

Review



# Mitochondrial Impairment: A Common Motif in Neuropsychiatric Presentation? The Link to the Tryptophan–Kynurenine Metabolic System

Masaru Tanaka <sup>1</sup>, Ágnes Szabó <sup>2,3</sup>, Eleonóra Spekker <sup>1</sup>, Helga Polyák <sup>2,3</sup>, Fanni Tóth <sup>1</sup> and László Vécsei <sup>1,2,\*</sup>

- <sup>1</sup> ELKH-SZTE Neuroscience Research Group, Danube Neuroscience Research Laboratory, Danube Neuroscience Research Laboratory, Eötvös Loránd Research Network, University of Szeged (ELKH-SZTE), Tisza Lajos krt. 113, H-6725 Szeged, Hungary
- <sup>2</sup> Department of Neurology, Albert Szent-Györgyi Medical School, University of Szeged, Semmelweis u. 6, H-6725 Szeged, Hungary
- <sup>3</sup> Doctoral School of Clinical Medicine, University of Szeged, Korányi fasor 6, H-6720 Szeged, Hungary
- \* Correspondence: vecsei.laszlo@med.u-szeged.hu; Tel.: +36-62-545-351

Abstract: Nearly half a century has passed since the discovery of cytoplasmic inheritance of human chloramphenicol resistance. The inheritance was then revealed to take place maternally by mitochondrial DNA (mtDNA). Later, a number of mutations in mtDNA were identified as a cause of severe inheritable metabolic diseases with neurological manifestation, and the impairment of mitochondrial functions has been probed in the pathogenesis of a wide range of illnesses including neurodegenerative diseases. Recently, a growing number of preclinical studies have revealed that animal behaviors are influenced by the impairment of mitochondrial functions and possibly by the loss of mitochondrial stress resilience. Indeed, as high as 54% of patients with one of the most common primary mitochondrial diseases, mitochondrial encephalomyopathy with lactic acidosis and strokelike episodes (MELAS) syndrome, present psychiatric symptoms including cognitive impairment, mood disorder, anxiety, and psychosis. Mitochondria are multifunctional organelles which produce cellular energy and play a major role in other cellular functions including homeostasis, cellular signaling, and gene expression, among others. Mitochondrial functions are observed to be compromised and to become less resilient under continuous stress. Meanwhile, stress and inflammation have been linked to the activation of the tryptophan (Trp)-kynurenine (KYN) metabolic system, which observably contributes to the development of pathological conditions including neurological and psychiatric disorders. This review discusses the functions of mitochondria and the Trp-KYN system, the interaction of the Trp-KYN system with mitochondria, and the current understanding of the involvement of mitochondria and the Trp-KYN system in preclinical and clinical studies of major neurological and psychiatric diseases.

**Keywords:** mitochondria; stress resilience; plasticity; stress; kynurenine; Alzheimer's disease; neurodegenerative; depression; anxiety; psychiatric

# 1. Introduction

Mitochondria are double membrane-bound cell organelles abundant in the cytosol of eukaryotes. The most prominent role of mitochondria is the production of high-energy storage molecule adenosine triphosphate (ATP) [1–3]. For ultimate energy production, mitochondria employ a variety of metabolic activities, including the tricarboxylic (TCA) cycle, oxidative phosphorylation (OXPHOS), ketogenesis/ketolysis, fatty acid oxidation, and glutamate metabolism [4–9]. Each component forms a complex metabolic network and dynamically adapts to the cellular environment to ensure the optimal energy supply. Mitochondrial malfunction can occur due to the defects of proteins directly or indirectly

**Citation:** Tanaka, M.; Szabó, Á.; Spekker, E.; Polyák, H.; Tóth, F.; Vécsei, L. Mitochondrial Impairment: A Common Motif in Neuropsychiatric Presentation? The Link to the Tryptophan–Kynurenine Metabolic System. *Cells* **2022**, *11*, 2607. https://doi.org/10.3390/ cells11162607

Academic Editors: Mounia Chami and Hemal H. Patel

Received: 7 July 2022 Accepted: 19 August 2022 Published: 21 August 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/). responsible for the OXPHOS or to the dysfunction of cellular mechanisms outside of mitochondria [10–13]. However, the role of mitochondria is not limited to cellular energy production. Other functions of mitochondria include calcium storage, subcellular signaling such as gene expression, autophagy, and apoptosis, among others [14,15].

Meanwhile, the tryptophan (Trp)–kynurenine (KYN) metabolic system plays a major role in Trp metabolism, as over 95% of Trp catabolizes into nicotinamide adenine dinucleotide (NADH). Accumulating evidence is revealing that the enzymes and the metabolic products of the Trp-KYN system actively influence the metabolism of mitochondria and participates in normal aging in organisms as well as the pathogenesis of mitochondrial diseases, neurodegenerative diseases, and psychiatric disorders [16,17]. The enzymes of the Trp-KYN system are activated by inflammation, oxidative stress, antioxidant system, and downstream bioactive metabolites [18,19].

Normal functions of mitochondria are typically compensated in mitochondrial diseases, a heterogenous group of chronic, genetic, and often inherited metabolic disorders caused by mitochondrial dysfunction, resulting in the impairment of cellular energy production and other crucial mitochondrial functions [20]. The prevalence of inherited mitochondrial diseases is estimated to occur one in 5000 live births, and they are the most common inborn errors of metabolism [21]. Primary mitochondrial disease (PMD) is caused by the pathogenic mutation of mitochondrial DNA (mtDNA) or nuclear DNA (nDNA) encoding either the proteins of OXPHOS or the proteins affecting the energy production of OXPHOS. Secondary mitochondrial diseases (SMDs) can be hereditary, caused by the genes for mtDNA transcription or expression, homeostasis, or metabolism [22]. Furthermore, mitochondrial dysfunction can be caused by acquired multifactorial diseases such as diabetes, cancer, heart or kidney disease, or neurodegenerative diseases [23,24].

Mitochondrial malfunction occurs in other conditions including normal aging in organisms, neurodegenerative diseases, and psychiatric disorders. Age-related physiological changes are strongly associated with mitochondrial malfunction with decreased mtDNA volume and mitochondrial integrity, which results from cumulative damage to mtDNAs by reactive chemical species [25]. Mitochondrial dysfunction also occurs in most neurodegenerative diseases such as Alzheimer's disease (AD), Parkinson's diseases (PD), Huntington's disease (HD), Friedreich's ataxia (FA), and amyotrophic lateral sclerosis (ALS) [26]. Psychiatric disorders include mood disorders such as major depressive disorder (MDD) and bipolar disorder (BD), schizophrenia (SCZ), autism spectrum disorder (ASD), and attention-deficit hyperactive disorder (ADHD) [27].

This review discusses the functions of mitochondria, the Trp-KYN system, the interaction of the Trp-KYN system with mitochondria, the mitochondrial environment upon the activation of the Trp-KYN system, and the link to neuropsychiatric presentation in clinical and preclinical settings in search for possible diagnostic biomarkers and novel interventional targets for mitochondria-associated diseases.

## 2. Mitochondria in the Central Nervous System

The brain accounts for only 2% of the body weight; however, it consumes as much as 20% of body's total oxygen supply. An estimated number of one to two million mitochondria is present per single neuron in the human substantia nigra [28]. Mitochondria take responsibility for the production of cellular energy and the proper conduction of neural circuits in the nervous system [29–33]. Mitochondria are multifunctional organelles maintaining calcium homeostasis and signaling to other organelles in the cell as well as with other mitochondria at distance [34,35]. Furthermore, mitochondria are highly plastic in morphology, functions, and cell cycle, depending on the tissue type and the need of cells [36]. Mitochondria can be even transferred from cell to cell [37].

#### 2.1. Mitochondrial Bioenergetics

Glucose, other sugars, and some amino acids are broken down in the cytosol to threecarbon molecule pyruvate which transfers into the mitochondria. Pyruvate is degraded to two carbon molecule acetyl coenzyme A (acetyl-CoA) which enters the second stage of cellular respiration, the TCA cycle that takes place in the matrix of mitochondria. Initially, Szent-Györgyi reported a cyclic chemical reaction between a four-carbon molecule (oxaloacetate) and two four-carbon molecules (fumarate and L-(-)-malate (the Szent-Györgyi cycle)). Later, Krebs revealed larger cyclic biochemical reactions in which a two-carbon molecule "triose" bonds with oxaloacetate to form a six-carbon molecule citrate, which is then oxidized to a five-carbon molecule (alpha-ketoglutarate) and four-carbon molecules (succinyl-CoA, succinate, fumarate, and malate), thus forming the TCA cycle (the Krebs cycle) [38]. "Triose" was eventually identified as a product of pyruvate and coenzyme A, acetyl-CoA [39,40]. The TCA cycle employs eight different enzymes, reproducing one molecule of oxaloacetate, two molecules of carbon dioxide, water, three molecules of NADH, and one molecule of flavin adenine dinucleotide (FADH<sub>2</sub>) and guanosine triphosphate (GTP). GTP is readily converted to ATP. In the TCA cycle, most of the high-energy storage molecule ATP is consumed by NAD<sup>+</sup> and FAD to form NADH and FADH<sub>2</sub> [29] (Figure 1a). The NAD<sup>+</sup> excess has been reported to improve mitochondrial function and thus prolong the life span of mice [30].

Of note, 16–18 carbon chain fatty acids transported by plasma albumin diffuse into the cytosol using a protein transporter. Consuming ATP, fatty acid is transformed to acyl coenzyme A (acyl-CoA) that crosses the inner membrane of mitochondria by carnitineacyl-CoA transferase. The beta-oxidation takes place in the mitochondrial matrix in which acetyl-CoA, water, and five ATP molecules are produced by shortening two carbon chains until an acyl-CoA molecule is reduced to an acetyl-CoA molecule [31].

Amino acids are recycled to produce new proteins, but when they are in excess, or cells are under starvation, amino acids are catabolized to supply energy. All essential amino acids except histidine, alanine, and cysteine (Cys) are involved in mitochondrial metabolic pathways. All essential amino acids are converted to pyruvate in the cytosol, which enters to mitochondria to fuel in the TCA cycle [36].

NADH and FADH<sup>2</sup> transfer their energy to the third stage of cellular respiration OXPHOS consisting of the electron transport, chemiosmosis, and ATP synthesis. The electron transport chain (ETC) generates a proton (H<sup>+</sup>) gradient across the inner membrane and the subsequent return of the H<sup>+</sup> to the matrix produces ATP from ADP by ATP synthetase. The ETC is a group of protein complexes composed of the NADH coenzyme Q reductase (Complex I), coenzyme Q, succinate dehydrogenase (Complex II), cytochrome bc1 complex (Complex III), and cytochrome c oxidase (Complex IV). NADH donates an electron to Complex I, generating three protons, while FADH<sup>2</sup> donates an electron to Complex I, generating three protons, while FADH<sup>2</sup> donates an electron to Complex II, generating two protons. ATP synthetase (Complex V) utilizes a proton gradient across the inner membrane to synthesize ATP from ADP and inorganic phosphate (Pi) [32]. In general, three to four protons are required to produce one ATP with 42% efficiency of energy conservation [33]. However, the cellular energy production can be altered under stressful condition or pathological processes according to the availability of substrates, enzyme activity, mitochondrial and cellular conditions, and adjacent biosystems including Trp-KYN metabolic system.



**Figure 1.** The tricarboxylic acid cycle (TCA) and its interface with the tryptophan (Trp)–kynurenine (KYN) metabolic system. (a) The TCAcycle is initiated with acetyl coenzyme A (acetyl-CoA) reacting with oxaloacetate to form citrate. Citrate is oxidized to alpha ( $\alpha$ )-ketoglutarate (2-oxoglutarate) with the formation of nicotinamide adenine dinucleotide (NADH).  $\alpha$ -ketoglutarate is oxidized to suc-

cinyl coenzyme A (succinyl-CoA) with the formation of NADH. Succinyl-CoA is converted to succinate with the formation of adenosine triphosphate (ATP). Succinate is oxidized to fumarate with the formation of flavin adenine dinucleotide (FADH<sub>2</sub>). Fumarate is hydrated to malate which is oxidized to oxaloacetate to end the cycle. (**b**) Cytosolic kynurenine aminotransferase (KAT) I catalyzes the reaction of an *S*-substituted L-Cys to pyruvate. KAT I also catalyzes the reaction of L-glutamine to  $\alpha$ -ketoglutarate (2-oxoglutarate). (**c**) Mitochondrial KAT II, KAT III, and KAT IV catalyze the reaction of  $\alpha$ -ketoglutarate catalyzes the reaction of L-glutamine to  $\alpha$ -ketoglutarate (2-oxoglutarate) to L-glutamate. (**d**) KAT II catalyzes the reaction of  $\alpha$ -ketoglutarate (2-oxoglutarate) to L-glutamate. (**d**) KAT II catalyzes the reaction of  $\alpha$ -ketoglutarate (2-oxoglutarate) to 2-oxoadipate which is eventually degraded to acetyl-CoA. (**e**) Mitochondrial KAT III catalyzes the reaction of an *S*-substituted L-Cys to pyruvate. (**f**) Mitochondrial KAT IV catalyzes the reaction of  $\alpha$ -ketoglutarate (2-oxoglutarate) and L-aspartate to  $\alpha$ -ketoglutarate (2-oxaloacetate) and L-glutamate.

# 2.2. Other Mitochondrial Functions

Mitochondria play an important role in cellular calcium homeostasis. The concentration of calcium ions in the intermembrane space is the same as that in the cytosol due to the high permeability of the outer mitochondria membrane. A higher concentration of mitochondrial calcium ions enhances ATP production; however, severe calcium overloads are associated with pathological conditions [34,35,41].

Mitochondria constantly communicate with other cellular organelles such the nucleus, the ER, lysozymes, and peroxisomes. The coordinated interaction of mitochondrial and nuclear factors is required for mitochondrial gene expression offered by mitochondrial ribonuclease P, ribosomal RNAs, transfer RNAs, introns, and a protein [42]. The nucleus sends signals to the mitochondria via anterograde regulation to modulate mitochondrial biogenesis upon stressful events. On the other hand, mitochondria constantly transmit information on mitochondrial status and cellular stress to the nucleus by retrograde signaling [43].

Mitochondria and the ER are at a close contact through the mitochondria-associated membrane to exchange information on energy production, calcium homeostasis, lipid transport, and apoptosis [44]. Lysosomes interact with mitochondria to transport amino acids, lipids, and calcium ions [45]. Mitochondria and peroxisomes function in concert in fatty acid metabolism. Mitochondria degrade long-chain fatty acids to supply acetyl-CoA and produce ATP, while peroxisome performs beta-oxidation to generate hydrogen per-oxide and anabolic metabolic metabolism such as plasmalogen and bile acid synthesis [46].

Mitochondria undergo division during mitosis, dividing equally between the cell soma to daughter cells in interaction with the ER and cytoskeleton [47]. The morphology, functions, and dynamics of mitochondria change upon tissue differentiation [48]. Mitochondria constantly divide and fuse, controlling their morphology and functions. The fusion takes place by initially merging the outer membrane and subsequently the inner membrane of two mitochondria. The continuous events of fusion and division generate mitochondrial networks [49]. Mitophagy refers to mitochondria autophagy in which double-membraned vesicle autophagosomes deliver mitochondria to lysosomes for destruction. Mitophagy is induced by prolonged fission, promoting the repair process, but may lead to mitochondrial degradation. MicroRNAs play an important role in regulation of protein expression responsible for autophagy [50].

Mitochondria also induce an immune response via the activation of the mitochondrial antiviral signaling protein which leads to the secretion of cytokines via the virally infected cells [51]. Furthermore, mitochondria induce mitochondrial apoptosis through mitochondrial outer membrane permeabilization which leads to the disruption of mitochondrial outer membrane and the release of intermembrane space proteins such as cytochrome c [52]. Therefore, mitochondria impairment may lead to multifarious consequences from ion homeostasis to entire organismal levels.

# 3. The Tryptophan-Kynurenine Metabolic System

The aromatic amino acid L-Trp is an essential component for the biosynthesis of proteins and a substrate for the production of neurotransmitters and hormones [53]. Acute Trp deprivation increases pain sensitivity, motor activity, acoustic startle, and aggression, while chronic Trp deficiency induces ataxia, cognitive impairment, and dysphoria [54,55]. A meta-analysis showed decreased levels of Trp in blood samples of patients with MDD and BD [56]. Trp is catabolized in the serotonin (5-HT) and KYN metabolic system. More than 95% of L-Trp enters the KYN system, producing several biomolecules. The main enzymes of the KYN system are tryptophan 2,3-dioxygenase (TDO), indoleamine 2,3-dioxygenases (IDOs), kynurenine formamidase (KFA), kynurenine 3-monooxygenase (KMO), kynurenine aminotransferases (KATs), and kynureninase (KYNU). The main metabolites are L-KYN, kynurenic acid (KYNA), 3-hydroxy-L-kynurenine (3-HK), quinolinic acid (QUIN), and NAD<sup>+</sup>. Those metabolites possess a wide range of biological properties such as oxidative, antioxidative, neurotoxic, neuroprotective, cognition-enhancing and impairing, and/or immunomodulating properties, and have attracted growing attention as potential biomarkers, therapeutic targets, cross-species markers, and indicators for environmental resilience [19,57,58] (Figure 2).



**Figure 2.** The tryptophan (Trp)–kynurenine (KYN) metabolic system and the subcellular location of the enzymes. More than 95% of L-Trp enters the KYN system producing multifarious biomolecules. The main enzymes of the KYN system are tryptophan 2,3-dioxygenase (TDO), indoleamine 2,3-dioxygenases (IDOs), kynurenine formamidase (KFA), kynurenine 3-monooxygenase (KMO), kynurenine aminotransferases (KATs), and kynureninase (KYNU). Most of the enzymes are located in the cytosol. However, KMO is located in the outer membrane of mitochondria; KAT II and KAT

III are in the inner membrane of mitochondria; and KAT IV is in the matrix of mitochondria and in the plasma membrane. The main metabolites are L-KYN, kynurenic acid (KYNA), 3-hydroxy-L-kynurenine (3-HK), quinolinic acid (QUIN), and nicotinamide adenine dinucleotide (NAD<sup>+</sup>) which exhibit a wide range of biological properties and the metabolites freely cross the mitochondrial membranes.

# 3.1. Tryptophan 2,3-Dioxygenase, Indoleamine 2,3-Dioxygenases, and Kynurenine Formamidase

TDO and IDOs are heme-containing enzymes that catalyze the oxidation of L-Trp to N-formyl-L-kynurenine. This is the first rate-limiting step of the Trp-KYN system, which regulates the systemic level of Trp in the body. IDO isoform 1 (IDO1) also catalyzes a stereoisomer D-Trp which is a product of the gut microbiome. TDO is a cytosolic enzyme encoded by the gene *tdo2*, stimulated by the stress hormone cortisol and the downward metabolite 3-HK [17,59]. Human *tdo2* gene polymorphisms are associated with ADHD, Tourette syndrome, MDD, ASD, and SCZ [60,61]. *tdo2-/-* mice exhibited less anxious behavior, increased exploratory activities, and increased cognitive function, with increased concentration of Trp, 5-HT, 5-hydroxyindoleacetic acid, and/or KYN (in the plasma, hippocampus, or midbrain) [62,63]. However, it was later reported that anxiolytic and exploratory behaviors are less prominent in *tdo2-/-* mice [64,65]. Thus, the behaviors of *tdo2-/-* mice remain inconclusive.

IDOs are cytosolic enzymes with two isoforms. IDO1 is encoded by the gene *ido1* and expressed in various parts of the body, including the brain, while IDO isoform 2 (IDO2) is encoded by the gene *ido2* and expressed widely in tissues such as kidney, liver, or anti-gen-presenting cells [66,67]. The two isoforms differentiate in kinetics, substrate specificity, and function. IDOs are upregulated by pro-inflammatory cytokines and lipopolysaccharide but are downregulated by anti-inflammatory cytokines and the antioxidant enzyme superoxide dismutase [17].

*ido1*<sup>-/-</sup> knockout (KO) mice showed diurnally hypolocomotive behavior with higher brain 5-HT levels and attenuated nociceptive sensation and depressive-like behavior [63,68]. Furthermore, Bacille Calmette-Guérin (BCG) elicited proinflammatory cytokines, but BCG-induced depressive-like behavior was not induced in *ido1*<sup>-/-</sup> KO mice [69]. However, another study reported that *ido1*<sup>-/-</sup> KO mice did not exhibit any significantly different behavior, compared to the wild type, in comprehensive behavioral assessments including the domain of cognitive function, negative valence system, motor function, social interaction, and pain sensitivity [70]. *Ido2*<sup>-/-</sup> KO mice increased exploratory activity during light phase [63]. *ido1* polymorphism is associated with the susceptibility of interferon  $\alpha$  treatment-induced depression in patients with chronic hepatitis C [71]. The common polymorphisms of *ido1* and *ido2* genes are associated with the outcome of a selective serotonin reuptake inhibitor (SSRI) citalopram treatment [72].

N-formyl-L-kynurenine hydrolyses to L-KYN spontaneously or enzymatically via KFA. The enzyme is encoded by the gene *afmid*, predominantly cytosolic, expressed in the liver and kidney, and participates in glyoxylate and dicarboxylate metabolism [73]. KFA is stimulated by the proinflammatory cytokine interferon- $\gamma$ . L-KYN is an antioxidant and an aryl hydrocarbon receptor (AHR) agonist [74]. A meta-analysis showed decreased levels of KYN and an increased KYN/Trp ratio in blood samples of patients with MDD [56]. No study regarding KFA polymorphism or gene KO has been reported (Table 1).

**Table 1.** The enzymes, genes, substrates, products, activities, knockout, and human gene variants of the tryptophan–kynurenine metabolic system.

Enzymes	Genes	Substrates	Products	Locations	Transgenic Models	Animal Traits	Human Gene Variants
TDO	tdo2	L-Trp	N-formyl-L-	Cytosol	tdo2-/-	- Anxiety (inconclusive)	ADHD, MDD, ASD, SCZ,
100			kynurenine				Tourette syndrome

						- Exploratory activities	
						(inconclusive)	
						- Cognitive function	
						- Locomotion	
IDO1	ido1	L-Trp,	N-tormyl-L-	Cytosol	ido1-/-	- Nociception	MDD
		D-Trp	kynurenine			- Depression	
			N-formyl-L-			-	
IDO2	ido2	L-Trp	kvnurenine	Cytosol	Ido2-/-	-	MDD
		N-formvl-L-					
KFA	afmid	kynurenine	L-KYN	Cytosol	-	-	-
		Rynarchine				Lower contextual	
						- Lower contextual	
						memory function	- Cognitive dysfunction
				Mitochondria		- more anxious-like be-	- Lower cognitive perfor-
КМО	kmo	L-KYN	3-HK	(outer mem-	kmo-/-	havior	mance
				brane)		- Higher horizontal ac-	- A trend effect on cogni-
						tivity upon a D-am-	tive function
						phetamine challenge	
		L-KYN	KYNA				
KAT I							
(Kynurenineox-		S-substituted L-Cys	thiol, NH4, pyruvat	te			
oglutarate trans-	kyat1			Cytosol	-	-	-
aminase 1)		3-phenylpyruvate	2-oxoglutarate				
,		L-glutamine	L-phenylalanine				
		Eglutamine	2 prierty talaitaile			Transitory hyperlose	
KAT II		L-KYN	KYNA			- mative estivity	
(kynurenine/α-				Inner mem-	1		
aminoadipate ami-	aadat	α-ketoglutarate L-glutamate 2-oxoglutarate L-glutamate	L-glutamate	brane of mi- tochondria	aadat-/- (aka kat2-/-)	- Transitory abnormal	-
notransferase)						motor coordination	
(KAT/AadAT)					- Increased cognitive		
· · · · · · · · · · · · · · · · · · ·		0	0			functions	
		L-KYN					
			KYNA				
		$\alpha$ -ketoglutarate					
			L-glutamate				
		3-HK					
KAT III		glvoxvlate	Glycine	Cytosol			
(kynurenineox-	kuat3	8,5,5,5,5	HiO	inner mem-	_	-	
oglutarate trans-	кушо		× A	brane of mi-			
aminase 3)			XA	tochondria			
		giyoxylate L-KYN					
			Glycine				
			H2O KYNA				
		S-substituted L-Cys					
		H <sub>2</sub> O	Thiol				

			NH4 <sup>+</sup>				
			pyruvate				
		L-KYN	KYNA				
KAT IV				Matrix of mi-			
(aspartate ami-	12	$\alpha$ -ketoglutarate	L-glutamate	tochondria			
notransferase,	got2			plasma mem-	-	-	-
mAspAT)		2-oxoglutarate	L-glutamate	brane			
		L-aspartate	oxaloacetate				
KYNU	kynu	L-KYN	AA 3-HAA	Cytosol	kynu-/-	-	- Vertebral, cardiac, renal, and limb defects syn- drome 1
	L-alaı	L-alanine 3-HK	(3-arylcarbonyl)-ala nine	-			- Essential hypertension
3-HAO	haao	3-HAA	ACMS	Cytosol	-	-	- Vertebral, cardiac, renal, and limb defects syn-
							drome 1

# 3.2. Kynurenine 3-Monooxygenase

KMO is encoded by the gene *kmo* and catalyzes the rate-limiting step of the redox reaction from L-KYN to 3-HK. KMO is located in the outer membrane of mitochondria and is expressed in many tissues of the body, including the brain glial cells and microglia [75]. KMO is stimulated by oxygen molecules, pro-inflammatory cytokines, and the downstream metabolite NADH, while it is inhibited by the superoxide dismutase and anti-inflammatory cytokines [17]. 3-HK generates free radicals to elicit excitotoxic injury. The oxidant molecule 3-HK may function as an antioxidant in certain conditions [76]. A meta-analysis showed a decreased ratio of KYNA/3-HK in blood samples of patients with MDD [56].

*kmo<sup>-/-</sup>* KO mice have been generated to study the transgenic effects on the metabolites of the Trp-KYN metabolites. The levels of 3-HK were lower in the liver, brain, and plasma; the levels of QUIN were greatly lower in the liver and plasma, while slightly lower in the brain; the levels of KYN, KYNA, and anthranilic acid (AA) were substantially higher, but depending on a tissue; and the levels of NAD<sup>+</sup> were not different, compared to the wild type [77]. *kmo<sup>-/-</sup>* KO mice showed lower contextual memory function, more anxious-like behavior, and higher horizontal activity upon a D-amphetamine challenge. The behaviors were associated with the elevated levels of KYNA in the brain, especially in the cerebellum. [78].

A small sample study reported that KMO single-nucleotide polymorphism (SNP) rs1053230 polymorphism was potentially associated with lower CSF KYNA concentrations in SCZ patients [79]. KMO polymorphism has been associated with cognitive dysfunction. The *KMO* rs2275163C>T C (risk) allele was related to the lower cognitive performance in healthy controls, and it is more prominent in SCZ patients. Furthermore, other KMO polymorphism showed a trend effect in cognitive function [80] (Table 1).

# 3.3. Kynurenine Aminotransferases

KATs belong to transferases, specifically transaminases, employing pyridoxal 5'phosphate (PLP). KATs typically catalyze substrates L-KYN and 2-oxoglutarate to 4-(2aminophenyl)-2,4-dioxobutanoate and L-glutamate, and then the unstable former product forms KYNA via intramolecular cyclization [81]. Kynurenine–oxoglutarate transaminase 1 (aka KAT I) encoded by the gene *kyat1*, kynurenine/ $\alpha$ -aminoadipate aminotransferase (KAT/AadAT, aka KAT II) encoded by the gene *aadat*, kynurenine–oxoglutarate transaminase 3 (KAT III) encoded by the gene *kyat3*, and aspartate aminotransferase (mAspAT, aka KAT IV) encoded by the gene *got2* are isofoms representing KATs. KAT I is located in the cytosol; KAT II is in the inner membrane of mitochondria; KATIII is in the inner membrane of mitochondria and the cytosol; and KAT IV is in the matrix of mitochondria and the plasma membrane [82–85]. KYNAs are allosterically regulated by  $\alpha$ ketoglutarate in cooperation with KYN [86].

KAT I also catalyzes the reaction of S-substituted L-Cys and H<sub>2</sub>O to a thiol, NH<sub>4</sub>, and pyruvate, as well as the reaction of 3-phenylpyruvate and L-glutamine to 2-oxoglutarate and L-phenylalanine (Figure 1b). 2-oxoglutarate is the same as  $\alpha$ -ketoglutarate [82]. Mitochondrial KAT II, KAT III, and KAT IV may compete for the substrate  $\alpha$ -ketoglutarate of the TCA cycle, supplying L-glutamate in glutamine metabolism (Figure 1c) [83–85]. KAT II also catalyzes the reaction of 2-oxoglutarate and L-2-aminoadipate to 2-oxoadipate and L-glutamate. 2-oxoadipate is an intermediate molecule of L-lysin catabolism, which is further degraded to glutaryl coenzyme A, and finally to acetyl-CoA, forming a side loop of the TCA cycle (Figure 1d) [83]. KAT III catalyzes the reactions of 3-HK and glyoxylate to glycine, H<sub>2</sub>O, and xanthurenic acid (XA); of glyoxylate and L-KYN to glycine, H<sub>2</sub>O, and KYNA; and of an S-substituted L-Cys and H<sub>2</sub>O to a thiol, NH<sub>4</sub><sup>+</sup>, and pyruvate (Figure 1e) [84]. Furthermore, mitochondrial KAT IV catalyzes the reaction of 2-oxoglutarate and L-aspartate to L-glutamate and oxaloacetate, thus bypassing the TCA cycle and participating in essential amino acid aspartate metabolism (Figure 1f) [85]. Thus, KAT enzymes relay molecules between the TCA cycle and glutamate, lysine, and aspartate metabolisms.

KYNA is a receptor antagonist of glutamate receptors including ionotropic  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), kainate, and N-methyl-D-aspartic acid (NMDA) receptors [87]. KYNA's action depends on its concentration at AMPA receptor and cognitive function [88,89]. KYNA's action at the  $\alpha$ -7 nicotinic acetylcholine receptor in vivo remains controversial [90]. KYNA is a G-protein-coupled receptor 35 (GPR35) ligand and an AHR agonist. A meta-analysis showed decreased ratios of KYNA/KYN, KYNA/QUIN, and KYNA/3-HK in patients with MDD; a decreased level of KYNA; and a decreased ratio of KYNA/QUIN in patients with BD [56]. XA is an AHR agonist and a glutamate vesicular transporter (VGLUT) inhibitor and may be a Group II metabotropic glutamate receptor agonist [91,92]. A meta-analysis revealed lower XA levels in the blood of patients with BP and a significantly lower level of XA was observed in the serum of SCZ patients [93,94].

The intracerebroventricular (i.c.v.) administration of KYNA was reported to enhance memory function at low doses and impair it at higher doses in the passive avoidance test of mice [95]. Furthermore, i.c.v. KYNA and synthetic KYNA analogues showed antide-pressant-like effects in modified forced swim test of mice [96,97].

*aadat*<sup>-/-</sup> (aka *kat*2<sup>-/-</sup>) KO mice exhibited transitory hyperlocomotive activity and abnormal motor coordination during postnatal day 17 to 26 with transiently reduced total brain KAT activity and KYNA levels during the first month, which returned to normal levels later [94]. Three-week-old *kat*2<sup>-/-</sup> KO mice showed significantly increased cognitive functions including object exploration and recognition, passive avoidance, and spatial discrimination with significantly reduced levels of KYNA in the hippocampus [98]. Intriguingly, homozygous *got*2<sup>-/-</sup> KO mice result in embryonal death in utero [99]. No study has been reported regarding KATs variants associated with psychiatric symptoms (Table 1).

## 3.4. Kynureninase

KYNU is a PLP-dependent hydrolase, encoded by the gene *kynu*, which catalyses L-KYN to AA and L-alanine and 3-HK to 3-hydroxyanthranilic acid (3-HAA) and (3-arylcarbonyl)-alanine. The enzyme also has Cys-conjugate-beta-lyase activity. KYNU is located in the cytosol [100]. However, there is a report that the enzyme has no activity for L-KYN and is inhibited by L-KYN and D-KYN. AA inhibits the TCA cycle and respiratory chain complexes I–III, interfering with mitochondrial function [101]. 3-HAA is also biosynthesized from AA by spontaneous hydroxylation. AA was once thought to be watersoluble vitamin L<sub>1</sub> and is possibly related to an endogenous anti-inflammatory derivative in the celluar environment, as the molecule is a pharmaceutical precursor of nonsteroidal anti-inflammatory agents such as mefenamic acid and diclofenac [102]. 3-HAA can be either an oxidant or an antioxidant depending on the cellular condition. AA and 3-HAA suppress pro-inflammatory cytokines and invoke anti-inflammatory cytokine interleukin [IL]-10 [103].

*kynu*<sup>-/-</sup> KO mice have been generated; however, no study has been reported regarding neurological and psychiatric diseases. Homozygous variant KYNU p.V57Efs\*21 and heterozygous KYNU variants p.Y156\* and p.F349Kfs\*4 were identified in patients with vertebral, cardiac, renal, and limb defects syndrome 1, as well as autosomal recessive congenital malformation, characterized by vertebral segmentation abnormalities, congenital cardiac defects, renal defects, and distal mild limb defects. The NADH levels of patients are significantly lower [104]. KYNU SNP rs2304705 has been associated with essential hypertension and a 50% reduction in enzyme activity, but the activity reduction was not observed in another SNP [105].

In humans, increased levels of pro-inflammatory cytokine IL-1, associated with reduced levels of hippocampal neurogenesis, have been reported in depressed patients and in animal models of depression [106,107]. Identifying the mechanisms by which inflammatory cytokines block neurogenesis in the human brain may provide insight that can be used to manage inflammation-associated mental health disorders, including diet, new diagnostic methods, and treatment therapies for depression [108–110]. Recently, building on previous evidence, a new theoretical model named the neurovisceral integration model of fear (NVI-f), conceptualized the anatomical–functional interplay between the prefrontal cortex and heart-related dynamics in human emotional learning [111]. The potential of a novel approach, the time-frequency decomposition of heart rate variability, has gained attention in the evaluation of the abnormal fear learning that characterizes several neurological and psychiatric disoders [112]. Moreover, several studies have suggested the effectiveness of non-invasive brain simulation to interfere and modulate the abnormal activity of neural circuits, such as amygdala-medial prefrontal cortex (PFC) and hippocampus, involved in the acquisition and consolidation of emotional memories, which are altered in psychiatric disorders, such as fear-related disorder including anxiety disorder, phobias, post-traumatic stress disorder (PTSD), or depression [113-116] (Table 1).

## 3.5. 3-Hydroxyanthranilate 3,4-Dioxygenase, and toward the Tricyclic Carboxylic Cycle

3-hydroxyanthranilate oxidase (3-HAO) is the most active enzyme of Trp-KYN metabolic system, encoded by gene *haao*. The non-heme iron-dependent enzyme 3-HAO catalyzes 3-HAA to 2-amino-3-carboxymuconate semialdehyde (ACMS), which spontaneously cyclizes to QUIN. The enzyme is located in the cytosol [117]. QUIN forms a highly reactive hydroxyl free radical and is a NMDA receptor agonist which elicits excitotoxicity [118]. The concentration QUIN is increased upon immune activation and decreased by immune suppressant dexamethasone [119,120]. QUIN inhibits around 35% succinate dehydrogenase, an enzyme involved in the TCA cycle and in the respiratory chain [121]. *haao* gene variants HAAO p.D162\* and HAAO p.W186\* were identified in patients with vertebral, cardiac, renal, and limb defects syndrome 1, and the NADH level of patients was significantly lower [104].

Quinolinate phosphoribosyltransferase (QPRT) catalizes QUIN to nicotinic acid mononucleotide (NaMN), which is converted by NaMN adenyltransferase to nicotinic acid adenine dinucleotide (NaAD). Finally, NAD synthetase converts NaAD<sup>+</sup> to NAD<sup>+</sup>. ACMS is catalized by 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase (AC-MSD) or picolinate carboxylase to 2-aminomuconic-6-semialdehyde (AMS) that is nonenzymatically cyclized to picolinic acid (PIC) [17]. The significantly lower levels of PIC were observed in ASD [122]. XA is converted to cinnabarinic acid (CA) by autoxidation. CA is

12 of 46

also produced from 3-HK or QUIN. CA is an AHR agonist, and the reduced concentration of CA in the PFC is linked to SCZ [123]. 2-aminomuconate semialdehyde dehydrogenase (AMSD) catalyzes the conversion of AMS to 2-aminomuconic acid, which is further degraded to acetyl-CoA that replenishes the TCA cycle [102] (Table 1).

Currently, researchers are focusing on finding scientific frameworks for understanding the relationship between the molecular regulation of higher-order neural circuits and neuropathological alterations, and how this may lead to PFC dysfunction and to the symptoms of mental illnesses and comorbidity [124]. The deficit in control and motor inhibition [125,126], but also in motor imagery or in the suppression of on-going action [127], which depend on aberrant neural activity in the PFC associated with serious impulsivity problems, are characterized by psychopathological and psychiatric conditions including MDD, SCZ, obsessive–compulsive disorder (OCD), and PD [128,129].

Furthermore, the intentional neglect of adaptive procees necessary for memory functioning and functional alterations in the PFC affects the memory and learning abilities of psychiatric and brain-damaged patients. The human ventromedial PFC is responsible for the capacity of associative learning [130–132]. Hypoactivation in the ventromedial PFC with hyperactivation in the dorsal anterior cingulate cortex are reported in patients with PTSD and SCZ [133,134]. This evidence suggests that PFC dysfunctions cause impairment of aversive learning and emotional memory circuits, which might be transversal across many psychiatric disorders in humans as well as in neurologic patients [135].

## 4. Diseases Linked to Mitochondrial Dysfunction

The activity of neurons depends on mitochondrial function to elicit membrane excitability, execute neurotransmission, and maintain neuroplasticity [136]. Mitochondria are located throughout the cytoplasm of neurons, but more mitochondria are found in energy-demanding areas such as in the sites of branching axons, synaptic contacts, and glial processes [137]. The volume fraction of mitochondria is the highest in the cortical layer IV and mitochondrial volume is higher in dendrites than axons in rats, suggesting most energy consumption takes place at the postsynaptic side [138]. The dynamics of mitochondria are governed by mitochondrial fission, fusion, mitophagy, motility, and anchoring. Furthermore, mitochondria play a crucial role in axon degeneration and regeneration [139,140].

Mitochondrial dysfunction can affect any part of the body, but the most vulnerable organs are those with high-energy requirements, such as the central nervous system (CNS), peripheral nervous system, heart, and musculoskeletal system [141]. Mitochondrial diseases often go misdiagnosed or undiagnosed due to a wide range of manifestation, including fatigue; muscle weakness; visual or hearing loss; seizures; strokes; dementia; severe constipation; diabetes; thyroid or adrenal dysfunction; heart, liver, or kidney failure; poor growth; developmental delays; learning disabilities; and ASD in children [142]. The conditions can appear in adolescence and in adulthood [143]. The mitochondrial dysfunction also exhibits psychiatric symptoms such as depression, cognitive impairment, psychosis, and anxiety [144]. Currently, there is no cure for mitochondrial diseases and the mainstay of the treatment remains symptom-relieving or progression-delaying measures, which vary from patient to patient and depend on the mitochondrial disease and its severity [145]. Nevertheless, novel treatment for mitochondrial diseases is under extensive research. The strategies include oxidative stress modulation, mitochondrial biogenesis augmentation, mitochondrial autophagy modulation, nitric oxide restoration, mitochondria genome modulation, nucleotides pool restoration, hypoxia, enzyme replacement, and mitochondrial augmentation [146].

# 4.1. Primary Mitochondrial Diseases

PMDs are a clinically heterogenous group of uncurable, chronic, and genetic conditions caused by the mutations of mtDNA. PMDs commonly affect the nervous system of developmental stage, predominantly affecting skeletal muscles, but presenting many nonspecific symptoms from muscle weakness to seizure [147]. The mutations of the genes may encode proteins functioning for OXPHOS, mtDNA replication and expression, mitochondrial dynamics, homeostasis, quality control, mitochondrial metabolism, metabolism of cofactors, and metabolism of toxic compounds, among others [148].

The severe forms of PMDs typically present early in life, but the milder forms tend to have later presentations [149]. Primary mitochondrial myopathy (PMM) causes progressive external ophthalmoplegia, frequently presented with diplopia, bilateral ptosis, or a head tilt. Progressive external ophthalmoplegia can be a part of a syndrome with facial muscle weakness or paralysis, swallowing difficulty, slurred speech, or breathing difficulty. Furthermore, PMM may show involvement of the muscles of the neck, shoulder, arms, hips, or legs, presenting cramping stiffness, weakness, pain, or paralysis of the affected muscles. Exercise intolerance is a common symptom [150]. Mitochondrial encephalomyopathy is characterized by neurological presentation in infancy or childhood, such as vision loss, sensorineural hearing loss, migraine, ataxia, or seizures. Other neurological manifestation includes dysphagia, dysarthria, myasthenia, or muscle rigidity. Some patients experience peripheral neuropathy. Developmental delays, failure to thrive, or short statue is a common finding in children [151]. In addition, many genetic disorders present mitochondrial myopathy or encephalomyopathy as a part of the main symptoms involved in multiple organ systems [152].

The diagnosis is made clinically but is very difficult and not always confirmed by a DNA mutation. The causative gene mutations can be located in both mtDNA and nDNA [153]. In total, 413 genes have been associated with PMDs. PMDs caused by mtDNA are estimated to have a prevalence of 1 in 5000 cases, while PMDs caused by nDNA are estimated to have prevalence of 1 in 35,000 [154]. The mutations can be either inherited or spontaneous. The mutations of nDNA can be inherited either autosomal dominantly or autosomal recessively. The mutations of mtDNA are considered to be inherited only maternally; however, a biparental mode of inheritance of mtDNA has been reported [155,156]. A variety of clinical manifestations in a single family may be due to the heteroplasmy of mtDNA or the different number of mutant mtDNAs in daughter cells as a result of mitosis.

Leigh syndrome is a neurodegenerative disorder, and it is the most prevalent mitochondrial disease in childhood. It is known that more than 75 genetical mutations appear in the basis of the disorder [157]. A recent study showed a reduction in the L-KYN and 3-HAA levels in blood with French Canadian variants of Leigh syndrome patients. In addition to this, the level of indoxyl sulfate increased in these patients which suggest a shift in Trp metabolism to the indol pathway [158]. Trp can metabolize not only into KYN or 5-HT, but also to indoxyl sulfate. Thus, Trp can transform to indole by tryptophanase, and indole metabolizes to indoxyl by cytochrome P450 2E1. Thereafter, sulfotransferase can convert indole to indoxyl sulfate [159].

Leber hereditary optic neuropathy is accompanied by a degeneration of retinal ganglion cells, causing a loss of vision [160]. The patients usually have some mutations in the genes, which encode complex I in the ETC. In patients with Leber hereditary optic neuropathy, decreased levels of Trp and glutamate have been found, suggesting the possible role of the Trp-KYN metabolic system in the pathomechanism of the disease [161,162].

# 4.2. Secondary Mitochondrial Dysfunction

SMD can be caused by genes not encoding proteins for OXPHOS or mitochondrial functions, secondary to other illnesses (such as cancer, sepsis, and infections, as well as metabolic, neuromuscular, neurodegenerative, and psychiatric diseases); by drugs such as tetracycline and valproate; by environmental factors (including alcohol, cigarette smoke, carbon monoxide, asbestos, and metals, as well as antiretroviral, tetracycline, valproate, and aminoglycosides therapy); or by normal aging [163–165]. Therefore, SMD can be inherited or acquired. The diagnosis of SMD is made based on clinical signs of mitochondrial dysfunction with negative or equivocal DNA tests. However, it is often

difficult to distinguish SMD from PMD, but important for the prognosis and treatment. Sometimes, the treatment of PMD is effective to SMD [22].

#### 4.3. Neurological Disesases Linked to Mitochondrial Dysfunction

The loss of stress resilience and the functional impairment of mitochondria have been linked to neuropsychiatric symptoms comorbidities of neurological diseases such as AD, PD, HD, ALS, FA, and Charcot–Marie–Tooth disease [166–168]. Preclinical animal research plays a major role in revealing the involvement of endogenous peptides, neurohormones, and metabolites including KYNs [169–175].

# 4.3.1. Alzheimer's Disease

AD is the most common chronic neurodegenerative disease with an insidious onset of progressive cytokine dysfunctions, particularly memory impairment, but it progresses to motor, sensory, and autonomic dysfunctions in later stages [176]. The age-related impairments in the ability to process contextual information and in the regulation of responses to threat are related to structural and physiological alterations in the PFC and medial temporal lobe [177]. Positron emission tomography with 2-deoxy-2-[fluorine-18]fluoro-D-glucose has shown the deficit of hubs in the theory of mind network in patients with mild cognitive impairment due to AD [178]. Brain autopsy and imaging studies reveal the atrophy of the brain including the frontal, temporal, parietal, entorhinal cortices, amygdala, and hippocampus [179]. The deposition of amyloid beta (A $\beta$ ) peptide and tau protein is a characteristic finding, but not limited to AD [180].

More than 170 genetically manipulated mouse models of AD have been created. Most transgenic mouse models of AD are designed to overexpress genes associated with early onset familial type of AD, such genes APP, PSEN-1, and PSEN-2 genes, and the mouse strains are characterized with the pathological deposition of A $\beta$  peptide [181]. Mitochondrial dysfunction including decreased mitochondrial respiration and pyruvate dehydrogenase protein was observed in the brains of triple transgenic mice of AD (3xTg-AD) at the age of 3 months, while mitochondrial A $\beta$  levels significantly increased in 3xTg-AD at the age of 9 months. Mitochondrial impairment was even detected in embryonal neurons of 3xTg-AD [182]. Mitochondrial impairments were reported in a transgenic mouse expressing human amyloid precursor protein with the Arctic mutation (TgAPParc) mice at the age of six months. The mitochondrial membrane potential was decreased; the amount of reactive oxygen species was increased; and oxidative DNA damage was increased. Mitochondrial abnormality is more prominent in TgAPParc mice at the age of 24 months [183]. Mitochondrial dysfunction was also observed in transgenic mice carrying the APPswe and PSEN1dE9 mutations, heterozygous sodium-dependent vitamin C transporter (SVCT2<sup>+/-</sup>) KO mice, and transgenic APP/PSEN1 mice with heterozygous SVCT2 expression at the age of 4 months [184]. However, familial AD accounts for less than 5% of AD. Recently, human A $\beta$ -knockin (KI) mice were engineered, which mimics a late-onset type.  $hA\beta$ -KI mice develop age-dependent phenotypic and behavioral alterations and may be more relevant to study polygenic and multifactorial pathogenesis of AD [185].

The ratio of KYN/TRP was increased in the plasma and CSF of patients with AD and an increased 3-HK/KYN ratio in samples from CSF positively correlated with amounts of t-tau and p-tau peptides, while plasma KYN and PIC inversely correlated with p-tau and t-tau, respectively [186–191]. The levels of KYNA were found to be decreased in the plasma of AD patients [188]. AD had strong effect sizes for shared deficits in complex I and IV in the peripheral blood, frontal cortex, cerebellum, and substantia nigra [192] (Table 2).

Neurological	Preclinical Models	Mitochondrial	Findings in Kynurenines	
Diseases		Involvement		
	>170 genetic models	_		
	(APP, PSEN-1, PSEN-2)			
		- decreased mitochondrial respiration		
	3xTg-AD	- decreased pyruvate dehydrogenase protein	- increased ratio of KYN/Trp	
		- increased mitochondrial $A\beta$ level	- decreased KYNA	
		- decreased mitochondrial membrane potential	- 3-HK/KYN positively correlated with t-tau and	
	TaAPPara	- increased reactive oxygen species	p-tau peptides	
Alzheimer's disease	IgAITaic	- increased oxidative DNA damage	- KYN and PIC negatively correlated with t-tau	
		- mitochondria impairments	and p-tau peptides	
	APPswe	-		
	PSEN1de9	-		
	SVCT2+/-	-		
	human Aβ-KI	-		
	PINK1	-		
	Parkin	-		
	Parkinson disease protein 7	-		
	CHCHD2	- fragmented mitochondria	- lower activities of KATT and KATTI	
		0	- decreased KYNA	
Parkinson's disease	complex I Park model	- neurodegeneration	- increased 3-HK	
	methyl-4-phenyl-1,2,3,6-tet-		- lower KINA/KIN ratio	
	rahydropyridine	-	- increased QUIN	
	Potonono		- nigner QUIN/KYNA ratio	
	Kotenone	-		
	6-hydroxydopamine	-		
	experimental autoim-	- depolarized fragmented mitochondria	- increased KYN/TRP ratio	
	mune/allergic encephalo-	- trafficking-impaired	- decreased NADH	
	myelitis (EAE)	tunicking imparicu	- higher 3-HK	
	Theiler's murine encephalo-		- higher QUIN/KYNA ratio	
Multiple sclerosis	myelitis virus-induced	-	- Trp, QUIN, KYNA depending on subtypes	
	chronic demyelination		- higher QUIN	
	cuprizone-induced demye-		- higher QUIN/KYN ratio	
	lination	- megamitochondria		
	R6/1	-	- lower Trp	
Huntington's dis-	D6/2		- higher KYN	
ease	K0/2	-	- higher KYN/Trp ratio	
	HTT+97CAG-CAA repeats	-	- higher 3-HK	

**Table 2.** The preclinical models, mitochondrial involvement, and findings in kynurenines in main neurological diseases.

	KI		- higher HAO activity	
(endogenous Hdh promoter)			- lower KYNA	
			- lower KAT activity	
	HdhQ111KI	- multiple mitochondria abnormality	- AA levels correlated with the number of CAG	
			repeats	
	FVB-C9orf72 BAC	-		
	Cu/Zn SOD1-G93A	-		
	TDP43-Q331K	-		
Amyotrophic lat- eral sclerosis	iPSC model of C9orf72-asso- ciated ALS	<ul> <li>swollen mitochondria</li> <li>cluster formation of mitochondria</li> <li>elongated spherical mitochondria</li> <li>mitochondrial fission and apoptosis</li> </ul>	- increased TRP, KYN, QUIN - decreased PIC - KYNA inconclusive	
	SOD1 G93A	-		
	BPA	- Drp1 translocation - mitochondrial RCS		
	BSSG	-		
Migraine	inflammatory soup	<ul> <li>small, fragmented mitochondria</li> <li>reduced mt DNA</li> <li>increased Drp1 fission protein</li> <li>decreased Mfn1 fusion protein</li> <li>valproic acid stabilized mitochondria</li> </ul>	- decreased L-KYN, KYNA, 3-HK, 3-HAA, 5- HIAA, QUIN - increased L-Trp, AA, XA	
	nitroglycerin-induced tri-		-	
	geminovascular activation	-		

# 4.3.2. Parkinson's Disease

PD is a progressive neurological disorder that affects the motor system with muscle rigidity, tremors, and changes in speech and gait. PD patients frequently experience non-motor symptoms (NMSs), such as sensory complaints, mental disorders, sleep disturbances, and autonomic dysfunction. NMSs often occur in PD due to the loss of dopamine-producing cells and the presence of Lewy bodies in the brain, having negative impacts on the quality of life and causing major challenges for disease management [193,194]. Human studies in healthy individuals have revealed that the modulation of autonomic nervous system responses is fundamental for behavioral regulation [195,196]. The pathogenesis of PD is considered to be largely due to the denervation of dopaminergic nigrostriatal nervous system and the aggregations of  $\alpha$ -synuclein [194].

In a familial form of PD, mutations have been identified in genes encoding mitochondrion-associated proteins such as mitochondrial phosphatase and tensin homologue (PTEN)-induced kinase 1 (PINK1), Parkinson juvenile disease protein 2 (parkin), protein deglycase DJ-1 (Parkinson disease protein 7), and coiled-coil-helix-coiled-coil-helix domain containing 2 (CHCHD2) [197]. PINK1 KO mice may use a prodromal model of PD, as the mice show olfactory and gain disturbances [198]. The Parkin KO mouse is a classic transgenic PD model, while there are few studies using DJ-1 KO rats [199]. Homozygous CHCHD2 KO mice mimic PD pathology in an age-dependent manner; they are indistinguishable at birth, but fragmented mitochondria in dopaminergic neurons compared to the wild type [200]. NADH ubiquinone oxidoreductase core subunit S2 (NDUFS2) is a subunit of complex I in neurons that produce dopamine. The mitochondrial complex I-Park model that lacks the gene encoding NDUFS2 shows neurodegeneration [201]. However, transgenic mice that lack the gene encoding another complex I subunit NDUFS4 do not show neurodegeneration in dopaminergic neurons [202]. Complex I inhibitors 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and rotenone are used for pharmacological models of PD, while 6-hydroxydopamine injections are applied for the oxidative stress model of PD [203]. Various types of  $\alpha$ -syn transgenic mice do not develop significant nigrostriatal degeneration [204].

Significantly lower activities of KAT I and KAT II with a decreasing tendency of plasma KYNA levels were observed in the plasma samples of PD patients. Increased 3-HK levels in CSF and a lower KYNA/KYN ratio, increased QUIN levels, and higher QUIN/KYNA ratio were reported in plasma of PD patients [205,206]. In addition, it is reported that the single nuclear polymorphism variants of IDO1 influence the age onset of PD [207]. PD had strong effect sizes for shared deficits in complex I and IV in the peripheral blood, frontal cortex, cerebellum, and substantia nigra [192] (Table 2).

# 4.3.3. Multiple Sclerosis

Multiple sclerosis (MS) is an autoimmune demyelinating neurodegenerative disease leading to the damage of neurons in the CNS. More common symptoms of MS range widely from motor and autonomic dysfunctions to psychobehavioral disturbances including pain, cognitive and emotional changes, and depression [208]. The neural lesion characteristic in MS is numerous plaques which are glial scars in the white matter and spinal cord [209]. The pathogenesis and progression of MS is ascribed at least partly to mitochondrial dysfunction, including reduced fidelity in gene expression, inadequate DNA repair, lower ATP supply, and increased reactive chemical species (RCS), among others [210]. Monitoring the redox status in patients with MS has been proposed to assess stress resilience and thus predictive biomarkers for therapeutic agents [211].

The most characterized animal models of MS are experimental autoimmune/allergic encephalomyelitis (EAE), Theiler's murine encephalomyelitis virus-induced chronic demyelination, and toxin-induced demyelination [212,213]. After three days, immunization in EAE abnormal mitochondrial morphology appears such as vacuolization, swelling, and crista dissolution [214]. Focal intra-axonal mitochondrial alteration proceeds focal axonal degeneration, leading to axon fragmentation, which is triggered by macrophage-derived RCS [215]. Furthermore, the mitochondria of the spinal cord axon are depolarized, fragmented, and trafficking impaired in EAE mice [216]. Cuprizone produces cellular megamitochondria, leading to ATP shortage, RCS production, and ER stress in oligodendrocytes [217]. In addition, cuprizone mobilizes iron molecules from ferritin by chelating copper, leading to iron-mediated lipid peroxidation. The ferroptosis also leads to the production of more RCS via the Fenton reaction [218,219]. Upon withdrawal of the toxin treatment, the mitochondria reverse to normal original morphology [219,220].

Up to only a half of polymorphic loci is identified in the nuclear genome in MS inheritance. The rest of inheritable polymorphic variants may lie in the mitochondrial genome and interaction of mitochondrial and nuclear genes. The allele m.9055\*G is found to be the mitochondrial variant associated with MS. The mitochondrial variants m.4216, m.4580, or m.13708 in biallelic combinations with nuclear gene variants of IL7R, CLEC16A, CD6, CD86, or PVT1 are found to be associated with MS [221]. Regarding KYNs, significantly decreased levels of KYNA were observed in the plasma and the brain tissue of mice treated with cuprizone [222].

The KYN/TRP ratio was significantly increased in the serum of MS patients. The QUIN levels were elevated, while NADH was decreased. 3-HK was found to be significantly higher in MS groups. The QUIN/KYNA ratio was higher in primary progressive MS, secondary progressive MS, and relapsing-remitting MS. KYNA levels were the highest in primary progressive MS, but lower in progressive MS [223]. The QUIN, neurofilament light, and neopterin levels were elevated in the CSF of MS patients compared to

controls [224]. Significantly elevated QUIN/KYN and QUIN/KYNA ratios were observed in the CSF of the relapsing subgroup of relapsing-remitting MS. Trp, KYNA, and QUIN levels were increased in primary progressive MS, while Trp and KYNA levels were decreased in secondary progressive MS [225]. KYNA levels were significantly increased in the plasma of MS patients [226].

# 4.3.4. Huntington's Disease

HD is a fatal autosomal-dominant disease characterized by progressive and irreversible motor dysfunctions, leading to coordination and gait difficulties, as well as cognitive and behavioral changes. The degeneration and neural loss of the striatum, particularly the caudate nuclei, targeting the cerebral cortex, pallidum, thalamus, brainstem, and cerebellum, is a specific neuropathological finding in HD [227]. An abundance of ballooned neurons in the cerebellum, thalamus, and brain stem is another characteristic finding [180]. Mutant huntingtin (HTT) protein is associated with ballooning cell death.

The R6/1 and R6/2 mice are the first transgenic mice generated with a gene containing the promoter and exon 1 of human HTT with 115 or 150 CAT repeats, respectively. The transgenic mouse exhibits cognitive and motor deficits, irregular gait, clasping, weight loss, and seizure, resulting in early death [228]. The R6/2 model exhibited age-dependent changes in mitochondrial respiration in different regions of the brain [229]. The bacterial artificial chromosome transgenic mouse model of HD which carries full-length mutant HTT with a mixture of 97 CAG-CAA repeats exhibits progressive motor dysfunction, synaptic dysfunction, late-onset neuropathology, and neural degeneration [230]. Many KI models of HD have been generated [231]. The KI mice have the advantage of carrying a certain mutation under the endogenous *Hdh* promoter. Some models develop behavioral, molecular, cellular, and neuropathological phenotypes at an early age [231]. The role of aggregates remains unclear. The development of aggregates inhibitors has been under extensive research; however, the aggregates may not play an important role in the pathogenesis [232]. The homozygous HdhQ111 KI mutant huntingtin was found to be associated with the outer mitochondrial membrane, directly induced mitochondrial permeability transition (MPT) pore opening, and significantly decreased the Ca<sup>2+</sup> threshold to trigger MPT pore opening [233]. KI mouse models exhibit the slow progression of behavioral abnormalities; thus, they may help reveal the pathomechanisms of HD and identify a new target for therapeutic intervention [234].

The lower levels of Trp and higher levels of KYN together with the higher KYN/Trp ratios were found in the serum of HD patients, suggesting the up-regulation of IDO activity [235]. The higher levels of 3-HK and QUIN and the higher activity of 3-HAO were observed in the striatum [236,237]. In contrast, the lower levels of KYNA and the lower activity of KATs were found in the plasma and the brain [238,239]. The levels of AA are found to be well correlated with the inflammatory status and the number of CAG repeats [240]. Thus, AA may be a potential prognostic biomarker for HD (Table 2).

## 4.3.5. Amyotrophic Lateral Sclerosis

ALS is a progressive neurodegenerative disease causing the dysfunction of neurons controlling voluntary muscles. ALS often begins with fasciculation, myasthenia, or dysarthria, progressing to the involvement of the muscles responsible for moving, speaking, eating, and breathing [241]. Mitochondrial impairment is an early pathological event in ALS, leading to the death of motor neurons. The dysfunction of mitochondria affects calcium homeostasis, mitochondrial respiration, ATP production, mitochondrial dynamics, and apoptotic signaling. This is caused by the accumulation of ALS-associated mutant proteins such as superoxide dismutase 1 (SOD1), transactive response (TAR) DNA binding protein 43 kDa (TDP-43), fused sarcoma, chromosome 9 open reading frame 72 (C9orf72) gene product, and the C9orf72 GGGGCC repeat expansion-associated glycine/arginine dipeptide repeat protein [242].

Current rodent ALS models include the Friend leukemia virus B (FVB)-C9orf72 bacterial artificial chromosome (BAC) mouse that carries C9orf72 mutations most associated with ALS, Cu/Zn SOD1-G93A mice that encode the human SOD1 protein containing the G93A mutation, and the TDP43-Q331K mouse model that mildly overexpresses human mutant TDP-43 [243]. The FVB-C9orf72 BAC mice develop paralysis and the loss of neuromuscular junction integrity, but the pathological manifestation depends on the mouse strain. The Cu/Zn SOD1-G93A mice show progressive motor dysfunction and loss of motor neurons, but it also depends on the mouse strain, and there is no evident upper motor neuron loss. The TDP43-Q331K mice develop progressive motor dysfunction with motor neuron and axon degeneration, but the progressive degeneration is limited in time, and it does not lead to death [243]. Morphological abnormalities have been observed in ALS models. Mitochondria were swollen in an induced pluripotent stem cell (iPSC) model of C9orf72-associated ALS [244]. The abnormal cluster formation of mitochondria was observed in the axon of SOD1 G93A transgenic mice [245]. Less elongated and more spherical mitochondria were isolated from the motor neuron of SOD1 G93A transgenic mice [246]. Expressions of wild-type or ALS TDP-43 mutants lead to abnormal morphology including aggregated, fragmented, and vacuolated mitochondria [247].

Environmental factors are considered to play a role in the pathogenesis of ALS and tabacco smoke has been linked to that of ALS. Exposure to bisphenol A (BPA) (a chemical used in the production of polycarbonate plastics) and beta-sitosterol beta-D-glucoside (BSSG) (an estrogen receptor-binding phytosterol) has been found to be neurotoxic to motor neurons; thus, those compounds have been applied to environmental models of ALS [243]. BPA induces neurotoxicity and neurodegeneration through alternations of mitochondrial functions, leading to fission and apoptosis via the translocation of dynamin-related protein 1 (Drp1) from the cytosol [248]. The neurotoxic effects of BSSG appear to be caused by mitochondrial production RCS associated with succinate oxidation (Complex II) [249].

Significantly increased TRP, KYN, and QUIN in serum and CSF, as well as significantly decreased PIC in serum, were observed in ALS. The neuronal and microglial expression of IDO and the levels of QUIN were increased in the motor cortex and spinal cord of ALS patients [250]. The levels of KYNA in ALS remain inconclusive, as studies showed significantly higher levels in the CSF, significantly lower levels in the serum, or no significant difference between healthy control and ALS patients [237]. Thus, the levels of KYNA may depend on the subgroup, the severity, and the stage of ALS, and further studies may reveal a potential role of KYNA measurement for biomarkers (Table 2).

## 4.3.6. Migraine

Migraine is a primary headache disorder, characterized mostly by a headache on one side of the head. The exact pathomechanism of the disease is not fully known, but morphological and biochemical studies have shown that the pathophysiology of migraine is linked to mitochondrial dysfunction [251]. Abnormal mitochondria have been identified in patients with migraine with aura [252] and with familial hemiplegic migraine [253]. In addition, increased levels of lactate were shown in the blood and cerebrospinal fluid of patients with migraines, which clearly suggests a defective oxidative function [254,255]. The activities of mitochondrial enzymes including monoamine oxidase, succinate dehydrogenase, NADH dehydrogenase, cyclooxygenase, and citrate synthetase were found to be reduced in the platelets of migraineurs [256,257]. Furthermore, the biochemical changes were restricted to enzymes of the respiratory chain encoded by mtDNA [252]. OXPHOS has been found to be impaired in the brain of patients with migraines during and between migraine attacks [258–260]. This impairment is seen as increased levels of ADP, decreased levels of organic phosphate, and a decreased phosphorylation potential [261]. Reyngoudt et al. found that brain ATP decreased by 16% between attacks in patients with migraine without aura compared with healthy controls [262].

The mitochondrial involvements in migraine have been reported in animal models of migraine. The rodent inflammatory soup model revealed that mitochondria were small and fragmented, and that the number of mtDNA was significantly reduced in the trigeminal neurons. Furthermore, fission protein Drp1 was increased, while fusion protein mitofusin (Mfn) 1 was decreased, suggesting that mitochondrial dynamics were under disturbance after repeated dural stimulation [263]. The same chronic migraine model also showed that the trigeminal nucleus caudalis decreased spare respiratory capacity, i.e., the amount of ATP to be produced by oxidative phosphorylation in case of a sudden increased demand [264]. Neuroprotective, antiepileptic, and migraine prophylactic agent valproic acid stabilized the mtDNA copy number, restored the ATP level, and maintained the mitochondrial membrane potential in a rat model of nitroglycerin-induced trigeminovascular activation [265].

In the serum of chronic migraineurs, the levels of L-KYN, KYNA, 3-HK, 3-HAA, 5-HIAA, and QUIN were decreased, while the levels of L-Trp, AA, and XA were significantly higher compared to healthy controls [266]. The similar results were observed in patients with episodic or chronic cluster headache [267]. The levels of L-Trp, L-KYN, KYNA, 3-HAA, 5-hydroxyindolacetic acid, PIC, and melatonin were decreased in the plasma of episodic migraineurs in the interictal period. The tendency was more prominent in those without aura. In addition, the levels of 3-HAA, 5-hydroxyindoleacetic acid, and melatonin were increased in the ictal period [268]. The expression of KAT II was decreased in the upper cervical spinal cord (C1-C2) in nitroglycerin-induced trigeminovascular activation of rats [269]. Preclinical and clinical findings suggest that mitochondrial dysfunction and KYN metabolites play a role in the pathomechanism of migraine [270]. Furthermore, KYNs are also involved in the pathogenesis of chronic pain and their adjacent position to serotonin metabolism is drawing close attention to development of anti-migraine drugs [271–274]. The acute administration of antidepressant SSRI citalopram altered Trp-KYN metabolism in patients with migraines [275]. In addition, gastrointestinal disorders have been linked to migraines through Trp-KYN metabolism [276] (Table 2).

## 4.4. Psychiatric Disorders Linked to Mitochondrial Dysfunction

A growing number of researchers cast more attention on the contribution of mitochondria in mental health, susceptibility by genetic variants, and its interaction with environmental factors. Clinical and preclinical studies are revealing evidence that the organelles play a key role in psychiatric disorders and neurodevelopmental disorder such as MDD, generalized anxiety disorder (GAD), PTSD, SCZ, ADHD, and ASD. Furthermore, preclinical animal research plays a major role in revealing the involvement of endogenous neurotransmitters, neurohormones, and metabolites [277–280].

# 4.4.1. Major Depressive Disorder

MDD is a mental disorder with at least two weeks of low mood, often accompanied by low self-esteem, loss of interest, low energy, and pain without a cause. The lifetime prevalence of MDD ranged from 2 to 21% [281]. The pathogenesis of depression has been linked to air pollution and depressive symptom frequently presents with cormobid conditions including anxiety, cognitive impairment, and chronic pain [282–285]. The monoamine hypothesis has prevailed for the pathogenesis of depression. The hypothesis holds that depression is caused by the depletion of 5-HT, norepinephrine, or dopamine in the CNS [286]. The atrophic lesion and synaptic impairment in the PFC and hippocampus and hypertrophy and increased synaptic activity in the nucleus accumbens, and amygdala are observed [287]. SSRIs are commonly used as a first-line treatment for MDD. However, only 42–53% of patients treated with SSRIs see an improvement, and medication for treatment-resistant depression remains a challenge. Novel treatment is under extensive study such as intermittent theta-burst stimulation [288,289]. Furthermore, psychotherapy is an effective treatment of choice, which can serve as a powerful measure for patients who cannot tolerate medication, and KYNs may be potentially useful as prognostic biomarkers [290].

A chronic mild stress (CMS) model of depression showed decreased ATP production, decreased hippocampal Na<sup>+</sup> and K<sup>+</sup>-ATPase activity, and anhedonia in the sucrose preference test [291]. The damaged structure, impaired respiration rate, and altered membrane potentials of mitochondria were observed in the hippocampus, hypothalamus, and the cortex of CMS mice which exhibit anhedonia in the sucrose preference test and depression-like behavior in the tail suspension test (TST) [292]. Furthermore, SSRI fluoxetine demonstrated Na<sup>+</sup> and K<sup>+</sup>-ATPase activity, mitochondrial respiration, and sucrose preference in the chronic unpredictable stress model [293]. Thus, mitochondrial dysfunction may be involved in depression-like behavior.

Transgenic models for depression have been generated by manipulating genes responsible for the metabolism of 5-HT. Tryptophan hydroxylase (TPH) is the rate-limiting enzyme in 5-HT biosynthesis. The  $Tph1^{-/-}$  mice produced a normal level of 5-HT in the brain and showed no significant change in behavior [294,295]. The  $Tph2^{-/-}$  mice exhibited depressive-like behavior in TST and anxiety-related behavior in the marble burying test [296]. However, it was reported that Tph2 null mutants ( $Tph2^{-/-}$ ) mice showed slightly reduced depression-like and anxiety-like behaviors, but significantly increased fear-conditioning responses. The behaviors, including impulsivity, aggressiveness, and emotional reactivity of  $Tph2^{-/-}$  mice, are sex-dependent [297]. The double KO  $Tph1/Tph2^{-/-}$  mice showed depressive-like behavior in TST and anxiety-related behavior in the marble burying test but antidepressive-like behavior in the forced swim test (FST) with reduced levels of 5-HT in the brain and periphery [298]. In addition, KI of the TPH2 variant (R439H) in mice showed depression-like behavior in TST [299].

MDD patients showed moderate effect sizes for similar abnormality patterns in the expression of complex I of samples from frontal cortex, cerebellum, and striatum [192]. The levels of Trp, KYN, and KYNA were decreased in the plasma of MDD patients, and the levels of QUIN were increased in MDD patients without antidepressant treatment. The immunoreactivity of QUIN was increased in the PFC and hippocampus of the postmortem brain tissues from MDD patients [300,301]. An increased risk of depression was reported following the activation of the Trp-KYN metabolic system in chronic illnesses [302]. Furthermore, KYNA may serve as a diagnostic and predictive biomarker in depression [303]. The serum KYNs were found to be correlated with depression in poststroke patients [304]. KYN analogues have been extensively researched in the search for novel antidepressants [305] (Table 3).

Psychiatric	Preclinical Models	Mitochondrial	Findings in Kynurenines	
Diseases	Treemieur Woucis	Involvement		
	CMS	- decreased ATP, ATPase activity		
	TST	- altered membrane potential		
	FST	-		
Major depressive disor- der	Tph1 <sup>-/-</sup>	-	- decreased Trp, KYN, KYNA - increased QUIN	
	Tph2-/-	-		
	Tph1/Tph2-/-	-		
	TPH2 variant (R439H) KI	-		

**Table 3.** The preclinical models, mitochondrial involvement, and findings in kynurenines in main psychiatric diseases.

Generalized anxiety dis- order	outbred Wistar rats	<ul> <li>reduced mitochondrial GTPase expression</li> <li>altered mitochondrial morphology and functions</li> </ul>	- decresed KYN
	social hierarchy	- NAc mitochondrial bioenergetic profiles	
	FKBP5-/-	-	
	PAC1R <sup>-/-</sup>	-	
	5-HT1AR-/-	-	
Post-traumatic stress	COMT-/-	-	
disorder	GAD6-/-	-	-
	GABAB1a-/-	-	
	CB1R-/-	-	
	single prolonged stress model	- abnormal apoptosis	
Bipolar disorder	Clock∆19	-	
	dominant negative mutant of mtDNA Polg1	-	- reduced KYNA - increased 3-HK/KYN, 3-HK/KYNA ratio - increased KYNA in CSF
	-	- complex I expression abnormality	
Substance use disorder	-	- reduced mitochondrial copy numbers	- higher 5-HT - lower KYN/5-HT ratio
Schizophrenia	DISC1	- affect mitochondrial transport, fission, and fusion	- higher KYN and KYN/TRP ratio - increased KYN, KYNA
Ĩ	hypertensive rats	-	- decreased KYNA/KYN ratio
	ND6 <sup>p25L</sup> KI	-	
	Shank3 <sup>Δc/Δc</sup>	-	
Autism spectrum disor-	Cntnap2 KO	-	- lowered KYNA
der	ADGRL3-/-	-	- higher KYN/KYNA ratio - higher KYN/Trp ration, KYN, QUIN
	valproate	-	
	polyinosinic–polycytidylic acid	- mitochondrial dysfunction	
Attention-deficit hyper-	Ptchd1-/-	-	- lowered Trp, KYNA, XA, 3-HAA
activity disorder	-	- higher mtDNA copy number	- higher Trp, KYN

# 4.4.2. Generalized Anxiety Disorder

GAD is a mental disorder characterized by excessive, uncontrollable, and irrational anxiety. GAD has a combined lifetime prevalence of 3.7% [306]. 5-HT, dopamine, norepinephrine, and gamma-aminobutyric acid (GABA) are linked to anxiety [307]. The amygdala in the middle of the brain which processes emotion, memory, and fear is involved in GAD [308]. Benzodiazepines such as alprazolam, clonazepam, and diazepam bring relief in 30 min; SSRIs are the first line of treatment in GAD; and cognitive behavioral therapy is the most effective form of psychotherapy [309].

An increasing number of preclinical studies are revealing that anxiety is linked to mitochondrial functions including bioenergetics, oxidative stress, neurosteroid production, biogenesis, and apoptosis [310]. Mitochondrial dysfunction is located in the nucleus accumbens (NAc) which interfaces motivation and action, playing a key role in motivation, aversion, reward, and reinforcement learning. The shell of NAc is considered to be part of the extended amygdala [310]. The outbred Wistar rats which exhibit anxiety-like behavior showed reduced expressions of the mitochondrial GTPase MFN 2 in the NAc, altered mitochondrial morphology and functions, and the morphology of medium spiny neurons (MSNs) projecting from the NAc. The behavioral, mitochondrial, and neuronal phenotypes were reversed by the viral overexpression of MFN2 [311]. Furthermore, more anxious rats are prone to become subordinate during a social encounter with less anxious rats and social hierarchy is linked to the mitochondrial bioenergetic profiles of the NAc. Thus, anxiety appears to directly influence social dominance mediated by mitochondrial functions [312].

The levels of KYN were reduced in endogenous anxiety and normalized after treatment in the plasma samples [313]. The levels of KYN were lower in people with Type D personality, the joint tendency towards negative affectivity and social inhibition [314].

# 4.4.3. Post-Traumatic Stress Disorder

PTSD is a behavioral and mental disorder that develops after experiencing a traumatic event. Individuals with PTSD suffer from flashbacks, nightmares, severe anxiety, and uncontrollable thoughts regarding the event [315]. The lifetime prevalence of PTSD ranges from 6.1 to 9.2% [316]. PTSD is considered to be caused by insufficient integration of a trauma memory into the hippocampal-cortical memory networks, forming fragmented, incomplete, and disorganized intrusive memories [317]. The primary treatment of PTSD is psychotherapy, and SSRIs such as sertraline and paroxetine are considered first-line therapy for PTSD [318].

Animal models are revealing the pathogenesis of PTSD. The genetic factors contributing to the development of PTSD include the stress response system, such as the hypothalamic–pituitary–adrenal (HPA) axis; neuroplasticity such as brain-derived neurotrophic factor (BDNF); and monoamine neurotransmission such as serotonergic, dopaminergic, glutamatergic, and GABA-ergic systems [319]. FK506-binding protein 51 (FKBP5) is a co-chaperone which modulates glucocorticoid receptor activity. FKBP5<sup>-/-</sup> mice prevents the age-induced impairment of stress resilience [320]. Chronic variate stress increases the bed nucleus of the stria terminalis pituitary adenylate cyclase activating polypeptide (PACAP) [321]. The pituitary adenylate cyclase 1 receptor type 1 KO (PAC1R<sup>-/-</sup>) mice show reduced anxiety [322]. The BDNF promotor IV-disrupted mutant Bdnf-e4 mice and BDNF Met-Val mutant mice showed impaired fear extinction [323,324].

Serotonin 1A receptor KO 5-HT1AR<sup>-/-</sup> mice show increased fear memory to contextual cues [325]. 5-HT transporter (5-HTT) gene KO 5-HTT<sup>-/-</sup> mice show impaired stress response and impaired fear extinction with abnormal corticolimbic structure [326]. Dopamine is degraded by catechol-O-methyltransferase (COMT). COMT gene KO COMT<sup>-/-</sup> mice showed an increased response to repeated stress exposures [327]. GABA is synthesized from L-glutamic acid by glutamic acid decarboxylase. The 65-kDa isozyme of glutamic acid decarboxylase 2 KO GAD6<sup>-/-</sup> mice shows increased generalized fear and impaired extinction of cued fear [328,329]. GABA receptor subunit B1a KO GABAB1a<sup>-/-</sup> mice shows a generalization of conditioned fear to nonconditioned stimuli [330]. Cannabinoids directly interact with GABAergic neurotransmission. Cannabinoid 1 receptor KO CB1R<sup>-/-</sup> mice show an increased response to repeated stress exposures [331].

Mitochondrial functions are linked to PTSD-like behavior in preclinical studies. Following exposure to a trauma, mice with PTSD-like symptoms exhibit reduced activities of mitochondrial electron transport in the cerebellum and the dysfunction of fatty acid oxidation in cerebellum and plasma. The activity of cerebellar mitochondrial electron transport complex is negatively correlated with PTSD-like symptoms [332]. Abnormal apoptosis has been observed in the brain areas closely associated with emotion and cognition, including the hippocampus, the amygdala, and the medial PFC in single prolonged stress model of PTSD [333]. In addition, the decreased kynurenine pathway potentiates resilience to the social defeat effect of a cocaine reward [334]. Early intervention with a glucocorticoid receptor antagonist a RU486 facilitates the correction of traumatic stressinduced fear and anxiety dysregulation [335]. No clinical study was reported regarding the peripheral or CSF samples of KYNs in patients with PTSD. KYN metabolites are monitored in clinical settings as evidence of inflammatory responses contributing to sleep deprivation and the formation of intrusive memories [336] (Table 3).

# 4.4.4. Bipolar Disorder

BD is a mental disorder characterized by mood oscillations with episodes of mania and depression. A large cross-sectional survey of 11 countries found the overall lifetime prevalence of BD was 2.4% [337]. Neuroimaging and postmortem studies have found abnormalities in a variety of brain regions, and the most commonly implicated regions include the ventral PFC and the amygdala [338,339]. Dysfunctions in emotional circuits located in these regions have been hypothesized as a mechanism for BD. The left side of the hippocampus regulates verbal and visual memory. This part of the brain also helps regulate how you emotionally respond to situations. When your mood shifts, your hippocampus changes shapes and shrinks [340]. Patients with BP showed diminished GABA neurotransmission. Thus, low GABA levels can result in excitatory toxicity [341].

There have been no established animal models of BD that exhibit both manic and depressive episodes. Typical current animal models of mania involve drug-induced hyperactivity or genetically modified animals that exhibit continuous hyperactivity. The targeting of circadian rhythm genes to disrupt mechanisms regulating the circadian rhythm has been widely used to create animal models for BD [342].

The most common model is the  $Clock \Delta 19$  mutant mouse. These mice carry a deletion at exon 19 of the *Clock* gene, resulting in a dominant-negative protein, unable to activate transcription [343]. Mutant mice exhibit mania-like behavior and altered sleep patterns [344]. The dominant negative mutant of mtDNA *Polg1* transgenic mice showed recurrent hypoactive periods [345]. The withdrawal of lithium provokes depression in mice, while antidepressant medications alleviate depressive symptoms [346]. Thus, the transgenic strain appears to be a good animal model for BD.

Meta-analysis revealed that BD showed moderate effect sizes for similar abnormality patterns in the expression of complex I of samples from frontal cortex, cerebellum, and striatum [192]. The dysfunctional mitochondrial hypothesis is one of the current hypotheses that attempt to explain the origin of mood disorders. Many studies have confirmed that mood stabilizers affect mitochondrial functions, even though the exact mechanism or localization of action is unknown [347].

Regarding the KYN system a case-control study showed that KYNA levels were reduced and the 3-HK/KYN and 3-HK/KYNA ratio was increased in BD compared to healthy control [348]. However, a meta-analysis reported no significant difference of TRP and KYN levels, KYN/TRP and KYNA/QUIN ratios in serum from BD patients [349] KYNA was significantly increased in CSF of BD patients [350] (Table 3).

# 4.4.5. Substance Use Disorders

Substance use disorders (SUDs) represent a type of mental disorder that affects the brain and behavior, leading to an inability to control the use of a drug or medication. The exact cause of SUDs is not known, but the known risk factors are the genes, the action of the drug, peer pressure, emotional distress, anxiety, depression, and environmental stress [351]. In addition to an impaired control, common substances are alcohol, sedatives, caffeine, hallucinogens, inhalants, stimulants, and tobacco, among others [352]. The main brain area associated with SUDs is the limbic system, comprising the cingulate gyrus, amygdala, hippocampus, PFC, ventral tegmental area, and the nucleus accumbens. The system is related to reward, emotion, and punishment [353].

The mitochondrial copy numbers were found to be reduced in blood samples of patients with opioid use disorder; however, the link between changes in the reward neural circuitry and the peripheral measurements remains unclear [354]. No clinical study was found regarding KYNs in patients with SUDs. Although there are few studies on this, growing attention is paid to a relationship between KYN metabolites and SUDs, the alteration of the Trp-KYN system, and a potential approach to SUDs (including ethanol, nicotine, cannabis, amphetamines, cocaine, and opioids) [355]. Furthermore, 5-HT concentration was significantly higher and the KYN/5-HT ratio was significantly lower in plasma of patients with cocaine use disorder in SUD-induced MDD compared to those with MDD, but there were no differences between SUD primary MDD and MDD. This may suggest that the Trp-KYN pathway participates less in SUD-induced MDD [356].

## 4.4.6. Schizophrenia

SCZ is a mental disorder characterized by abnormally interpret reality, hallucinations, delusions, apathy, lack of social functioning, and extremely disordered thinking and behavior. Cognitive symptoms, including concentration and attention difficulties, as well as memory impairments, can be subtle [357]. Cognitive deterioration in patients with SCZ has been linked to vitamin D deficiency which may directly affect processing speed [358]. SCZ is generally considered to be a neurodegenerative disorder with neurodevelopmental antecedents. The underlying changes occur before the onset of symptoms arising from the interaction between genes and the environment, leading to deficits in the neural circuitry in the age of 18–25 [359]. Maternal infections, malnutrition, and complication during pregnancy and parturition are risk factors [360]. In total, 30–50% of SCZ patients develop antipsychotic-resistant SCZ which is associated with a high level of dissociation, a loss of integrity between memories, and perceptions of reality [361]. Many people with SCZ have hypertension, disturbance of lipid metabolism, and other mental disorders (including SUDs, MDD, GAD, and OCD) [362–364].

Disrupted in schizophrenia 1 (DISC1), encoded by the *DISC1* gene, is a protein which plays a role in presynaptic regulation of dopamine. DISC1 alterations increase the risk of SCZ [365]. DISC1 also plays various roles in many other cellular functions, including mitochondrial transport, fission, and fusion. The dynamic processes of mitochondrial transport, fission, and fusion determine mitochondrial morphology, localization, and network [366]. DISC1 mouse models display abnormal changes relevant to SCZ. The neuro-anatomical changes include displaced dentate granule neurons, altered axonal targeting, reduced dendrite growth, and dendritic spine density. The behavioral abnormalities include the impairment of working memory [367]. Furthermore, the spontaneously hypertensive rat strain haves been proposed to an animal model of SCZ, which exhibits abnormal behaviors resembling cognitive, psychotic, and negative symptoms [368].Neuro-degenerative changes in SCZ are caused by a series of malfunction including mitochondrial impairment, oxidative stress responses, and the activation of immune responses, leading to chronic low-grade inflammation [17]. Clinical studies linked mitochondrial impairment with increased risk of SCZ and suggested that abnormal mitochondrial dynam-

ics contribute to compromising normal neural connectivity in the brain [369,370]. Furthermore, meta-analysis revealed that SCZ showed moderate effect sizes for similar abnormality patterns in the expression of complex I in samples from the frontal cortex, cerebellum, and striatum [192].

Regarding the Trp-KYN metabolic system, KYN and the ratios of KYN/TRP were higher in the serum of SCZ patients [371]. Meta-analyses showed increased KYN and KYNA levels in CSF samples of SCZ patients and increased levels of KYNA in plasma, CSF, brain tissue, or saliva, respectively [53,256]. Thus, the KYN system is activated in SCZ, and elevated KYNA levels are considered to contribute to the impairment of cognitive function. However, another meta-analysis reported that KYNA levels and the KYNA/3-HK ratio were not altered, and the KYNA/KYN ratio was decreased in SCZ, suggesting the presence of differential patterns between SCZ and mood disorders [53] (Table 3).

## 4.4.7. Autism Spectrum Disorder

ASD is a neurodevelopmental disorder characterized by persistent deficits in social interaction, restricted-repetitive patterns of behavior, and the loss of interests or activities [372]. These social impairments may be related to the interpretation of social signals [373]. Potentially threatening situations, such as the proximity of others, can trigger a number of physiological responses that help to regulate the distance between themselves and others during social interaction, showing the critical role of social signal interpretation in social interaction. Individuals with ASD have social impairments, potentially due to the lack of social signal interpretation, resulting in an inability to interpret these signals to guide appropriate behaviors. The prevalence of mitochondrial diseases is higher in the population of ASD than in general population and up to a half of children with ASD showed evidence of mitochondrial dysfunction [374]. However, most mitochondrial disease-associated ASD is not associated with genetic abnormalities, suggesting secondary mitochondrial impairment, such as environmental factors [375].

Animal models of ASD include prenatal exposure to valproate during pregnancy, inbred strains of mice expressing autism traits, and genetical modification targeting autism-related genes (including mtDNA) [376]. The insertion of mtDNA ND6 gene missense mutation (ND6<sup>P25L</sup>) exhibits ASD endophenotypes, including autism-like behaviors and electroencephalographic profiles, and correlates with mitochondrial respiration and increased RCS of the brain, suggesting a link to mitochondrial dysfunction [377]. The hemideletion of the Src-homology 3 and the multiple ankyrin repeat domain 3 (SHANK3) gene is found in patients with Phelan–McDermid syndrome, demonstrating ASD-like behaviors. The homozygous Shank3<sup>Δc/Δc</sup> mice with C-terminal 508 deletions exhibit significant impairments in social novelty preference, stereotyped behavior, and gait [377]. The contactin-associated protein-like 2 (CNTNAP2) encodes a neurexin that regulates the interactions of neurons and glial cells. The Cntnap2 knockout mice show ASD-like behavior [378]. The adhesion of G protein-coupled receptor L3 (ADGRL3) encodes latrophilins. AD-*GRL3<sup>-/-</sup>* mice show hyperactivity and less depression-like behavior. Preclinical studies also showed links between the pathogenesis of ASD and maternal immune activation, maternal microbiota profile, and exposure to nutritional and toxic metals during mid-fatal development [379-382].

The alteration of the Trp-KYN metabolic system was also observed in patients with ASD. The levels of KYNA were significantly lower, and the ratio of KYN/KYNA was significantly higher in the serum of children with ASD [383]. The ratio of KYN/Trp and the levels of KYN and QUIN were significantly higher in blood samples of ASD patients, but there was no significant difference in KYNA and the levels of PIC were significantly lower in ASD patients [121] (Table 3).

#### 4.4.8. Attention-Deficit Hyperactivity Disorder

ADHD is a behavioral and neurodevelopmental disorder characterized by inattention, hyperactivity, and impulsivity, which are pervasive, impairing, and otherwise ageinappropriate [384]. ADHD is associated with SUDs, alcoholism, and other mental disorders, including MDD, GAD, and ASD [385]. Furthermore, multidirectional relationships between stress, anxiety, and inflammation in the pathogenesis of ADHD are discussed recently [386].

Patched domain-containing protein (*Ptchd*) is a membrane protein with a patched domain. The deletion of the *Ptchd* gene has been identified in patients with intellectual disability and ASD. The *Ptchd1* KO mice exhibit ADHD-like behaviors. No changes in Trp and 3-HK were found, but significant increased levels of KYN, KYNA, AA, and 3-HK were observed in the serum of the *Ptchd1* KO mice. Meanwhile, significantly increased levels of AA, 3-HK, and 3-HAA were observed in the frontal cortex of *Ptchd1* KO mice, but there were no changes in KYNA levels [387]. A clinical study showed that lower concentrations of Trp, KYNA, and XA, 3-HAA were found in the serum of patients with ADHD, and that higher levels of Trp and KYN were associated with higher scores of ADHD symptoms [388].

Mitochondria may be sensitive to psychological stress in early life [389]. People who experienced childhood trauma appear to possess a larger number of mitochondrial genomes per cell [389]. Indeed, mtDNA copy number was observed to be higher in the peripheral blood of ADHD patients, which suggests a possible link to mitochondrial impairments in the pathogenesis of ADHD [390] (Table 3).

### 5. Conclusions and Future Perspective

This review article recapitulated the involvement of mitochondria with an emphasis on its connection to the Trp-KYN metabolic system in clinical manifestations of neuropsychiatric symptoms and advances in preclinical research in major neurological and psychiatric disorders. Growing evidence has revealed that mitochondria have a close link to KYN metabolism and that mitochondrial dysfunction and the activation of the KYN system contribute to the pathogenesis of neuropsychiatric disorders. Extensive clinical and preclinical research has helped delineate the multifunctional facets, compartmentalization, and dynamic nature of mitochondria, including cell differentiation, cell-type determination, cell movement, and pattern formation. The pathological changes in functions, morphologies, and dynamics have been probed in mitochondrial diseases, as well as diseases linked to mitochondrial dysfunction. For example, functional magnetic resonance imaging, measurements of fibroblast mitochondrial spare respiratory capacity, the NAD<sup>+</sup>/NADH ratio, Complex II levels, and a combination of the detection of amyloid and/or tau protein and signs of neuronal injury on brain imaging or cerebrospinal fluid sampling are emerging techniques used to assess mitochondrial functions.

However, little is known about the reversibility, plasticity, and/or resilience of mitochondrial functions, integrity, dynamics, and/or network formation. The measurement of such parameters is of particular importance. Revealing the link between mitochondria and the KYN metabolic system may be a promising option for this direction of research. The development of the mitochondrial stress test, for example, which assesses recoverability, may help with the early detection of mitochondria-related diseases and the possible application of prophylactic measures. For this purpose, engineering fine preclinical models high in construct, face, and predictive validity is an essential step. Two-hit models consisting of a certain genetic susceptibility and environmental trigger with pharmacological agents which initiate neuropsychiatric manifestations can help develop preventive measures, understand the pathomechanism, make accurate diagnoses, delay disease progression, and choose the most appropriate therapeutic option.

**Author Contributions:** Conceptualization, M.T.; writing—original draft preparation, M.T., E.S. and H.P.; writing—review and editing, M.T., Á.S., E.S, H.P., F.T. and L.V.; visualization; Á.S.; project

administration, L.V.; funding acquisition, M.T. and L.V. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by National Research, Development and Innovation Office – NKFIH 138125, MTA-JSP-050609, and ELKH-SZTE. Helga Polyák was supported by the New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research Development and Innovation Fund [ÚNKP-21-3] and EFOP-3.6.3-VEKOP-16-2017-00009.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest regarding this review article.

## Abbreviations

AA	anthranilic acid
Acetyl-CoA	acetyl coenzyme A
ACMS	2-amino-3-carboxymuconate semialdehyde
ACMSD	2-amino-3-carboxymuconate-6-semialdehyde decarboxylase
acyl-CoA	acyl coenzyme A
AD	Alzheimer's disease
ADHD	attention-deficit hyperactive disorder
AHR	aryl hydrocarbon receptor
ALS	amyotrophic lateral sclerosis
AMPA	lpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
AMS	2-aminomuconic-6-semialdehyde
AMSD	2-aminomuconate semialdehyde dehydrogenase
Αβ	amyloid beta
ASD	autism spectrum disorder
ATP	adenosine triphosphate
BAC	bacterial artificial chromosome
BCG	Bacillus Calmette-Guérin
BD	bipolar disorder
BPA	bisphenol A
BSSG	beta-sitosterol beta-d-glucoside
CA	cinnabarinic acid
CMS	chronic mild stress
C9orf72	chromosome 9 open reading frame 72
Cys	cysteine
EAE	experimental autoimmune/allergic encephalomyelitis
ER	endoplasmic reticulum
ETC	electron transport chain
FA	Friedreich's ataxia
FADH <sub>2</sub>	flavin adenine dinucleotide
FVB	Friend leukemia virus B
GABA	gamma-aminobutyric acid
GAD	generalized anxiety disorder

GTP	guanosine triphosphate
GPR35	G-protein-coupled receptor 35
H⁺	proton
HD	Huntington's disease
5-HT	serotonin
i.c.v.	intracerebroventricular
IDO	indoleamine 2,3-dioxygenase
IDO1	indoleamine 2.3-dioxygenase isoform 1
IDO2	indoleamine 2.3-dioxygenase isoform 2
3-HAA	3-bydroxyanthranilic acid
3-НК	3-hydroxy-I -kynurenine
3-HAO	3-hydroxyapthrapilate ovidace
UTT	huntingtin
1111 5 UTT	
3-H11	5-rii transporter
iPSC	induced pluripotent stem cells
KAT	kynurenine aminotransferase
KFA	kynurenine formamidase
KI	knock-in
КМО	kynurenine 3-monooxygenase
КО	knockout
KYN	kynurenine
KYNA	Kynurenic acid
KYNU	kynureninase
MDD	major depressive disorder
MELAS	mitochondrial encephalomyopathy with lactic acidosis and stroke-like epi- sodes
MPT	mitochondrial permeability transition
MS	multiple sclerosis
mtDNA	mitochondrial DNA
MTF	mitofusin
NaAD	nicotinic acid adenine dinucleotide
NADH	nicotinamide adenine dinucleotide
NaMN	nicotinic acid mononucleotide
nDNA	nuclear DNA
NDUFS2	NADH:Ubiquinone Oxidoreductase Core Subunit S2
NMDA	N-methyl-D-aspartic acid
	obsessive compulsive disorder
OXPHOS	ovidative phosphorylation
parkin	Parkinson juvenile disease protein 2
Pi	inorganic phosphate
PINK1	phosphatase and tensin homolog (PTEN)-induced kinase1
PD	Parkinson's diseases
PFC	prefrontal cortex
PIC	picolinic acid
PLP	pyridoxal 5'-phosphate
PMD	primary mitochondrial disease
PMM	primary mitochondrial myopathy

PTSD	post-traumatic stress disorder
QPRT	quinolinate phosphoribosyltransferase
QUIN	quinolinic acid
RCS	reactive chemical species
SCZ	schizophrenia
SNP	single-nucleotide polymorphism
SMD	secondary mitochondrial dysfunction
SOD1	superoxide dismutase 1
SSRI	selective serotonin reuptake inhibitor
succinyl-CoA	succinyl coenzyme A
SUD	substance use disorder
TDP-43	transactive response (TAR) DNA binding protein 43 kDa
TDO	tryptophan 2,3-dioxygenase
TPH	tryptophan hydroxylase
Trp	tryptophan
TST	tail suspension test
ХА	xanthurenic acid

## References

- 1. Fiske, C.H.; Subbarow, Y. Phosphorus Compounds of Muscle and Liver. *Science* **1929**, *70*, 381–382. https://doi.org/10.1126/science.70.1816.381-a.
- 2. Lohmann, K. Über die Pyrophosphatfraktion im Muskel. Naturwissenschaften 1929, 17, 624–625. https://doi.org/10.1007/BF01506215.
- 3. Maruyama, K. The Discovery of Adenosine Triphosphate and the Establishment of its Structure. *J. Hist. Biol.* **1991**, *24*, 145–154. https://doi.org/10.1007/BF00130477.
- 4. Krebs, H.A.; Johnson, W.A. Metabolism of ketonic acids in animal tissues. *Biochem. J.* **1937**, *31*, 645–660. https://doi.org/10.1042/bj0310645.
- Friedkin, M.; Lehninger, A.L. Esterification of inorganic phosphate coupled to electron transport between dihydrodiphosphopyridine nucleotide and oxygen. J. Biol. Chem. 1949, 178, 611–644. https://doi.org/10.1016/S0021-925856879-4.
- 6. Belitser, V.A.; Tsibakova, E.T. About phosphorilation mechanism coupled with respiration. *Biokhimiya* **1939**, *4*, 516–534.
- 7. Gano, L.B.; Patel, M.; Rho, J.M. Ketogenic diets, mitochondria, and neurological diseases. J. Lipid Res. 2014, 55, 2211–2228. https://doi.org/10.1194/jlr.R048975.
- 8. Knoop, F. Der Abbau aromatischer Fettsäuren im Tierkörper. Beitr. Chem. Physiol. Pathol. 1904, 6, 150–162.
- 9. Klingman, D.; Handler, P. Partial purification and properties of renal glutaminase. J. Biol. Chem. 1958, 232, 369–380.
- 10. Ernster, L.; Ikkos, D.; Luft, R. Enzymatic activities of human skeletal muscle mitochondria: A tool in clinical metabolic research. *Nature* **1959**, *184*, 1851–1854. https://doi.org/10.1038/1841851a0.
- Holt, I.J.; Harding, A.E.; Morgan-Hughes, J.A. Deletions of muscle mitochondrial DNA in patients with mitochondrial myopathies. *Nature* 1988, 331, 717–719. https://doi.org/10.1038/331717a0.
- Wallace, D.C.; Singh, G.; Lott, M.T.; Hodge, J.A.; Schurr, T.G.; Lezza, A.M.; Elsas, L.J., 2nd; Nikoskelainen, E.K. Mitochondrial DNA mutation associated with Leber's hereditary optic neuropathy. *Science* 1988, 242, 1427–1430. https://doi.org/10.1126/science.3201231.
- 13. Bourgeron, T.; Rustin, P.; Chretien, D.; Birch-Machin, M.; Bourgeois, M.; Viegas-Péquignot, E.; Munnich, A.; Rötig, A. Mutation of a nuclear succinate dehydrogenase gene results in mitochondrial respiratory chain deficiency. *Nat. Genet.* **1995**, *11*, 144–149. https://doi.org/10.1038/ng1095-144.
- 14. Carafoli, E. The interplay of mitochondria with calcium: An historical appraisal. *Cell Calcium* **2012**, *52*, 1–8. https://doi.org/10.1016/j.ceca.2012.02.007.
- Diogo, C.V.; Yambire, K.F.; Fernández Mosquera, L.; Branco, F.T.; Raimundo, N. Mitochondrial adventures at the organelle society. *Biochem. Biophys. Res. Commun.* 2018, 500, 87–93. https://doi.org/10.1016/j.bbrc.2017.04.124.
- 16. Tanaka, M.; Toldi, J.; Vécsei, L. Exploring the Etiological Links behind Neurodegenerative Diseases: Inflammatory Cytokines and Bioactive Kynurenines. *Int. J. Mol. Sci.* 2020, *21*, 2431. https://doi.org/10.3390/ijms21072431.
- 17. Tanaka, M.; Tóth, F.; Polyák, H.; Szabó, Á.; Mándi, Y.; Vécsei, L. Immune Influencers in Action: Metabolites and Enzymes of the Tryptophan-Kynurenine Metabolic Pathway. *Biomedicines* **2021**, *9*, 734. https://doi.org/10.3390/biomedicines9070734.
- Tanaka, M.; Bohár, Z.; Vécsei, L. Are Kynurenines Accomplices or Principal Villains in Dementia? Maintenance of Kynurenine Metabolism. *Molecules* 2020, 25, 564. https://doi.org/10.3390/molecules25030564.
- 19. Tanaka, M.; Vécsei, L. Monitoring the kynurenine system: Concentrations, ratios or what else? *Adv. Clin. Exp. Med.* **2021**, *30*, 775–778. https://doi.org/10.17219/acem/139572.
- Stenton, S.L.; Prokisch, H. Genetics of mitochondrial diseases: Identifying mutations to help diagnosis. *EBioMedicine* 2020, 56, 102784. https://doi.org/10.1016/j.ebiom.2020.102784.

- Koene, S.; Wortmann, S.B.; de Vries, M.C.; Jonckheere, A.I.; Morava, E.; de Groot, I.J.; Smeitink, J.A. Developing outcome measures for pediatric mitochondrial disorders: Which complaints and limitations are most burdensome to patients and their parents? *Mitochondrion* 2013, 13, 15–24. https://doi.org/10.1016/j.mito.2012.11.002.
- 22. Niyazov, D.M.; Kahler, S.G.; Frye, R.E. Primary Mitochondrial Disease and Secondary Mitochondrial Dysfunction: Importance of Distinction for Diagnosis and Treatment. *Mol. Syndromol.* **2016**, *7*, 122–137. https://doi.org/10.1159/000446586.
- Cenini, G.; Lloret, A.; Cascella, R. Oxidative Stress and Mitochondrial Damage in Neurodegenerative Diseases: From Molecular Mechanisms to Targeted Therapies. Oxid. Med. Cell Longev. 2020, 2020, 1270256. https://doi.org/10.1155/2020/1270256.
- Rigotto, G.; Basso, E. Mitochondrial Dysfunctions: A Thread Sewing Together Alzheimer's Disease, Diabetes, and Obesity. Oxid. Med. Cell Longev. 2019, 2019, 7210892. https://doi.org/10.1155/2019/7210892.
- Giorgi, C.; Marchi, S.; Simoes, I.C.M.; Ren, Z.; Morciano, G.; Perrone, M.; Patalas-Krawczyk, P.; Borchard, S.; Jędrak, P.; Pierzynowska, K.; et al. Mitochondria and Reactive Oxygen Species in Aging and Age-Related Diseases. *Int. Rev. Cell Mol. Biol.* 2018, 340, 209–344. https://doi.org/10.1016/bs.ircmb.2018.05.006.
- 26. Wang, Y.; Xu, E.; Musich, P.R.; Lin, F. Mitochondrial dysfunction in neurodegenerative diseases and the potential countermeasure. *CNS Neurosci. Ther.* **2019**, *25*, 816–824. https://doi.org/10.1111/cns.13116.
- Daniels, T.E.; Olsen, E.M.; Tyrka, A.R. Stress and Psychiatric Disorders: The Role of Mitochondria. *Annu. Rev. Clin. Psychol.* 2020, 16, 165–186. https://doi.org/10.1146/annurev-clinpsy-082719-104030.
- Raichle, M.E.; Gusnard, D.A. Appraising the brain's energy budget. Proc. Natl. Acad. Sci. USA 2002, 99, 10237–10239. https://doi.org/10.1073/pnas.172399499.
- 29. Kay, J.; Weitzman, P.D. Krebs' Citric Acid Cycle: Half a Century and Still Turning; Biochemical Society: London, UK, 1987; p. 25.
- Zhang, H.; Ryu, D.; Wu, Y.; Gariani, K.; Wang, X.; Luan, P.; D'Amico, D.; Ropelle, E.R.; Lutolf, M.P.; Aebersold, R.; et al. NAD+ repletion improves mitochondrial and stem cell function and enhances life span in mice. *Science* 2016, 352, 1436–1443. https://doi.org/10.1126/science.aaf2693.
- 31. Stryer, L. Fatty acid metabolism. In Biochemistry, 4th ed.; W.H. Freeman and Company: New York, NY, USA, 1995; pp. 603–628.
- 32. Ahmad, M.; Wolberg, A.; Kahwaji, C.I. Biochemistry, Electron Transport Chain.. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2022. Available online: https://www.ncbi.nlm.nih.gov/books/NBK526105/ (accessed on 6 July 2022).
- 33. Klingenberg, M. The ADP and ATP transport in mitochondria and its carrier. *Biochim. Biophys. Acta* 2008, 1778, 1978–2021. https://doi.org/10.1016/j.bbamem.2008.04.011.
- 34. Finkel, T.; Menazza, S.; Holmström, K.M.; Parks, R.J.; Liu, J.; Sun, J.; Liu, J.; Pan, X.; Murphy, E. The ins and outs of mitochondrial calcium. *Circ. Res.* 2015, *116*, 1810–1819. https://doi.org/10.1161/CIRCRESAHA.116.305484.
- 35. Xu, Z.; Zhang, D.; He, X.; Huang, Y.; Shao, H. Transport of Calcium Ions into Mitochondria. *Curr. Genom.* 2016, 17, 215–219. https://doi.org/10.2174/1389202917666160202215748.
- 36. Palego, L.; Betti, L.; Rossi, A.; Giannaccini, G. Tryptophan Biochemistry: Structural, Nutritional, Metabolic, and Medical Aspects in Humans. J. Amino Acids 2016, 2016, 8952520. https://doi.org/10.1155/2016/8952520.
- 37. Liu, D.; Gao, Y.; Liu, J.; Huang, Y.; Yin, J.; Feng, Y.; Shi, L.; Meloni, B.P.; Zhang, C.; Zheng, M.; et al. Intercellular mitochondrial transfer as a means of tissue revitalization. *Signal. Transduct. Target Ther.* **2021**, *6*, 65. https://doi.org/10.1038/s41392-020-00440-z.
- Krebs, H.A. The citric acid cycle and the Szent-Györgyi cycle in pigeon breast muscle. *Biochem. J.* 1940, 34, 775–779. https://doi.org/10.1042/bj0340775.
- 39. Lipmann, F.; Kaplan, N.O. A common factor in the enzymatic acetylation of sulfanilamide and of choline. *J. Biol. Chem.* **1946**, 162, 743–744.
- 40. Lipmann, F. Development of the Acetylation Problem: A personal Account. Available online: https://www.no-belprize.org/prizes/medicine/1953/lipmann/lecture/ (accessed on 6 July 2022).
- 41. Romero-Garcia, S.; Prado-Garcia, H. Mitochondrial calcium: Transport and modulation of cellular processes in homeostasis and cancer (Review). *Int. J. Oncol.* 2019, *54*, 1155–1167. https://doi.org/10.3892/ijo.2019.4696.
- 42. Xia, M.; Zhang, Y.; Jin, K.; Lu, Z.; Zeng, Z.; Xiong, W. Communication between mitochondria and other organelles: a brandnew perspective on mitochondria in cancer. *Cell Biosci.* **2019**, *9*, 27. https://doi.org/10.1186/s13578-019-0289-8.
- 43. Soledad, R.B.; Charles, S.; Samarjit, D. The secret messages between mitochondria and nucleus in muscle cell biology. *Arch. Biochem. Biophys.* **2019**, *666*, 52–62. https://doi.org/10.1016/j.abb.2019.03.019.
- Yeo, A.J.; Chong, K.L.; Gatei, M.; Zou, D.; Stewart, R.; Withey, S.; Wolvetang, E.; Parton, R.G.; Brown, A.D.; Kastan, M.B.; et al. Impaired endoplasmic reticulum-mitochondrial signaling in ataxia-telangiectasia. *iScience* 2020, 24, 101972. https://doi.org/10.1016/j.isci.2020.101972.
- 45. Todkar, K.; Ilamathi, H.S.; Germain, M. Mitochondria and Lysosomes: Discovering Bonds. *Front. Cell Dev. Biol.* 2017, 7, 106. https://doi.org/10.3389/fcell.2017.00106.
- 46. Demarquoy, J.; Le Borgne, F. Crosstalk between mitochondria and peroxisomes. *World J. Biol. Chem.* 2015, *6*, 301–319. https://doi.org/10.4331/wjbc.v6.i4.301.
- 47. Mishra, P.; Chan, D.C. Mitochondrial dynamics and inheritance during cell division, development and disease. *Nat. Rev. Mol. Cell Biol.* **2014**, *15*, 634–646. https://doi.org/10.1038/nrm3877.
- Noguchi, M.; Kasahara, A. Mitochondrial dynamics coordinate cell differentiation. *Biochem. Biophys. Res. Commun.* 2018, 27, 59– 64. https://doi.org/10.1016/j.bbrc.2017.06.094.
- 49. Hales, K.G. Mitochondrial Fusion and Division. Nat. Educ. 2010, 3, 12.

- 50. Thomas, R.L.; Gustafsson, A.B. Mitochondrial autophagy An essential quality control mechanism for myocardial homeostasis. *Circ. J.* 2013, 77, 2449–2454. https://doi.org/10.1253/circj.cj-13-0835.
- Refolo, G.; Vescovo, T.; Piacentini, M.; Fimia, G.M.; Ciccosanti, F. Mitochondrial Interactome: A Focus on Antiviral Signaling Pathways. Front. Cell Dev. Biol. 2020, 8, 8. https://doi.org/10.3389/fcell.2020.00008.
- 52. Vringer, E.; Tait, S.W.G. Mitochondria and Inflammation: Cell Death Heats Up. Front. Cell Dev. Biol. 2019, 7, 100. https://doi.org/10.3389/fcell.2019.00100.
- 53. Barik, S. The Uniqueness of Tryptophan in Biology: Properties, Metabolism, Interactions and Localization in Proteins. *Int. J. Mol. Sci.* **2020**, *21*, 8776. https://doi.org/10.3390/ijms21228776.
- 54. van Donkelaar, E.L.; Blokland, A.; Ferrington, L.; Kelly, P.A.; Steinbusch, H.W.; Prickaerts, J. Mechanism of acute tryptophan depletion: Is it only serotonin? *Mol. Psychiatry* **2011**, *16*, 695–713. https://doi.org/10.1038/mp.2011.9.
- 55. Blankfield, A. A Brief Historic Overview of Clinical Disorders Associated with Tryptophan: The Relevance to Chronic Fatigue Syndrome (CFS) and Fibromyalgia (FM). *Int. J. Tryptophan. Res.* **2012**, *5*, 27–32. https://doi.org/10.4137/IJTR.S10085.
- 56. Marx, W.; McGuinness, A.J.; Rocks, T.; Ruusunen, A.; Cleminson, J.; Walker, A.J.; Gomes-da-Costa, S.; Lane, M.; Sanches, M.; Diaz, A.P.; et al. The kynurenine pathway in major depressive disorder, bipolar disorder, and schizophrenia: A meta-analysis of 101 studies. *Mol. Psychiatry* 2021, 26, 4158–4178. https://doi.org/10.1038/s41380-020-00951-9.
- 57. Török, N.; Tanaka, M.; Vécsei, L. Searching for Peripheral Biomarkers in Neurodegenerative Diseases: The Tryptophan-Kynurenine Metabolic Pathway. *Int. J. Mol. Sci.* **2020**, *21*, 9338. https://doi.org/10.3390/ijms21249338.
- Jamshed, L.; Debnath, A.; Jamshed, S.; Wish, J.V.; Raine, J.C.; Tomy, G.T.; Thomas, P.J.; Holloway, A.C. An Emerging Cross-Species Marker for Organismal Health: Tryptophan-Kynurenine Pathway. *Int. J. Mol. Sci.* 2022, 23, 6300. https://doi.org/10.3390/ijms23116300.
- 59. UniProtKB-P48775 (T23O\_HUMAN). Available online: https://www.uniprot.org/uniprot/P48775 (accessed on 6 July 2022).
- 60. Comings, D.E. Clinical and molecular genetics of ADHD and Tourette syndrome. Two related polygenic disorders. *Ann. N. Y. Acad. Sci.* **2001**, *931*, 50–83. https://doi.org/10.1111/j.1749-6632.2001.tb05773.x.
- 61. Nabi, R.; Serajee, F.J.; Chugani, D.C.; Zhong, H.; Huq, A.H. Association of tryptophan 2,3 dioxygenase gene polymorphism with autism. *Am. J. Med. Genet. B. Neuropsychiatr. Genet.* 2004, 125B, 63–68. https://doi.org/10.1002/ajmg.b.20147.
- Kanai, M.; Funakoshi, H.; Takahashi, H.; Hayakawa, T.; Mizuno, S.; Matsumoto, K.; Nakamura, T. Tryptophan 2,3-dioxygenase is a key modulator of physiological neurogenesis and anxiety-related behavior in mice. *Mol. Brain* 2009, 27, 8. https://doi.org/10.1186/1756-6606-2-8.
- 63. Too, L.K.; Li, K.M.; Suarna, C.; Maghzal, G.J.; Stocker, R.; McGregor, I.S.; Hunt, N.H. Behavioral and cognitive data in mice with different tryptophan-metabolizing enzymes knocked out. *Data Brief* **2016**, *6*, 275–287. https://doi.org/10.1016/j.dib.2016.08.071.
- Too, L.K.; Li, K.M.; Suarna, C.; Maghzal, G.J.; Stocker, R.; McGregor, I.S.; Hunt, N.H. Deletion of TDO2, IDO-1 and IDO-2 dif-64. ferentially affects mouse behavior and cognitive function. Behav. Brain Res. 2016, 1, 102-117. https://doi.org/10.1016/j.bbr.2016.06.018.
- 65. Hattori, S.; Takao, K.; Funakoshi, H.; Miyakawa, T. Comprehensive behavioral analysis of tryptophan 2,3-dioxygenase (Tdo2) knockout mice. *Neuropsychopharmacol. Rep.* **2018**, *38*, 52–60. https://doi.org/10.1002/npr2.12006.
- 66. UniProtKB-P14902 (I23O1\_HUMAN). Available online: https://www.uniprot.org/uniprot/P14902 (accessed on on 6 July 2022).
- UniProtKB-Q6ZQW0 (I23O2\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q6ZQW0 (accessed on on 6 July 2022).
- Kim, H.; Chen, L.; Lim, G.; Sung, B.; Wang, S.; McCabe, M.F.; Rusanescu, G.; Yang, L.; Tian, Y.; Mao, J. Brain indoleamine 2,3dioxygenase contributes to the comorbidity of pain and depression. *J. Clin. Investig.* 2012, 122, 2940–2954. https://doi.org/10.1172/JCI61884.
- O'Connor, J.C.; Lawson, M.A.; André, C.; Briley, E.M.; Szegedi, S.S.; Lestage, J.; Castanon, N.; Herkenham, M.; Dantzer, R.; Kelley, K.W. Induction of IDO by bacille Calmette-Guérin is responsible for development of murine depressive-like behavior. J. Immunol. 2009, 182, 3202–3212. https://doi.org/10.4049/jimmunol.0802722.
- 70. Hirata, N.; Hattori, S.; Shoji, H.; Funakoshi, H.; Miyakawa, T. Comprehensive behavioral analysis of indoleamine 2,3-dioxygenase knockout mice. *Neuropsychopharmacol. Rep.* **2018**, *38*, 133–144. https://doi.org/10.1002/npr2.12019.
- 71. Smith, A.K.; Simon, J.S.; Gustafson, E.L.; Noviello, S.; Cubells, J.F.; Epstein, M.P.; Devlin, D.J.; Qiu, P.; Albrecht, J.K.; Brass, C.A.; et al. Association of a polymorphism in the indoleamine-2,3-dioxygenase gene and interferon-α-induced depression in patients with chronic hepatitis C. *Mol. Psychiatry* 2012, *17*, 781–789. https://doi.org/10.1038/mp.2011.67.
- Cutler, J.A.; Rush, A.J.; McMahon, F.J.; Laje, G. Common genetic variation in the indoleamine-2,3-dioxygenase genes and antidepressant treatment outcome in major depressive disorder. *J. Psychopharmacol.* 2012, 26, 360–367. https://doi.org/10.1177/0269881111434622.
- 73. UniProtKB-Q63HM1 (KFA\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q63HM1 (accessed on on 6 July 2022).
- 74. Ramírez Ortega, D.; Ugalde Muñiz, P.E.; Blanco Ayala, T.; Vázquez Cervantes, G.I.; Lugo Huitrón, R.; Pineda, B.; González Esquivel, D.F.; Pérez de la Cruz, G.; Pedraza Chaverrí, J.; Sánchez Chapul, L.; et al. On the Antioxidant Properties of L-Kynurenine: An Efficient ROS Scavenger and Enhancer of Rat Brain Antioxidant Defense. *Antioxidants* 2022, *11*, 31. https://doi.org/10.3390/antiox11010031.
- 75. UniProtKB-O15229 (KMO\_HUMAN). Available online: https://www.uniprot.org/uniprot/O15229 (accessed on on 6 July 2022).

- Mor, A.; Tankiewicz-Kwedlo, A.; Krupa, A.; Pawlak, D. Role of Kynurenine Pathway in Oxidative Stress during Neurodegenerative Disorders. *Cells* 2021, 10, 1603. https://doi.org/10.3390/cells10071603.
- 77. Giorgini, F.; Huang, S.Y.; Sathyasaikumar, K.V.; Notarangelo, F.M.; Thomas, M.A.; Tararina, M.; Wu, H.Q.; Schwarcz, R.; Muchowski, P.J. Targeted deletion of kynurenine 3-monooxygenase in mice: A new tool for studying kynurenine pathway metabolism in periphery and brain. J. Biol. Chem. 2013, 288, 36554–36566. https://doi.org/10.1074/jbc.M113.503813.
- Erhardt, S.; Pocivavsek, A.; Repici, M.; Liu, X.C.; Imbeault, S.; Maddison, D.C.; Thomas, M.A.R.; Smalley, J.L.; Larsson, M.K.; Muchowski, P.J.; et al. Adaptive and Behavioral Changes in Kynurenine 3-Monooxygenase Knockout Mice: Relevance to Psychotic Disorders. *Biol. Psychiatry* 2017, *82*, 756–765. https://doi.org/10.1016/j.biopsych.2016.12.011.
- Holtze, M.; Saetre, P.; Engberg, G.; Schwieler, L.; Werge, T.; Andreassen, O.A.; Hall, H.; Terenius, L.; Agartz, I.; Jönsson, E.G.; et al. Kynurenine 3-monooxygenase polymorphisms: Relevance for kynurenic acid synthesis in patients with schizophrenia and healthy controls. *J. Psychiatry Neurosci.* 2012, *37*, 53–57. https://doi.org/10.1503/jpn.100175.
  - Wonodi, I.; McMahon, R.P.; Krishna, N.; Mitchell, B.D.; Liu, J.; Glassman, M.; Hong, L.E.; Gold, J.M. Influence of kynurenine 3-monooxygenase (KMO) gene polymorphism on cognitive function in schizophrenia. *Schizophr. Res.* 2014, 160, 80–87. https://doi.org/10.1016/j.schres.2014.10.026.
- Han, Q.; Cai, T.; Tagle, D.A.; Robinson, H.; Li, J. Substrate specificity and structure of human aminoadipate aminotransferase/kynurenine aminotransferase II. *Biosci. Rep.* 2008, 28, 205–215. https://doi.org/10.1042/BSR20080085.
- UniProtKB-Q16773 (KAT1\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q16773 (accessed on on 6 July 2022).
- UniProtKB-Q4W5N8 (Q4W5N8\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q4W5N8 (accessed on on 6 July 2022).
- 84. UniProtKB-Q6YP21 (KAT3\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q6YP21 (accessed on on 6 July 2022).
- 85. UniProtKB-P00505 (AATM\_HUMAN). Available online: https://www.uniprot.org/uniprot/P00505 (accessed on on 6 July 2022).
- Okada, K.; Angkawidjaja, C.; Koga, Y.; Takano, K.; Kanaya, S. Characteristic features of kynurenine aminotransferase allosterically regulated by (alpha)-ketoglutarate in cooperation with kynurenine. *PLoS ONE* 2012, 7, e40307. https://doi.org/10.1371/journal.pone.0040307.
- 87. Herédi, J.; Berkó, A.M.; Jankovics, F.; Iwamori, T.; Iwamori, N.; Ono, E.; Horváth, S.; Kis, Z.; Toldi, J.; Vécsei, L.; et al. Astrocytic and neuronal localization of kynurenine aminotransferase-2 in the adult mouse brain. *Brain Struct. Funct.* 2017, 222, 1663–1672.
- 88. Prescott, C.; Weeks, A.M.; Staley, K.J.; Partin, K.M. Kynurenic acid has a dual action on AMPA receptor responses. *Neurosci. Lett.* **2006**, 402, 108–112. https://doi.org/10.1016/j.neulet.2006.03.051.
- 89. Rózsa, E.; Robotka, H.; Vécsei, L.; Toldi, J. The Janus-face kynurenic acid. J. Neural. Transm. 2008, 115, 1087–1091. https://doi.org/10.1007/s00702-008-0052-5.
- 90. Stone, T.W. Does kynurenic acid act on nicotinic receptors? An assessment of the evidence. J. Neurochem. 2020, 152, 627–649. https://doi.org/10.1111/jnc.14907.
- Copeland, C.S.; Neale, S.A.; Salt, T.E. Actions of Xanthurenic acid, a putative endogenous Group II metabotropic glutamate receptor agonist, on sensory transmission in the thalamus. *Neuropharmacology* 2013, 66, 133–142. https://doi.org/10.1016/j.neuropharm.2012.03.009.
- 92. Bartlett, R.D.; Esslinger, C.S.; Thompson, C.M.; Bridges, R.J. Substituted quinolines as inhibitors of L-glutamate transport into synaptic vesicles. *Neuropharmacology* **1998**, *37*, 839–846. https://doi.org/10.1016/s0028-390800080-x.
- Fazio, F.; Lionetto, L.; Curto, M.; Iacovelli, L.; Cavallari, M.; Zappulla, C.; Ulivieri, M.; Napoletano, F.; Capi, M.; Corigliano, V.; et al. Xanthurenic Acid Activates mGlu2/3 Metabotropic Glutamate Receptors and is a Potential Trait Marker for Schizophrenia. *Sci. Rep.* 2015, *8*, 17799. https://doi.org/10.1038/srep17799.
- Yu, P.; Di Prospero, N.A.; Sapko, M.T.; Cai, T.; Chen, A.; Melendez-Ferro, M.; Du, F.; Whetsell, W.O., Jr.; Guidetti, P.; Schwarcz, R.; et al. Biochemical and phenotypic abnormalities in kynurenine aminotransferase II-deficient mice. *Mol. Cell. Biol.* 2004, 24, 6919–6930. https://doi.org/10.1128/MCB.24.16.6919-6930.2004.
- 95. Martos, D.; Tuka, B.; Tanaka, M.; Vécsei, L.; Telegdy, G. Memory Enhancement with Kynurenic Acid and Its Mechanisms in Neurotransmission. *Biomedicines* 2022, *10*, 849. https://doi.org/10.3390/biomedicines10040849.
- 96. Tanaka, M.; Bohár, Z.; Martos, D.; Telegdy, G.; Vécsei, L. Antidepressant-like effects of kynurenic acid in a modified forced swim test. *Pharmacol. Rep.* 2020, 72, 449–455. https://doi.org/10.1007/s43440-020-00067-5.
- Tanaka, M.; Szabó, Á.; Lőrinczi, B.; Szatmári, I.; Fülöp, F.; Vécsei, L. Antidepressant-like Effects of Kynurenic Acid Analogues. In Proceedings of the 1st International Electronic Conference on Biomedicine, Online, 1–26 June 2021;. https://doi.org/10.3390/ECB2021-10301.
- Potter, M.C.; Elmer, G.I.; Bergeron, R.; Albuquerque, E.X.; Guidetti, P.; Wu, H.Q.; Schwarcz, R. Reduction of endogenous kynurenic acid formation enhances extracellular glutamate, hippocampal plasticity, and cognitive behavior. *Neuropsychopharmacology* 2010, *35*, 1734–1742. https://doi.org/10.1038/npp.2010.39.
- 99. Vécsei, L. (University of Szeged, Szeged, Hungary); 2020 Personal communication.
- 100. UniProtKB-Q16719 (KYNU\_HUMAN). Available online: https://www.uniprot.org/uniprot/Q16719 (accessed on on 6 July 2022).

- Issa, F.; Kirch, D.G.; Gerhardt, G.A.; Bartko, J.J.; Suddath, R.L.; Freedman, R.; Wyatt, R.J. A Multidimensional Approach to Analysis of Cerebrospinal Fluid Biogenic Amines in Schizophrenia: II. Correlations with Psychopathology. *Psychiatry Res.* 1994, 52, 251–258. https://doi.org/10.1016/0165-178190070-1.
- Badawy, A.A. Kynurenine Pathway of Tryptophan Metabolism: Regulatory and Functional Aspects. Int. J. Tryptophan Res. 2017, 15, 1178646917691938. https://doi.org/10.1177/1178646917691938.
- Bala, S.; Kamboj, S.; Saini, V.; Prasad, D.N. Anti-inflammatory, analgesic evaluation and molecular docking studies of N-phenyl anthranilic acid-based 1,3,4-oxadiazole analogues. J. Chem. 2013, 2013, 412053. https://doi.org/10.1155/2013/412053.
- 104. Shi, H.; Enriquez, A.; Rapadas, M.; Martin, E.M.M.A.; Wang, R.; Moreau, J.; Lim, C.K.; Szot, J.O.; Ip, E.; Hughes, J.N.; et al. NAD Deficiency, Congenital Malformations, and Niacin Supplementation. N. Engl. J. Med. 2017, 10, 377, 544-552. https://doi.org/10.1056/NEJMoa1616361.
- 105. Zhang, Y.; Shen, J.; He, X.; Zhang, K.; Wu, S.; Xiao, B.; Zhou, X.; Phillips, R.S.; Gao, P.; Jeunemaitre, X.; et al. A rare variant at the KYNU gene is associated with kynureninase activity and essential hypertension in the Han Chinese population. *Circ. Cardiovasc. Genet.* 2011, 4, 687–694. https://doi.org/10.1161/CIRCGENETICS.110.959064.
- 106. Farooq, R.K.; Asghar, K.; Kanwal, S.; Zulqernain, A. Role of inflammatory cytokines in depression: Focus on interleukin-1β. *Biomed. Rep.* 2017, *6*, 15–20. https://doi.org/10.3892/br.2016.807.
- 107. Wang, Y.; Qu, P.; Sun, Y.; Li, Z.; Liu, L.; Yang, L. Association between increased inflammatory cytokine expression in the lateral habenular nucleus and depressive-like behavior induced by unpredictable chronic stress in rats. *Exp. Neurol.* 2022, 349, 113964. <u>https://doi.org/10.1016/j.expneurol.2021.113964</u>.
- 108. Hepsomali, P.; Coxon, C. Inflammation and diet: Focus on mental and cognitive health. *Adv. Clin. Exp. Med.* 2022 [published online ahead of print, 2022 Aug 11]. https://doi.org/10.17219/acem/152350.
- 109. Liu, Y.; Zhou, L.J.; Wang, J.; Li, D.; Ren, W.J.; Peng, J.; Wei, X.; Xu, T.; Xin, W.J.; Pang, R.P.; et al. TNF-α Differentially Regulates Synaptic Plasticity in the Hippocampus and Spinal Cord by Microglia-Dependent Mechanisms after Peripheral Nerve Injury. J. Neurosci. 2017, 37, 871–881. https://doi.org/10.1523/JNEUROSCI.2235-16.2016.
- Boes, A.D.; Kelly, M.S.; Trapp, N.T.; Stern, A.P.; Press, D.Z.; Pascual-Leone, A. Noninvasive Brain Stimulation: Challenges and Opportunities for a New Clinical Specialty. *J. Neuropsychiatry Clin. Neurosci.* 2018, 30, 173–179. https://doi.org/10.1176/appi.neuropsych.17110262.
- 111. Battaglia, S.; Thayer, J.F. Functional interplay between central and autonomic nervous systems in human fear conditioning. *Trends Neurosci.* 2022, 45, 504–506. https://doi.org/10.1016/j.tins.2022.04.003.
- Battaglia, S.; Orsolini, S.; Borgomaneri, S.; Barbieri, R.; Diciotti, S.; di Pellegrino, G. Characterizing cardiac autonomic dynamics of fear learning in humans. *Psychophysiology* 2022, e14122. https://doi.org/10.1111/psyp.14122.
- 113. Borgomaneri, S.; Battaglia, S.; Garofalo, S.; Tortora, F.; Avenanti, A.; di Pellegrino, G. State-Dependent TMS over Prefrontal Cortex Disrupts Fear-Memory Reconsolidation and Prevents the Return of Fear. *Curr. Biol.* **2020**, *30*, 3672–3679.e4. https://doi.org/10.1016/j.cub.2020.06.091.
- Borgomaneri, S.; Battaglia, S.; Sciamanna, G.; Tortora, F.; Laricchiuta, D. Memories are not written in stone: Re-writing fear memories by means of non-invasive brain stimulation and optogenetic manipulations. *Neurosci. Biobehav. Rev.* 2021, 127, 334– 352. https://doi.org/10.1016/j.neubiorev.2021.04.036.
- 115. Borgomaneri, S.; Battaglia, S.; Avenanti, A.; Pellegrino, G.D. Don't Hurt Me No More: State-dependent Transcranial Magnetic Stimulation for the treatment of specific phobia. J. Affect Disord. 2021, 286, 78–79. https://doi.org/10.1016/j.jad.2021.02.076.
- 116. Gonzalez-Escamilla, G.; Dörfel, D.; Becke, M.; Trefz, J.; Bonanno, G.A.; Groppa, S. Associating Flexible Regulation of Emotional Expression with Psychopathological Symptoms. *Front. Behav. Neurosci.* 2022, 16, 924305. https://doi.org/10.3389/fnbeh.2022.924305.
- 117. UniProtKB-P46952 (3HAO\_HUMAN). Available online: https://www.uniprot.org/uniprot/P46952 (accessed on 6 July 2022).
- 118. Lugo-Huitrón, R.; Ugalde Muñiz, P.; Pineda, B.; Pedraza-Chaverrí, J.; Ríos, C.; Pérez-de la Cruz, V. Quinolinic acid: An endogenous neurotoxin with multiple targets. *Oxid. Med. Cell. Longev.* **2013**, 2013, 104024. https://doi.org/10.1155/2013/104024.
- Majláth, Z.; Toldi, J.; Vécsei, L. The potential role of kynurenines in Alzheimer's disease: Pathomechanism and therapeutic possibilities by influencing the glutamate receptors. *J. Neural. Transm.* 2014, 121, 881–889. https://doi.org/10.1007/s00702-013-1135-5.
- Zhang, W.; Egashira, N.; Masuda, S. Recent Topics on The Mechanisms of Immunosuppressive Therapy-Related Neurotoxicities. Int. J. Mol. Sci. 2019, 20, 3210. https://doi.org/10.3390/ijms20133210.
- 121. Schuck, P.F.; Tonin, A.; da Costa Ferreira, G.; Rosa, R.B.; Latini, A.; Balestro, F.; Perry, M.L.; Wannmacher, C.M.; de Souza Wyse, A.T.; Wajner, M. In vitro effect of quinolinic acid on energy metabolism in brain of young rats. *Neurosci. Res.* 2007, *57*, 277–288. https://doi.org/10.1016/j.neures.2006.10.013.
- 122. Lim, C.K.; Essa, M.M.; de Paula Martins, R.; Lovejoy, D.B.; Bilgin, A.A.; Waly, M.I.; Al-Farsi, Y.M.; Al-Sharbati, M.; Al-Shaffae, M.A.; Guillemin, G.J. Altered kynurenine pathway metabolism in autism: Implication for immune-induced glutamatergic activity. *Autism Res.* 2016, 9, 621–631. https://doi.org/10.1002/aur.1565.
- 123. Ulivieri, M.; Wierońska, J.M.; Lionetto, L.; Martinello, K.; Cieslik, P.; Chocyk, A.; Curto, M.; Di Menna, L.; Iacovelli, L.; Traficante, A.; et al. The Trace Kynurenine, Cinnabarinic Acid, Displays Potent Antipsychotic-Like Activity in Mice and Its Levels Are Reduced in the Prefrontal Cortex of Individuals Affected by Schizophrenia. *Schizophr. Bull.* 2020, 46, 1471–1481. https://doi.org/10.1093/schbul/sbaa074.

- 124. Lesiewska, N.; Borkowska, A.; Junik, R.; Kamińska, A.; Jaracz, K.; Bieliński, M. Consequences of diabetes and pre-diabetes and the role of biochemical parameters of carbohydrate metabolism for the functioning of the prefrontal cortex in obese patients. *Front. Biosci.* 2022, 27, 76. https://doi.org/10.31083/j.fbl2703076.
- 125. Borgomaneri, S.; Serio, G.; Battaglia, S. Please, don't do it! Fifteen years of progress of non-invasive brain stimulation in action inhibition. *Cortex* **2020**, *132*, 404–422. https://doi.org/10.1016/j.cortex.2020.09.002.
- 126. Battaglia, S.; Cardellicchio, P.; Di Fazio, C.; Nazzi, C.; Fracasso, A.; Borgomaneri, S. The Influence of Vicarious Fear-Learning in "Infecting" Reactive Action Inhibition. *Front. Behav. Neurosci.* **2022**, *16*, 946263. https://doi.org/10.3389/fnbeh.2022.94626.
- 127. Sellitto, M.; Terenzi, D.; Starita, F.; di Pellegrino, G.; Battaglia, S. The Cost of Imagined Actions in a Reward-Valuation Task. *Brain Sci.* **2022**, *12*, 582. https://doi.org/10.3390/brainsci12050582.
- 128. Battaglia, S.; Harrison, B.J.; Fullana, M.A. Does the human ventromedial prefrontal cortex support fear learning, fear extinction or both? A commentary on subregional contributions. *Mol. Psychiatry* **2022**, *27*, 784–786. https://doi.org/10.1038/s41380-021-01326-4.
- 129. Garofalo, S.; Timmermann, C.; Battaglia, S.; Maier, M.E.; di Pellegrino, G. Mediofrontal Negativity Signals Unexpected Timing of Salient Outcomes. J. Cogn. Neurosci. 2017, 29, 718–727. https://doi.org/10.1162/jocn\_a\_01074.
- 130. Gamboa, O.L.; Chuan-Peng, H.; Salas, C.E.; Yuen, K.S.L. Obliviate! Reviewing Neural Fundamentals of Intentional Forgetting from a Meta-Analytic Perspective. *Biomedicines* **2022**, *10*, 1555. https://doi.org/10.3390/biomedicines10071555.
- 131. Battaglia, S.; Garofalo, S.; di Pellegrino, G.; Starita, F. Revaluing the Role of vmPFC in the Acquisition of Pavlovian Threat Conditioning in Humans. J. Neurosci. 2020, 40, 8491–8500. https://doi.org/10.1523/JNEUROSCI.0304-20.202.
- Battaglia, S.; Serio, G.; Scarpazza, C.; D'Ausilio, A.; Borgomaneri, S. Frozen in (e)motion: How reactive motor inhibition is influenced by the emotional content of stimuli in healthy and psychiatric populations. *Behav. Res. Ther.* 2021, 146, 103963. https://doi.org/10.1016/j.brat.2021.103963.
- 133. Milad, M.R.; Pitman, R.K.; Ellis, C.B.; Gold, A.L.; Shin, L.M.; Lasko, N.B.; Zeidan, M.A.; Handwerger, K.; Orr, S.P.; Rauch, S.L. Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biol. Psychiatry* 2009, 66, 1075–1082. https://doi.org/10.1016/j.biopsych.2009.06.026.
- Holt, D.J.; Lebron-Milad, K.; Milad, M.R.; Rauch, S.L.; Pitman, R.K.; Orr, S.P.; Cassidy, B.S.; Walsh, J.P.; Goff, D.C. Extinction memory is impaired in schizophrenia. *Biol. Psychiatry* 2009, 65, 455–463. https://doi.org/10.1016/j.biopsych.2008.09.017.
- 135. Battaglia, S.; Fabius, J.H.; Moravkova, K.; Fracasso, A.; Borgomaneri, S. The Neurobiological Correlates of Gaze Perception in Healthy Individuals and Neurologic Patients. *Biomedicines* **2022**, *10*, 627. https://doi.org/10.3390/biomedicines10030627.
- Flippo, K.H.; Strack, S. Mitochondrial dynamics in neuronal injury, development and plasticity. J. Cell Sci. 2017, 130, 671–681. https://doi.org/10.1242/jcs.171017.
- Sheng, Z.H. The Interplay of Axonal Energy Homeostasis and Mitochondrial Trafficking and Anchoring. *Trends Cell Biol.* 2017, 27, 403–416. https://doi.org/10.1016/j.tcb.2017.01.005.
- Santuy, A.; Turégano-López, M.; Rodríguez, J.R.; Alonso-Nanclares, L.; DeFelipe, J.; Merchán-Pérez, A. A Quantitative Study on the Distribution of Mitochondria in the Neuropil of the Juvenile Rat Somatosensory Cortex. *Cereb. Cortex* 2018, 28, 3673–3684. https://doi.org/10.1093/cercor/bhy159.
- Lee, A.; Hirabayashi, Y.; Kwon, S.K.; Lewis, T.L., Jr.; Polleux, F. Emerging roles of mitochondria in synaptic transmission and neurodegeneration. *Curr. Opin. Physiol.* 2018, *3*, 82–93. https://doi.org/10.1016/j.cophys.2018.03.009.
- Wang, B.; Huang, M.; Shang, D.; Yan, X.; Zhao, B.; Zhang, X. Mitochondrial Behavior in Axon Degeneration and Regeneration. Front. Aging Neurosci. 2021, 13, 650038. https://doi.org/10.3389/fnagi.2021.650038.
- Carelli, V.; Chan, D.C. Mitochondrial DNA: Impacting central and peripheral nervous systems. *Neuron* 2014, 84, 1126–1142. https://doi.org/10.1016/j.neuron.2014.11.022.
- 142. Liang, C.; Ahmad, K.; Sue, C.M. The broadening spectrum of mitochondrial disease: Shifts in the diagnostic paradigm. *Biochim. Biophys. Acta* 2014, *1840*, 1360–1367. https://doi.org/10.1016/j.bbagen.2013.10.040.
- 143. Frye, R.E. Mitochondrial Dysfunction in Autism Spectrum Disorder: Unique Abnormalities and Targeted Treatments. *Semin. Pediatr. Neurol.* **2020**, *35*, 100829. https://doi.org/10.1016/j.spen.2020.100829.
- 144. Scaglia, F. The role of mitochondrial dysfunction in psychiatric disease. *Dev. Disabil. Res. Rev.* 2010, *16*, 136–143. https://doi.org/10.1002/ddrr.115.
- Khan, N.A.; Govindaraj, P.; Meena, A.K.; Thangaraj, K. Mitochondrial disorders: Challenges in diagnosis & treatment. *Indian J. Med. Res.* 2015, 141, 13–26. https://doi.org/10.4103/0971-5916.154489.
- 146. Falk, M.J.; Sondheimer, N. Mitochondrial genetic diseases. *Curr. Opin. Pediatr.* 2010, 22, 711–716. https://doi.org/10.1097/MOP.0b013e3283402e21.
- 147. Almannai, M.; El-Hattab, A.W.; Ali, M.; Soler-Alfonso, C.; Scaglia, F. Clinical trials in mitochondrial disorders, an update. *Mol. Genet. Metab.* 2020, *131*, 1–13. https://doi.org/10.1016/j.ymgme.2020.10.002.
- McCormick, E.M.; Zolkipli-Cunningham, Z.; Falk, M.J. Mitochondrial disease genetics update: Recent insights into the molecular diagnosis and expanding phenotype of primary mitochondrial disease. *Curr. Opin. Pediatr.* 2018, 30, 714–724. https://doi.org/10.1097/MOP.0000000000686.
- Kanungo, S.; Morton, J.; Neelakantan, M.; Ching, K.; Saeedian, J.; Goldstein, A. Mitochondrial disorders. Ann. Transl. Med. 2018, 6, 475. https://doi.org/10.21037/atm.2018.12.13.
- 150. Montano, V.; Gruosso, F.; Simoncini, C.; Siciliano, G.; Mancuso, M. Clinical features of mtDNA-related syndromes in adulthood. *Arch. Biochem. Biophys.* **2021**, 697, 108689. https://doi.org/10.1016/j.abb.2020.108689.

- 151. Fan, H.C.; Lee, H.F.; Yue, C.T.; Chi, C.S. Clinical Characteristics of Mitochondrial Encephalomyopathy, Lactic Acidosis, and Stroke-Like Episodes. *Life* **2021**, 11, 1111. https://doi.org/10.3390/life1111111.
- 152. Tetsuka, S.; Ogawa, T.; Hashimoto, R.; Kato, H. Clinical features, pathogenesis, and management of stroke-like episodes due to MELAS. *Metab. Brain Dis.* 2021, *36*, 2181–2193. https://doi.org/10.1007/s11011-021-00772-x.
- 153. Schlieben, L.D.; Prokisch, H. The Dimensions of Primary Mitochondrial Disorders. Front. Cell Dev. Biol. 2020, 8, 600079. https://doi.org/10.3389/fcell.2020.600079.
- 154. Gorman, G.S.; Schaefer, A.M.; Ng, Y.; Gomez, N.; Blakely, E.L.; Alston, C.L.; Feeney, C.; Horvath, R.; Yu-Wai-Man, P.; Chinnery, P.F.; et al. Prevalence of nuclear and mitochondrial DNA mutations related to adult mitochondrial disease. *Ann. Neurol.* 2015, 77, 753–759. https://doi.org/10.1002/ana.24362.
- 155. Beekman, M.; Dowling, D.K.; Aanen, D.K. The costs of being male: Are there sex-specific effects of uniparental mitochondrial inheritance? *Philos. Trans. R. Soc. Lond. Bol. Sci.* 2014, *369*, 20130440. https://doi.org/10.1098/rstb.2013.0440.
- 156. Luo, S.; Valencia, C.A.; Zhang, J.; Lee, N.C.; Slone, J.; Gui, B.; Wang, X.; Li, Z.; Dell, S.; Brown, J.; et al. Biparental Inheritance of Mitochondrial DNA in Humans. Proc. Natl. Acad. Sci. USA 2018, 115, 13039–13044. https://doi.org/10.1073/pnas.1810946115.
- 157. Ruhoy, I.S.; Saneto, R.P. The genetics of Leigh syndrome and its implications for clinical practice and risk management. *Appl. Clin. Genet.* **2014**, *7*, 221–234. https://doi.org/10.2147/TACG.S46176.
- 158. Thompson Legault, J.; Strittmatter, L.; Tardif, J.; Sharma, R.; Tremblay-Vaillancourt, V.; Aubut, C.; Boucher, G.; Clish, C.B.; Cyr, D.; Daneault, C.; et al. A Metabolic Signature of Mitochondrial Dysfunction Revealed through a Monogenic Form of Leigh Syndrome. *Cell Rep.* 2015, *13*, 981–989. https://doi.org/10.1016/j.celrep.2015.09.054.
- 159. Agus, A.; Planchais, J.; Sokol, H. Gut Microbiota Regulation of Tryptophan Metabolism in Health and Disease. *Cell Host Microbe* **2018**, 23, 716–724. https://doi.org/10.1016/j.chom.2018.05.003.
- Sundaramurthy, S.; SelvaKumar, A.; Ching, J.; Dharani, V.; Sarangapani, S.; Yu-Wai-Man, P. Leber hereditary optic neuropathynew insights and old challenges. *Graefes. Arch. Clin. Exp. Ophthalmol.* 2021, 259, 2461–2472. https://doi.org/10.1007/s00417-020-04993-1.
- Stenton, S.L.; Sheremet, N.L.; Catarino, C.B.; Andreeva, N.A.; Assouline, Z.; Barboni, P.; Barel, O.; Berutti, R.; Bychkov, I.; Caporali, L.; et al. Impaired complex I repair causes recessive Leber's hereditary optic neuropathy. *J. Clin. Investig.* 2021, 131, e138267. https://doi.org/10.1172/JCI138267.
- Klivenyi, P.; Karg, E.; Rozsa, C.; Horvath, R.; Komoly, S.; Nemeth, I.; Turi, S.; Vécsei, L. alpha-Tocopherol/lipid ratio in blood is decreased in patients with Leber's hereditary optic neuropathy and asymptomatic carriers of the 11,778 mtDNA mutation. J. Neurol. Neurosurg. Psychiatry 2001, 70, 359–362. https://doi.org/10.1136/jnnp.70.3.359.
- Khajuria, K.; Khajuria, V.; Sawhney, V. Secondary Mitochondrial Dysfunction. Int. J. Pharm. Pharm. Sci. 2021, 13, 14–19. https://doi.org/10.22159/ijpps.2021v13i3.40335.
- 164. Barahona, A.J.; Bursac, Z.; Veledar, E.; Lucchini, R.; Tieu, K.; Richardson, J.R. Relationship of Blood and Urinary Manganese Levels with Cognitive Function in Elderly Individuals in the United States by Race/Ethnicity, NHANES 2011–2014. *Toxics* 2022, 10, 191. https://doi.org/10.3390/toxics10040191.
- 165. Jopowicz, A.; Wiśniowska, J.; Tarnacka, B. Cognitive and Physical Intervention in Metals' Dysfunction and Neurodegeneration. *Brain Sci.* 2022, *12*, 345. https://doi.org/10.3390/brainsci12030345.
- Chaturvedi, R.K.; Beal, M.F. Mitochondrial diseases of the brain. Free Radic. Biol. Med. 2013, 63, 1–29. https://doi.org/10.1016/j.freeradbiomed.2013.03.018.
- Tanaka, M.; Vécsei, L. Editorial of Special Issue "Crosstalk between Depression, Anxiety, and Dementia: Comorbidity in Behavioral Neurology and Neuropsychiatry". *Biomedicines* 2021, 9, 517. https://doi.org/10.3390/biomedicines9050517.
- Tanaka, M.; Vécsei, L. Editorial of Special Issue 'Dissecting Neurological and Neuropsychiatric Diseases: Neurodegeneration and Neuroprotection'. Int. J. Mol. Sci. 2022, 23, 6991. https://doi.org/10.3390/ijms23136991.
- 169. Telegdy, G.; Tanaka, M.; Schally, A.V. Effects of the LHRH antagonist Cetrorelix on the brain function in mice. *Neuropeptides* **2009**, 43, 229–234. https://doi.org/10.1016/j.npep.2009.03.001.
- 170. Telegdy, G.; Tanaka, M.; Schally, A.V. Effects of the growth hormone-releasing hormone (GH-RH) antagonist on brain functions in mice. *Behav. Brain Res.* 2011, 224, 155–1558. https://doi.org/10.1016/j.bbr.2011.05.036.
- Palotai, M.; Telegdy, G.; Tanaka, M.; Bagosi, Z.; Jászberényi, M. Neuropeptide AF induces anxiety-like and antidepressant-like behavior in mice. *Behav. Brain Res.* 2014, 274, 264–269. https://doi.org/10.1016/j.bbr.2014.08.007.
- Tanaka, M.; Kádár, K.; Tóth, G.; Telegdy, G. Antidepressant-like effects of urocortin 3 fragments. *Brain Res. Bull.* 2011, 84, 414–418. https://doi.org/10.1016/j.brainresbull.2011.01.016.
- 173. Mirchandani-Duque, M.; Barbancho, M.A.; López-Salas, A.; Alvarez-Contino, J.E.; García-Casares, N.; Fuxe, K.; Borroto-Escuela, D.O.; Narváez, M. Galanin and Neuropeptide Y Interaction Enhances Proliferation of Granule Precursor Cells and Expression of Neuroprotective Factors in the Rat Hippocampus with Consequent Augmented Spatial Memory. *Biomedicines* 2022, 10, 1297. https://doi.org/10.3390/biomedicines10061297.
- 174. Simon, C.; Soga, T.; Ahemad, N.; Bhuvanendran, S.; Parhar, I. Kisspeptin-10 Rescues Cholinergic Differentiated SHSY-5Y Cells from α-Synuclein-Induced Toxicity In Vitro. *Int. J. Mol. Sci.* 2022, 23, 5193. https://doi.org/10.3390/ijms23095193.
- 175. Naito, M.; Iwakoshi-Ukena, E.; Moriwaki, S.; Narimatsu, Y.; Kato, M.; Furumitsu, M.; Miyamoto, Y.; Esumi, S.; Ukena, K. Immunohistochemical Analysis of Neurotransmitters in Neurosecretory Protein GL-Producing Neurons of the Mouse Hypothalamus. *Biomedicines* 2022, 10, 454. https://doi.org/10.3390/biomedicines10020454.

- 176. Dias, F.L.; Silva, R.M.; Moraes, E.N.; Caramelli, P. Clinical and autonomic profile of patients with Alzheimer's disease and mixed dementia patients. *Rev. Assoc. Médica Bras.* 2013, 59, 435–441. https://doi.org/10.1016/j.ramb.2013.04.004.
- 177. Battaglia, S.; Garofalo, S.; di Pellegrino, G. Context-dependent extinction of threat memories: Influences of healthy aging. *Sci. Rep.* **2018**, *8*, 12592. https://doi.org/10.1038/s41598-018-31000-9.
- 178. Orso, B.; Lorenzini, L.; Arnaldi, D.; Girtler, N.; Brugnolo, A.; Doglione, E.; Mattioli, P.; Biassoni, E.; Massa, F.; Peira, E.; et al. The Role of Hub and Spoke Regions in Theory of Mind in Early Alzheimer's Disease and Frontotemporal Dementia. *Biomedicines* 2022, 10, 544. https://doi.org/10.3390/biomedicines10030544.
- 179. DeTure, M.A.; Dickson, D.W. The neuropathological diagnosis of Alzheimer's disease. *Mol. Neurodegener.* 2019, 14, 32. https://doi.org/10.1186/s13024-019-0333-5.
- 180. Dugger, B.N.; Dickson, D.W. Pathology of Neurodegenerative Diseases. *CSH Perspect. Biol.* 2017, 9, a028035. https://doi.org/10.1101/cshperspect.a028035.
- 181. Dai, M.H.; Zheng, H.; Zeng, L.D.; Zhang, Y. The genes associated with early-onset Alzheimer's disease. *Oncotarget* 2017, 9, 15132–15143. https://doi.org/10.18632/oncotarget.23738.
- 182. Yao, J.; Irwin, R.W.; Zhao, L.; Nilsen, J.; Hamilton, R.T.; Brinton, R.D. Mitochondrial bioenergetic deficit precedes Alzheimer's pathology in female mouse model of Alzheimer's disease. Proc. Natl. Acad. Sci. USA 2009, 106, 14670–14675. https://doi.org/10.1073/pnas.0903563106.
- Rönnbäck, A.; Pavlov, P.F.; Mansory, M.; Gonze, P.; Marlière, N.; Winblad, B.; Graff, C.; Behbahani, H. Mitochondrial dysfunction in a transgenic mouse model expressing human amyloid precursor protein (APP) with the Arctic mutation. *J. Neurochem.* 2016, 136, 497–502. https://doi.org/10.1111/jnc.13410.
- Dixit, S.; Fessel, J.P.; Harrison, F.E. Mitochondrial dysfunction in the APP/PSEN1 mouse model of Alzheimer's disease and a novel protective role for ascorbate. *Free Radic. Biol. Med.* 2017, 112, 515–523. https://doi.org/10.1016/j.freeradbiomed.2017.08.021.
- 185. Mehla, J.; Lacoursiere, S.G.; Lapointe, V.; McNaughton, B.L.; Sutherland, R.J.; McDonald, R.J.; Mohajerani, M.H. Age-dependent behavioral and biochemical characterization of single APP knock-in mouse (APPNL-G-F/NL-G-F) model of Alzheimer's disease. *Neurobiol. Aging* 2019, 75, 25–37. https://doi.org/10.1016/j.neurobiolaging.2018.10.026.
- Guillemin, G.J.; Brew, B.J.; Noonan, C.E.; Takikawa, O.; Cullen, K.M. Indoleamine 2,3 hippocampus dioxygenase and quinolinic acid immunoreactivity in Alzheimer's disease. *Neuropathol. Appl. Neurobiol.* 2005, 31, 395–404. https://doi.org/10.1111/j.1365-2990.2005.00655.x.
- 187. Bonda, D.; Mailankot, M.; Stone, J.G.; Garrett, M.R.; Staniszewska, M.; Castellani, R.J.; Siedlak, S.L.; Zhu, X.; Lee, H.; Perry, G.; et al. Indoleamine 2,3-dioxygenase and 3-hydroxykynurenine modifications are found in the neuropathology of Alzheimer's disease. *Redox Rep.* 2010, 15, 161–168. https://doi.org/10.1179/174329210X12650506623645.
- Gulaj, E.; Pawlak, K.; Bien, B.; Pawlak, D. Kynurenine and its metabolites in Alzheimer's disease patients. Adv. Med. Sci. 2010, 55, 204–211. https://doi.org/10.2478/v10039-010-0023-6.
- Almulla, A.F.; Supasitthumrong, T.; Amrapala, A.; Tunvirachaisakul, C.; Jaleel, A.K.A.; Oxenkrug, G.; Al-Hakeim, H.K.; Maes, M. The Tryptophan Catabolite or Kynurenine Pathway in Alzheimer's Disease: A Systematic Review and Meta-Analysis. J. Alzheimer's Dis. 2022, in press. https://doi.org/10.3233/JAD-220295.
- Schwarcz, M.J.; Guillemin, G.J.; Teipel, S.J.; Buerger, K.; Hampel, H. Increased 3-hydroxykynurenine serum concentrations differentiate Alzheimer's disease patients from controls. *Eur. Arch. Psychiatry Clin. Neurosci.* 2013, 263, 345. https://doi.org/10.1007/s00406-012-0384-x.
- 191. Jacobs, K.; Lim, C.K.; Blennow, K.; Zetterberg, H.; Chatterjee, P.; Martins, R.N.; Brew, B.; Guillemin, G.; Lovejoy, D. Correlation between plasma and CSF concentrations of kynurenine pathway metabolites in Alzheimer's disease and relationship to amyloid-β and tau. *Neurobiol. Aging* 2019, *80*, 11–20. https://doi.org/10.1016/j.neurobiolaging.2019.03.015.
- Holper, L.; Ben-Shachar, D.; Mann, J.J. Multivariate meta-analyses of mitochondrial complex I and IV in major depressive disorder, bipolar disorder, schizophrenia, Alzheimer disease, and Parkinson disease. *Neuropsychopharmacology* 2019, 44, 837–849. https://doi.org/10.1038/s41386-018-0090-0.
- 193. Park, A.; Stacy, M. Non-motor symptoms in Parkinson's disease. J. Neurol. 2009, 256 (Suppl. 3), 293–298. https://doi.org/10.1007/s00415-009-5240-1.
- 194. Spillantini, M.G.; Schmidt, M.L.; Lee, V.M.; Trojanowski, J.Q.; Jakes, R.; Goedert, M. Alpha-synuclein in Lewy bodies. *Nature* **1997**, *388*, 839–840. https://doi.org/10.1038/42166.
- Ellena, G.; Battaglia, S.; Làdavas, E. The spatial effect of fearful faces in the autonomic response. *Exp. Brain. Res.* 2020, 238, 2009–2018. https://doi.org/10.1007/s00221-020-05829-4.
- Borgomaneri, S.; Vitale, F.; Battaglia, S.; Avenanti, A. Early Right Motor Cortex Response to Happy and Fearful Facial Expressions: A TMS Motor-Evoked Potential Study. *Brain Sci.* 2021, *11*, 1203. https://doi.org/10.3390/brainsci11091203.
- Pickrell, A.M.; Youle, R.J. The roles of PINK1, parkin, and mitochondrial fidelity in Parkinson's disease. *Neuron* 2015, 85, 257–273. https://doi.org/10.1016/j.neuron.2014.12.007.
- Moisoi, N.; Fedele, V.; Edwards, J.; Martins, L.M. Loss of PINK1 enhances neurodegeneration in a mouse model of Parkinson's disease triggered by mitochondrial stress. *Neuropharmacology* 2014, 77, 350–357. https://doi.org/10.1016/j.neuropharm.2013.10.009.
- 199. Chia, S.J.; Tan, E.-K.; Chao, Y.-X. Historical Perspective: Models of Parkinson's Disease. Int. J. Mol. Sci. 2020, 21, 2464. https://doi.org/10.3390/ijms21072464.

- 200. Kee, T.R.; Espinoza Gonzalez, P.; Wehinger, J.L.; Bukhari, M.Z.; Ermekbaeva, A.; Sista, A.; Kotsiviras, P.; Liu, T.; Kang, D.E.; Woo, J.A. Mitochondrial CHCHD2: Disease-Associated Mutations, Physiological Functions, and Current Animal Models. *Front. Aging Neurosci.* 2021, 13, 660843. https://doi.org/10.3389/fnagi.2021.66084.
- 201. González-Rodríguez, P.; Zampese, E.; Stout, K.A.; Guzman, J.N.; Ilijic, E.; Yang, B.; Tkatch, T.; Stavarache, M.A.; Wokosin, D.L.; Gao, L.; et al. Disruption of mitochondrial complex I induces progressive parkinsonism. *Nature* 2021, 599, 650–656. https://doi.org/10.1038/s41586-021-04059-0.
- 202. Kim, H.W.; Choi, W.S.; Sorscher, N.; Park, H.J.; Tronche, F.; Palmiter, R.D.; Xia, Z. Genetic reduction of mitochondrial complex I function does not lead to loss of dopamine neurons in vivo. *Neurobiol. Aging* 2015, 36, 2617–2627. https://doi.org/10.1016/j.neurobiolaging.2015.05.008.
- 203. Latchoumycandane, C.; Anantharam, V.; Jin, H.; Kanthasamy, A.; Kanthasamy, A. Dopaminergic neurotoxicant 6-OHDA induces oxidative damage through proteolytic activation of PKCδ in cell culture and animal models of Parkinson's disease. *Toxicol. Appl. Pharmacol.* 2011, 256, 314–323. https://doi.org/10.1016/j.taap.2011.07.021.
- 204. Blesa, J.; Przedborski, S. Parkinson's disease: Animal models and dopaminergic cell vulnerability. *Front. Neuroanat.* 2014, *8*, 155. https://doi.org/10.3389/fnana.2014.00155.
- 205. Hartai, Z.; Klivenyi, P.; Janaky, T.; Penke, B.; Dux, L.; Vécsei, L. Kynurenine metabolism in plasma and in red blood cells in Parkinson's disease. J. Neurol. Sci. 2005, 239, 31–35. https://doi.org/10.1016/j.jns.2005.07.006.
- 206. Lewitt, P.A.; Li, J.; Lu, M.; Beach, T.G.; Adler, C.H.; Guo, L. Arizona Parkinson's Disease Consortium. 3-hydroxykynurenine and other Parkinson's disease biomarkers discovered by metabolomic analysis. *Mov. Disord.* 2013, 28, 1653–1660. https://doi.org/10.1002/mds.25555.
- 207. Török, N.; Maszlag-Török, R.; Molnár, K.; Szolnoki, Z.; Somogyvári, F.; Boda, K.; Tanaka, M.; Klivényi, P.; Vécsei, L. Single Nucleotide Polymorphisms of Indoleamine 2,3-Dioxygenase 1 Influenced the Age Onset of Parkinson's Disease. *Preprints* 2020, 2020090470. https://doi.org/10.20944/preprints202010.0172.v2.
- 208. Boeschoten, R.E.; Braamse, A.M.J.; Beekman, A.T.F.; Cuijpers, P.; van Oppen, P.; Dekker, J.; Uitdehaag, B.M.J. Prevalence of Depression and Anxiety in Multiple Sclerosis: A Systematic Review and Meta-Analysis. J. Neurol. Sci. 2017, 372, 331–341. https://doi.org/10.1016/j.jns.2016.11.067.
- Huang, W.J.; Chen, W.W.; Zhang, X. Multiple sclerosis: Pathology, diagnosis and treatments. *Exp. Ther. Med.* 2017, 13, 3163–3166. https://doi.org/10.3892/etm.2017.4410.
- 210. Barcelos, I.P.D.; Troxell, R.M.; Graves, J.S. Mitochondrial Dysfunction and Multiple Sclerosis. *Biology* 2019, *8*, 37. https://doi.org/10.3390/biology8020037.
- Tanaka, M.; Vécsei, L. Monitoring the Redox Status in Multiple Sclerosis. *Biomedicines* 2020, *8*, 406. https://doi.org/10.3390/bio-medicines8100406.
- 212. Procaccini, C.; De Rosa, V.; Pucino, V.; Formisano, L.; Matarese, G. Animal models of Multiple Sclerosis. *Eur. J. Pharmacol.* 2015, 759, 182–191. https://doi.org/10.1016/j.ejphar.2015.03.042.
- Pukoli, D.; Polyák, H.; Rajda, C.; Vécsei, L. Kynurenines and Neurofilament Light Chain in Multiple Sclerosis. *Front. Neurosci.* 2021, 15, 658202. https://doi.org/10.3389/fnins.2021.658202.
- Qi, X.; Lewin, A.S.; Sun, L.; Hauswirth, W.W.; Guy. J. Mitochondrial protein nitration primes neurodegeneration in experimental autoimmune encephalomyelitis. J. Biol. Chem. 2006, 281, 31950–31962. https://doi.org/10.1074/jbc.M603717200.
- Nikić, I.; Merkler, D.; Sorbara, C.; Brinkoetter, M.; Kreutzfeldt, M.; Bareyre, F.M.; Brück, W.; Bishop, D.; Misgeld, T.; Kerschensteiner, M. A reversible form of axon damage in experimental autoimmune encephalomyelitis and multiple sclerosis. *Nat. Med.* 2011, *17*, 495–499. https://doi.org/10.1038/nm.2324.
- 216. Sadeghian, M.; Mastrolia, V.; Rezaei Haddad, A.; Mosley, A.; Mullali, G.; Schiza, D.; Sajic, M.; Hargreaves, I.; Heales, S.; Duchen, M.R.; et al. Mitochondrial dysfunction is an important cause of neurological deficits in an inflammatory model of multiple sclerosis. *Sci. Rep.* 2016, 14, 33249. https://doi.org/10.1038/srep33249.
- 217. Wakabayashi, T.; Asano, M.; Kurono, C. Mechanism of the formation of megamitochondria induced by copper-chelating agents. II. Isolation and some properties of megamitochondria from the cuprizone-treated mouse liver. *Acta Pathol. Jpn.* **1975**, *25*, 39–49.
- Jhelum, P.; Santos-Nogueira, E.; Teo, W.; Haumont, A.; Lenoël, I.; Stys, P.K.; David, S. Ferroptosis Mediates Cuprizone-Induced Loss of Oligodendrocytes and Demyelination. *J. Neurosci.* 2020, 40, 9327–9341. https://doi.org/10.1523/JNEUROSCI.1749-20.2020.
- Praet, J.; Guglielmetti, C.; Berneman, Z.; Van der Linden, A.; Ponsaerts, P. Cellular and molecular neuropathology of the cuprizone mouse model: Clinical relevance for multiple sclerosis. *Neurosci. Biobehav. Rev.* 2014, 47, 485–505. https://doi.org/10.1016/j.neubiorev.2014.10.004.
- 220. Flatmark, T.; Kryvi, H.; Tangerås, A. Induction of megamitochondria by cuprizone (biscyclohexanone oxaldihydrazone). Evidence for an inhibition of the mitochondrial division process. *Eur. J. Cell Biol.* **1980**, *23*, 141–148.
- Kozin, M.; Kulakova, O.; Kiselev, I.; Baulina, N.; Boyko, A.; Favorova, O. Mitonuclear interactions influence multiple sclerosis risk. *Gene* 2020, 758, 144962. https://doi.org/10.1016/j.gene.2020.144962.
- 222. Polyák, H.; Cseh, E.K.; Bohár, Z.; Rajda, C.; Zádori, D.; Klivényi, P.; Toldi, J.; Vécsei, L. Cuprizone markedly decreases kynurenic acid levels in the rodent brain tissue and plasma. *Heliyon* **2021**, *7*, e06124. https://doi.org/10.1016/j.heliyon.2021.e06124.
- 223. Lim, C.K.; Bilgin, A.; Lovejoy, D.B.; Tan, V.; Bustamante, S.; Taylor, B.V.; Bessede, A.; Brew, B.J.; Guillemin, G.J. Kynurenine pathway metabolomics predicts and provides mechanistic insight into multiple sclerosis progression. *Sci. Rep.* 2017, *7*, 41473. https://doi.org/10.1038/srep41473.

- 224. Rajda, C.; Galla, Z.; Polyák, H.; Maróti, Z.; Babarczy, K.; Pukoli, D.; Vécsei, L. Cerebrospinal Fluid Neurofilament Light Chain Is Associated with Kynurenine Pathway Metabolite Changes in Multiple Sclerosis. *Int. J. Mol. Sci.* 2020, 21, 2665. https://doi.org/10.3390/ijms21082665.
- 225. Aeinehband, S.; Brenner, P.; Ståhl, S.; Bhat, M.; Fidock, M.D.; Khademi, M.; Olsson, T.; Engberg, G.; Jokinen, J.; Erhardt, S.; et al. Cerebrospinal fluid kynurenines in multiple sclerosis; relation to disease course and neurocognitive symptoms. *Brain Behav. Immun.* **2016**, *51*, 47–55. https://doi.org/10.1016/j.bbi.2015.07.016.
- 226. Hartai, Z.; Klivenyi, P.; Janaky, T.; Penke, B.; Dux, L.; Vécsei, L. Kynurenine metabolism in multiple sclerosis. Acta Neurol. Scand. 2005, 112, 93–96. https://doi.org/10.1111/j.1600-0404.2005.00442.x.
- 227. Reiner, A.; Dragatsis, I.; Dietrich, P. Genetics and neurophatology of huntington's disease. *Int. Rev. Neurobiol.* 2011, 98, 325–372. https://doi.org/10.1016/B978-0-12-381328-2.00014-6.
- 228. Mangiarini, L.; Sathasivam, K.; Seller, M.; Cozens, B.; Harper, A.; Hetherington, C.; Lawton, M.; Trottier, Y.; Lehrach, H.; Davies, S.W.; et al. Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice. *Cell* 1996, *87*, 493–506. https://doi.org/10.1016/s0092-867481369-0.
- Burtscher, J.; Di Pardo, A.; Maglione, V.; Schwarzer, C.; Squitieri, F. Mitochondrial Respiration Changes in R6/2 Huntington's Disease Model Mice during Aging in a Brain Region Specific Manner. *Int. J. Mol. Sci.* 2020, 21, 5412. https://doi.org/10.3390/ijms21155412.
- Gray, M.; Shirasaki, D.I.; Cepeda, C.; André, V.M.; Wilburn, B.; Lu, X.H.; Tao, J.; Yamazaki, I.; Li, S.H.; Sun, Y.E.; et al. Fulllength human mutant huntingtin with a stable polyglutamine repeat can elicit progressive and selective neuropathogenesis in BACHD mice. J. Neurosci. 2008, 28, 6182–6195. https://doi.org/10.1523/JNEUROSCI.0857-08.2008.
- 231. Menalled, L.B. Knock-in mouse models of Huntington's disease. *NeuroRx* 2005, 2, 465–470. https://doi.org/10.1602/neurorx.2.3.465.
- Arrasate, M.; Mitra, S.; Schweitzer, E.S.; Segal, M.R.; Finkbeiner, S. Inclusion body formation reduces levels of mutant huntingtin and the risk of neuronal death. *Nature* 2004, 431, 805–810. https://doi.org/10.1038/nature02998.
- Choo, Y.S.; Johnson, G.V.; MacDonald, M.; Detloff, P.J.; Lesort, M. Mutant huntingtin directly increases susceptibility of mitochondria to the calcium-induced permeability transition and cytochrome c release. *Hum. Mol. Genet.* 2004, 13, 1407–1420. https://doi.org/10.1093/hmg/ddh162.
- Kaye, J.; Reisine, T.; Finkbeiner, S. Huntington's disease mouse models: Unraveling the pathology caused by CAG repeat expansion. *Fac. Rev.* 2021, 10, 77. https://doi.org/10.12703/r/10-77.
- 235. Widner, B.; Leblhuber, F.; Walli, J.; Tilz, G.P.; Demel, U.; Fuchs, D. Degradation of tryptophan in neurodegenerative disorders. *Adv. Exp. Med. Biol.* **1999**, 467, 133–138. https://doi.org/10.1007/978-1-4615-4709-9\_19.
- 236. Guidetti, P.; Luthi-Carter, R.E.; Augood, S.J.; Schwarcz, R. Neostriatal and cortical quinolinate levels are increased in early grade Huntington's disease. *Neurobiol. Dis.* **2004**, *17*, 455–461. https://doi.org/10.1016/j.nbd.2004.07.006.
- 237. Pearson, S.J.; Reynolds, G.P. Increased brain concentrations of a neurotoxin, 3-hydroxykynurenine, in Huntington's disease. *Neurosci. Lett.* **1992**, 144, 199–201. https://doi.org/10.1016/0304-394090749-w.
- Iłzecka, J.; Kocki, T.; Stelmasiak, Z.; Turski, W.A. Endogenous protectant kynurenic acid in amyotrophic lateral sclerosis. *Acta Neurol. Scand.* 2003, 107, 412–418. https://doi.org/10.1034/j.1600-0404.2003.00076.x.
- Beal, M.F.; Matson, W.R.; Swartz, K.J.; Gamache, P.H.; Bird, E.D. Kynurenine pathway measurements in Huntington's disease striatum: Evidence for reduced formation of kynurenic acid. J. Neurochem. 1990, 55, 1327–1339. https://doi.org/10.1111/j.1471-4159.1990.tb03143.x.
- Forrest, C.M.; Mackay, G.M.; Stoy, N.; Spiden, S.L.; Taylor, R.; Stone, T.W.; Darlington, L.G. Blood levels of kynurenines, interleukin-23 and soluble human leucocyte antigen-G at different stages of Huntington's disease. J. Neurochem. 2010, 112, 112–122. https://doi.org/10.1111/j.1471-4159.2009.06442.x.
- 241. Gubbay, S.; Kahana, E.; Zilber, N.; Cooper, G.; Pintov, S.; Leibowitz, Y. Amyotrophic lateral sclerosis. A study of its presentation and prognosis. J. Neurol. 1985, 232, 295–300. https://doi.org/10.1007/BF00313868.
- 242. Smith, E.F.; Shaw, P.J.; De Vos, K.J. The role of mitochondria in amyotrophic lateral sclerosis. *Neurosci. Lett.* **2019**, *710*, 132933. https://doi.org/10.1016/j.neulet.2017.06.052.
- 243. Morrice, J.R.; Gregory-Evans, C.Y.; Shaw, C.A. Animal models of amyotrophic lateral sclerosis: A comparison of model validity. *Neural Regen. Res.* 2018, 13, 2050–2054. https://doi.org/10.4103/1673-5374.241445.
- 244. Dafinca, R.; Scaber, J.; Ababneh, N.; Lalic, T.; Weir, G.; Christian, H.; Vowles, J.; Douglas, A.G.; Fletcher-Jones, A.; Browne, C.; et al. C9orf72 Hexanucleotide Expansions Are Associated with Altered Endoplasmic Reticulum Calcium Homeostasis and Stress Granule Formation in Induced Pluripotent Stem Cell-Derived Neurons from Patients with Amyotrophic Lateral Sclerosis and Frontotemporal Dementia. *Stem Cells* 2016, 34, 2063–2078. https://doi.org/10.1002/stem.2388.
- 245. Magrané, J.; Cortez, C.; Gan, W.B.; Manfredi, G. Abnormal mitochondrial transport and morphology are common pathological denominators in SOD1 and TDP43 ALS mouse models. *Hum. Mol. Genet.* 2014, 23, 1413–1424. https://doi.org/10.1093/hmg/ddt528.
- 246. De Vos, K.J.; Chapman, A.L.; Tennant, M.E.; Manser, C.; Tudor, E.L.; Lau, K.F.; Brownlees, J.; Ackerley, S.; Shaw, P.J.; McLoughlin, D.M.; et al. Familial amyotrophic lateral sclerosis-linked SOD1 mutants perturb fast axonal transport to reduce axonal mitochondria content. *Hum. Mol. Genet.* 2007, *16*, 2720–2728. https://doi.org/10.1093/hmg/ddm226.

- 247. Wang, W.; Li, L.; Lin, W.L.; Dickson, D.W.; Petrucelli, L.; Zhang, T.; Wang, X. The ALS disease-associated mutant TDP-43 impairs mitochondrial dynamics and function in motor neurons. *Hum. Mol. Genet.* **2013**, *22*, 4706–4719. https://doi.org/10.1093/hmg/ddt319.
- 248. Agarwal, S.; Yadav, A.; Tiwari, S.K.; Seth, B.; Chauhan, L.K.; Khare, P.; Ray, R.S.; Chaturvedi, R.K. Dynamin-related Protein 1 Inhibition Mitigates Bisphenol A-mediated Alterations in Mitochondrial Dynamics and Neural Stem Cell Proliferation and Differentiation. J. Biol. Chem. 2016, 291, 15923–15939. https://doi.org/10.1074/jbc.M115.709493.
- 249. Panov, A.; Kubalik, N.; Brooks, B.R.; Shaw, C.A. In vitro effects of cholesterol β-D-glucoside, cholesterol and cycad phytosterol glucosides on respiration and reactive oxygen species generation in brain mitochondria. J. Membr. Biol. 2010, 237, 71–77. https://doi.org/10.1007/s00232-010-9307-9.
- 250. Chen, Y.; Stankovic, R.; Cullen, K.; Meininger, V.; Garner, B.; Coggan, S.; Grant, R.; Brew, B.J.; Guillemin, G.J. The kynurenine pathway and inflammation in amyotrophic lateral sclerosis. *Neurotox. Res.* **2010**, *18*, 132–142. https://doi.org/10.1007/s12640-009-9129-7.
- 251. Sparaco, M.; Feleppa, M.; Lipton, R.B.; Rapoport, A.M.; Bigal, M.E. Mitochondrial dysfunction and migraine: Evidence and hypotheses. *Cephalalgia* 2006, *26*, 361–372. https://doi.org/10.1111/j.1468-2982.2005.01059.x.
- 252. Montagna, P.; Sacquegna, T.; Martinelli, P.; Cortelli, P.; Bresolin, N.; Moggio, M.; Baldrati, A.; Riva, R.; Lugaresi, E. Mitochondrial Abnormalities in Migraine. Preliminary Findings. *Headache J. Head Face Pain* 1988, 28, 477–480. https://doi.org/10.1111/j.1526-4610.1988.hed2807477.x.
- 253. Uncini, A.; Lodi, R.; Di Muzio, A.; Silvestri, G.; Servidei, S.; Lugaresi, A.; Iotti, S.; Zaniol, P.; Barbiroli, B. Abnormal brain and muscle energy metabolism shown by 31P-MRS in familial hemiplegic migraine. *J. Neurol. Sci.* 1995, 129, 214–222. https://doi.org/10.1016/0022-510x(94)00283-t.
- Skinhøj, E. Hemodynamic Studies within the Brain during Migraine. Arch. Neurol. 1973, 29, 95–98. https://doi.org/10.1001/archneur.1973.00490260039007.
- 255. Okada, H.; Araga, S.; Takeshima, T.; Nakashima, K. Plasma lactic acid and pyruvic acid levels in migraine and tension-type headache. *Headache* **1998**, *38*, 39–42. https://doi.org/10.1046/j.1526-4610.1998.3801039.x.
- 256. Littlewood, J.; Glover, V.; Sandler, M.; Peatfield, R.; Petty, R.; Clifford Rose, F. Low platelet monoamine oxidase activity in headache: No correlation with phenolsulphotransferase, succinate dehydrogenase, platelet preparation method or smoking. J. Neurol. Neurosurg. Psychiatry 1984, 47, 338–343. https://doi.org/10.1136/jnnp.47.4.338.
- 257. Sangiorgi, S.; Mochi, M.; Riva, R.; Cortelli, P.; Monari, L.; Pierangeli, G.; Montagna, P. Abnormal platelet mitochondrial function in patients affected by migraine with and without aura. *Cephalalgia* **1994**, *14*, 21–23. https://doi.org/10.1046/j.1468-2982.1994.1401021.x.
- 258. Barbiroli, B.; Montagna, P.; Cortelli, P.; Funicello, R.; Iotti, S.; Monari, L.; Pierangeli, G.; Zaniol, P.; Lugaresi, E. Abnormal brain and muscle energy metabolism shown by 31P magnetic resonance spectroscopy in patients affected by migraine with aura. *Neurology* **1992**, *42*, 1209–1214. https://doi.org/10.1212/wnl.42.6.1209.
- 259. Schulz, U.G.; Blamire, A.M.; Corkill, R.G.; Davies, P.; Styles, P.; Rothwell, P.M. Association between cortical metabolite levels and clinical manifestations of migrainous aura: An MR-spectroscopy study. *Brain* **2007**, *130 Pt 12*, 3102–3110. https://doi.org/10.1093/brain/awm165.
- Reyngoudt, H.; Achten, E.; Paemeleire, K. Magnetic resonance spectroscopy in migraine: What have we learned so far? *Cephalalgia* 2012, 32, 845–859. https://doi.org/10.1177/0333102412452048.
- Gross, E.C.; Lisicki, M.; Fischer, D.; Sándor, P.S.; Schoenen, J. The metabolic face of migraine From pathophysiology to treatment. *Nat. Rev. Neurol.* 2019, 15, 627–643. https://doi.org/10.1038/s41582-019-0255-4.
- 262. Reyngoudt, H.; Paemeleire, K.; Descamps, B.; De Deene, Y.; Achten, E. 31P-MRS demonstrates a reduction in high-energy phosphates in the occipital lobe of migraine without aura patients. *Cephalalgia* 2011, 31, 1243–1253. https://doi.org/10.1177/0333102410394675.
- Dong, X.; Guan, X.; Chen, K.; Jin, S.; Wang, C.; Yan, L.; Shi, Z.; Zhang, X.; Chen, L.; Wan, Q. Abnormal mitochondrial dynamics and impaired mitochondrial biogenesis in trigeminal ganglion neurons in a rat model of migraine. *Neurosci. Lett.* 2017, 636, 127– 133. https://doi.org/10.1016/j.neulet.2016.10.054.
- 264. Fried, N.T.; Moffat, C.; Seifert, E.L.; Oshinsky, M.L. Functional mitochondrial analysis in acute brain sections from adult rats reveals mitochondrial dysfunction in a rat model of migraine. *Am. J. Physiol. Cell Physiol.* 2014, 307, C1017–C1030. https://doi.org/10.1152/ajpcell.00332.2013.
- 265. Li, R.; Liu, Y.; Chen, N.; Zhang, Y.; Song, G.; Zhang, Z. Valproate Attenuates Nitroglycerin-Induced Trigeminovascular Activation by Preserving Mitochondrial Function in a Rat Model of Migraine. *Med. Sci. Monit.* 2016, 22, 3229–3237. https://doi.org/10.12659/msm.900185.
- 266. Curto, M.; Lionetto, L.; Negro, A.; Capi, M.; Perugino, F.; Fazio, F.; Giamberardino, M.A.; Simmaco, M.; Nicoletti, F.; Martelletti, P. Altered serum levels of kynurenine metabolites in patients affected by cluster headache. *J. Headache Pain* 2015, 17, 27. https://doi.org/10.1186/s10194-016-0620-2.
- Curto, M.; Lionetto, L.; Negro, A.; Capi, M.; Fazio, F.; Giamberardino, M.A.; Simmaco, M.; Nicoletti, F.; Martelletti, P. Altered kynurenine pathway metabolites in serum of chronic migraine patients. *J. Headache Pain* 2015, 17, 47. https://doi.org/10.1186/s10194-016-0638-5.

- 268. Tuka, B.; Nyári, A.; Cseh, E.K.; Körtési, T.; Veréb, D.; Tömösi, F.; Kecskeméti, G.; Janáky, T.; Tajti, J.; Vécsei, L. Clinical relevance of depressed kynurenine pathway in episodic migraine patients: Potential prognostic markers in the peripheral plasma during the interictal period. *J. Headache Pain* 2021, 22, 60. https://doi.org/10.1186/s10194-021-01239-1.
- Nagy-Grócz, G.; Tar, L.; Bohár, Z.; Fejes-Szabó, A.; Laborc, K.F.; Spekker, E.; Vécsei, L.; Párdutz, Á. The modulatory effect of anandamide on nitroglycerin-induced sensitization in the trigeminal system of the rat. *Cephalalgia* 2016, 36, 849–861. https://doi.org/10.1177/0333102415613766.
- Spekker, E.; Tanaka, M.; Szabó, Á.; Vécsei, L. Neurogenic Inflammation: The Participant in Migraine and Recent Advancements in Translational Research. *Biomedicines* 2022, 10, 76. https://doi.org/10.3390/biomedicines10010076.
- Tanaka, M.; Török, N.; Vécsei, L. Are 5-HT1 receptor agonists effective anti-migraine drugs? *Expert. Opin. Pharmacother.* 2021, 22, 1221–1225. https://doi.org/10.1080/14656566.2021.1910235.
- 272. Tanaka, M.; Török, N.; Tóth, F.; Szabó, Á.; Vécsei, L. Co-Players in Chronic Pain: Neuroinflammation and the Tryptophan-Kynurenine Metabolic Pathway. *Biomedicines* 2021, *9*, 897. https://doi.org/10.3390/biomedicines9080897.
- 273. Ciapała, K.; Mika, J.; Rojewska, E. The Kynurenine Pathway as a Potential Target for Neuropathic Pain Therapy Design: From Basic Research to Clinical Perspectives. *Int. J. Mol. Sci.* **2021**, *22*, 11055. https://doi.org/10.3390/ijms222011055.
- Jovanovic, F.; Candido, K.D.; Knezevic, N.N. The Role of the Kynurenine Signaling Pathway in Different Chronic Pain Conditions and Potential Use of Therapeutic Agents. *Int. J. Mol. Sci.* 2020, 21, 6045. https://doi.org/10.3390/ijms21176045.
- 275. Gecse, K.; Édes, A.E.; Nagy, T.; Demeter, A.K.; Virág, D.; Király, M.; Dalmadi Kiss, B.; Ludányi, K.; Környei, Z.; Denes, A.; et al. Citalopram Neuroendocrine Challenge Shows Altered Tryptophan and Kynurenine Metabolism in Migraine. *Cells* 2022, 11, 2258. https://doi.org/10.3390/cells1114225.
- 276. Fila, M.; Chojnacki, J.; Pawlowska, E.; Szczepanska, J.; Chojnacki, C.; Blasiak, J. Kynurenine Pathway of Tryptophan Metabolism in Migraine and Functional Gastrointestinal Disorders. *Int. J. Mol. Sci.* **2021**, *22*, 10134. https://doi.org/10.3390/ijms221810134.
- 277. Tanaka, M.; Schally, A.V.; Telegdy, G. Neurotransmission of the antidepressant-like effects of the growth hormone-releasing hormone antagonist MZ-4-71. *Behav. Brain Res.* 2012, 228, 388–391. https://doi.org/10.1016/j.bbr.2011.12.02.
- Tanaka, M.; Telegdy, G. Neurotransmissions of antidepressant-like effects of neuromedin U-23 in mice. *Behav. Brain Res.* 2014, 259, 196–199. https://doi.org/10.1016/j.bbr.2013.11.005.
- 279. Castillo-Mariqueo, L.; Giménez-Llort, L. Impact of Behavioral Assessment and Re-Test as Functional Trainings That Modify Survival, Anxiety and Functional Profile (Physical Endurance and Motor Learning) of Old Male and Female 3xTg-AD Mice and NTg Mice with Normal Aging. *Biomedicines* 2022, 10, 973. https://doi.org/10.3390/biomedicines10050973
- Tanaka, M.; Telegdy, G. Antidepressant-like Effects of Neuropeptide SF (NPSF). arXiv 2020, arXiv:200513256. https://doi.org/10.48550/arXiv.2005.13256.
- Gutiérrez-Rojas, L.; Porras-Segovia, A.; Dunne, H.; Andrade-González, N.; Cervilla, J.A. Prevalence and correlates of major depressive disorder: A systematic review. *Braz. J. Psychiatry* 2020, 42, 657–672. https://doi.org/10.1590/1516-4446-2020-0650.
- 282. Gładka, A.; Zatoński, T.; Rymaszewska, J. Association between the long-term exposure to air pollution and depression. *Adv. Clin. Exp. Med.* **2022**. https://doi.org/10.17219/acem/149988.
- 283. Chen, C. Recent advances in the study of the comorbidity of depressive and anxiety disorders. *Adv. Clin. Exp. Med.* **2022**, *31*, 355–358. https://doi.org/10.17219/acem/147441.
- Carrera-González, M.; Cantón-Habas, V.; Rich-Ruiz, M. Aging, depression and dementia: The inflammatory process. *Adv. Clin. Exp. Med.* 2022, 31, 469–473. https://doi.org/10.17219/acem/149897.
- 285. Tanaka, M. Crosstalk between Depression, Anxiety, and Dementia: Comorbidity in Behavioral Neurology and Neuropsychiatry, 1st ed.; MDPI: Basel, Switzerland, 2022; pp. 1–266.
- 286. Delgado, P.L. Depression: The case for a monoamine deficiency. J. Clin. Psychiatry 2000, 61 (Suppl. 6), 7–11.
- 287. Balogh, L.; Tanaka, M.; Török, N.; Vécsei, L.; Taguchi, S. Crosstalk between Existential Phenomenological Psychotherapy and Neurological Sciences in Mood and Anxiety Disorders. *Biomedicines* **2021**, *9*, 340. https://doi.org/10.3390/biomedicines9040340.
- 288. Wohleb, E.; Franklin, T.; Iwata, M.; Duman, R.S. Integrating neuroimmune systems in the neurobiology of depression. *Nat. Rev. Neurosci.* **2016**, *17*, 497–511. https://doi.org/10.1038/nrn.2016.69.
- 289. Adeel, M.; Chen, C.-C.; Lin, B.-S.; Chen, H.-C.; Liou, J.-C.; Li, Y.-T.; Peng, C.-W. Safety of Special Waveform of Transcranial Electrical Stimulation (TES): In Vivo Assessment. *Int. J. Mol. Sci.* 2022, 23, 6850. https://doi.org/10.3390/ijms23126850.
- Taliaz, D.; Spinrad, A.; Barzilay, R.; Barnett-Itzhaki, Z.; Averbuch, D.; Teltsh, O.; Schurr, R.; Darki-Morag, S.; Lerer, B. Optimizing prediction of response to antidepressant medications using machine learning and integrated genetic, clinical, and demographic data. *Transl. Psychiatry* 2021, *11*, 381. https://doi.org/10.1038/s41398-021-01488-3.
- 291. Gamaro, G.D.; Streck, E.L.; Matté, C.; Prediger, M.E.; Wyse, A.T.; Dalmaz, C. Reduction of hippocampal Na+, K+-ATPase activity in rats subjected to an experimental model of depression. *Neurochem. Res.* 2003, 28, 1339–1344. https://doi.org/10.1023/a:1024988113978.
- 292. Gong, Y.; Chai, Y.; Ding, J.H.; Sun, X.L.; Hu, G. Chronic mild stress damages mitochondrial ultrastructure and function in mouse brain. *Neurosci. Lett.* 2011, 488, 76–80. https://doi.org/10.1016/j.neulet.2010.11.006.
- Wen, L.; Jin, Y.; Li, L.; Sun, S.; Cheng, S.; Zhang, S.; Zhang, Y.; Svenningsson, P. Exercise prevents raphe nucleus mitochondrial overactivity in a rat depression model. *Physiol. Behav.* 2014, 132, 57–65. https://doi.org/10.1016/j.physbeh.2014.04.050.
- Trowbridge, S.; Narboux-Nême, N.; Gaspar, P. Genetic models of serotonin (5-HT) depletion: What do they tell us about the developmental role of 5-HT? *Anat. Rec.* 2011, 294, 1615–1623. https://doi.org/10.1002/ar.21248.

- 295. Crane, J.D.; Palanivel, R.; Mottillo, E.P.; Bujak, A.L.; Wang, H.; Ford, R.J.; Collins, A.; Blümer, R.M.; Fullerton, M.D.; Yabut, J.M.; et al. Inhibiting peripheral serotonin synthesis reduces obesity and metabolic dysfunction by promoting brown adipose tissue thermogenesis. *Nat. Med.* 2015, 21, 166–172. https://doi.org/10.1038/nm.3766.
- 296. Mosienko, V.; Bert, B.; Beis, D.; Matthes, S.; Fink, H.; Bader, M.; Alenina, N. Exaggerated aggression and decreased anxiety in mice deficient in brain serotonin. *Transl. Psychiatry* 2012, 2, e122. https://doi.org/10.1038/tp.2012.44.
- 297. Gutknecht, L.; Popp, S.; Waider, J.; Sommerlandt, F.M.; Göppner, C.; Post, A.; Reif, A.; van den Hove, D.; Strekalova, T.; Schmitt, A.; et al. Interaction of brain 5-HT synthesis deficiency, chronic stress and sex differentially impact emotional behavior in Tph2 knockout mice. *Psychopharmacology* 2015, 232, 2429–2441. https://doi.org/10.1007/s00213-015-3879-0.
- Wang, Q.; Timberlake, M.A., 2nd; Prall, K.; Dwivedi, Y. The recent progress in animal models of depression. Prog. Neuropsychopharmacol. Biol. Psychiatry 2017, 77, 99–109. https://doi.org/10.1016/j.pnpbp.2017.04.008.
- Beaulieu, J.M.; Zhang, X.; Rodriguiz, R.M.; Sotnikova, T.D.; Cools, M.J.; Wetsel, W.C.; Gainetdinov, R.R.; Caron, M.G. Role of GSK3 beta in behavioral abnormalities induced by serotonin deficiency. *Proc. Natl. Acad. Sci. USA* 2008, 105, 1333–1338. https://doi.org/10.1073/pnas.0711496105.
- 300. Ogawa, S.; Fujii, T.; Koga, N.; Hori, H.; Teraishi, T.; Hattori, K.; Noda, T.; Higuchi, T.; Motohashi, N.; Kunugi, H. Plasma L-tryptophan concentration in major depressive disorder: New data and meta-analysis. J. Clin. Psychiatry 2014, 75, e906–e915. https://doi.org/10.4088/JCP.13r08908.
- 301. Ogyu, K.; Kubo, K.; Noda, Y.; Iwata, Y.; Tsugawa, S.; Omura, Y.; Wada, M.; Tarumi, R.; Plitman, E.; Moriguchi, S.; et al. Kynurenine pathway in depression: A systematic review and meta-analysis. *Neurosci. Biobehav. Rev.* 2018, 90, 16–25. https://doi.org/10.1016/j.neubiorev.2018.03.023.
- Tanaka, M.; Török, N.; Vécsei, L. Novel Pharmaceutical Approaches in Dementia. In *NeuroPsychopharmacotherapy*; Riederer, P., Laux, G., Nagatsu, T., Le, W., Riederer, C., Eds.; Springer: Cham, Switzerland, 2021. https://doi.org/10.1007/978-3-319-56015-1\_444-1.
- 303. Hunt, C.; Macedo e Cordeiro, T.; Suchting, R.; de Dios, C.; Cuellar Leal, V.A.; Soares, J.C.; Dantzer, R.; Teixeira, A.L.; Selvaraj, S. Effect of immune activation on the kynurenine pathway and depression symptoms—A systematic review and meta-analysis. *Neurosci. Biobeha. Rev.* 2020, *118*, 514. https://doi.org/j.neubiorev.2020.08.010.
- 304. Erabi, H.; Okada, G.; Shibasaki, C.; Setoyama, D.; Kang, D.; Takamura, M.; Yoshino, A.; Fuchikami, M.; Kurata, A.; Kato, T.A.; et al. Kynurenic acid is a potential overlapped biomarker between diagnosis and treatment response for depression from metabolome analysis. *Sci. Rep.* 2020, *10*, 16822. https://doi.org/10.1038/s41598-020-73918-z.
- 305. Carrillo-Mora, P.; Pérez-De la Cruz, V.; Estrada-Cortés, B.; Toussaint-González, P.; Martínez-Cortéz, J.A.; Rodríguez-Barragán, M.; Quinzaños-Fresnedo, J.; Rangel-Caballero, F.; Gamboa-Coria, G.; Sánchez-Vázquez, I.; et al. Serum Kynurenines Correlate with Depressive Symptoms and Disability in Poststroke Patients: A Cross-sectional Study. *Neurorehabilit. Neural Repair* 2020, 34, 154596832095367. https://doi.org/10.1177/1545968320953671.
- 306. Ruscio, A.M.; Hallion, L.S.; Lim, C.C.W.; Aguilar-Gaxiola, S.; Al-Hamzawi, A.; Alonso, J.; Andrade, L.H.; Borges, G.; Bromet, E.J.; Bunting, B.; et al. Cross-sectional Comparison of the Epidemiology of DSM-5 Generalized Anxiety Disorder Across the Globe. JAMA Psychiatry 2017, 74, 465–475. https://doi.org/10.1001/jamapsychiatry.2017.0056.
- Martin, E.I.; Ressler, K.J.; Binder, E.; Nemeroff, C.B. The neurobiology of anxiety disorders: Brain imaging, genetics, and psychoneuroendocrinology. Psychiatr. *Clin. N. Am.* 2009, 32, 549–575. https://doi.org/10.1016/j.psc.2009.05.004.
- Strawn, J.R.; Geracioti, L.; Rajdev, N.; Clemenza, K.; Levine, A. Pharmacotherapy for generalized anxiety disorder in adult and pediatric patients: An evidence-based treatment review. *Expert. Opin. Pharmacother.* 2018, 19, 1057–1070. https://doi.org/10.1080/14656566.2018.1491966.
- 309. Bandelow, B.; Michaelis, S.; Wedekind, D. Treatment of anxiety disorders. Dialogues *Clin. Neurosci.* 2017, 19(2), 93-107. https://doi.org/10.31887/DCNS.2017.19.2/bbandelow.
- Chen, A.P.; Chen, L.; Kim, T.A.; Xiong, Q. Integrating the Roles of Midbrain Dopamine Circuits in Behavior and Neuropsychiatric Disease. *Biomedicines* 2021, 9, 647. https://doi.org/10.3390/biomedicines9060647.
- Gebara, E.; Zanoletti, O.; Ghosal, S.; Grosse, J.; Schneider, B.L.; Knott, G.; Astori, S.; Sandi, C. Mitofusin-2 in the Nucleus Accumbens Regulates Anxiety and Depression-like Behaviors Through Mitochondrial and Neuronal Actions. *Biol. Psychiatry* 2021, 89, 1033–1044. https://doi.org/10.1016/j.biopsych.2020.12.003.
- Hollis, F.; van der Kooij, M.A.; Zanoletti, O.; Lozano, L.; Cantó, C.; Sandi, C. Mitochondrial function in the brain links anxiety with social subordination. *Proc. Natl. Acad. Sci. USA* 2015, *112*, 15486–15491. https://doi.org/10.1073/pnas.1512653112.
- Orlikov, A.B.; Prakhye, I.B.; Ryzov, I.V. Kynurenine in blood plasma and DST in patients with endogenous anxiety and endogenous depression. *Biol. Psychiatry* 1994, 36, 97–102. https://doi.org/10.1016/0006-322391189-4.
- 314. Altmaier, E.; Emeny, R.T.; Krumsiek, J.; Lacruz, M.E.; Lukaschek, K.; Häfner, S.; Kastenmüller, G.; Römisch-Margl, W.; Prehn, C.; Mohney, R.P.; et al. Metabolomic profiles in individuals with negative affectivity and social inhibition: A population-based study of Type D personality. *Psychoneuroendocrinology* 2013, *38*, 1299–1309. https://doi.org/10.1016/j.psyneuen.2012.11.014.
- 315. National Institute of Mental Health. Transforming the Understanding and Treatment of Mental Illnesses. Post-Traumatic Stress Disorder. Available online: https://www.nimh.nih.gov/health/publications/post-traumatic-stress-disorder-ptsd (accessed on 15 June 2022).
- 316. Sareen, J.; Stein, M.B.; Friedman, M. Posttraumatic Stress Disorder in Adults: Epidemiology, Pathophysiology, Clinical Manifestations, Course, Assessment, and Diagnosis. UpToDate 2022. Available online: https://www.uptodate.com/contents/post-

traumatic-stress-disorder-in-adults-epidemiology-pathophysiology-clinical-manifestations-course-assessment-and-diagnosis/print#:~:text=PTSD%20prevalence%20%E2%80%94%20The%20lifetime%20prevalence,percent%20%5B6%2C7%5D (accessed on 15 June 2022).

- Bedard-Gilligan, M.; Zoellner, L.A.; Feeny, N.C. Is Trauma Memory Special? Trauma Narrative Fragmentation in PTSD: Effects of Treatment and Response. *Clin. Psychol. Sci.* 2017, *5*, 212–225. https://doi.org/10.1177/2167702616676581.
- 318. Ehret, M. Treatment of posttraumatic stress disorder: Focus on pharmacotherapy. *Ment. Health Clin.* 2019, *9*, 373–382. https://doi.org/10.9740/mhc.2019.11.373.
- Richter-Levin, G.; Stork, O.; Schmidt, M.V. Animal models of PTSD: A challenge to be met. *Mol. Psychiatry* 2019, 24, 1135–1156. https://doi.org/10.1038/s41380-018-0272-5.
- 320. Sabbagh, J.J.; O'Leary, J.C., 3rd; Blair, L.J.; Klengel, T.; Nordhues, B.A.; Fontaine, S.N.; Binder, E.B.; Dickey, C.A. Age-associated epigenetic upregulation of the FKBP5 gene selectively impairs stress resiliency. *PLoS ONE* 2014, 9, e107241. https://doi.org/10.1371/journal.pone.0107241.
- 321. King, S.B.; Lezak, K.R.; O'Reilly, M.; Toufexis, D.J.; Falls, W.A.; Braas, K.; May, V.; Hammack, S.E. The Effects of Prior Stress on Anxiety-Like Responding to Intra-BNST Pituitary Adenylate Cyclase Activating Polypeptide in Male and Female Rats. *Neuro*psychopharmacology 2017, 42, 1679–1687. https://doi.org/10.1038/npp.2017.16.
- Otto, C.; Martin, M.; Wolfer, D.P.; Lipp, H.P.; Maldonado, R.; Schütz, G. Altered emotional behavior in PACAP-type-I-receptordeficient mice. *Brain Res. Mol. Brain Res.* 2001, 92, 78–84. https://doi.org/10.1016/s0169-328x(01)00153-x.
- 323. Hill, J.L.; Hardy, N.F.; Jimenez, D.V.; Maynard, K.R.; Kardian, A.S.; Pollock, C.J.; Schloesser, R.J.; Martinowich, K. Loss of promoter IV-driven BDNF expression impacts oscillatory activity during sleep, sensory information processing and fear regulation. *Transl. Psychiatry* 2016, *6*, e873. https://doi.org/10.1038/tp.2016.153.
- 324. Soliman, F.; Glatt, C.E.; Bath, K.G.; Levita, L.; Jones, R.M.; Pattwell, S.S.; Jing, D.; Tottenham, N.; Amso, D.; Somerville, L.H.; et al. A genetic variant BDNF polymorphism alters extinction learning in both mouse and human. *Science* 2010, 327, 863–866. https://doi.org/10.1126/science.1181886.
- Klemenhagen, K.C.; Gordon, J.A.; David, D.J.; Hen, R.; Gross, C.T. Increased fear response to contextual cues in mice lacking the 5-HT1A receptor. *Neuropsychopharmacology* 2006, *31*, 101–111. https://doi.org/10.1038/sj.npp.1300774.
- Wellman, C.L.; Izquierdo, A.; Garrett, J.E.; Martin, K.P.; Carroll, J.; Millstein, R.; Lesch, K.P.; Murphy, D.L.; Holmes, A. Impaired stress-coping and fear extinction and abnormal corticolimbic morphology in serotonin transporter knock-out mice. *J. Neurosci.* 2007, 27, 684–691. https://doi.org/10.1523/JNEUROSCI.4595-06.2007.
- 327. O'Tuathaigh, C.M.; Clarke, G.; Walsh, J.; Desbonnet, L.; Petit, E.; O'Leary, C.; Tighe, O.; Clarke, N.; Karayiorgou, M.; Gogos, J.A.; et al. Genetic vs. pharmacological inactivation of COMT influences cannabinoid-induced expression of schizophrenia-related phenotypes. *Int. J. Neuropsychopharmacol.* 2012, *15*, 1331–1342. https://doi.org/10.1017/S1461145711001581.
- 328. Bergado-Acosta, J.R.; Sangha, S.; Narayanan, R.T.; Obata, K.; Pape, H.C.; Stork, O. Critical role of the 65-kDa isoform of glutamic acid decarboxylase in consolidation and generalization of Pavlovian fear memory. *Learn. Mem.* 2008, 15, 163–171. https://doi.org/10.1101/lm.705408.
- Sangha, S.; Narayanan, R.T.; Bergado-Acosta, J.R.; Stork, O.; Seidenbecher, T.; Pape, H.C. Deficiency of the 65 kDa isoform of glutamic acid decarboxylase impairs extinction of cued but not contextual fear memory. J. Neurosci. 2009, 29, 15713–15720. https://doi.org/10.1523/JNEUROSCI.2620-09.2009.
- 330. Shaban, H.; Humeau, Y.; Herry, C.; Cassasus, G.; Shigemoto, R.; Ciocchi, S.; Barbieri, S.; van der Putten, H.; Kaupmann, K.; Bettler, B.; et al. Generalization of amygdala LTP and conditioned fear in the absence of presynaptic inhibition. *Nat. Neurosci.* 2006, 9, 1028–1035. https://doi.org/10.1038/nn1732.
- Fride, E.; Suris, R.; Weidenfeld, J.; Mechoulam, R. Differential response to acute and repeated stress in cannabinoid CB1 receptor knockout newborn and adult mice. *Behav. Pharmacol.* 2005, *16*, 431–440. https://doi.org/10.1097/00008877-200509000-00016.
- 332. Preston, G.; Emmerzaal, T.; Kirdar, F.; Schrader, L.; Henckens, M.; Morava, E.; Kozicz, T. Cerebellar mitochondrial dysfunction and concomitant multi-system fatty acid oxidation defects are sufficient to discriminate PTSD-like and resilient male mice. *Brain Behav. Immun. Health* 2020, *6*, 100104. https://doi.org/10.1016/j.bbih.2020.100104.
- Jia, Y.; Han, Y.; Wang, X.; Han, F. Role of apoptosis in the Post-traumatic stress disorder model-single prolonged stressed rats. *Psychoneuroendocrinology* 2018, 95, 97–105. https://doi.org/10.1016/j.psyneuen.2018.05.015.
- 334. Giménez-Gómez, P.; Ballestín, R.; Gil de Biedma-Elduayen, L.; Vidal, R.; Ferrer-Pérez, C.; Reguilón, M.D.; O'Shea, E.; Miñarro, J.; Colado, M.I.; Rodríguez-Arias, M. Decreased kynurenine pathway potentiate resilience to social defeat effect on cocaine reward. *Neuropharmacology* 2021, 197, 108753. https://doi.org/10.1016/j.neuropharm.2021.
- 335. Lin, C.-C.; Cheng, P.-Y.; Hsiao, M.; Liu, Y.-P. Effects of RU486 in Treatment of Traumatic Stress-Induced Glucocorticoid Dysregulation and Fear-Related Abnormalities: Early versus Late Intervention. Int. J. Mol. Sci. 2022, 23, 5494. https://doi.org/10.3390/ijms23105494.
- 336. Kim, T.D.; Lee, S.; Yoon, S. Inflammation in Post-Traumatic Stress Disorder (PTSD): A Review of Potential Correlates of PTSD with a Neurological Perspective. *Antioxidants* 2020, 9, 107. https://doi.org/10.3390/antiox9020107.
- 337. Merikangas, K.R.; Jin, R.; He, J.P.; Kessler, R.C.; Lee, S.; Sampson, N.A.; Viana, M.C.; Andrade, L.H.; Hu, C.; Karam, E.G.; et al. Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Arch. Gen. Psychiatry* 2011, 68, 241–251. https://doi.org/10.1001/archgenpsychiatry.2011.12.
- Drevets, W.C.; Price, J.L.; Furey, M.L. Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Struct. Funct.* 2008, 213, 93–118. https://doi.org/10.1007/s00429-008-0189-x.

- Womer, F.Y.; Kalmar, J.H.; Wang, F.; Blumberg, H.P. A Ventral Prefrontal-Amygdala Neural System in Bipolar Disorder: A View from Neuroimaging Research. Acta Neuropsychiatr. 2009, 21, 228–238.
- Cao, B.; Passos, I.C.; Mwangi, B.; Amaral-Silva, H.; Tannous, J.; Wu, M.J.; Zunta-Soares, G.B.; Soares, J.C. Hippocampal subfield volumes in mood disorders. *Mol. Psychiatry* 2017, 22, 1352–1358. https://doi.org/10.1038/mp.2016.262.
- Brady, R.O., Jr.; McCarthy, J.M.; Prescot, A.P.; Jensen, J.E.; Cooper, A.J.; Cohen, B.M.; Renshaw, P.F.; Ongür, D. Brain gammaaminobutyric acid (GABA) abnormalities in bipolar disorder. *Bipolar Disord*. 2013, 15, 434–439. https://doi.org/10.1111/bdi.12074.
- 342. Beyer, D.K.E.; Freund, N. Animal models for bipolar disorder: From bedside to the cage. Int. J. Bipolar Disord. 2017, 5, 35. https://doi.org/10.1186/s40345-017-0104-6.
- 343. Kristensen, M.; Nierenberg, A.A.; Østergaard, S.D. Face and predictive validity of the Clock∆19 mouse as an animal model for bipolar disorder: A systematic review. *Mol. Psychiatry* 2018, 23, 70–80. https://doi.org/10.1038/mp.2017.192.
- 344. van Enkhuizen, J.; Minassian, A.; Young, J.W. Further evidence for Clock∆19 mice as a model for bipolar disorder mania using cross-species tests of exploration and sensorimotor gating. *Behav. Brain Res.* 2013, 249, 44–54. https://doi.org/10.1016/j.bbr.2013.04.023.
- 345. Kasahara, T.; Takata, A.; Kato, T.M.; Kubota-Sakashita, M.; Sawada, T.; Kakita, A.; Mizukami, H.; Kaneda, D.; Ozawa, K.; Kato, T. Depression-like episodes in mice harboring mtDNA deletions in paraventricular thalamus. *Mol. Psychiatry* 2016, 21, 39–48. https://doi.org/10.1038/mp.2015.156.
- Kato, T. Neurobiological basis of bipolar disorder: Mitochondrial dysfunction hypothesis and beyond. Schizophr. Res. 2017, 187, 62–66. https://doi.org/10.1016/j.schres.2016.10.037.
- Giménez-Palomo, A.; Dodd, S.; Anmella, G.; Carvalho, A.F.; Scaini, G.; Quevedo, J.; Pacchiarotti, I.; Vieta, E.; Berk, M. The Role of Mitochondria in Mood Disorders: From Physiology to Pathophysiology and to Treatment. *Front. Psychiatry* 2021, 12, 546801. https://doi.org/10.3389/fpsyt.2021.546801.
- 348. Birner, A.; Platzer, M.; Bengesser, S.A.; Dalkner, N.; Fellendorf, F.T.; Queissner, R.; Pilz, R.; Rauch, P.; Maget, A.; Hamm, C.; et al. Increased breakdown of kynurenine towards its neurotoxic branch in bipolar disorder. *PLoS ONE* 2017, 12, e0172699. https://doi.org/10.1371/journal.pone.0172699.
- Arnone, D.; Saraykar, S.; Salem, H.; Teixeira, A.L.; Dantzer, R.; Selvaraj, S. Role of Kynurenine pathway and its metabolites in mood disorders: A systematic review and meta-analysis of clinical studies. *Neurosci. Biobehav. Rev.* 2018, 92, 477–485. https://doi.org/10.1016/j.neubiorev.2018.05.031.
- 350. Hiles, S.A.; Baker, A.L.; de Malmanche, T.; Attia, J. A meta-analysis of differences in IL-6 and IL-10 between people with and without depression: Exploring the causes of heterogeneity. *Brain Behav. Immun.* **2012**, *26*, 1180–1188. https://doi.org/10.1016/j.bbi.2012.06.001.
- 351. Substance Abuse and Mental Health Services Administration (US); Office of the Surgeon General (US). Chapter 2, the Neurobiology of Substance Use, Misuse, and Addiction. *Facing Addiction in America: The Surgeon General's Report on Alcohol, Drugs, and Health [Internet]*; US Department of Health and Human Services: Washington, DC, USA, 2016. Available online: https://www.ncbi.nlm.nih.gov/books/NBK424849/ (accessed on 15 June 2022).
- 352. National Institute on Drug Abuse. Advancing Addiction Science. Available online: https://www.drugabuse.gov/publications/media-guide/science-drug-use-addiction-basics (accessed on 15 June 2022).
- 353. Uhl, G.R.; Koob, G.F.; Cable, J. The neurobiology of addiction. Ann. N. Y. Acad. Sci. 2019, 451, 5–28. https://doi.org/10.1111/nyas.13989.
- 354. Calarco, C.A.; Fox, M.E.; Van Terheyden, S.; Turner, M.D.; Alipio, J.B.; Chandra, R.; Lobo, M.K. Mitochondria-Related Nuclear Gene Expression in the Nucleus Accumbens and Blood Mitochondrial Copy Number After Developmental Fentanyl Exposure in Adolescent Male and Female C57BL/6 Mice. Front. Psychiatry 2021, 12, 737389. https://doi.org/10.3389/fpsyt.2021.737389.
- 355. Morales-Puerto, N.; Giménez-Gómez, P.; Pérez-Hernández, M.; Abuin-Martínez, C.; Gil de Biedma-Elduayen, L.; Vidal, R.; Gutiérrez-López, M.D.; O'Shea, E.; Colado, M.I. Addiction and the kynurenine pathway: A new dancing couple? *Pharmacol. Ther.* 2021, 223, 107807. https://doi.org/10.1016/j.pharmthera.2021.107807.
- 356. Fonseca, F.; Mestre-Pintó, J.I.; Gómez-Gómez, Á.; Martinez-Sanvisens, D.; Rodríguez-Minguela, R.; Papaseit, E.; Pérez-Mañá, C.; Langohr, K.; Valverde, O.; Pozo, Ó.J.; et al. The Tryptophan System in Cocaine-Induced Depression. J. Clin. Med. 2020, 9, 4103. https://doi.org/10.3390/jcm9124103.
- 357. Salazar de Pablo, G.; Woods, S.W.; Drymonitou, G.; de Diego, H.; Fusar-Poli, P. Prevalence of Individuals at Clinical High-Risk of Psychosis in the General Population and Clinical Samples: Systematic Review and Meta-Analysis. *Brain Sci.* **2021**, *11*, 1544. https://doi.org/10.3390/brainsci11111544.
- 358. Gaebler, A.J.; Finner-Prével, M.; Sudar, F.P.; Langer, F.H.; Keskin, F.; Gebel, A.; Zweerings, J.; Mathiak, K. The Interplay between Vitamin D, Exposure of Anticholinergic Antipsychotics and Cognition in Schizophrenia. *Biomedicines* 2022, 10, 1096. https://doi.org/10.3390/biomedicines10051096.
- Nyatega, C.O.; Qiang, L.; Adamu, M.J.; Younis, A.; Kawuwa, H.B. Altered Dynamic Functional Connectivity of Cuneus in Schizophrenia Patients: A Resting-State fMRI Study. *Appl. Sci.* 2021, *11*, 11392. https://doi.org/10.3390/app112311392.
- Tanaka, M.; Spekker, E.; Szabó, Á.; Polyák, H.; Vécsei, L. Modelling the neurodevelopmental pathogenesis in neuropsychiatric disorders. Bioactive kynurenines and their analogues as neuroprotective agents—In celebration of 80th birthday of Professor Peter Riederer. J. Neural. Transm. 2022, 129, 627–642. https://doi.org/10.1007/s00702-022-02513-5.
- Panov, G. Dissociative Model in Patients with Resistant Schizophrenia. Front. Psychiatry 2022, 13, 845493. https://doi.org/10.3389/fpsyt.2022.845493.

- 362. Avan, R.; Sahebnasagh, A.; Hashemi, J.; Monajati, M.; Faramarzi, F.; Henney, N.C.; Montecucco, F.; Jamialahmadi, T.; Sahebkar, A. Update on Statin Treatment in Patients with Neuropsychiatric Disorders. *Life* 2021, 11, 1365. https://doi.org/10.3390/life11121365.
- Correia, B.S.B.; Nani, J.V.; Waladares Ricardo, R.; Stanisic, D.; Costa, T.B.B.C.; Hayashi, M.A.F.; Tasic, L. Effects of Psychostimulants and Antipsychotics on Serum Lipids in an Animal Model for Schizophrenia. *Biomedicines* 2021, 9, 235. https://doi.org/10.3390/biomedicines9030235.
- 364. Rog, J.; Błażewicz, A.; Juchnowicz, D.; Ludwiczuk, A.; Stelmach, E.; Kozioł, M.; Karakula, M.; Niziński, P.; Karakula-Juchnowicz, H. The Role of GPR120 Receptor in Essential Fatty Acids Metabolism in Schizophrenia. *Biomedicines* 2020, *8*, 243. https://doi.org/10.3390/biomedicines8080243.
- Dahoun, T.; Trossbach, S.V.; Brandon, N.J.; Korth, C.; Howes, O.D. The impact of Disrupted-in-Schizophrenia 1 (DISC1) on the dopaminergic system: A systematic review. *Transl. Psychiatry* 2017, 7, e1015. https://doi.org/10.1038/tp.2016.282.
- 366. Norkett, R.; Modi, S.; Birsa, N.; Atkin, T.A.; Ivankovic, D.; Pathania, M.; Trossbach, S.V.; Korth, C.; Hirst, W.D.; Kittler, J.T. DISC1-dependent Regulation of Mitochondrial Dynamics Controls the Morphogenesis of Complex Neuronal Dendrites. J. Biol. Chem. 2016, 291, 613–629. https://doi.org/10.1074/jbc.M115.699447.
- 367. Kvajo, M.; McKellar, H.; Drew, L.J.; Lepagnol-Bestel, A.M.; Xiao, L.; Levy, R.J.; Blazeski, R.; Arguello, P.A.; Lacefield, C.O.; Mason, C.A.; et al. Altered axonal targeting and short-term plasticity in the hippocampus of Disc1 mutant mice. *Proc. Natl. Acad. Sci. USA* 2011, 108, E1349–E1358. https://doi.org/10.1073/pnas.1114113108.
- 368. Niigaki, S.T.; Peres, F.F.; Ferreira, L.; Libanio, T.; Gouvea, D.A.; Levin, R.; Almeida, V.; Silva, N.D.; Diana, M.C.; Suiama, M.A.; Calzavara, M.B.; Abilio, V.C. Young spontaneously hypertensive rats (SHRs) display prodromal schizophrenia-like behavioral abnormalities. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 2019, 2(90), 169-176. https://doi.org/10.1016/j.pnpbp.2018.11.020.
- Park, C.; Park, S.K. Molecular links between mitochondrial dysfunctions and schizophrenia. *Mol. Cells* 2012, 33, 105–110. https://doi.org/10.1007/s10059-012-2284-3.
- Rajasekaran, A.; Venkatasubramanian, G.; Berk, M.; Debnath, M. Mitochondrial dysfunction in schizophrenia: Pathways, mechanisms and implications. *Neurosci. Biobehav. Rev.* 2015, 48, 10–21. https://doi.org/10.1016/j.neubiorev.2014.11.005.
- 371. Okusaga, O.; Fuchs, D.; Reeves, G.; Giegling, I.; Hartmann, A.M.; Konte, B.; Friedl, M.; Groer, M.; Cook, T.B.; Stearns-Yoder, K.A.; et al. Kynurenine and Tryptophan Levels in Patients with Schizophrenia and Elevated Antigliadin Immunoglobulin G Antibodies. *Psychosom. Med.* 2016, 78, 931–939. https://doi.org/10.1097/PSY.00000000000352.
- 372. Wang, A.K.; Miller, B.J. Meta-analysis of Cerebrospinal Fluid Cytokine and Tryptophan Catabolite Alterations in Psychiatric Patients: Comparisons Between Schizophrenia, Bipolar Disorder, and Depression. *Schizophr. Bull.* 2018, 44, 75–83. https://doi.org/10.1093/schbul/sbx035.
- 373. Faras, H.; Al Ateeqi, N.; Tidmarsh, L. Autism spectrum disorders. Ann. Saudi. Med. 2010, 30, 295–300. https://doi.org/10.4103/0256-4947.65261.
- 374. Candini, M.; Battaglia, S.; Benassi, M.; di Pellegrino, G.; Frassinetti, F. The physiological correlates of inter-personal space. *Sci. Rep.* 2021, *11*, 2611. https://doi.org/10.1038/s41598-021-82223-2.
- Rossignol, D.A.; Frye, R.E. Mitochondrial dysfunction in autism spectrum disorders: A systematic review and meta-analysis. *Mol. Psychiatry* 2012, 17, 290–314. https://doi.org/10.1038/mp.2010.136.
- 376. Thabault, M.; Turpin, V.; Maisterrena, A.; Jaber, M.; Egloff, M.; Galvan, L. Cerebellar and Striatal Implications in Autism Spectrum Disorders: From Clinical Observations to Animal Models. *Int. J. Mol. Sci.* 2022, 23, 2294. https://doi.org/10.3390/ijms23042294.
- Agarwala, S.; Ramachandra, N.B. Role of CNTNAP2 in autism manifestation outlines the regulation of signaling between neurons at the synapse. *Egypt. J. Med. Hum. Genet.* 2021, 22, 22. https://doi.org/10.1186/s43042-021-00138-z.
- 378. Yardeni, T.; Cristancho, A.G.; McCoy, A.J.; Schaefer, P.M.; McManus, M.J.; Marsh, E.D.; Wallace, D.C. An mtDNA mutant mouse demonstrates that mitochondrial deficiency can result in autism endophenotypes. *Proc. Natl. Acad. Sci. USA* 2021, 118, e2021429118. https://doi.org/10.1073/pnas.2021429118.
- 379. Lee, G.A.; Lin, Y.-K.; Lai, J.-H.; Lo, Y.-C.; Yang, Y.-C.S.H.; Ye, S.-Y.; Lee, C.-J.; Wang, C.-C.; Chiang, Y.-H.; Tseng, S.-H. Maternal Immune Activation Causes Social Behavior Deficits and Hypomyelination in Male Rat Offspring with an Autism-Like Microbiota Profile. *Brain Sci.* 2021, 11, 1085. https://doi.org/10.3390/brainsci11081085.
- 380. Abuaish, S.; Al-Otaibi, N.M.; Abujamel, T.S.; Alzahrani, S.A.; Alotaibi, S.M.; AlShawakir, Y.A.; Aabed, K.; El-Ansary, A. Fecal Transplant and Bifidobacterium Treatments Modulate Gut Clostridium Bacteria and Rescue Social Impairment and Hippocampal BDNF Expression in a Rodent Model of Autism. *Brain Sci.* 2021, 11, 1038. https://doi.org/10.3390/brainsci11081038.
- Frye, R.E.; Cakir, J.; Rose, S.; Delhey, L.; Bennuri, S.C.; Tippett, M.; Palmer, R.F.; Austin, C.; Curtin, P.; Arora, M. Early life metal exposure dysregulates cellular bioenergetics in children with regressive autism spectrum disorder. *Transl. Psychiatry* 2020, 10, 223. https://doi.org/10.1038/s41398-020-00905-3.
- Bryn, V.; Verkerk, R.; Skjeldal, O.H.; Saugstad, O.D.; Ormstad, H. Kynurenine Pathway in Autism Spectrum Disorders in Children. *Neuropsychobiology* 2017, 76, 82–88. https://doi.org/10.1159/000488157.
- Mahone, E.M.; Denckla, M.B. Attention-Deficit/Hyperactivity Disorder: A Historical Neuropsychological Perspective. J. Int. Neuropsychol. Soc. 2017, 23, 916–929. https://doi.org/10.1017/S1355617717000807.
- Kyaga, S.; Landén, M.; Boman, M.; Hultman, C.M.; Långström, N.; Lichtenstein, P. Mental illness, suicide and creativity: 40year prospective total population study. J. Psychiatr. Res. 2013, 47, 83–90. https://doi.org/10.1016/j.jpsychires.2012.09.010.

- Saccaro, L.F.; Schilliger, Z.; Perroud, N.; Piguet, C. Inflammation, Anxiety, and Stress in Attention-Deficit/Hyperactivity Disorder. *Biomedicines* 2021, 9, 1313. https://doi.org/10.3390/biomedicines9101313.
- 386. Murakami, Y.; Imamura, Y.; Saito, K.; Sakai, D.; Motoyama, J. Altered kynurenine pathway metabolites in a mouse model of human attention-deficit hyperactivity/autism spectrum disorders: A potential new biological diagnostic marker. *Sci. Rep.* 2019, 9, 13182. https://doi.org/10.1038/s41598-019-49781-y.
- 387. Aarsland, T.I.; Landaas, E.T.; Hegvik, T.A.; Ulvik, A.; Halmøy, A.; Ueland, P.M.; Haavik, J. Serum concentrations of kynurenines in adult patients with attention-deficit hyperactivity disorder (ADHD): A case-control study. *Behav. Brain Funct.* 2015, 11, 36. https://doi.org/10.1186/s12993-015-0080-x.
- 388. Picard, M.; McEwen, B.S. Psychological Stress and Mitochondria: A Systematic Review. *Psychosom. Med.* 2018, *80*, 141–153. https://doi.org/10.1097/PSY.00000000000545.
- Jiang, S.; Postovit, L.; Cattaneo, A.; Binder, E.B.; Aitchison, K.J. Epigenetic Modifications in Stress Response Genes Associated with Childhood Trauma. *Front. Psychiatry* 2019, 10, 808. https://doi.org/10.3389/fpsyt.2019.00808.
- Kim, J.I.; Lee, S.Y.; Park, M.; Kim, S.Y.; Kim, J.W.; Kim, S.A.; Kim, B.N. Peripheral Mitochondrial DNA Copy Number is Increased in Korean Attention-Deficit Hyperactivity Disorder Patients. *Front. Psychiatr.* 2019, 10, 506. https://doi.org/10.3389/fpsyt.2019.00506.