

Article

Açai Berry Mitigates Vascular Dementia-Induced Neuropathological Alterations Modulating Nrf-2/Beclin1 Pathways



Daniela Impellizzeri ^{1,†}, Ramona D'Amico ^{1,†}, Roberta Fusco ¹, Tiziana Genovese ¹, Alessio Filippo Peritore ¹, Enrico Gugliandolo ², Rosalia Crupi ², Livia Interdonato ¹, Davide Di Paola ¹, Rosanna Di Paola ^{2,*}, Salvatore Cuzzocrea ^{1,3,*}, Rosalba Siracusa ^{1,‡} and Marika Cordaro ^{4,‡}

- ¹ Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Viale Ferdinando Stagno D'Alcontres 31, 98166 Messina, Italy
- ² Department of Veterinary Sciences, University of Messina, 98168 Messina, Italy
- ³ Department of Pharmacological and Physiological Science, Saint Louis University School of Medicine, Saint Louis, MO 63104, USA
- Department of Biomedical, Dental and Morphological and Functional Imaging, University of Messina,
 Via Consolare Valeria, 98125 Messina, Italy
- * Correspondence: dipaolar@unime.it (R.D.P.); salvator@unime.it (S.C.)
- + These authors equally contributed to the work.
- ‡ These authors shared senior authorship.

Abstract: The second-most common cause of dementia is vascular dementia (VaD). The majority of VaD patients experience cognitive impairment, which is brought on by oxidative stress and changes in autophagic function, which ultimately result in neuronal impairment and death. In this study, we examine a novel method for reversing VaD-induced changes brought on by açai berry supplementation in a VaD mouse model. The purpose of this study was to examine the impact of açai berries on the molecular mechanisms underlying VaD in a mouse model of the disease that was created by repeated ischemia–reperfusion (IR) of the whole bilateral carotid artery. Here, we found that açai berry was able to reduce VaD-induced behavioral alteration, as well as hippocampal death, in CA1 and CA3 regions. These effects are probably due to the modulation of nuclear factor erythroid 2-related factor 2 (Nrf-2) and Beclin-1, suggesting a possible crosstalk between these molecular pathways. In conclusion, the protective effects of açai berry could be a good supplementation in the future for the management of vascular dementia.

Keywords: vascular dementia; oxidative stress; autophagy; açai berry

1. Introduction

In the Western world, vascular dementia (VaD) is the second-most common cause of organic acquired cognitive dysfunction; in several Asian nations, it is likely the main cause [1].

There are no viable pharmaceuticals approved for the treatment of VaD in any country despite the recent investments made in experimental and clinical neuroscience [2]. Smoking, not exercising, eating poorly, and being exposed to pollutants are all known to worsen the neuronal environment and increase oxidative stress, which can lead to the cognitive loss associated with aging and neurodegenerative disorders. Important brain areas like the hippocampus should be more vulnerable to oxidative and inflammatory stresses since these conditions might impair synaptic plasticity and memory by causing dendritic alteration and neuronal death. Furthermore, the brain is also subjected to an altered expression of two important neuronal markers of wellbeing: MAP-2 and β -Tubulin.

Citation: Impellizzeri, D.; D'Amico, R.; Fusco, R.; Genovese, T.; Peritore, A.F.; Gugliandolo, E.; Crupi, R.; Interdonato, L.; Di Paola, D.; Di Paola, R.; et al. Açai Berry Mitigates Vascular Dementia-Induced Neuropathological Alterations Modulating Nrf-2/Beclin1 Pathways. *Cells* **2022**, *11*, 2616. https://doi.org/10.3390/cells 11162616

Academic Editors: Alessia Remigante and Rossana Morabito

Received: 28 July 2022 Accepted: 20 August 2022 Published: 22 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/license s/by/4.0/). As it promotes microtubule stiffness, MAP2—which is found in cell bodies and dendrites—is regarded as a microtubule stabilizer. When expression is changed, MAP2 builds up into granules, causing neurotoxicity and neuronal degeneration. During cortical development, the tubulin family primarily expresses itself in post-mitotic neurons with a particular geographic and temporal expression pattern. Microtubules' primary building blocks, tubulin, are essential for the mechanisms of the central nervous system's (CNS) development, including neuronal migration and axonal guidance. The altered expression of these proteins may result in the death of neurons. The loss of MAP2 and β -Tubulin have been strongly associated with VaD [3,4].

Additionally, defective autophagy has been linked to increased oxidative stress in the brain, altering protein "quality control," accumulating undesired proteins and organelles in brain cells [5].

A fundamental leucine zipper redox-sensitive transcription factor called nuclear factor erythroid 2-related factor 2 (Nrf2) regulates the redox status of the cell under damaging stressors. Under normal circumstances, Kelch-like ECH-associated protein 1(Keap1), Nrf2's inhibitor, anchors Nrf2 in the cytoplasm. However, during oxidative stress, Nrf2 separates from Keap1, moves into the nucleus, and interacts with the antioxidant response element (ARE) to control the production of antioxidant genes such heme oxygenase-1 (HO-1). Additionally, recent studies have shown that Nrf2 could control the initiation of autophagy, furthering its protective benefits [6–8]. In fact, inclusion bodies, which are pathogenic markers of many neurodegenerative disorders, are known to accumulate excessively as a result of dysfunctional autophagy. Numerous studies have demonstrated that entire fruits and dietary polyphenols such as anthocyanins, stilbenes, melatonin, kaemferol, quercetin, and resveratrol protect neurons by a modulation of autophagy process [9–20].

Scientists have recently developed an interest in açai seeds. The berry known as the açai berry has a variety of beneficial nutritional properties as well as some possible medicinal uses. This fruit has a tart flavor and is produced by the *Euterpe oleracea* palm, which can only be found in the Amazon. For millennia, Amazonian Indians have employed the açai fruit, which is regarded as a high-energy meal, as a food source and a natural remedy for a number of illnesses [21–29].

The pulp of the açai fruit has been extensively researched due to its high bioactive nutritional and phytochemical content. Açai berry pulp composition contains a wide range of physiologically active phytochemicals as well as significant amounts of monoand polyunsaturated fatty acids, which are uncommon in most fruits and other berries. Phytochemicals such proanthocyanidins, anthocyanins, and other flavonoids are present in açai pulp. Additionally, phytochemical analyses revealed that the açai berry contains a significant amount of luteolin, quercetin, dihydrokaempferol, and chrysoerial, among other polyphenolics, as well as a variety of anthocyanins, including cyanidin, delphinidin, malvidin, pelargonidin, and peonidin. Açai fruit pulp included five different types of carotenoids, including carotene, lycopene, astaxanthin, lutein, and zeaxanthin. [30].

Numerous pharmacological benefits of açai berry extract and its bioactive components include anti-inflammatory and anti-anxiety properties through modulation of oxidative stress, inflammation, autophagy, and Nrf2 expression in the hippocampus and frontal cortex [31–39]. However, more data are needed to confirm the neuroprotective, anti-inflammatory and antioxidant effect. For this reason, we used a consolidated murine model of VaD to investigate the potential beneficial effects of açai supplementation and the molecular way by which it acts.

2. Materials and Methods

2.1. Animals

CD1 male mice (8-week-old, 18–24 g) were acquired from Envigo (Milan, Italy) and located in a controlled environment. All animal experiments complied with the new Italian regulations (D. Lgs 2014/26), EU regulations (EU Directive 2010/63) and the ARRIVE guidelines.

2.2. Experimental Design and Groups

Mice were subjected to temporary bilateral carotid occlusion as previously described by Cordaro et al. [40]. Briefly, following anesthesia, the bilateral carotid arteries underwent three cycles of ligation and release lasting a total of 10 min each. After the threading was taken out, the incision was stitched up. After 15 days of induction, the mice were sacrificed, and their brains were removed and processed for histological and biochemical analysis [40,41].

Mice were arbitrarily divided into groups:

Group 1: Sham = mice were subjected to the surgery without carotid arteries ligation and treated daily orally with saline (vehicle) for 15 days.

Group 2: Sham + açai berry = same as above, but açai berry at a dose of 500 mg/kg, dissolved in saline, was administered orally once a day for 15 days (data not shown).

Group 3: VaD = mice were subjected to the VaD surgery described above and treated with vehicle.

Group 4: VaD + açai berry = same as above, but açai berry at a dose of 500 mg/kg was administered daily orally for 15 days.

A total of six samples was used for each technique. Freeze-dried açai extract was dissolved in distilled water. This substance (Cas Number: 879496-95, 906351-38-0) was purchased from Farmalabor Srl (Canosa di Puglia, Barletta, Italy) (see Supplementary File for technical datasheet). The dose and the route of administration of açai berry were chosen based on our previous study [42]. No significant difference was found between the sham and sham + açai berry, so only data regarding the sham groups are shown.

2.3. Behavioral Tests

2.3.1. Open Field Test (OFT)

Locomotor activity and anxiety-like behavior were monitored by the OFT. Each mouse was carefully placed in the center of the box after training, and any action was counted as a line crossing if a mouse moved all four paws out of one square and into another [43,44].

2.3.2. Novel Object Recognition (NOR) Test

The spontaneous inclination of mice to spend time investigating a novel object or a familiar one was examined with the NOR test. After a training session, mice were switched out in the box for a five-minute exam, during which the examiner, at random, switched out one of the familiar objects for a new one. The amount of time the mouse spent investigating each item was measured [41,45].

2.3.3. Social Interaction Test (SIT)

A three-chambered device was used to create the social interaction test as previously described. The number of active interactions and the duration of contacts were measured [41].

2.4. Histopathological Evaluation

Brains were stained with hematoxylin/eosin (H/E) or cresyl-violet and blindly analyzed using light microscopy LeicaDM6 connected to an imaging system (LasX

Navigator)(Leica Microsystem, Buccinasco, Italy) by two investigators without knowledge of the experimental groups [46–48]. The severity of brain injury and numbers of dead cells was assessed as previously described [40,41].

2.5. Immunohistochemical Evaluation

Brain sections were incubated with anti-NRF2 antibody (1:250, Santa Cruz Biotechnology) as previously described by Cordaro et al. [49,50]. At the end of the protocol, the digital photos were analyzed by two observers blinded to the treatment as previously made in our laboratory [51–55].

2.6. Terminal Deoxynucleotidyl Nick-End Labeling (TUNEL) Assay

TUNEL staining for apoptotic cell nuclei and DAPI staining for all cell nuclei were performed in brain sections as described previously [56–59].

2.7. Immunofluorescence Evaluation

Brain sections were incubated with the following primary antibodies: HO-1 (Santa Cruz Biotechnology; 1:50 in PBS, v/v) or β -Tubulin (Santa Cruz Biotechnology Heidelberg Germany; 1:50) or MAP-2 (Santa Cruz Biotechnology; 1:50), as previously described [60–65]. After the incubation, sections were washed with PBS and incubated with secondary antibody FITC-conjugated anti-mouse Alexa Fluor-488 (1:2000 v/v Molecular Probes, UK) and 40,60-diamidino-2-phenylindole (DAPI; Hoechst, Frankfurt; Germany). Sections were observed and photographed using a Leica DM6 microscope (Leica Microsystems SpA, Milan, Italy)[66–70].

2.8. RT-qPCR

Total RNA was isolated from the hippocampus using TRIzol Reagent (Invitrogen, USA) following the manufacturer's recommendations. Reverse transcription of cDNA was performed according to the manufacturer's instructions as previously described [71]. sequences The (5'-3')following primer were used: Beclin-1 (F) GCTGTAGCCAGCCTCTGAAA AATGGCTCCTGTGAGTTCCTG; (R) LC3B (F) GGGACCCTAACCCCATAGGA TCTCCCCCTTGTATCGCTCT; (R) P62 (F) ACTGCTCAGGAGGAGACGAT (R) CCGGGGATCAGCCTCTGTAG; (F) Bcl-2 GCGTCAACAGGGAGATGTCA (R) GCATGCTGGGGGCCATATAGT; Bax (F) CTGGATCCAAGACCAGGGTG (R) GTGAGGACTCCAGCCACAAA β-Actin (F) ACACTCTCCCAGAAGGAGGG (R) TTTATAGGACGCCACAGCGG. Relative expression of mRNA was calculated by the delta-delta CT method [72,73]. All values for the mRNA species were normalized to β -actin.

2.9. Western Blot Analysis of Cytosolic and Nuclear Extracts

Extracts from cytosol and nucleus were prepared as previously described and incubated with the antibodies anti-NRF-2 (1–500, SCB, Heidelberg, Germany, #sc-365949), anti-heme oxygenase 1 (HO-1; 1-500, SCB, Heidelberg, Germany, #sc-136960), anti- β -tubulin (1–500, SCB, Heidelberg, Germany, #sc-166729), and anti-MAP2 (1–500, SCB, Heidelberg, Germany, #sc-74421) in 1× PBS, 5% w/v non-fat dried milk and 0.1% Tween-20 at 4 °C overnight [46–48,74–83]. For the cytosolic and nuclear fraction, blots were also probed with β -actin and lamin A/C protein to ensure that they were filled with equivalent amounts of proteins (1:500; Santa Cruz Biotechnology). Signals were detected as previously described in our works [49,84–86].

2.10. Materials

All compounds were purchased from Sigma-Aldrich (Milan, Italy). All solutions used for in vivo infusions were prepared using nonpyrogenic saline (0.9% NaCl; Baxter Healthcare Ltd., Thetford, Norfolk, UK).

2.11. Statistical Evaluation

In this study, the data are expressed as the average ± SEM and represent at least three experiments carried out on different days. For in vivo studies, N represents the number of animals used. The number of animals used for in vivo studies was determined by G*Power 3.1 software (Die Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany). The images used in the histology, immunofluorescence, and immunohistochemistry came from at least three independent investigations. For multiple comparisons, a one-way ANOVA was employed, then a Bonferroni post hoc analysis; 0.05 was a significant *p*-value.

3. Results

3.1. Açai Berry Improve Behavioral Changes Vascular Dementia-Induced

We observed that after the unfamiliar mouse was placed in the three-chamber test arena, the number of contacts (Figure 1A) considerably increased while the total duration of contacts (Figure 1B) decreased compared to the control group. On the other hand, after oral açai administration for 15 days at the dose of 500 mg/kg, we noticed that the behavior was more similar to the control group than the vehicle group. Additionally, we assessed alterations in cognitive function using the novel object recognition test (Figure 1C,D). Controls and VaD animals did not significantly differ in their exploration of items during training; however, 15 days after carotid artery ligation, VaD animals dramatically decreased their preference for the novel object, indicating compromised cognitive function. However, after taking açai orally for 15 days at a dose of 500 mg/kg, we found that the behavior was more like the control group than the vehicle group.



Figure 1. Açai berry improves vascular-dementia-induced behavioral changes. (**A**) Number of contacts and (**B**) total contact duration measured during social interaction test; (**C**) time spent to explore familiar or novel object, and (**D**) total exploration time measured during novel object recognition test; (**E**) number of crossing and (**F**) time in the center measured during open field test. Values are means ± SEM of six mice for all groups. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD; ## *p* < 0.01 vs. VaD.

3.2. Açai Berry Limits Histological Changes Vascular Dementia-Induced

At 15 days after the induction of vascular dementia, all slices were stained with H&E to observe the severity of the hippocampal areas' damage. As showed in Figure 2B (for CA1 region) and 2E (for CA3 region), the hippocampal regions of the animals that had suffered VaD damage exhibited disorderly and rigidly labeled neurons. After daily oral administration of açai, brain sections showed a clear reorganization of the hippocampal CA1 and CA3 regions with an increased number of hippocampal neurons (Figure 2C for CA1 region, 2F for CA3 region). No alteration was observed in sham animals (Figure 2A for CA1 region, 2D for CA3 region; see histological score G).



Figure 2. Açai berry limits vascular-dementia-induced histological changes. Hippocampal CA1 figure of Sham (**A**), VaD (**B**), Açai Berry (**C**); Hippocampal CA3 figure of Sham (**D**), VaD (**E**), Açai Berry (**F**); Histological score (**G**). Arrows indicate altered neurons and loss of hippocampal architecture. Figures are representative from least three experiments. Values are means ± SEM of six mice for all groups. See manuscript for further details. *** p < 0.001 vs. sham; ### p < 0.001 vs. VaD.

3.3. Açai Berry Limits Neuronal Death in the Hippocampus

Next, we investigated whether açai berry could attenuate neuronal death in the hippocampal structure, which is an essential feature in VaD patients [87]. For this, we used cresil violet and TUNEL staining to assess the effects of açai berry administration on the loss of pyramidal neurons in the hippocampus, especially in the CA1 and CA3 subregion. As showed in Figure 3B,E and in Figure 4B,E, the vehicle group was characterized by an intense neuronal loss due to the death of pyramidal neurons in both CA1 and CA3 regions compared to the control group, Figure 3A,D and in Figure 4A,D. On the other hand, daily administration of açai berry at the dose of 500 mg/kg was able to significantly reduce neuronal death in both regions (Figure 3C,F and Figure 4C,F; see relative score, Figures 3G and 4G).



Figure 3. Açai berry limits neuronal hippocampal deaths. Hippocampal CA1 figure of Sham (A), VaD (B), Açai Berry (C); Hippocampal CA3 figure of Sham (D), VaD (E), Açai Berry (F); count of dead cells (G). Arrows indicate death neurons in both hippocampal sections. Figures are representative from at least three experiments. Values are means ± SEM of six mice for all groups. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.



Hippocampal CA1

Figure 4. Açai berry reduces hippocampal TUNEL positive staining. Hippocampal CA1 figure of Sham (**A**,**A1**,**A2**), VaD (**B**,**B1**,**B2**), Açai Berry (**C**,**C1**,**C2**); Hippocampal CA3 figure of Sham (**D**,**D1**,**D2**), VaD (**E**,**E1**,**E2**), Açai Berry (**F**,**F1**,**F2**); count of dead cells (**G**). Arrows indicate TUNEL positive cells, merged with DAPI. Figures are representative from at least three experiments. Values are means ± SEM of six mice for all groups. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.

3.4. Açai Berry Modulates Nrf-2 Pathways

In our work, by immunohistochemistry staining and western blot analysis, we found an increase in NRF-2 expression after VaD induction (see Figure 5B for CA1, Figure 5E for CA3, Figure 5G for quantification, and Figure 5H,I for western blot) compared to the control group (see Figure 5A for CA1, Figure 5D for CA3, Figure 5G for quantification, and Figure 5H,I for western blot). As expected, after açai berry administration, the physiological response was significant (see Figure 5C for CA1, Figure 5F for CA3, Figure 6B,B1,B2 for CA1 region and Figure 6E,E1,E2 for CA3 region of VaD group, Figure 6C,C1,C2 for CA1 region and Figure 6F,F1,F2 for CA3 region of Açai Berry group, and

Figure 6H,I for western blot).

CA1 CA3 Sham B VaD F С VaD+Açai G Vad + Acai 500 Nrf-2 61 kDa н CA1 CA3 I Sham Sha A/0 200 VaD W// VaD VaD+Aça I Jumber of Nrf-2

Figure 5. Açai berry improve NRF-2 expression. Hippocampal CA1 figure of Sham (**A**), VaD (**B**), Açai Berry (**C**); Hippocampal CA3 figure of Sham (**D**), VaD (**E**), Açai Berry (**F**); Number of Nrf-2 positive pixel (**G**); (**H**) western blot of NRF-2 expression and (**I**) relative densitometric analysis. Arrows indicate Nrf-2 positive cells. Figures are representative from at least three experiments. Values are means ± SEM of six mice for all groups. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.



Hippocampal CA1

Figure 6. Açai berry increase HO-1 expression. Hippocampal CA1 figure of Sham (**A**,**A1**,**A2**), VaD (**B**,**B1**,**B2**), Açai Berry (**C**,**C1**,**C2**); Hippocampal CA3 figure of Sham (**D**,**D1**,**D2**), VaD (**E**,**E1**,**E2**), Açai Berry (**F**,**F1**,**F2**); (**G**) western blot of HO-1 and (**H**) relative densitometric analysis. Arrows indicate HO-1-DAPI merged cells in both hippocampal sections. Values are means ± SEM of six mice for all groups. Figures are representative from at least three experiments. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.

3.5. Açai Berry Modulates Apoptotic and Autophagic Pathways

To further investigated neuronal death, we made RT-qPCR for Bax, and Bcl-2 and we found that after VaD induction there was a significant increase in Bax expression and a significant decrease in Bcl-2 expression compared to sham group (Figure 7A,B). Vice versa, the daily oral administration of açai berry at a dose of 500 mg/kg was able to restore both apoptotic markers to almost physiological values (Figure 7A,B). Autophagy and

apoptosis interact in a complex way because, depending on the biological context, it can either promote cell survival or cell death [88]. In our study, açai berry supplementation increased autophagy in the hippocampus by activating Beclin1 (Figure 7C) decreasing LC3B turnover (Figure 7D), which led also to a decrease in p62 accumulation (Figure 7E).





3.6. Açai Berry Mitigates Brain Structure Change VaD-Induced

It is well-known that reduced levels of tubulin were found in patients with VaD [3]. In our study, we detected by immunofluorescence a significant decrease of b-tubulin in animals subjected to the operation compared to Sham group in both hippocampal regions. This decrease was significantly reduced after daily administration of açai berry at the dose of 500 mg/kg (see Figure 8A,A1,A2 for CA1 region and Figure 8D,D1,D2 for CA3 region of Sham group, Figure 8B,B1,B2 for CA1 region and Figure 8E,E1,E2 for CA3 region of VaD group, Figure 8C,C1,C2 for CA1 region and Figure 8F,F1,F2 for CA3 region of Açai Berry group). The same result was also observed with the immunofluorescence analysis of MAP-2. In this case as well, we observed a significant reduction of MAP-2 expression after VaD induction compared to the control group, which was significant restored after açai berry administration (see Figure 9A,A1,A2 for CA1 region and Figure 9D,D1,D2 for

13 of 21

CA3 region of sham group, Figure 9B,B1,B2 for CA1 region and Figure 9E,E1,E2 for CA3 region of VaD group, Figure 9C,C1,C2 for CA1 region and Figure 9F,F1,F2 for CA3 region of Açai Berry group). These results were also confirmed by western blots analysis of β -tubulin and MAP-2 (see Figure 8G,H and Figure 9G,H, respectively).



Figure 8. Açai berry mitigates β-tubulin alteration. Hippocampal CA1 figure of Sham (**A**,**A1**,**A2**), VaD (**B**,**B1**,**B2**), Açai Berry (**C**,**C1**,**C2**); Hippocampal CA3 figure of Sham (**D**,**D1**,**D2**), VaD (**E**,**E1**,**E2**), Açai Berry (**F**,**F1**,**F2**); β-tubulin western blot (**G**) and relative densitometric analysis (**H**). Arrows indicate the alteration or absence of β-tubulin. Values are means ± SEM of six mice for all groups. Figures are representative from at least three experiments. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.



Figure 9. Açai berry mitigates MAP-2 alteration. Hippocampal CA1 figure of Sham (**A**,**A1**,**A2**), VaD (**B**,**B1**,**B2**), Açai Berry (**C**,**C1**,**C2**); Hippocampal CA3 figure of Sham (**D**,**D1**,**D2**), VaD (**E**,**E1**,**E2**), Açai Berry (**F**,**F1**,**F2**); MAP-2 western blot (**G**), and relative densitometric analysis (**H**) Arrows indicate the alteration or absence of MAP-2. Values are means ± SEM of 6 mice for all groups. Figures are representative from at least three experiments. See manuscript for further details. *** *p* < 0.001 vs. sham; ### *p* < 0.001 vs. VaD.

4. Discussion

The pathologic disease known as VaD, which affects the elderly, is characterized by disturbed cerebral blood flow. Loss of judgment, reason, and especially cognitive and memory abilities are its symptoms [89]. Advanced age, previous stroke episodes, hypertension, higher blood pressure, diabetes mellitus, smoking, high hematocrit, changes in hemostasis, dyslipoproteinemia, high alcohol consumption, aspirin use, psychological stress in early life, occupational exposure to pesticides, herbicides, liquid

plastic or rubber, and premorbid personality traits made the subjects more susceptible to developing VaD [90]. Neuronal degeneration, axonal injury, and white matter injury in VaD are caused by a number of processes. They frequently share vascular dysfunctioninduced common molecular and regulatory pathways of neuronal survival and death. Of these, oxidative stress, apoptosis, and autophagy pathways typically demonstrate different processes of their own. Along with age and other risk factors acting in concert, a crosstalk between these signaling pathways contributes to myelinated axonal damage and cerebrovascular diseases. The diseased characteristics cause changes in the health of the neuronal population as well as a series of harmful events and white matter destruction. Cognitive impairment and brain atrophy are the results of these diseased traits.

Although numerous papers describe the pathologic mechanisms behind vascular dementia, it is uncertain how the signaling pathways involved in vascular neuropathologies and cerebrovascular dysfunction interact with one another [91]. Accumulated data from numerous studies show that supplementing the diet with fruits, nuts, and vegetables of vibrant colors enhances the motor, memory, and cognitive abilities of both people and animals [5,66–70]. The antioxidant defense system may be compromised by increased oxidative stress brought on by a poor diet. On the other hand, a nutritious diet that is well-balanced and rich in a variety of nutrients, such as numerous servings of fruits and vegetables, low dosages of omega-3 fatty acids, tea, coffee, and wine, may provide neuroprotection [92,93].

The new food that is discussed in this paper is a fruit from the Euterpe genus of tropical palm trees that is native to South America and is commonly referred to as "Acai." Researchers have been investigating *Euterpe oleracea* due to its high antioxidant content when compared to other fruits and berries. The composition of açai pulp has also been studied, and it contains a number of phytochemicals that have physiological activity. Numerous studies have demonstrated the neuroprotective properties of açai berries [30]. The causes of many of these diseases are multifactorial, including oxidative stress, chronic neuroinflammation, excitotoxicity, mitochondrial dysfunction, alteration in autophagy, abnormal protein accumulation in brain tissues, and other cellular aetiologias. These diseases are brought on by a combination of aging, genetic disorders, and exposure to one or more environmental factors. Açai berry extracts have been shown in studies to have neuroprotective effects by, among other things, restoring calcium homeostasis and mitochondrial function, preventing the formation of toxic protein aggregates, and demonstrating antioxidant and anti-inflammatory activities. Additionally, açai fruit has anticonvulsant and antidepressant effects that may be beneficial for those with certain neurological conditions [94–101].

Keeping this aim in our mind, we decided to explore the neuroprotective effects of açai berry administration in a murine model of VaD. Depressive, anxious, or apathetic mood swings are common in people with vascular dementia. A person with vascular dementia may start acting strangely as their condition worsens. For instance, they might become angrier, hostile, or perplexed [102,103]. In our study, using different behavioral tests, we found that açai supplementation was able to reduce VaD-induced behavioral decline. It is well-known that a strong relationship exists between cognitive impairment and neuronal loss in the hippocampus. For this, we investigated by different staining what happen in CA1 and CA3. In particular, we found that both hippocampal sections showed, after VaD induction, a significant increase in altered and dead neurons, compared to Sham animals. After açai berry supplementation as well as a decrease in dead neurons in both hippocampal regions.

Nrf2 is the master regulator of redox status and controls the transcription of a panel of antioxidative and anti-inflammatory genes. By cooperating with NF- κ B (nuclear factor κ B), Nrf2 also coordinates cellular oxidative and inflammatory balance. Nrf2 deficiency is linked to aging, and mounting evidence supports Nrf2's role in slowing the VaD process. By reducing neuroinflammation and oxidative stress, targeting Nrf2 has become a popular strategy for treating neurodegenerative diseases [104–106]. In our study, we found that açai berry supplementation was able to improve physiological anti-oxidant defense, improving Nrf-2 expression as well as HO-1, reducing oxidative stress. Additionally, recent studies suggest that autophagy and the Kelch-like ECH-associated protein 1 (Keap1-Nrf2) signaling pathway cooperate to avoid oxidative injury in the brain, and that p62 phosphorylation activates Nrf2 in response to oxidative stress. This is particularly intriguing because recent articles have demonstrated dynamic Beclin-1 alterations in the hippocampus of male mice with VaD. Beclin-1 is a crucial regulator of several trafficking pathways, including autophagy and receptor recycling in microglia. [71,107]. Additionally, it has also been demonstrated that Beclin 1 regulates signaling through the neuroprotective TGF- β pathway by decreasing TGF- β 1 receptor II (TBRII). TGF- β 1 plays a key function in hippocampus synaptic plasticity, memory, and neuronal survival. It also regulates brain homeostasis [108–110]. In our study, we found that acai berry restored autophagic flux at physiological levels. Interesting studies have shown that the altered expression of MAP-2 and β -tubulin leads to neuronal death; such results could be used in the future target for neuroprotective therapy targeting [111–114]. In our work, we found that açai berry stops the process of cell death and neurodegeneration and restores the microtubule network.

5. Conclusions

In conclusion, we demonstrate for the first time that açai berry supplementation at a dose of 500 mg/kg was able to enhance the physiological anti-oxidant defense, suggesting a possible protective role during VaD events.

Supplementary Materials: The following are available online at www.mdpi.com/article/10.3390/cells11162616/s1, Supplementary File: Technical Datasheet of EUTERPE OLERACEA P.E. 10% POLYPHENOLS.

Author Contributions: Conceptualization, R.S. and M.C.; Formal analysis, L.I. and D.D.P.; Funding acquisition, R.D.P. and S.C.; Methodology, D.I., R.D. and E.G.; Software, R.F. and T.G.; Supervision, R.D.P.; Validation, A.F.P. and R.C.; Writing—original draft, R.S.; Writing—review & editing, M.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The animal study protocol was approved by the Institutional Review Board of University of Messina.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Orgogozo, J.M.; Abadie, E. Vascular dementia: European perspectives. *Alzheimer Dis. Assoc. Disord.* 1999, 13 (Suppl. 3), S192– S200. https://doi.org/10.1097/00002093-199912003-00027.
- Baskys, A.; Hou, A.C. Vascular dementia: Pharmacological treatment approaches and perspectives. *Clin. Interv. Aging* 2007, 2, 327.
- Santiago-Mujika, E.; Luthi-Carter, R.; Giorgini, F.; Kalaria, R.N.; Mukaetova-Ladinska, E.B. Tubulin and Tubulin Posttranslational Modifications in Alzheimer's Disease and Vascular Dementia. *Front. Aging Neurosci.* 2021, 13, 730107. https://doi.org/10.3389/fnagi.2021.730107.
- Sabbatini, M.; Catalani, A.; Consoli, C.; Marletta, N.; Tomassoni, D.; Avola, R. The hippocampus in spontaneously hypertensive rats: An animal model of vascular dementia? *Mech. Ageing Dev.* 2002, 123, 547–559. https://doi.org/10.1016/s0047-6374(01)00362-1.
- Poulose, S.M.; Bielinski, D.F.; Carey, A.; Schauss, A.G.; Shukitt-Hale, B. Modulation of oxidative stress, inflammation, autophagy and expression of Nrf2 in hippocampus and frontal cortex of rats fed with acai-enriched diets. *Nutr. Neurosci.* 2017, 20, 305–315. https://doi.org/10.1080/1028415X.2015.1125654.

- Jiang, T.; Harder, B.; Rojo de la Vega, M.; Wong, P.K.; Chapman, E.; Zhang, D.D. p62 links autophagy and Nrf2 signaling. *Free Radic. Biol. Med.* 2015, 88, 199–204. https://doi.org/10.1016/j.freeradbiomed.2015.06.014.
- Jo, C.; Gundemir, S.; Pritchard, S.; Jin, Y.N.; Rahman, I.; Johnson, G.V. Nrf2 reduces levels of phosphorylated tau protein by inducing autophagy adaptor protein NDP52. *Nat. Commun.* 2014, *5*, 3496. https://doi.org/10.1038/ncomms4496.
- Shah, S.Z.A.; Zhao, D.; Hussain, T.; Sabir, N.; Mangi, M.H.; Yang, L. p62-Keap1-NRF2-ARE Pathway: A Contentious Player for Selective Targeting of Autophagy, Oxidative Stress and Mitochondrial Dysfunction in Prion Diseases. *Front. Mol. Neurosci.* 2018, 11, 310. https://doi.org/10.3389/fnmol.2018.00310.
- 9. Zhang, L.; Wang, H.; Fan, Y.; Gao, Y.; Li, X.; Hu, Z.; Ding, K.; Wang, Y.; Wang, X. Fucoxanthin provides neuroprotection in models of traumatic brain injury via the Nrf2-ARE and Nrf2-autophagy pathways. *Sci. Rep.* **2017**, *7*, 46763. https://doi.org/10.1038/srep46763.
- Xiao, X.T.; He, S.Q.; Wu, N.N.; Lin, X.C.; Zhao, J.; Tian, C. Green Tea Polyphenols Prevent Early Vascular Aging Induced by High-Fat Diet via Promoting Autophagy in Young Adult Rats. *Curr. Med. Sci.* 2022, *Epub ahead of print*. https://doi.org/10.1007/s11596-022-2604-6.
- 11. Arias, C.; Salazar, L.A. Autophagy and Polyphenols in Osteoarthritis: A Focus on Epigenetic Regulation. *Int. J. Mol. Sci.* 2021, 23, 421. https://doi.org/10.3390/ijms23010421.
- Brimson, J.M.; Prasanth, M.I.; Malar, D.S.; Thitilertdecha, P.; Kabra, A.; Tencomnao, T.; Prasansuklab, A. Plant Polyphenols for Aging Health: Implication from Their Autophagy Modulating Properties in Age-Associated Diseases. *Pharmaceuticals* 2021, 14, 982. https://doi.org/10.3390/ph14100982.
- 13. Garcia-Aguilar, A.; Palomino, O.; Benito, M.; Guillen, C. Dietary Polyphenols in Metabolic and Neurodegenerative Diseases: Molecular Targets in Autophagy and Biological Effects. *Antioxidants* **2021**, *10*, 142. https://doi.org/10.3390/antiox10020142.
- 14. Musial, C.; Siedlecka-Kroplewska, K.; Kmiec, Z.; Gorska-Ponikowska, M. Modulation of Autophagy in Cancer Cells by Dietary Polyphenols. *Antioxidants* **2021**, *10*, 123. https://doi.org/10.3390/antiox10010123.
- 15. Abbaszadeh, F.; Fakhri, S.; Khan, H. Targeting apoptosis and autophagy following spinal cord injury: Therapeutic approaches to polyphenols and candidate phytochemicals. *Pharmacol. Res.* **2020**, *160*, 105069. https://doi.org/10.1016/j.phrs.2020.105069.
- Wu, M.; Luo, Q.; Nie, R.; Yang, X.; Tang, Z.; Chen, H. Potential implications of polyphenols on aging considering oxidative stress, inflammation, autophagy, and gut microbiota. *Crit. Rev. Food Sci. Nutr.* 2021, 61, 2175–2193. https://doi.org/10.1080/10408398.2020.1773390.
- 17. Zhou, H.; Chen, Y.; Huang, S.W.; Hu, P.F.; Tang, L.J. Regulation of autophagy by tea polyphenols in diabetic cardiomyopathy. *J. Zhejiang Univ. Sci. B* 2018, *19*, 333–341. https://doi.org/10.1631/jzus.B1700415.
- Shabalala, S.; Muller, C.J.F.; Louw, J.; Johnson, R. Polyphenols, autophagy and doxorubicin-induced cardiotoxicity. *Life Sci.* 2017, 180, 160–170. https://doi.org/10.1016/j.lfs.2017.05.003.
- Zhang, P.W.; Tian, C.; Xu, F.Y.; Chen, Z.; Burnside, R.; Yi, W.J.; Xiang, S.Y.; Xie, X.; Wu, N.N.; Yang, H.; et al. Green Tea Polyphenols Alleviate Autophagy Inhibition Induced by High Glucose in Endothelial Cells. *Biomed. Environ. Sci.* 2016, 29, 524– 528. https://doi.org/10.3967/bes2016.069.
- 20. Pallauf, K.; Rimbach, G. Autophagy, polyphenols and healthy ageing. *Ageing Res. Rev.* 2013, 12, 237–252. https://doi.org/10.1016/j.arr.2012.03.008.
- Maciel-Silva, F.W.; Buller, L.S.; MBB Gonçalves, M.L.; Rostagno, M.A.; Forster-Carneiro, T. Sustainable development in the Legal Amazon: Energy recovery from açaí seeds. *Biofuels Bioprod. Biorefin.* 2021, 15, 1174–1189.
- Melo, P.S.; Massarioli, A.P.; Lazarini, J.G.; Soares, J.C.; Franchin, M.; Rosalen, P.L.; Alencar, S.M. Simulated gastrointestinal digestion of Brazilian acai seeds affects the content of flavan-3-ol derivatives, and their antioxidant and anti-inflammatory activities. *Heliyon* 2020, 6, e05214. https://doi.org/10.1016/j.heliyon.2020.e05214.
- 23. Melo, P.S.; Selani, M.M.; Gonçalves, R.H.; de Oliveira Paulino, J.; Massarioli, A.P.; de Alencar, S.M. Açaí seeds: An unexplored agro-industrial residue as a potential source of lipids, fibers, and antioxidant phenolic compounds. *Ind. Crops Prod.* **2021**, *161*, 113204.
- Rodrigues, R.B.; Lichtenthaler, R.; Zimmermann, B.F.; Papagiannopoulos, M.; Fabricius, H.; Marx, F.; Maia, J.G.; Almeida, O. Total oxidant scavenging capacity of *Euterpe oleracea* Mart. (acai) seeds and identification of their polyphenolic compounds. *J. Agric. Food Chem.* 2006, *54*, 4162–4167. https://doi.org/10.1021/jf058169p.
- de Moura, R.S.; Pires, K.M.; Santos Ferreira, T.; Lopes, A.A.; Nesi, R.T.; Resende, A.C.; Sousa, P.J.; da Silva, A.J.; Porto, L.C.; Valenca, S.S. Addition of acai (*Euterpe oleracea*) to cigarettes has a protective effect against emphysema in mice. *Food Chem. Toxicol.* 2011, 49, 855–863. https://doi.org/10.1016/j.fct.2010.12.007.
- Lee, J.Y.; Kim, N.; Choi, Y.J.; Nam, R.H.; Lee, S.; Ham, M.H.; Suh, J.H.; Choi, Y.J.; Lee, H.S.; Lee, D.H. Anti-inflammatory and Anti-tumorigenic Effects of Acai Berry in Helicobacter felis-infected mice. *J. Cancer Prev.* 2016, 21, 48–54. https://doi.org/10.15430/JCP.2016.21.1.48.
- Moura, R.S.; Ferreira, T.S.; Lopes, A.A.; Pires, K.M.; Nesi, R.T.; Resende, A.C.; Souza, P.J.; Silva, A.J.; Borges, R.M.; Porto, L.C.; et al. Effects of *Euterpe oleracea* Mart. (ACAI) extract in acute lung inflammation induced by cigarette smoke in the mouse. *Phytomedicine* 2012, *19*, 262–269. https://doi.org/10.1016/j.phymed.2011.11.004.
- Poulose, S.M.; Fisher, D.R.; Larson, J.; Bielinski, D.F.; Rimando, A.M.; Carey, A.N.; Schauss, A.G.; Shukitt-Hale, B. Anthocyaninrich acai (*Euterpe oleracea* Mart.) fruit pulp fractions attenuate inflammatory stress signaling in mouse brain BV-2 microglial cells. *J. Agric. Food Chem.* 2012, 60, 1084–1093. https://doi.org/10.1021/jf203989k.

- Santos, I.B.; de Bem, G.F.; da Costa, C.A.; de Carvalho, L.; de Medeiros, A.F.; Silva, D.L.B.; Romao, M.H.; de Andrade Soares, R.; Ognibene, D.T.; de Moura, R.S.; et al. Acai seed extract prevents the renin-angiotensin system activation, oxidative stress and inflammation in white adipose tissue of high-fat diet-fed mice. *Nutr. Res.* 2020, 79, 35–49. https://doi.org/10.1016/j.nutres.2020.05.006.
- 30. ALNasser, M.N.; Mellor, I.R. Neuroprotective activities of acai berries (*Euterpe* sp.): A review. J. Herbmed Pharmacol. 2022, 11, 166–181.
- Kang, J.; Li, Z.; Wu, T.; Jensen, G.S.; Schauss, A.G.; Wu, X. Anti-oxidant capacities of flavonoid compounds isolated from acai pulp (*Euterpe oleracea* Mart.). Food Chem. 2010, 122, 610–617.
- 32. de Moraes Arnoso, B.J.; Magliaccio, F.M.; de Araujo, C.A.; de Andrade Soares, R.; Santos, I.B.; de Bem, G.F.; Fernandes-Santos, C.; Ognibene, D.T.; de Moura, R.S.; Resende, A.C.; et al. Acai seed extract (ASE) rich in proanthocyanidins improves cardiovascular remodeling by increasing antioxidant response in obese high-fat diet-fed mice. *Chem. Biol. Interact.* 2022, 351, 109721. https://doi.org/10.1016/j.cbi.2021.109721.
- Bellucci, E.R.B.; Dos Santos, J.M.; Carvalho, L.T.; Borgonovi, T.F.; Lorenzo, J.M.; Silva-Barretto, A.C.D. Acai extract powder as natural antioxidant on pork patties during the refrigerated storage. *Meat Sci.* 2022, 184, 108667. https://doi.org/10.1016/j.meatsci.2021.108667.
- 34. da Silva, T.V.N.; Torres, M.F.; Sampaio, L.A.; Hamoy, M.; Monserrat, J.M.; Barbas, L.A.L. Dietary *Euterpe oleracea* Mart. attenuates seizures and damage to lipids in the brain of Colossoma macropomum. *Fish Physiol. Biochem.* **2021**, 47, 1851–1864. https://doi.org/10.1007/s10695-021-01010-y.
- Kim, K.J.; Kim, Y.; Jin, S.G.; Kim, J.Y. Acai berry extract as a regulator of intestinal inflammation pathways in a Caco-2 and RAW 264.7 co-culture model. J. Food Biochem. 2021, e13848. https://doi.org/10.1111/jfbc.13848.
- Silva, M.; Costa, J.H.; Pacheco-Fill, T.; Ruiz, A.; Vidal, F.C.B.; Borges, K.R.A.; Guimaraes, S.J.A.; Azevedo-Santos, A.P.S.; Buglio, K.E.; Foglio, M.A.; et al. Acai (*Euterpe oleracea* Mart.) Seed Extract Induces ROS Production and Cell Death in MCF-7 Breast Cancer Cell Line. *Molecules* 2021, 26, 3546. https://doi.org/10.3390/molecules26123546.
- Martins, G.R.; Guedes, D.; Marques de Paula, U.L.; de Oliveira, M.; Lutterbach, M.T.S.; Reznik, L.Y.; Servulo, E.F.C.; Alviano, C.S.; Ribeiro da Silva, A.J.; Alviano, D.S. Acai (*Euterpe oleracea* Mart.) Seed Extracts from Different Varieties: A Source of Proanthocyanidins and Eco-Friendly Corrosion Inhibition Activity. *Molecules* 2021, 26, 3433. https://doi.org/10.3390/molecules26113433.
- de Bem, G.F.; Okinga, A.; Ognibene, D.T.; da Costa, C.A.; Santos, I.B.; Soares, R.A.; Silva, D.L.B.; da Rocha, A.P.M.; Isnardo Fernandes, J.; Fraga, M.C.; et al. Anxiolytic and antioxidant effects of *Euterpe oleracea* Mart. (acai) seed extract in adult rat offspring submitted to periodic maternal separation. *Appl. Physiol. Nutr. Metab.* 2020, 45, 1277–1286. https://doi.org/10.1139/apnm-2020-0099.
- Remigante, A.; Spinelli, S.; Straface, E.; Gambardella, L.; Caruso, D.; Falliti, G.; Dossena, S.; Marino, A.; Morabito, R. Acai (*Euterpe oleracea*) Extract Protects Human Erythrocytes from Age-Related Oxidative Stress. *Cells* 2022, *11*, 2391. https://doi.org/10.3390/cells11152391.
- Cordaro, M.; D'Amico, R.; Fusco, R.; Peritore, A.F.; Genovese, T.; Interdonato, L.; Franco, G.; Arangia, A.; Gugliandolo, E.; Crupi, R.; et al. Discovering the Effects of Fisetin on NF-kappaB/NLRP-3/NRF-2 Molecular Pathways in a Mouse Model of Vascular Dementia Induced by Repeated Bilateral Carotid Occlusion. *Biomedicines* 2022, 10, 1448. https://doi.org/10.3390/biomedicines10061448.
- Siracusa, R.; Impellizzeri, D.; Cordaro, M.; Crupi, R.; Esposito, E.; Petrosino, S.; Cuzzocrea, S. Anti-Inflammatory and Neuroprotective Effects of Co-UltraPEALut in a Mouse Model of Vascular Dementia. *Front. Neurol.* 2017, *8*, 233. https://doi.org/10.3389/fneur.2017.00233.
- Genovese, T.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Peritore, A.F.; Crupi, R.; Interdonato, L.; Gugliandolo, E.; Cuzzocrea, S.; Paola, R.D.; et al. Acai (*Euterpe oleraceae* Mart.) Seeds Regulate NF-kappaB and Nrf2/ARE Pathways Protecting Lung against Acute and Chronic Inflammation. *Cell Physiol. Biochem.* 2022, 56, 1–20. https://doi.org/10.33594/000000529.
- 43. Prut, L.; Belzung, C. The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: A review. *Eur. J. Pharmacol.* **2003**, *463*, 3–33. https://doi.org/10.1016/S0014-2999(03)01272-X.
- Crupi, R.; Cambiaghi, M.; Spatz, L.; Hen, R.; Thorn, M.; Friedman, E.; Vita, G.; Battaglia, F. Reduced adult neurogenesis and altered emotional behaviors in autoimmune-prone B-cell activating factor transgenic mice. *Biol. Psychiatry* 2010, 67, 558–566. https://doi.org/10.1016/j.biopsych.2009.12.008.
- 45. Pan, Z.; Cui, M.; Dai, G.; Yuan, T.; Li, Y.; Ji, T.; Pan, Y. Protective Effect of Anthocyanin on Neurovascular Unit in Cerebral Ischemia/Reperfusion Injury in Rats. *Front. Neurosci.* **2018**, *12*, 947. https://doi.org/10.3389/fnins.2018.00947.
- Di Paola, D.; Iaria, C.; Capparucci, F.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Aflatoxin B1 Toxicity in Zebrafish Larva (*Danio rerio*): Protective Role of *Hericium erinaceus*. *Toxins* 2021, 13, 710. https://doi.org/10.3390/toxins13100710.
- Di Paola, D.; Natale, S.; Gugliandolo, E.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Assessment of 2-Pentadecyl-2-oxazoline Role on Lipopolysaccharide-Induced Inflammation on Early Stage Development of Zebrafish (*Danio rerio*). *Life* 2022, *12*, 128. https://doi.org/10.3390/life12010128.
- Di Paola, D.; Capparucci, F.; Abbate, J.M.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Genovese, T.; Impellizzeri, D.; et al. Environmental Risk Assessment of Oxaliplatin Exposure on Early Life Stages of Zebrafish (*Danio rerio*). *Toxics* 2022, 10, 81. https://doi.org/10.3390/toxics10020081.

- Cordaro, M.; Impellizzeri, D.; Gugliandolo, E.; Siracusa, R.; Crupi, R.; Esposito, E.; Cuzzocrea, S. Adelmidrol, a Palmitoylethanolamide Analogue, as a New Pharmacological Treatment for the Management of Inflammatory Bowel Disease. *Mol. Pharmacol.* 2016, 90, 549–561. https://doi.org/10.1124/mol.116.105668.
- Fusco, R.; Siracusa, R.; D'Amico, R.; Cordaro, M.; Genovese, T.; Gugliandolo, E.; Peritore, A.F.; Crupi, R.; Di Paola, R.; Cuzzocrea, S.; et al. Mucosa-Associated Lymphoid Tissue Lymphoma Translocation 1 Inhibitor as a Novel Therapeutic Tool for Lung Injury. *Int. J. Mol. Sci.* 2020, *21*, 7761.
- Fusco, R.; Siracusa, R.; Peritore, A.F.; Gugliandolo, E.; Genovese, T.; D'Amico, R.; Cordaro, M.; Crupi, R.; Mandalari, G.; Impellizzeri, D.; et al. The Role of Cashew (*Anacardium occidentale* L.) Nuts on an Experimental Model of Painful Degenerative Joint Disease. *Antioxidants* 2020, 9, 511. https://doi.org/10.3390/antiox9060511.
- 52. Peritore, A.F.; Siracusa, R.; Fusco, R.; Gugliandolo, E.; D'Amico, R.; Cordaro, M.; Crupi, R.; Genovese, T.; Impellizzeri, D.; Cuzzocrea, S.; et al. Ultramicronized Palmitoylethanolamide and Paracetamol, a New Association to Relieve Hyperalgesia and Pain in a Sciatic Nerve Injury Model in Rat. *Int. J. Mol. Sci.* 2020, *21*, 3509. https://doi.org/10.3390/ijms21103509.
- Fusco, R.; Cordaro, M.; Siracusa, R.; D'Amico, R.; Genovese, T.; Gugliandolo, E.; Peritore, A.F.; Crupi, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Biochemical Evaluation of the Antioxidant Effects of Hydroxytyrosol on Pancreatitis-Associated Gut Injury. *Antioxidants* 2020, *9*, 781. https://doi.org/10.3390/antiox9090781.
- Sawant, S.; Gokulan, R.; Dongre, H.; Vaidya, M.; Chaukar, D.; Prabhash, K.; Ingle, A.; Joshi, S.; Dange, P.; Joshi, S.; et al. Prognostic role of Oct4, CD44 and c-Myc in radio-chemo-resistant oral cancer patients and their tumourigenic potential in immunodeficient mice. *Clin. Oral. Investig.* 2016, 20, 43–56. https://doi.org/10.1007/s00784-015-1476-6.
- Siracusa, R.; Paterniti, I.; Cordaro, M.; Crupi, R.; Bruschetta, G.; Campolo, M.; Cuzzocrea, S.; Esposito, E. Neuroprotective Effects of Temsirolimus in Animal Models of Parkinson's Disease. *Mol. Neurobiol.* 2018, 55, 2403–2419. https://doi.org/10.1007/s12035-017-0496-4.
- Gugliandolo, E.; D'Amico, R.; Cordaro, M.; Fusco, R.; Siracusa, R.; Crupi, R.; Impellizzeri, D.; Cuzzocrea, S.; Di Paola, R. Effect of PEA-OXA on neuropathic pain and functional recovery after sciatic nerve crush. *J. Neuroinflamm.* 2018, 15, 264. https://doi.org/10.1186/s12974-018-1303-5.
- Genovese, T.; Siracusa, R.; D'Amico, R.; Cordaro, M.; Peritore, A.F.; Gugliandolo, E.; Crupi, R.; Trovato Salinaro, A.; Raffone, E.; Impellizzeri, D.; et al. Regulation of Inflammatory and Proliferative Pathways by Fotemustine and Dexamethasone in Endometriosis. *Int. J. Mol. Sci.* 2021, *22*, 5998. https://doi.org/10.3390/ijms22115998.
- Peritore, A.F.; Crupi, R.; Scuto, M.; Gugliandolo, E.; Siracusa, R.; Impellizzeri, D.; Cordaro, M.; D'Amico, R.; Fusco, R.; Di Paola, R.; et al. The Role of Annexin A1 and Formyl Peptide Receptor 2/3 Signaling in Chronic Corticosterone-Induced Depression-Like behaviors and Impairment in Hippocampal-Dependent Memory. CNS Neurol. Disord. Drug Targets 2020, 19, 27–43. https://doi.org/10.2174/1871527319666200107094732.
- Fusco, R.; Cordaro, M.; Siracusa, R.; Peritore, A.F.; Gugliandolo, E.; Genovese, T.; D'Amico, R.; Crupi, R.; Smeriglio, A.; Mandalari, G.; et al. Consumption of *Anacardium occidentale* L. (Cashew Nuts) Inhibits Oxidative Stress through Modulation of the Nrf2/HO-1 and NF-kB Pathways. *Molecules* 2020, 25, 4426. https://doi.org/10.3390/molecules25194426.
- Cordaro, M.; Siracusa, R.; Crupi, R.; Impellizzeri, D.; Peritore, A.F.; D'Amico, R.; Gugliandolo, E.; Di Paola, R.; Cuzzocrea, S. 2-Pentadecyl-2-Oxazoline Reduces Neuroinflammatory Environment in the MPTP Model of Parkinson Disease. *Mol. Neurobiol.* 2018, 55, 9251–9266. https://doi.org/10.1007/s12035-018-1064-2.
- Impellizzeri, D.; Cordaro, M.; Bruschetta, G.; Crupi, R.; Pascali, J.; Alfonsi, D.; Marcolongo, G.; Cuzzocrea, S. 2-pentadecyl-2oxazoline: Identification in coffee, synthesis and activity in a rat model of carrageenan-induced hindpaw inflammation. *Pharmacol. Res.* 2016, 108, 23–30. https://doi.org/10.1016/j.phrs.2016.04.007.
- Cordaro, M.; Siracusa, R.; Fusco, R.; D'Amico, R.; Peritore, A.F.; Gugliandolo, E.; Genovese, T.; Scuto, M.; Crupi, R.; Mandalari, G.; et al. Cashew (*Anacardium occidentale* L.) Nuts Counteract Oxidative Stress and Inflammation in an Acute Experimental Model of Carrageenan-Induced Paw Edema. *Antioxidants* 2020, 9, 660. https://doi.org/10.3390/antiox9080660.
- 63. D'Amico, R.; Fusco, R.; Cordaro, M.; Siracusa, R.; Peritore, A.F.; Gugliandolo, E.; Crupi, R.; Scuto, M.; Cuzzocrea, S.; Di Paola, R.; et al. Modulation of NLRP3 Inflammasome through Formyl Peptide Receptor 1 (Fpr-1) Pathway as a New Therapeutic Target in Bronchiolitis Obliterans Syndrome. *Int. J. Mol. Sci.* 2020, *21*, 2144. https://doi.org/10.3390/ijms21062144.
- 64. Esposito, E.; Impellizzeri, D.; Bruschetta, G.; Cordaro, M.; Siracusa, R.; Gugliandolo, E.; Crupi, R.; Cuzzocrea, S. A new comicronized composite containing palmitoylethanolamide and polydatin shows superior oral efficacy compared to their association in a rat paw model of carrageenan-induced inflammation. *Eur. J. Pharmacol.* 2016, 782, 107–118. https://doi.org/10.1016/j.ejphar.2016.03.033.
- 65. Impellizzeri, D.; Siracusa, R.; Cordaro, M.; Peritore, A.F.; Gugliandolo, E.; Mancuso, G.; Midiri, A.; Di Paola, R.; Cuzzocrea, S. Therapeutic potential of dinitrobenzene sulfonic acid (DNBS)-induced colitis in mice by targeting IL-1beta and IL-18. *Biochem. Pharmacol.* **2018**, *155*, 150–161. https://doi.org/10.1016/j.bcp.2018.06.029.
- 66. Di Paola, R.; Fusco, R.; Impellizzeri, D.; Cordaro, M.; Britti, D.; Morittu, V.M.; Evangelista, M.; Cuzzocrea, S. Adelmidrol, in combination with hyaluronic acid, displays increased anti-inflammatory and analgesic effects against monosodium iodoacetate-induced osteoarthritis in rats. *Arthritis Res. Ther.* 2016, *18*, 291. https://doi.org/10.1186/s13075-016-1189-5.
- Fusco, R.; Cordaro, M.; Genovese, T.; Impellizzeri, D.; Siracusa, R.; Gugliandolo, E.; Peritore, A.F.; D'Amico, R.; Crupi, R.; Cuzzocrea, S.; et al. Adelmidrol: A New Promising Antioxidant and Anti-Inflammatory Therapeutic Tool in Pulmonary Fibrosis. *Antioxidants* 2020, *9*, 601. https://doi.org/10.3390/antiox9070601.

- Fusco, R.; Gugliandolo, E.; Siracusa, R.; Scuto, M.; Cordaro, M.; D'Amico, R.; Evangelista, M.; Peli, A.; Peritore, A.F.; Impellizzeri, D.; et al. Formyl Peptide Receptor 1 Signaling in Acute Inflammation and Neural Differentiation Induced by Traumatic Brain Injury. *Biology* 2020, *9*, 238. https://doi.org/10.3390/biology9090238.
- 69. Impellizzeri, D.; Siracusa, R.; Cordaro, M.; Crupi, R.; Peritore, A.F.; Gugliandolo, E.; D'Amico, R.; Petrosino, S.; Evangelista, M.; Di Paola, R.; et al. N-Palmitoylethanolamine-oxazoline (PEA-OXA): A new therapeutic strategy to reduce neuroinflammation, oxidative stress associated to vascular dementia in an experimental model of repeated bilateral common carotid arteries occlusion. *Neurobiol. Dis.* 2019, 125, 77–91. https://doi.org/10.1016/j.nbd.2019.01.007.
- Cordaro, M.; Cuzzocrea, S.; Crupi, R. An Update of Palmitoylethanolamide and Luteolin Effects in Preclinical and Clinical Studies of Neuroinflammatory Events. *Antioxidants* 2020, 9, 216. https://doi.org/10.3390/antiox9030216.
- Deng, M.; Huang, L.; Zhong, X.; Huang, M. Dynamic Changes of Beclin-1 in the Hippocampus of Male Mice with Vascular Dementia at Different Time Points. J. Mol. Neurosci. 2020, 70, 1611–1618. https://doi.org/10.1007/s12031-020-01591-y.
- 72. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **2001**, *25*, 402–408. https://doi.org/10.1006/meth.2001.1262.
- 73. Lacruz, R.S.; Smith, C.E.; Smith, S.M.; Hu, P.; Bringas, P., Jr.; Sahin-Toth, M.; Moradian-Oldak, J.; Paine, M.L. Chymotrypsin C (caldecrin) is associated with enamel development. *J. Dent. Res.* **2011**, *90*, 1228–1233. https://doi.org/10.1177/0022034511418231.
- Paola, D.D.; Capparucci, F.; Natale, S.; Crupi, R.; Cuzzocrea, S.; Spano, N.; Gugliandolo, E.; Peritore, A.F. Combined Effects of Potassium Perchlorate and a Neonicotinoid on Zebrafish Larvae (*Danio rerio*). *Toxics* 2022, 10, 203. https://doi.org/10.3390/toxics10050203.
- Di Paola, D.; Capparucci, F.; Lanteri, G.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Combined Toxicity of Xenobiotics Bisphenol A and Heavy Metals on Zebrafish Embryos (*Danio rerio*). *Toxics* 2021, 9, 344. https://doi.org/10.3390/toxics9120344.
- 76. Di Paola, D.; Natale, S.; Iaria, C.; Crupi, R.; Cuzzocrea, S.; Spano, N.; Gugliandolo, E.; Peritore, A.F. Environmental Co-Exposure to Potassium Perchlorate and Cd Caused Toxicity and Thyroid Endocrine Disruption in Zebrafish Embryos and Larvae (*Danio rerio*). *Toxics* 2022, 10, 198. https://doi.org/10.3390/toxics10040198.
- 77. Di Paola, D.; Abbate, J.M.; Iaria, C.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Environmental Risk Assessment of Dexamethasone Sodium Phosphate and Tocilizumab Mixture in Zebrafish Early Life Stage (*Danio rerio*). *Toxics* 2022, 10, 279. https://doi.org/10.3390/toxics10060279.
- Di Paola, D.; Capparucci, F.; Lanteri, G.; Crupi, R.; Marino, Y.; Franco, G.A.; Cuzzocrea, S.; Spano, N.; Gugliandolo, E.; Peritore, A.F. Environmental Toxicity Assessment of Sodium Fluoride and Platinum-Derived Drugs Co-Exposure on Aquatic Organisms. *Toxics* 2022, 10, 272. https://doi.org/10.3390/toxics10050272.
- 79. Di Paola, D.; Natale, S.; Iaria, C.; Cordaro, M.; Crupi, R.; Siracusa, R.; D'Amico, R.; Fusco, R.; Impellizzeri, D.; Cuzzocrea, S.; et al. Intestinal Disorder in Zebrafish Larvae (*Danio rerio*): The Protective Action of N-Palmitoylethanolamide-oxazoline. *Life* 2022, 12, 125. https://doi.org/10.3390/life12010125.
- Genovese, T.; Impellizzeri, D.; D'Amico, R.; Fusco, R.; Peritore, A.F.; Di Paola, D.; Interdonato, L.; Gugliandolo, E.; Crupi, R.; Di Paola, R.; et al. Role of Bevacizumab on Vascular Endothelial Growth Factor in Apolipoprotein E Deficient Mice after Traumatic Brain Injury. *Int. J. Mol. Sci.* 2022, 23, 4162. https://doi.org/10.3390/ijms23084162.
- Cordaro, M.; Siracusa, R.; D'Amico, R.; Genovese, T.; Franco, G.; Marino, Y.; Di Paola, D.; Cuzzocrea, S.; Impellizzeri, D.; Di Paola, R.; et al. Role of Etanercept and Infliximab on Nociceptive Changes Induced by the Experimental Model of Fibromyalgia. *Int. J. Mol. Sci.* 2022, 23, 6139. https://doi.org/10.3390/ijms23116139.
- D'Amico, R.; Gugliandolo, E.; Cordaro, M.; Fusco, R.; Genovese, T.; Peritore, A.F.; Crupi, R.; Interdonato, L.; Di Paola, D.; Cuzzocrea, S.; et al. Toxic Effects of Endocrine Disruptor Exposure on Collagen-Induced Arthritis. *Biomolecules* 2022, *12*, 564. https://doi.org/10.3390/biom12040564.
- D'Amico, R.; Gugliandolo, E.; Siracusa, R.; Cordaro, M.; Genovese, T.; Peritore, A.F.; Crupi, R.; Interdonato, L.; Di Paola, D.; Cuzzocrea, S.; et al. Toxic Exposure to Endocrine Disruptors Worsens Parkinson's Disease Progression through NRF2/HO-1 Alteration. *Biomedicines* 2022, 10, 1073. https://doi.org/10.3390/biomedicines10051073.
- Fusco, R.; D'Amico, R.; Cordaro, M.; Gugliandolo, E.; Siracusa, R.; Peritore, A.F.; Crupi, R.; Impellizzeri, D.; Cuzzocrea, S.; Di Paola, R. Absence of formyl peptide receptor 1 causes endometriotic lesion regression in a mouse model of surgically-induced endometriosis. *Oncotarget* 2018, *9*, 31355–31366. https://doi.org/10.18632/oncotarget.25823.
- 85. Di Paola, R.; Impellizzeri, D.; Fusco, R.; Cordaro, M.; Siracusa, R.; Crupi, R.; Esposito, E.; Cuzzocrea, S. Ultramicronized palmitoylethanolamide (PEA-um((R))) in the treatment of idiopathic pulmonary fibrosis. *Pharmacol. Res.* **2016**, *111*, 405–412. https://doi.org/10.1016/j.phrs.2016.07.010.
- Fusco, R.; Siracusa, R.; D'Amico, R.; Peritore, A.F.; Cordaro, M.; Gugliandolo, E.; Crupi, R.; Impellizzeri, D.; Cuzzocrea, S.; Di Paola, R. Melatonin Plus Folic Acid Treatment Ameliorates Reserpine-Induced Fibromyalgia: An Evaluation of Pain, Oxidative Stress, and Inflammation. *Antioxidants* 2019, *8*, 628. https://doi.org/10.3390/antiox8120628.
- Kang, H.B.; Kim, S.H.; Uhm, S.H.; Kim, D.K.; Lee, N.S.; Jeong, Y.G.; Sung, N.Y.; Kim, D.S.; Han, I.J.; Yoo, Y.C.; et al. Perilla frutescens Leaf Extract Attenuates Vascular Dementia-Associated Memory Deficits, Neuronal Damages, and Microglial Activation. *Curr. Issues Mol. Biol.* 2022, 44, 257–272. https://doi.org/10.3390/cimb44010019.
- Young, M.M.; Kester, M.; Wang, H.G. Sphingolipids: Regulators of crosstalk between apoptosis and autophagy. J. Lipid Res. 2013, 54, 5–19. https://doi.org/10.1194/jlr.R031278.
- 89. Iadecola, C. The pathobiology of vascular dementia. Neuron 2013, 80, 844-866. https://doi.org/10.1016/j.neuron.2013.10.008.

- 90. Skoog, I. Status of risk factors for vascular dementia. Neuroepidemiology 1998, 17, 2–9. https://doi.org/10.1159/000026147.
- 91. Wang, X.X.; Zhang, B.; Xia, R.; Jia, Q.Y. Inflammation, apoptosis and autophagy as critical players in vascular dementia. *Eur. Rev. Med. Pharmacol. Sci.* 2020, 24, 9601–9614. https://doi.org/10.26355/eurrev_202009_23048.
- Barichella, M.; Cereda, E.; Pezzoli, G. Major nutritional issues in the management of Parkinson's disease. *Mov. Disord.* 2009, 24, 1881–1892. https://doi.org/10.1002/mds.22705.
- Seidl, S.E.; Santiago, J.A.; Bilyk, H.; Potashkin, J.A. The emerging role of nutrition in Parkinson's disease. *Front. Aging Neurosci.* 2014, 6, 36. https://doi.org/10.3389/fnagi.2014.00036.
- Chen, W.W.; Zhang, X.; Huang, W.J. Role of neuroinflammation in neurodegenerative diseases (Review). Mol. Med. Rep. 2016, 13, 3391–3396. https://doi.org/10.3892/mmr.2016.4948.
- 95. de Almeida Magalhaes, T.S.S.; de Oliveira Macedo, P.C.; Converti, A.; Neves de Lima, A.A. The Use of *Euterpe oleracea* Mart. As a New Perspective for Disease Treatment and Prevention. *Biomolecules* **2020**, *10*, 813. https://doi.org/10.3390/biom10060813.
- Denzer, I.; Munch, G.; Friedland, K. Modulation of mitochondrial dysfunction in neurodegenerative diseases via activation of nuclear factor erythroid-2-related factor 2 by food-derived compounds. *Pharmacol. Res.* 2016, 103, 80–94. https://doi.org/10.1016/j.phrs.2015.11.019.
- Kovacs, G.G. Molecular Pathological Classification of Neurodegenerative Diseases: Turning towards Precision Medicine. *Int. J. Mol. Sci.* 2016, 17, 189. https://doi.org/10.3390/ijms17020189.
- Lewerenz, J.; Maher, P. Chronic Glutamate Toxicity in Neurodegenerative Diseases-What is the Evidence? *Front. Neurosci.* 2015, 9, 469. https://doi.org/10.3389/fnins.2015.00469.
- Machado, A.K.; Andreazza, A.C.; da Silva, T.M.; Boligon, A.A.; do Nascimento, V.; Scola, G.; Duong, A.; Cadona, F.C.; Ribeiro, E.E.; da Cruz, I.B. Neuroprotective Effects of Acai (*Euterpe oleracea* Mart.) against Rotenone In Vitro Exposure. Oxid. Med. Cell Longev. 2016, 2016, 8940850. https://doi.org/10.1155/2016/8940850.
- Torma, P.D.; Brasil, A.V.; Carvalho, A.V.; Jablonski, A.; Rabelo, T.K.; Moreira, J.C.; Gelain, D.P.; Flores, S.H.; Augusti, P.R.; Rios, A.O. Hydroethanolic extracts from different genotypes of acai (*Euterpe oleracea*) presented antioxidant potential and protected human neuron-like cells (SH-SY5Y). *Food Chem.* 2017, 222, 94–104. https://doi.org/10.1016/j.foodchem.2016.12.006.
- 101. Xie, C.; Kang, J.; Li, Z.; Schauss, A.G.; Badger, T.M.; Nagarajan, S.; Wu, T.; Wu, X. The acai flavonoid velutin is a potent antiinflammatory agent: Blockade of LPS-mediated TNF-alpha and IL-6 production through inhibiting NF-kappaB activation and MAPK pathway. J. Nutr. Biochem. 2012, 23, 1184–1191. https://doi.org/10.1016/j.jnutbio.2011.06.013.
- 102. Gauthier, S.; Ferris, S. Outcome measures for probable vascular dementia and Alzheimer's disease with cerebrovascular disease. *Int. J. Clin. Pract. Suppl.* **2001**, 29–39.
- 103. Bathgate, D.; Snowden, J.S.; Varma, A.; Blackshaw, A.; Neary, D. Behaviour in frontotemporal dementia, Alzheimer's disease and vascular dementia. *Acta Neurol. Scand.* 2001, 103, 367–378. https://doi.org/10.1034/j.1600-0404.2001.2000236.x.
- 104. Yang, Y.; Jiang, S.; Yan, J.; Li, Y.; Xin, Z.; Lin, Y.; Qu, Y. An overview of the molecular mechanisms and novel roles of Nrf2 in neurodegenerative disorders. *Cytokine Growth Factor Rev.* 2015, 26, 47–57. https://doi.org/10.1016/j.cytogfr.2014.09.002.
- 105. Liu, D.D.; Yuan, X.; Chu, S.F.; Chen, C.; Ren, Q.; Luo, P.; Lin, M.Y.; Wang, S.S.; Zhu, T.B.; Ai, Q.D.; et al. CZ-7, a new derivative of Claulansine F, ameliorates 2VO-induced vascular dementia in rats through a Nrf2-mediated antioxidant responses. *Acta Pharmacol. Sin.* 2019, 40, 425–440. https://doi.org/10.1038/s41401-018-0078-7.
- 106. Zhang, D.; Xiao, Y.; Lv, P.; Teng, Z.; Dong, Y.; Qi, Q.; Liu, Z. Edaravone attenuates oxidative stress induced by chronic cerebral hypoperfusion injury: Role of ERK/Nrf2/HO-1 signaling pathway. *Neurol. Res.* 2018, 40, 1–10. https://doi.org/10.1080/01616412.2017.1376457.
- O'Brien, C.E.; Bonanno, L.; Zhang, H.; Wyss-Coray, T. Beclin 1 regulates neuronal transforming growth factor-beta signaling by mediating recycling of the type I receptor ALK5. *Mol. Neurodegener.* 2015, *10*, 69. https://doi.org/10.1186/s13024-015-0065-0.
- 108. Diniz, L.P.; Matias, I.; Siqueira, M.; Stipursky, J.; Gomes, F.C.A. Astrocytes and the TGF-beta1 Pathway in the Healthy and Diseased Brain: A Double-Edged Sword. *Mol. Neurobiol.* **2019**, *56*, 4653–4679. https://doi.org/10.1007/s12035-018-1396-y.
- 109. Caraci, F.; Gulisano, W.; Guida, C.A.; Impellizzeri, A.A.; Drago, F.; Puzzo, D.; Palmeri, A. A key role for TGF-beta1 in hippocampal synaptic plasticity and memory. *Sci. Rep.* **2015**, *5*, 11252. https://doi.org/10.1038/srep11252.
- 110. Kandasamy, M.; Anusuyadevi, M.; Aigner, K.M.; Unger, M.S.; Kniewallner, K.M.; de Sousa, D.M.B.; Altendorfer, B.; Mrowetz, H.; Bogdahn, U.; Aigner, L. TGF-beta Signaling: A Therapeutic Target to Reinstate Regenerative Plasticity in Vascular Dementia? *Aging Dis.* 2020, 11, 828–850. https://doi.org/10.14336/AD.2020.0222.
- 111. Liu, H.X.; Zhang, J.J.; Zheng, P.; Zhang, Y. Altered expression of MAP-2, GAP-43, and synaptophysin in the hippocampus of rats with chronic cerebral hypoperfusion correlates with cognitive impairment. *Brain Res. Mol. Brain Res.* 2005, 139, 169–177. https://doi.org/10.1016/j.molbrainres.2005.05.014.
- 112. Gomazkov, O.; Lagunin, A. Vascular dementia: Molecular targets of neuroprotective therapy. Biol. Bull. Rev. 2017, 7, 528-536.
- 113. Bertelli, S.; Remigante, A.; Zuccolini, P.; Barbieri, R.; Ferrera, L.; Picco, C.; Gavazzo, P.; Pusch, M. Mechanisms of Activation of LRRC8 Volume Regulated Anion Channels. *Cell Physiol. Biochem.* **2021**, *55*, 41–56. https://doi.org/10.33594/000000329.
- 114. Remigante, A.; Morabito, R.; Spinelli, S.; Trichilo, V.; Loddo, S.; Sarikas, A.; Dossena, S.; Marino, A. d-Galactose Decreases Anion Exchange Capability through Band 3 Protein in Human Erythrocytes. *Antioxidants* 2020, 9, 689. https://doi.org/10.3390/antiox9080689.