

Table S6. Clinical characteristics of the patient cohort.

Variable	Total %	PCNA_V2 expres- sion	<i>p value</i>	PCNA_V4 ex- pression	<i>p value</i>
Age					
0-9	40 (57.14%)	13 (18.5%) High 27 (38.5%) Low	0.400	17 24.2%) High 23 (23.8%) Low	0.240
>9	30 (42.85%)	7 (10%) High 23 (32.8%) Low		17 (24.2%) High 13 (18.5%) Low	
Sex					
Men	34 (48.6%)	12 (17.1%) High 22 (31.4%) Low	0.347	27 (38.6) High 7 (10.0%) Low	0.094
Women	36 (51.4%)	9 (12.9%) High 27 (38.6%) Low		22 (31.4%) High 14 (20.0%) Low	
Genetic findings: T(9;22), T(1;19), T(12;21)					
Yes	7 (10%)	3 (4.2%) High 4 (5.7%) Low	0.433	3 (4.2%) High 4 (5.7%) Low	0.433
Cerebrospinal fluid (blasts %)					
No blast	64 (91.4%)	19 (27.1%) High 45 (64.3%) Low	0.444	45 (64.3%) High 19 (27.1%) Low	0.444
< 5 blast	2 (2.9%%)	0 (0%) High 2 (2.9%) Low		2 (2.9%) High 0 (0%) Low	
> 5 blast	4 (5.7%)	2 (2.9%) High 2 (2.9%) Low		2 (2.9%) High 2 (2.9%) Low	
Treatment response					
Good	45 (64.3%)	16 (22.9%) High 29 (41.4%) Low	0.243	32 (45.7%) High 13 (18.6%) Low	0.894
Bad	23 (32.9%)	5 (7.1%) High 18 (25.7%) Low		16 (22.9%) High 7 (10%) Low	
Hypodiploidy					
Yes	23 (32.9%)	7 (10%) High 16 (22.9%) Low	0.955	15 (21.4%) High 8 (11.4%) Low	0.541
No	47 (67.1%)	14 (20.0%) High 33 (47.1%) Low		34 (48.6%) High 13 (16.6%) Low	
Blast %					
<50	33 (47.1%)	9 (12.9%) High 24 (34.3%) Low	0.638	20 (28.6%) High 13 (18.6%) Low	0.105
>50	37 (52.9%)	12 (17.1%) High 25 (35.7%) Low		29 (41.4%) High 8 (11.4%) Low	
Relapse					
Yes	22 (31.4%)	4 (5.7%) High 18 (25.7%) Low	0.118	10 (14.3%) High 12 (17.1%) Low	0.002
No	48 (68.6%)	17 (24.3%) High 31 (44.3%) Low		39 (55.7%) High 9 (12.9%) Low	
Status					
Alive	59 (84.3%)	19 (27.1%) High 40 (57.1%) Low	0.351	43 (61.4%) High 16 (22.9%) Low	0.223
Died	11 (15.7%)	2 (2.9%) High 9 (12.9%) Low		6 (8.6%) High 5 (7.1%) Low	

