

Supplementary Information

Article

Persisting effect of cytokines and toll-like receptor agonists on indoleamine-2,3-dioxygenase-1 in human periodontal ligament stem/stromal cells


Christian Behm^{1,2}, Alice Blufstein¹, Johannes Gahn¹, Barbara Kubin¹, Andreas Moritz¹, Xiaohui Rausch-Fan¹ and Oleh Andrukhov^{1,*}

¹ Division of Conservative Dentistry and Periodontology, University Clinic of Dentistry, Medical University of Vienna, Sensengasse 2a, 1090 Vienna, Austria; christian.behm@meduniwien.ac.at (CB); alice.blufstein@meduniwien.ac.at (AB); Johannes.gahn@gmail.com (JG); andreas.moritz@meduniwien.ac.at (AM); xiaohui.rausch-fan@meduniwien.ac.at (XRF)

² Division of Orthodontics, University Clinic of Dentistry, Medical University of Vienna, Sensengasse 2a, 1090 Vienna, Austria; christian.behm@meduniwien.ac.at (CB)

* Correspondence: oleh.andrukhov@meduniwien.ac.at; Tel.: +43(0)1-40070-2620

Supplementary Figure 1



Supplementary Figure 1. Representative dot plots (b) show the percentage of IDO-1 positive hPDLSCs compared to the unstimulated control (a). Primary hPDLSCs were stimulated with 10 ng/ml TNF- α , 5 ng/ml IL-1 β , 100 ng/ml IFN- γ , 1 μ g/ml Poly I/C or 1 μ g/ml Pam3CSK4. After 48 hours incubation, IDO-1 protein levels were determined by intracellular immunostaining followed by flow cytometry analysis.