

Article

Greenhouse Gas Emissions Response to Fertilizer Application and Soil Moisture in Dry Agricultural Uplands of Central Kenya

Peter Semba Mosongo^{1,2}, David E. Pelster^{3,†} , Xiaoxin Li¹, Gokul Gaudel^{1,2}, Yuying Wang¹, Suying Chen¹, Wenyan Li^{1,2}, David Mburu⁴ and Chunsheng Hu^{1,2,*}

- ¹ Key Laboratory of Agricultural Water Resources, Hebei Key Laboratory of Soil Ecology, Center for Agricultural Resources Research, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, 286 Huaizhong Road, Shijiazhuang 050021, China; petersemba@gmail.com (P.S.M.); xiaoxin_li@sjziam.ac.cn (X.L.); gokul.gaudel@gmail.com (G.G.); wangyuying@sjziam.ac.cn (Y.W.); csy@sjziam.ac.cn (S.C.); liwenyan215@gmail.com (W.L.)
- ² University of Chinese Academy of Sciences, Beijing 100049, China
- ³ Mazingira Centre, International Livestock Research Institute (ILRI), Nairobi 30709-00100, Kenya; david.pelster@agr.gc.ca
- ⁴ College of Agriculture and Natural Resources, Jomo Kenyatta University of Agriculture and Technology (JKUAT), Nairobi 62000-00200, Kenya; david.mburu@jkuat.ac.ke
- * Correspondence: cshu@sjziam.ac.cn
- † Current address: Agriculture and Agri-Food Canada, 2560 Blvd Hochelaga, Quebec, QC G1V 2J3, Canada.

Abstract: In sub-Saharan Africa, agriculture can account for up to 66% of anthropogenic greenhouse gas (GHG) emissions. Unfortunately, due to the low number of studies in the region there is still much uncertainty on how management activities can affect these emissions. To help reduce this uncertainty, we measured GHG emissions from three maize (*Zea mays*) growing seasons in central Kenya. Treatments included: (1) a no N application control (C); (2) split (30% at planting and 70% 1 month after planting) mineral nitrogen (N) applications (Min—100 kg N ha⁻¹); (3) split mineral N + irrigation (equivalent to 10 mm precipitation every three days—MI); (4) split mineral N + 40 kg N ha⁻¹ added as manure (MM—total N = 140 kg ha⁻¹); and (5) split mineral + intercropping with faba beans (*Phaseolus vulgaris*—MB). Soil CO₂ fluxes were lower in season 1 compared to seasons 2 and 3 with fluxes highest in Min ($p = 0.02$) in season 2 and lowest in C ($p = 0.02$) in season 3. There was uptake of CH₄ in these soils that decreased from season 1 to 3 as the mean soil moisture content increased. Cumulative N₂O fluxes ranged from 0.25 to 2.45 kg N₂O-N ha⁻¹, with the highest fluxes from MI during season 3 ($p = 0.01$) and the lowest from C season 1 ($p = 0.03$). The average fertilizer induced emission factor ($0.36 \pm 0.03\%$) was roughly one-third the default value of 1%. Soil moisture was a critical factor controlling GHG emissions in these central Kenya highlands. Under low soil moisture, the soils were CH₄ sinks and minimal N₂O sources.

Keywords: nitrogen fertilization; greenhouse gases; sub-Saharan Africa; small scale farmers



Citation: Mosongo, P.S.; Pelster, D.E.; Li, X.; Gaudel, G.; Wang, Y.; Chen, S.; Li, W.; Mburu, D.; Hu, C. Greenhouse Gas Emissions Response to Fertilizer Application and Soil Moisture in Dry Agricultural Uplands of Central Kenya. *Atmosphere* **2022**, *13*, 463. <https://doi.org/10.3390/atmos13030463>

Academic Editor: Antonio Donateo

Received: 7 January 2022

Accepted: 9 March 2022

Published: 13 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Increased atmospheric greenhouse gas (GHG) concentrations have been linked with increased mean global temperatures over the past century, with a third of the anthropogenic emissions coming from agriculture [1]. In Africa, agriculture accounts for up to 66% of anthropogenic GHG emissions, and although much of these emissions come from livestock, croplands are still responsible for approximately 15% of agricultural GHG emissions [2].

Sub-Saharan Africa (SSA), with a current population of 1.136 billion of whom 30% face acute food shortages annually [3], is home to one of the fastest growing populations in the world. According to the Food and Agricultural Organization (FAO) (2017), there are 4 million Kenyans facing acute food shortages. In response to these shortages, the Kenyan

government, hoping to emulate successful projects in Malawi and the Millennium Villages, has increased incentives for smallholder farmers to purchase and apply synthetic fertilizers through the National Accelerated Agricultural Input Access Program (NAAIAP) [4] to staple crops (e.g., maize (*Zea mays*)), with the aim of improving yields [3]. As a result, Kenya experienced a 4.7% annual increase in fertilizer use between 2015 and 2018 [5].

However, increased nitrogen (N) fertilization increases anthropogenic nitrous oxide (N_2O) emissions [5,6] as N addition tends to induce nitrification and denitrification [6]. Nitrification occurs under aerobic soil conditions when ammonium (NH_4^+) is oxidized to nitrite (NO_2^-) then nitrate (NO_3^-), with N_2O sometimes released as a by-product [7]. Denitrification (i.e., the reduction of NO_3^- to N_2O and then to N_2) on the other hand occurs under anaerobic conditions with about 10% of the N being lost as N_2O [8]. These processes are regulated by soil substrate availability and soil physical properties (e.g., temperature, pH, moisture content, oxygen availability) [9]. Soil water content influences N_2O emission by affecting soil oxygen concentrations and by altering gas diffusivity rates [6,10]. Irrigation, which increases the soil water content, therefore has a direct influence on N_2O emissions. However, by affecting plant growth, irrigation may also indirectly reduce N_2O emissions as it may increase plant N requirements [11,12]. Different fertilizer sources may also affect N_2O fluxes, with manure sometimes producing less N_2O than synthetic inorganic fertilizers because of the higher proportion of organic N [13], although the addition of the labile C together with the N often increases N_2O fluxes from organic fertilizers particularly if initial soil C concentrations are low [14,15].

Methane (CH_4) and carbon dioxide, CO_2 , are major GHGs emitted from agricultural croplands. Although, CH_4 is mainly from animal production systems such as manure and animal waste handling, a significant amount is emitted from croplands [16]. Methanogenic bacteria anaerobically produce CH_4 in deep layers of soils with limited oxygen which later diffuses to upper layers and is consequently released to the environment. In other circumstances, CH_4 produced is oxidized to CO_2 by methanotrophic bacteria and the balance between production and consumption leads to negative or positive CH_4 flux. CO_2 from soil and plant respiration has been of global concern especially after it was shown that the ability of terrestrial sinks is greatly reduced under increasing emissions [17]. Fossil fuels and other human activities have increasingly contributed to CO_2 emissions accounting for 75% of total CO_2 emissions. Agricultural activities and land use change together account for the rest [18] where deforestation is the major factor [19].

Currently, Kenya has 8 million smallholder farms that account for approximately 78% of agricultural production [16]. However, there are few empirical studies that have measured GHG fluxes in SSA in general and on smallholder farms in particular [14,20]. This leaves the impact of SSA smallholder agriculture on climate highly uncertain [21], resulting in large uncertainties for emission rates and a lack of clarity of what constitutes appropriate mitigation measures.

To address this lack of information on how irrigation, increased use of organic and inorganic fertilizer on smallholder Kenyan farms may impact GHG fluxes, we measured GHG fluxes for three growing seasons from maize plots receiving different amounts and types of N on an experimental farm set up to represent a typical smallholder farm in the uplands of central Kenya. We hypothesized that fertilized treatments would have higher emissions than the 0-N control, that combining manure with mineral fertilizer would decrease emissions compared with mineral fertilizer alone, and that irrigation in combination with mineral fertilizer would increase the emissions compared with mineral fertilizer alone.

2. Materials and Methods

2.1. Site Description

The study site was located on the campus of the Jomo Kenyatta University of Agriculture and Technology at $1^\circ 05' \text{ S}$ and $37^\circ 0' \text{ E}$ in Kiambu County, Central Kenya Figure 1. The area has a subtropical highland climate (cwb under the Koppen classification) with

two rainy seasons per year. The first rainy season runs from March to May (Long rains) and the second from August to October (Short rains). The presence of two rainy seasons means that there are also two cropping seasons per year; the first cropping season runs from March to September and the other from November through February. Average annual rainfall and air temperature (from 2016 to 2018) at the Kenya Agriculture and Livestock Research Organization Thika weather station (18 km southwest of the experimental site) were 825 mm and 20.4 °C, respectively. Rainfall and air temperature were measured using an automated weather station (Campbell Scientific Inc., ET107, Logan, UT, USA) installed at the site in November 2016. The field had been pasture with natural vegetation and scattered bushes similar to that of a savannah ecosystem prior to 2014, after which it was converted into a long-term trial examining the effects of different fertilizers types and rates on maize yields. Since 2014, the field has been continually cropped with maize, with two seasons annually. Prior to greenhouse gas experiment, initial soil conditions were measured (Table 1).




Figure 1. Map showing the study area.

Table 1. Initial soil conditions measured prior to the start of greenhouse gas measurements.

Soil Type	Chromic Vertisol
Silt content	26%
Clay content	62%
Available nitrogen (N)	374 mg/kg
SOM	2.98%
Total nitrogen	0.55%

We measured fluxes from five treatments, replicated in three blocks. The treatments used were as follows: 1. Control plot with no added N fertilizer (C); 2. Plots fertilized with diammonium phosphate (DAP) during planting with an additional topdress of urea 1 month after planting (30% of the fertilizer was applied at planting while 70% was applied as topdress), for a total of 100 kg N ha⁻¹ (Min); 3. The same as Min, but with irrigation equivalent to 10 mm precipitation every three days (MI); 4. Min + 40 kg N ha⁻¹ added as manure (MM)—total applied N = 140 kg; and 5. The same as Min but intercropped (MB) with faba beans (*Phaseolus vulgaris*). The plots were arranged as a randomized complete block design with the plots measuring 8 × 10 m and a buffer zone of 1 m between the plots and between the blocks.

Farm management activities were kept as similar as possible between growing seasons although there were minor differences due to weather and other technical factors. Planting was completed by hand along rows demarcated by sisal rope with an alternating inter-row spacing of 40 and 70 cm. Two seeds were placed in planting holes (approximately 2.5 cm deep) that were made in the ground 30 cm apart using a sharp stick. Intercropped plots

had similar spacing but with two alternating rows of maize and beans after each “wide” spacing. The plots were thinned (i.e., the smaller maize plant removed) prior to topdressing, and then weeded using handheld hoes. In cases where germination of both seeds failed, both plants in the adjacent holes were retained so that a constant plant density per plot was achieved. During the first season, DAP fertilizer and dry solid cattle manure (total N = 3.06%, P = 0.339%, K = 2.31%, and TC = 22.52%) were applied on 4 November 2016 with the maize planted the following day. Urea was applied on 22 December and the maize was harvested on 28 February 2017. Due to intense rains that caused waterlogging in some plots, planting for the second cropping season was delayed. The plots were seeded on 22 April 2017 with fertilizer and manure applied on 3 and 11 May 2017, respectively. Urea was applied on 17 June 2017 and the plots harvested on 12 September 2017. For the third growing season, plots were planted on 5 April 2018, with fertilizer applied the following day (basal application), topdressed on 28 May 2018, and harvested between 7 and 10 August 2018. Overhead irrigation was completed periodically (4 times in season 1 and 3 times for seasons 2 and 3) for all plots when the maize showed signs of extreme water stress (i.e., to avert crop failure) while the treatment with supplemented irrigation (MI) was watered by a watering can routinely without plants having to show water stress at an interval of 3–4 days with each equal to 10 mm of rainfall.

2.2. Chamber Design

Greenhouse gas fluxes were estimated using manually sampled non-flow through, non-steady state chamber methodology [22]. Briefly, six opaque plastic bases were installed 7.5 cm into the ground in each treatment immediately after tillage and left in the field for the entire growing season. The height of the chamber was measured from the outside to ensure slight changes caused by soil compaction were captured. During deployment, vented and ventilated opaque plastic lids ($25 \times 35 \times 12.5$ cm) were fastened to the bases with clamps to ensure an airtight seal as shown in Figure 2. All openings were lined with a rubber septum to ensure the chamber was airtight.




Figure 2. Schematic diagram showing the chamber design. Illustration (a) shows a three-dimensional hollow base and a top with a top while illustration (b) shows a transverse section of the base and the top cover.

2.3. Gas Sample Collection

Gas samples were collected between 9:00 am and 11:00 am to minimize temperature changes during chamber deployment and because fluxes measured during this time better represent daily mean fluxes [23]. The top of the chamber was also lined with a reflective duct tape to reduce temperature changes. Gas samples were drawn at 10 min intervals beginning at T_0 (immediately after closing the gas chamber) and ending at T_4 (30 min after closing the gas chamber). During each sampling, 20 mL of gas was drawn with a propylene syringe fitted with leuer-lock from each of the two replicates per plot and all 40 mL was injected into a 20 mL pre-evacuated glass vial for storage [24]. Air temperature was measured directly using a thermometer inserted into an air-tight port on top of the chamber. Sampling occurred twice a week during the rainy season and for the two weeks following fertilization and weekly during the rest of the growing seasons [25]. Sampling

was conducted from 8 November 2016 to 28 February 2017 for growing season 1, 5 May to 15 September 2017 for growing season 2, and 13 April to 24 July 2018 for growing season 3.

2.4. Greenhouse Gas Measurements

Gas concentrations of the three greenhouse gases, N₂O, CO₂ and CH₄, were determined using SRI gas chromatographs (8610C; SRI) at the Mazingira Centre at the International Livestock Research Institute (ILRI) in Kabete, Nairobi Kenya. Flame ionization detector (FID) was used in the case of CH₄ and CO₂ (after the sample passed through a methanizer) while an electron capture detector (ECD) was used for N₂O. Hayesep-D packed columns were used in operating the chromatograph and N₂ (flow rate of 30 mL min⁻¹) used as a carrier gas for both detector lines. Oven temperature was at 65 °C while that of the methanizer and ECD was set at 350 °C [21]. The peak areas from the GC were converted to concentrations by comparing peak areas of the samples to areas of standard gases with known CO₂, CH₄, and N₂O concentrations.

2.5. Flux Calculations

Gas fluxes were calculated using Equation (1) after estimating a linear rate of change in the mixing ratios.

$$Fr = \Delta (C:Rt) \times M/22.41 \times h \times P/(273 + T) \times 1013 \times 60/100 \quad (1)$$

where *Fr* is the flux rate (mg m⁻² h⁻¹), *C* = calculated concentration (CH₄/CO₂—ppmv; N₂O—ppbv), *Rt* is the relative time, *M* is the molecular mass of each of the gases (g), *h* is chamber height (cm), *T* is temperature in chamber during sampling (°C), 22.41 is ideal gas volume at room temperature, *P* is air pressure (mb), 60 is time conversion factor from minutes to hours, 100 is chamber length conversion factor, 1013 is air pressure at standard conditions (atm).

To calculate daily flux rate, the equation below was used.

$$DFr = Fr \times 100 \times 24 \text{ (kg ha}^{-1} \text{ day}^{-1}\text{)} \quad (2)$$

where *DFr* is the daily flux rate, *Fr* is the hourly flux rate and the 100 is to convert from mg m⁻² to kg ha⁻¹.

2.6. Soil Analysis

Soil samples from each plot were collected (once every 4 weeks for seasons 1 and 3, bi-weekly for season 2) using a regular 5 cm wide soil auger to a depth of 10 cm. Plastic zip-lock bags were used to store the samples, which were then transported in a cooler box to a laboratory for nitrate (NO₃⁻) and ammonium (NH₄⁺) extraction within 24 h. For the extraction, 20 g of fresh soil was mixed with 100 mL of 2 M KCl by shaking for 30 min at 175 rpm on an orbital shaker. The mixture was then allowed to settle for one hour, after which it was centrifuged at 3000 rpm for 10 min. The supernatant was then filtered into a vial and refrigerated (4 °C) until it could be analyzed. NO₃⁻ and NH₄⁺ concentrations were determined by first reducing the NO₃⁻ in the extract to nitrite by use of cadmium granules coated by copper. Sulphanilic acid and 5-amino-2-naphthalene sulphonic acid were used in diazotization to produce a colored dye which was quantitatively measured using a photometric analyzer (Aquakem 200: Thermo Scientific, Waltham, MA, USA). Standards of known concentrations analyzed alongside the samples were used to calculate the concentration of NO₃⁻-N and NH₄⁺-N in the soil. Soil NO₃⁻ and NH₄⁺ intensities were then calculated using the time-weighted sum of soil NO₃⁻-N and NH₄⁺-N concentrations [25].

Soil moisture content was estimated by drying approximately 30 g of the soil at 105 °C for 24 h with the weight change from the fresh soil weight expressed as a percentage of the final dry soil's weight [21]. Soil carbon (%) was calculated from measured soil organic matter using van Bemmelen factor (0.58)

2.7. Statistical Analysis

Cumulative flux data (model residuals) were checked for normality using SPSS (Shapiro–Wilk) and was normal at 95% confidence interval without transformation. This normality test was used since our sample size was <50 hence it had high accuracy of detecting non-normality of data and guided the choice of further methods of analysis. A general linear model was used where treatments were fixed factors, seasons were set as random factors and cumulative flux as the dependent variable. Least Significant Difference (LSD) was used as a post hoc test as all variances were equal. Regression models and Pearson's correlation were used to check the significance of measured variables (soil moisture content, NO_3^- and NH_4^+ concentrations and intensities, and plant biomass) to emissions. All analyses were completed using Windows SPSS version 20.0 at significance level of <0.05 unless stated otherwise.

3. Results

3.1. Environmental Factors

The first cropping season (November 2016–March 2017) received a total of 75 mm of precipitation and was the driest of the three seasons. Season 2 (April–August 2017) received a total of 303 mm of rainfall while the third season (April–August 2018) received 599 mm, the highest amount of precipitation for all the three seasons (Figure 3). The large amount of precipitation during season 3 caused flooding of some of the plots and delayed planting. Much of the precipitation in seasons 2 and 3 was during the first 2 months. Air temperatures were similar across the three seasons, with average daily maximums of 24 °C and average daily minimums of 13 °C. February was the hottest month while July–August was the coolest (Figure 3).

Soil moisture content showed no difference between treatments during the three growing seasons although it did correlate with rainfall events as higher moisture content was observed soon after rainfall episodes. The range was from 19 to 42% for season 1, 26–42% for season 2, and 23–61% for season 3 for all the treatments. The average soil moisture content was lowest in season 1 < season 2 < season 3 (Figure 3) ($p = 0.05$). During heavy rainfall episodes in seasons 2 and 3, ponded water formed on MB treatment and remained on the soil surface for 2–3 days. Initial soil C concentrations were highest in the control treatment and lowest in the MB treatment (Table 2). Total N concentrations in the soil followed an inverse pattern to the soil C concentrations with the C treatment having the lowest total N concentrations and the MB (along with the Min) treatment having the highest total N concentration which is in contrast to C:N ratios in undisturbed soils [26] (Table 2).

Table 2. Soil organic carbon (%) and soil nitrogen (TN) measured using samples collected from maize plots with different fertilizer treatments at the research farm at Jomo Kenyatta University Kenya during growing season 2.

	Total Soil Nitrogen	Soil Organic Carbon
C	0.13 ± 0.005 ^a	1.85 ± 0.15 ^a
Min	0.16 ± 0.006 ^a	1.76 ± 0.17 ^a
MI	0.15 ± 0.005 ^b	1.71 ± 0.05 ^b
MM	0.14 ± 0.011 ^b	1.72 ± 0.04 ^c
MB	0.16 ± 0.005 ^a	1.61 ± 0.06 ^d

Note: C = control (no N fertilizer applied); Min = mineral fertilizer (100 kg N ha⁻¹) applied; MI = Min + irrigation; MM = Min + additional 40 kg N ha⁻¹ added as cattle manure; MB = maize intercropped with Fava beans. Different letters indicate significance at 0.05.




Figure 3. Average daily rainfall and gravimetric soil water content measured during gas sampling to a depth of 20 cm during three maize growing seasons: Season 1 (November 2016–February 2017), season 2 (April–September 2017), and season 3 (April 2018). Vertical arrows indicate supplemented overhead irrigation.

For the fertilized plots, the soil NO_3^- concentrations increased following fertilization (mean of $17.5 \text{ mg N kg}^{-1}$ soil) and decreased gradually through the growing season to a mean concentration of approximately 10 mg N kg^{-1} soil at the end of the growing season. Soil NH_4^+ concentrations averaged 1.7 mg N kg^{-1} soil at the start of the season and 1 mg N kg^{-1} soil at the end of the season with no discernible pattern (Figure 4). For both the NO_3^- and the NH_4^+ concentrations, the increases after fertilization were greater during season 2 than in either season 1 or 3.




Figure 4. Soil nitrate (mg kg^{-1} soil), (a–c) for seasons 1, 2, and 3, respectively, and soil ammonium (mg kg^{-1} soil) (d–f) for seasons 1, 2, and 3, respectively, measured during gas sampling to a depth of 20 cm.

3.2. CH_4 Fluxes

Growing season 1 began with all treatments having an average emission rate of 0.3 mg h^{-1} except for the Min treatment which had -1.2 mg h^{-1} uptake. Average flux rates across all treatments were low (about $0.01 \text{ mg CH}_4\text{-C m}^{-2} \text{ h}^{-1}$) with 69% of the fluxes being negative (i.e., uptake) (Figure 5). Growing season 2 also had primarily negative fluxes with 59% of measurements being uptake, and fluxes averaging $-0.01 \text{ mg CH}_4\text{-C m}^{-2} \text{ h}^{-1}$. Season 3 had the lowest number of negative fluxes (52%) with emission rates as high as $0.12 \text{ mg CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ whereas uptake rates were quite low peaking at $0.03 \text{ mg CH}_4\text{-C m}^{-2} \text{ h}^{-1}$. Generally, high emission rates were recorded between the first week of May and the first week of June when many of the plots were flooded while the rest of the season had average emissions of -0.02 – $0.02 \text{ mg CH}_4\text{-C m}^{-2} \text{ h}^{-1}$.




Figure 5. CH₄ (CH₄-C mg ha⁻¹ h⁻¹) flux rate measured from spontaneous gas measurements for three different seasons: season 1 (a), season 2 (b), and season 3 (c).

Cumulative CH₄ fluxes ranged from -1.83 to -0.04 kg CH₄-C ha⁻¹ for season 1, -1.6 to 0.33 kg CH₄-C ha⁻¹ for growing season 2, and -0.19 to 0.24 for growing season 3 with the Min treatment having the highest uptake in two of the three seasons (Figure 6). Soils were CH₄ sinks and the rate of uptake changed from season 1 to season 3 with the highest uptake recorded in season 1. During the second season, four out of five treatments had negative flux rates measured over a period of 4.5 months (Figure 5). The order of cumulative fluxes for the three combined seasons was MM > MB > MI > C > Min. There was a significant correlation between mean seasonal soil moisture content and cumulative seasonal fluxes ($p = 0.038$). However, there were no correlations between soil moisture content and instantaneous CH₄ fluxes for any of the three seasons.




Figure 6. CO₂ (CO₂-C mg ha⁻¹ h⁻¹) flux rates from maize plots with different nitrogen (N) fertilizer treatments and a no N control (C) over three crop growing seasons. Continuous line arrows indicate basal fertilizer application while dashed line arrows indicate top dressing.

3.3. CO₂ Fluxes

Relatively low peaks in CO₂-C emission rates were observed at the beginning and end of each of the three growing seasons (Figure 6). Initial fluxes for growing season 1 ranged between 20 and 40 mg CO₂-C m⁻² h⁻¹ then rose steadily to a maximum of 160 mg CO₂-C m⁻² h⁻¹ in between 14 and 22 November 2016. The flux rates then decreased, remaining below 100 mg CO₂-C m⁻² h⁻¹ for the rest of the season. Season 2 followed a similar trend but had higher measurements initially, increasing to almost 500 mg CO₂-C m⁻² h⁻¹ in mid-May before decreasing rapidly and returning to fluxes of about 20 mg CO₂-C m⁻² h⁻¹ for most of the growing season. Growing season 3 showed a different pattern, with emissions initially ranging from about 80 to 250 mg CO₂-C m⁻² h⁻¹ before decreasing, only to increase again in late April and again in early June when they peaked at about 400 mg CO₂-C m⁻² h⁻¹ before returning to an average of 25 mg CO₂-C m⁻² h⁻¹ at the end of the season (Figure 6).

The lowest cumulative seasonal flux measured was 1246 kg ha⁻¹ in the C treatment during growing season 3 while the highest was 2054 kg ha⁻¹ in Min during growing season 2. Cumulative flux for the three seasons ranged from 4100 to 5543 kg CO₂-C ha⁻¹. Cumulative CO₂ fluxes tended to be lower in the MB treatment than in the other treatments with no consistent differences between the other treatments (Figure 7). There were no significant correlations between flux rates and either soil moisture ($p = 0.431$) content or yield ($p = 0.503$)




Figure 7. Cumulative CH₄ (CH₄-C kg ha⁻¹), CO₂ (CO₂-C kg ha⁻¹), and N₂O (N₂O-N kg ha⁻¹) with standard errors fluxes over the three growing seasons sampled from different plots with different fertilizer and manure/irrigation treatments. Bars with different letters indicate significant differences among the treatments (Duncan’s test 0.05 significance level).

3.4. N₂O Fluxes

The N₂O flux rates during season 1 increased shortly after the initial fertilizer application (4 November 2016), with the highest rates measured in late November (Figure 8). Rates remained elevated until early December 2016, when they decreased to less than 10 µg N₂O-N m⁻² h⁻¹ until the end of the sampling period. Season 2 N₂O flux rates followed a similar pattern, peaking shortly after fertilization before returning to baseline rates approximately 1 month later (Figure 8). However, during season 2, N₂O flux rates increased in the MM and MI treatments to almost 100 µg N₂O-N m⁻² hr⁻¹ on 17 August 2017. The first measurement of season 3, which was after basal fertilization, captured a declining peak with MB and MI maintaining relatively higher flux rates as compared to the other treatments. Emissions in season 3 had peaks other than those observed immedi-

ately after fertilization, with high emissions observed a few days after high precipitation episodes. The flux rates for all treatments except C increased after topdressing in late May, with the flux rates declining over the subsequent weeks to baseline and remaining low until the end of the season. Cumulative fluxes ranged between 0.25 and 0.48 kg ha⁻¹ during season 1, 0.89–1.69 kg ha⁻¹ during season 2, and 0.72–2.45 kg ha⁻¹ during season 3. Cumulatively, growing season 1 had the lowest emissions while growing season 3 had the highest (Figure 7). Cumulative emissions from MI were 2.24 kg ha⁻¹ greater than emissions from C treatment ($p = 0.046$) while the other treatments were similar. The emission factor was highest in MM treatment for seasons 1 and 2 whereas it was the second highest in season 3 (Figure 7).




Figure 8. N₂O (N₂O-N μg ha⁻¹ h⁻¹) flux rate from maize plots with different nitrogen (N) fertilizer treatments and a no N control (C) for crop growing season 1 (a), season 2 (b) and, season 3 (c). Continuous line arrows indicate basal fertilizer application while dashed line arrows indicate top.

There was a correlation between soil moisture content and cumulative N₂O emission ($F = 58.2, p = 0.02$). All the treatments had Emission Factors (EF) less than 1% during all three growing seasons (Table 3). Instantaneous flux rates showed no correlation with soil nitrate concentrations, yield, nor soil moisture content for all seasons. Nitrate intensity was not significantly correlated to cumulative emissions at $R^2 = 0.216$ ($p = 0.213$), 0.120 ($p = 0.173$) for seasons 1 and 2, respectively, whereas there was a significant correlation in season 3 ($R^2 = 0.520; p = 0.021$).

Table 3. N₂O Emission Factor which is a percentage of fertilizer applied released as N₂O emission (IPCC, 2006) for different fertilizer treatments and management depicting small scale farming systems in Central Kenya.

	Season 1	Season 2	Season 3	Mean
Min	0.23 ± 0.02	0.49 ± 0.01	0.08 ± 0.02	0.27 ± 0.12
MI	0.13 ± 0.03	0.39 ± 0.01	0.17 ± 0.03	0.23 ± 0.08
MM	0.14 ± 0.06	0.68 ± 0.03	0.93 ± 0.15	0.58 ± 0.23
MB	0.06 ± 0.06	0.12 ± 0.31	0.84 ± 0.26	0.34 ± 0.25

4. Discussion

4.1. CH₄ Emissions

CH₄ emissions are important in estimation of total GHG emissions since CH₄ is produced anaerobically by methanogenic microbes and later it may be oxidized to CO₂ by methanotrophic bacteria under aerobic conditions. Thus, CH₄ flux is important since it influences CO₂ flux [27]. CH₄ is also an important greenhouse gas since it has a global warming potential (GWP) of 25–28 in relation to CO₂ [28]. Methane flux rates were similar to a previous study completed in the region [13] where emissions from maize fields with different fertilizer and manure treatments ranged from -2.6 to 5.8 kg ha⁻¹ over a 6-month period. The oxidation of CH₄ during the two drier seasons (i.e., seasons 1 and 2) was consistent with prior studies that showed how upland soils tend to be CH₄ sinks when dry, and that the sink strength is much diminished (or the soils may even become sources) when wet, as seen in season 3 [14,29,30]. Increased soil moisture content can reduce diffusion of atmospheric CH₄ into the soil, thus reducing the ability of the soil to act as a CH₄ sink. However, fertilizer addition can also inhibit CH₄ oxidation, especially where fertilization results in high NH₄ concentrations [31]. However, the low NH₄ concentrations measured in this study suggest that fertilization had little effect on total CH₄ emissions. The steady increase in cumulative CH₄ flux as the mean soil moisture content increased from season 1 to season 3 suggests that soil moisture was the main driver determining the CH₄ fluxes in this study.

A number of studies have reported CH₄ oxidation from various ecosystems ranging from grasslands to croplands [32,33]. Whereas these studies were carried out in different regions, the crop growing season of [34] occurred in spring when prevailing atmospheric conditions were similar to the conditions of this study. The mechanism of CH₄ oxidation/emission greatly depends on SWC, temperature and soil conditions. SWC was positively correlated with CH₄ fluxes. Extremes of soil water content limited methanotrophic activities suppressing the CH₄ fluxes. Optimal SWC meant vibrant methanotrophic activity which led to higher CH₄ oxidation in season 2 compared with season 3 whose lower oxidation and emissions rates could be explained by limited gas diffusion at higher SWC [30,35]. Our study showed that atmospheric temperature was weakly correlated with CH₄ oxidation explaining only 20% of the CH₄ oxidation in agreement with [36] who also found that temperature was a minor co-factor in CH₄ oxidation.

4.2. CO₂ Emissions

Cumulative soil CO₂ emissions recorded in this study were at the low end of an earlier study carried out in maize with agroforestry [29] where emissions ranged from 1.8 to 2.3 Mg CO₂-C ha⁻¹ over a 99 d period, and also at the low end of the range of studies in Kenya and Tanzania where emissions ranged from 2.8 to 15.9 Mg CO₂ ha⁻¹ yr⁻¹ [14,21]. However, the emissions were at the high end of another study from Zimbabwe where 122d cumulative CO₂-C emissions ranged from 0.7 to 1.6 Mg CO₂ ha⁻¹ [13]. The flux rates were generally low at the beginning of the seasons (average of 70 mg CO₂ m⁻² h⁻¹) likely owing to the low root respiration of the young plants and increased during the plant growing stage (with average peaks of above 200 mg CO₂ m⁻² h⁻¹). The last sampling shows the lowest observed flux rate (50 mg CO₂ m⁻² h⁻¹) since the soil tended to be dry and the plants were at senescence stage with minimal root respiration [37]. The slight increase in soil respiration at the end of the first season can be attributed to increased microbial activity following the incorporation of the crop residues along with the arrival of some light rains (and increased soil moisture content) that occurred at the end of the season. Together, microbial activity and plant root respiration were the regulators of CO₂ emissions in this experiment.

We expected the MM treatment to experience the highest soil CO₂-C emission rates for all the three seasons because of the additional C applied as manure. While this was the case during season 3, this was not the case in seasons 1 and 2. Interestingly, treatment C had the highest initial soil C concentrations (Table 2) as well as similar or even higher

emission rates as compared to the other treatments which had C inputs. Over time, it is expected that these soil C concentrations will decrease compared to the other treatments as there are no external C additions to the plots, and as the lower biomass yield (Table 4) will mean less C returned to the soil via residue incorporation. Reductions in soil C in the control plots may explain why soil CO₂-C fluxes decreased steadily from seasons 1 to 3 in the control plots, whereas the other plots did not show this steady decline in emissions.

This further supports the observation that plots with higher root respiration/biomass have more emissions [38]. Soil C:N ratio reduced with increase in N input which is consistent with earlier studies [39] where <100 kg N input resulted in C:N of 14 whereas >200 kg of N resulted in a ratio of <11 consistent with our findings. Additional N through fertilizations adds plant available N which at times is not utilized by the plant and ends up being mineralized. This coupled with additional nitrogen in the form of plant residue increases soil nitrogen and consequently reduces the ratio as seen in Table 2. The consistently low emissions from the MB treatment, meanwhile, could be related to the wider spacing of maize plants in MB as two consecutive rows were alternated with two lines of beans. This may have increased water loss through evaporation as beans have less ground cover thereby reducing the soil moisture and slowing organic matter decomposition and root respiration. Additionally, the lower plant density may have reduced root densities resulting in lower root respiration, while lower biomass results in reduced C returned into the soil (Table 2), which could also cause lower CO₂ emissions.

Table 4. Mean (± 1 SD) above ground biomass (g) of 3 plants from maize plots near Nairobi Kenya with different fertilizer treatments taken during the mid-growing period of each of the three seasons. Different letters indicate significance at 0.05.

	Season 1	Season 2	Season 3
C	80.0 \pm 0.64 a	81.7 \pm 0.36 a	79.7 \pm 1.30 a
Min	95.0 \pm 0.11 a	85.8 \pm 1.30 b	82.3 \pm 0.80 a
MI	94.2 \pm 2.80 a	88.8 \pm 1.30 a	85.6 \pm 0.59 a
MM	94.3 \pm 1.50 b	91.9 \pm 2.30 a	90.3 \pm 0.38 a
MB	84.5 \pm 0.95 a	81.8 \pm 1.70 ab	89.3 \pm 2.40 a

4.3. N₂O Emissions

Cumulative N₂O emissions were consistent with earlier studies where measured cumulative emissions from maize fields in Kenya ranged from 0.1 to 4.1 kg N₂O-N ha⁻¹ over a 4-month period [40] and where cumulative emissions from a number of crop types in Kenya and Tanzania ranged from 0.4 to 3.9 kg N₂O-N ha⁻¹ yr⁻¹ [18]. However, with the exception of growing season 1 (with its overall low emissions), the emissions were higher than other studies from east Africa [13,41] where emissions were 0.62–0.81 kg N₂O-N ha⁻¹ (over a 6-month period) and 0.1–0.5 kg N₂O-N ha⁻¹ (over a 5-month period), respectively.

The lower emission rates observed in the control were consistent with expected results where fertilized plots were expected to have higher emission rates than the unfertilized plot [42]. The start of each season had high emission rates that were caused by the high N concentrations due to the added fertilizer and the increased soil moisture content due to high rainfall (seasons 2 and 3) or irrigation in the case of season 1. During season 1, the N₂O released was likely from nitrification as the low amount of NO₃⁻, relatively higher NH₄⁺, and drier soil conditions were ideal for nitrification to take place [9]. The first few days of each season were also characterized by soil disturbances caused by ploughing and planting which may have created greater soil aeration and thus enhancing emission rates [43]. Soil NO₃⁻ in MB treatment was low (Figure 4), which might be due to uptake or rapid immobilization [7] by either the growing crop or the microbial community, likely reducing the emissions. Waterlogging which caused ponded water on the soil surface and increased soil moisture content at the start of seasons 1 and 2 (Figure 3) observed in the site with MB treatment affected most suggest the possible reason of low N₂O emission from the MB treatment as compared to others.

The strong correlation between cumulative N_2O emission and NO_3^- intensities in fertilized plots indicates a direct influence of fertilizer application on N_2O emissions consistent with our expectations [44,45]. High NO_3^- concentrations in the soils provided the N substrate required for the production of N_2O through microbial denitrification [46]. The positive correlation between mean soil moisture content and cumulative emissions was likely because denitrification requires anaerobic conditions, and soils will have more anaerobic microsites at higher soil moisture content resulting in high N_2O emissions [47].

Emission rates between different treatments were not consistent across growing seasons as the seasons were short and each season presented different weather conditions. However, a trend could be deduced from the average emissions over the three seasons. Zero N input resulted in lower emissions whereas optimal soil moisture through supplemented irrigation and increased N levels through manure addition in the soil led to higher emissions from the MI and MM treatments, consistent with previous research [9]. Similar levels of emissions in MB and Min treatments showed that type of plant cover did not affect the emission rates.

Lower precipitation in season 1 and the resulting low soil moisture content resulted in lower emissions for season 1 as compared to seasons 2 and 3, likely because denitrification, which is generally the primary source of N_2O emissions from agricultural soils [48,49], generally requires soil water content above 60% WFPS [50]. Consistent with previous studies, emissions increased during or slightly after precipitation episodes [14,51] as soil NO_3^- concentrations tend to increase after re-wetting of soils due to temporary mobilization of N that accumulated over the dry period [52]. This is characterized by an increase in soil microbial populations some of which are involved in biogeochemical cycles [53]. The high emission rates experienced at the start of seasons 2 and 3 was due to fertilization and consequently, higher inorganic N levels. High rainfall during this period might cause anaerobic conditions suitable for denitrification. The increase in N_2O emissions after the first (fertilization peak) was likely due to the combined effect of nitrification and denitrification which may have occurred simultaneously [54,55]. The reduced soil moisture content also allows for more rapid gas diffusion that results in less opportunity for N_2O to be reduced to N_2 and hence greater N_2O emissions [10]. Season 3 emissions were similar to season 2 with high emission rates observed during the periods after basal fertilization and top dressing as well as additional peaks seen around 11 May due to rainfall episodes. Consistently higher soil moisture content during season 3 likely caused slightly lower emissions than season 2, perhaps because the higher soil water content would result in more complete denitrification to N_2 [9,56].

Although there was no significant difference among different treatments for instantaneous fluxes, cumulative fluxes provided an overall picture from which significant differences could be seen. Spontaneous greenhouse gas measurements were noisy in some cases and did not provide a distinct trend between different treatments. The erratic nature of spontaneous emissions could be attributed to the extreme weather conditions especially high rainfall which caused a variation in distal and proximal emission drivers hence affecting the stability of emissions [9,57]. Increased moisture content affects all the three gases which were measured differently, and it was a key factor which was constantly changing. High rainfall episodes which were far apart made the soil shift from extremely dry to extremely moist. This also caused shifts to occur from denitrification to nitrification which produce different amounts of N_2O [58]. The decomposition rate which is critical in CO_2 emissions may have also been affected by the moisture/rainfall. Oxidation of CH_4 which is reliant on oxygen levels in the soil was also greatly influenced by the erratic rainfall. These factors together may have caused the erratic spontaneous fluxes. Long-term effects can easily be observed on specific treatments making long-term experiments more suitable to study field emission experiments in the region as indicated by the cumulative fluxes.

5. Conclusions

This study shows that agricultural soils from the dry uplands of central Kenya were CH₄ sinks when conditions were dry. However, under flooded conditions they have the capacity to become sources. Cumulative soil N₂O fluxes were correlated in two out of the three growing seasons with mean soil moisture content where emissions increased with N input. Precipitation was a factor where low precipitation led to low emissions, moderate precipitation had high emissions, while high precipitation caused moderate to high emissions. Therefore, as the region moves to intensify crop production through increased N input and irrigation, CH₄ and CO₂ fluxes are likely to have minimal increases while N₂O emissions are expected to increase but will remain below <1% of N applied if application rates remain below 100 kg N ha⁻¹ season⁻¹. However, more long-term studies need to be conducted for more conclusive studies on annual emissions and estimation of regional emission rates.

Author Contributions: Experimental design: P.S.M., X.L. and S.C. Data collection: P.S.M. Laboratory analysis: P.S.M. and W.L. Statistical analysis: P.S.M. and G.G. Interpretation of results: D.E.P., P.S.M. and Y.W. Supervision of experiments: C.H., D.E.P. and D.M. Writing of first draft of the manuscript: P.S.M. and D.E.P. Preparation of final draft: All authors. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by International Partnership Program of Chinese Academy of Sciences, grant number 151542KYSB20200017.

Data Availability Statement: Not applicable.

Acknowledgments: The authors greatly acknowledge Mazingira Center, ILRI for providing the facilities for laboratory analysis of the samples. We also acknowledge Sino-Africa Joint Research Project for funding this study (SAJC201603).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Crippa, M.; Solazzo, E.; Guizzardi, D.; Monforti-Ferrario, F.; Tubiello, F.N.; Leip, A. Food systems are responsible for a third of global anthropogenic GHG emissions. *Nat. Food* **2021**, *2*, 198–209. [[CrossRef](#)]
2. Tubiello, F.N.; Salvatore, M.; Ferrara, A.; Rossi, S.; Biancalani, R.; Federici, S.; Jacobs, H.; Flammini, A. Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks. *ESS Work. Pap.* **2014**, *2*, 4–89.
3. Sánchez, P.A. Tripling crop yields in tropical Africa. *Nat. Geosci.* **2010**, *3*, 299–300. [[CrossRef](#)]
4. Ariga, J.; Jayne, T.S. Fertilizer in Kenya: Factors Driving the Increase in Usage by Smallholder Farmers. In *Yes, Africa Can: Success Stories from a Dynamic Continent*; World Bank Publications: Washington, DC, USA, 2011; pp. 269–288.
5. FAO. *World Fertilizer Trends and Outlook to 2022*; FAO: Rome, Italy, 2019.
6. Davidson, E.A.; Keller, M.; Erickson, H.E.; Verchot, L.V.; Veldkamp, E. Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides. *Bioscience* **2000**, *50*, 667. [[CrossRef](#)]
7. Norton, J.; Ouyang, Y. Controls and Adaptive Management of Nitrification in Agricultural Soils. *Front. Microbiol.* **2019**, *10*, 1931. [[CrossRef](#)] [[PubMed](#)]
8. Scheer, C.; Fuchs, K.; Pelster, D.E.; Butterbach-bahl, K. ScienceDirect Estimating global terrestrial denitrification from measured N₂O:(N₂O + N₂) product ratios. *Curr. Opin. Environ. Sustain.* **2020**, *47*, 72–80. [[CrossRef](#)]
9. Butterbach-Bahl, K.; Baggs, E.M.; Dannenmann, M.; Kiese, R.; Zechmeister-Boltenstern, S. Nitrous oxide emissions from soils: How well do we understand the processes and their controls? *Philos. Trans. R. Soc. B Biol. Sci.* **2013**, *368*, 20130122. [[CrossRef](#)]
10. Balaine, N.; Clough, T.J.; Beare, M.H.; Thomas, S.M.; Meenken, E.D.; Ross, J.G. Changes in Relative Gas Diffusivity Explain Soil Nitrous Oxide Flux Dynamics. *Soil Sci. Soc. Am. J.* **2013**, *77*, 1496–1505. [[CrossRef](#)]
11. Abegunde, V.O.; Sibanda, M.; Obi, A. The Dynamics of Climate Change Adaptation in Sub-Saharan Africa: A Review of Climate-Smart Agriculture among Small-Scale Farmers. *Climate* **2019**, *7*, 132. [[CrossRef](#)]
12. Ntinyari, W.; Gweyi-Onyango, J.P. Greenhouse Gases Emissions in Agricultural Systems and Climate Change Effects in Sub-Saharan Africa. In *African Handbook of Climate Change Adaptation*; Oguge, N., Ayal, D., Adeleke, L., da Silva, I., Eds.; Springer International Publishing: Cham, Switzerland, 2021; pp. 1081–1105.
13. Mapanda, F.; Wuta, M.; Nyamangara, J.; Rees, R.M. Effects of organic and mineral fertilizer nitrogen on greenhouse gas emissions and plant-captured carbon under maize cropping in Zimbabwe. *Plant Soil* **2011**, *343*, 67–81. [[CrossRef](#)]
14. Pelster, D.; Rufino, M.; Rosenstock, T.; Mango, J.; Saiz, G.; Diaz-Pines, E.; Baldi, G.; Butterbach-Bahl, K. Smallholder farms in eastern African tropical highlands have low soil greenhouse gas fluxes. *Biogeosciences* **2017**, *14*, 187–202. [[CrossRef](#)]

15. Zhou, M.; Zhu, B.; Wang, S.; Zhu, X.; Vereecken, H.; Brüggemann, N. Stimulation of N₂O emission by manure application to agricultural soils may largely offset carbon benefits: A global meta-analysis. *Glob. Chang. Biol.* **2017**, *23*, 4068–4083. [[CrossRef](#)] [[PubMed](#)]
16. Chen-Rui, W.; Yi, S.; Xiao-Ming, Y.; Jie, W.; Jin, Y. Advances of study on atmospheric methane oxidation (consumption) in forest soil. *J. For. Res.* **2003**, *14*, 230–238. [[CrossRef](#)]
17. Liu, S.; Ji, C.; Wang, C.; Chen, J.; Jin, Y.; Zou, Z.; Li, S.; Niu, S.; Zou, J. Climatic role of terrestrial ecosystem under elevated CO₂: A bottom-up greenhouse gases budget. *Ecol. Lett.* **2018**, *21*, 1108–1118. [[CrossRef](#)]
18. Menon, S.; Denman, K.L.; Brasseur, G.; Chidthaisong, A.; Ciais, P.; Cox, P.M.; Dickinson, R.E.; Hauglustaine, D.; Heinze, C.; Holland, E.; et al. *Couplings between Changes in the Climate System and Biogeochemistry*; Lawrence Berkeley National Lab.(LBNL): Berkeley, CA, USA, 2007.
19. Houghton, R. Balancing the Global Carbon Budget. *Annu. Rev. Earth Planet. Sci.* **2007**, *35*, 313–347. [[CrossRef](#)]
20. Kim, D.-G.; Thomas, A.D.; Pelster, D.; Rosenstock, T.S.; Sanz-Cobena, A. Greenhouse gas emissions from natural ecosystems and agricultural lands in sub-Saharan Africa: Synthesis of available data and suggestions for further research. *Biogeosciences* **2016**, *13*, 4789–4809. [[CrossRef](#)]
21. Rosenstock, T.S.; Mpanda, M.; Pelster, D.E.; Butterbach-Bahl, K.; Rufino, M.C.; Thiong’O, M.; Mutuo, P.; Abwanda, S.; Rioux, J.; Kimaro, A.A.; et al. Greenhouse gas fluxes from agricultural soils of Kenya and Tanzania. *J. Geophys. Res. Biogeosci.* **2016**, *121*, 1568–1580. [[CrossRef](#)]
22. Rochette, P. Towards a standard non-steady-state chamber methodology for measuring soil N₂O emissions. *Anim. Feed Sci. Technol.* **2011**, *166–167*, 141–146. [[CrossRef](#)]
23. Alves, B.J.; Smith, K.A.; Flores, R.; Cardoso, A.; Oliveira, W.R.; Jantalia, C.P.; Urquiaga, S.; Boddey, R.M. Selection of the most suitable sampling time for static chambers for the estimation of daily mean N₂O flux from soils. *Soil Biol. Biochem.* **2012**, *46*, 129–135. [[CrossRef](#)]
24. Arias-Navarro, C.; Díaz-Pinés, E.; Kiese, R.; Rosenstock, T.S.; Rufino, M.; Stern, D.; Neufeldt, H.; Verchot, L.; Butterbach-Bahl, K. Gas pooling: A sampling technique to overcome spatial heterogeneity of soil carbon dioxide and nitrous oxide fluxes. *Soil Biol. Biochem.* **2013**, *67*, 20–23. [[CrossRef](#)]
25. Barton, L.; Wolf, B.; Rowlings, D.W.; Scheer, C.; Kiese, R.; Grace, P.R.; Stefanova, K.; Butterbachbahl, K. Sampling frequency affects estimates of annual nitrous oxide fluxes. *Sci. Rep.* **2015**, *5*, 15912. [[CrossRef](#)] [[PubMed](#)]
26. Troeh, F.R.; Thompson, L.M. Soils and Soil Fertility. *Soils Soil Fertil.* **2005**, *7*, 5.
27. Le Mer, J.; Roger, P. Production, oxidation, emission and consumption of methane by soils: A review. *Eur. J. Soil Biol.* **2001**, *37*, 25–50. [[CrossRef](#)]
28. Johansson, D.J.A.; Persson, U.M.; Azar, C. The Cost of Using Global Warming Potentials: Analysing the Trade off between CO₂, CH₄ and N₂O. *Clim. Chang.* **2006**, *77*, 291–309. [[CrossRef](#)]
29. Baggs, E.; Chebii, J.; Ndufa, J. A short-term investigation of trace gas emissions following tillage and no-tillage of agroforestry residues in western Kenya. *Soil Tillage Res.* **2006**, *90*, 69–76. [[CrossRef](#)]
30. Butterbach-Bahl, K.; Papen, H. Four years continuous record of CH₄-exchange between the atmosphere and untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany. *Plant Soil* **2002**, *240*, 77–90. [[CrossRef](#)]
31. Mosier, A.R.; Pendall, E.; Morgan, J.A. Effect of water addition and nitrogen fertilization on the fluxes of CH₄, CO₂, NO_x, and N₂O following five years of elevated CO₂ in the Colorado Shortgrass Steppe. *Atmos. Chem. Phys.* **2003**, *3*, 1703–1708. [[CrossRef](#)]
32. Ussiri, D.A.; Lal, R.; Jarecki, M.K. Nitrous oxide and methane emissions from long-term tillage under a continuous corn cropping system in Ohio. *Soil Tillage Res.* **2009**, *104*, 247–255. [[CrossRef](#)]
33. Ruan, L.; Robertson, G. Initial nitrous oxide, carbon dioxide, and methane costs of converting conservation reserve program grassland to row crops under no-till vs. conventional tillage. *Glob. Chang. Biol.* **2013**, *19*, 2478–2489. [[CrossRef](#)]
34. Rong, Y.; Ma, L.; Johnson, D.A. Methane uptake by four land-use types in the agro-pastoral region of northern China. *Atmos. Environ.* **2015**, *116*, 12–21. [[CrossRef](#)]
35. Steinkamp, R.; Butterbach-Bahl, K.; Papen, H. Methane oxidation by soils of an N limited and N fertilized spruce forest in the Black Forest, Germany. *Soil Biol. Biochem.* **2001**, *33*, 145–153. [[CrossRef](#)]
36. Merino, A.; Pérez-Batallón, P.; Macías, F. Responses of soil organic matter and greenhouse gas fluxes to soil management and land use changes in a humid temperate region of southern Europe. *Soil. Biol. Biochem.* **2004**, *36*, 917–925. [[CrossRef](#)]
37. Czaban, J.; Gajda, A.; Wróblewska, B. The motility of bacteria from rhizosphere and different zones of winter wheat roots. *Pol. J. Environ. Stud.* **2007**, *16*, 301–308.
38. Rochette, P.; Angers, D.A.; Flanagan, L.B. Maize Residue Decomposition Measurement Using Soil Surface Carbon Dioxide Fluxes and Natural Abundance of Carbon-13. *Soil Sci. Soc. Am. J.* **1999**, *63*, 1385–1396. [[CrossRef](#)]
39. Deng, X.; Ma, W.; Ren, Z.; Zhang, M.; Grieneisen, M.L. Geoderma Spatial and temporal trends of soil total nitrogen and C/N ratio for croplands of East China. *Geoderma* **2020**, *361*, 114035. [[CrossRef](#)]
40. Millar, N.; Ndufa, J.K.; Cadisch, G.; Baggs, L. Nitrous oxide emissions following incorporation of improved-fallow residues in the humid tropics. *Glob. Biogeochem. Cycles* **2004**, *18*, 1–9. [[CrossRef](#)]
41. Hickman, J.E.; Palm, C.A.; Mutuo, P.; Melillo, J.M.; Tang, J. Nitrous oxide (N₂O) emissions in response to increasing fertilizer addition in maize (*Zea mays* L.) agriculture in western Kenya. *Nutr. Cycl. Agroecos.* **2014**, *100*, 177–187. [[CrossRef](#)]
42. IPCC. *Summary for Policymakers*; IPCC: Geneva, Switzerland, 2014.

43. Chatskikh, D.; Olesen, J.E. Soil tillage enhanced CO₂ and N₂O emissions from loamy sand soil under spring barley. *Soil Tillage Res.* **2007**, *97*, 5–18. [[CrossRef](#)]
44. Snyder, C.; Bruulsema, T.; Jensen, T.; Fixen, P. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agric. Ecosyst. Environ.* **2009**, *133*, 247–266. [[CrossRef](#)]
45. Shcherbak, I.; Millar, N.; Robertson, G.P. Global metaanalysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 9199–9204. [[CrossRef](#)]
46. Firestone, E.; Davidson, M.K. Microbial basis of NO and N₂O production and consumption in soil. Exchange of trace gases between terrestrial ecosystems and the atmosphere. In *Microbiological Basis of NO and N₂O Production and Consumption in Soil*; Andreae, D.S., Schimel, M.O., Eds.; Wiley: New York, NY, USA, 1989; pp. 7–21.
47. Ssiri, D.; Lal, R. Global sources of nitrous oxide. In *Soil Emission of Nitrous Oxide and Its Mitigation*; Springer: Berlin/Heidelberg, Germany, 2013; pp. 131–175.
48. Li, X.; Sørensen, P.; Olesen, J.E.; Petersen, S.O. Evidence for denitrification as main source of N₂O emission from residue-amended soil. *Soil Biol. Biochem.* **2016**, *92*, 153–160. [[CrossRef](#)]
49. Opdyke, M.R.; Ostrom, N.E.; Ostrom, P.H. Evidence for the predominance of denitrification as a source of N₂O in temperate agricultural soils based on isotopologue measurements. *Glob. Biogeochem. Cycles* **2009**, *23*, 1–10. [[CrossRef](#)]
50. Linn, D.M.; Doran, J.W. Effect of Water-Filled Pore Space on Carbon Dioxide and Nitrous Oxide Production in Tilled and Nontilled Soils. *Soil Sci. Soc. Am. J.* **1984**, *48*, 1267–1272. [[CrossRef](#)]
51. de Carvalho, A.M.; de Oliveira, W.R.D.; Ramos, M.L.G.; Coser, T.R.; de Oliveira, A.D.; Pulrolnik, K.; Souza, K.W.; Vilela, L.; Marchão, R.L. Soil N₂O fluxes in integrated production systems, continuous pasture and Cerrado. *Nutr. Cycl. Agroecos.* **2017**, *108*, 69–83. [[CrossRef](#)]
52. Birch, H.F. Nitrification in soils after different periods of dryness. *Plant Soil* **1960**, *12*, 81–96. [[CrossRef](#)]
53. Leitner, S.; Minixhofer, P.; Inselsbacher, E.; Keiblinger, K.M.; Zimmermann, M.; Zechmeister-Boltenstern, S. Short-term soil mineral and organic nitrogen fluxes during moderate and severe drying–rewetting events. *Appl. Soil Ecol.* **2017**, *114*, 28–33. [[CrossRef](#)]
54. Chen, Z.; Ding, W.; Xu, Y.; Müller, C.; Yu, H.; Fan, J. Increased N₂O emissions during soil drying after waterlogging and spring thaw in a record wet year. *Soil Biol. Biochem.* **2016**, *101*, 152–164. [[CrossRef](#)]
55. Kong, Q.; Wang, Z.-B.; Niu, P.-F.; Miao, M.-S. Greenhouse gas emission and microbial community dynamics during simultaneous nitrification and denitrification process. *Bioresour. Technol.* **2016**, *210*, 94–100. [[CrossRef](#)]
56. Guo, X.; Drury, C.F.; Yang, X.; Reynolds, W.D.; Fan, R. The Extent of Soil Drying and Rewetting Affects Nitrous Oxide Emissions, Denitrification, and Nitrogen Mineralization. *Soil Sci. Soc. Am. J.* **2014**, *78*, 194–204. [[CrossRef](#)]
57. Oertel, C.; Matschullat, J.; Zurba, K.; Zimmermann, F.; Erasmi, S. Greenhouse gas emissions from soils—A review. *Geochemistry* **2016**, *76*, 327–352. [[CrossRef](#)]
58. Klemetsson, L.; Svensson, B.; Rosswall, T. Relationships between soil moisture content and nitrous oxide production during nitrification and denitrification. *Biol. Fertil. Soils* **1988**, *6*, 106–111. [[CrossRef](#)]