

## Article

# Impact of Shale Gas Exploration and Exploitation Activities on the Quality of Ambient Air—The Case Study of Wysin, Poland

Janusz Jarosławski \*, Izabela Pawlak, Jakub Guzikowski and Aleksander Pietruczuk 

Institute of Geophysics Polish Academy of Sciences, 01-452 Warszawa, Poland; izap@igf.edu.pl (I.P.); jguzikowski@igf.edu.pl (J.G.); alek@igf.edu.pl (A.P.)

\* Correspondence: januszj@igf.edu.pl; Tel.: +48-48-22-6915-869

**Abstract:** The continuous two-year monitoring of a set of air pollutants, as well as gases directly related to shale gas exploration processes (methane, non-methane hydrocarbons, carbon dioxide), was carried out at Stary Wiec village in the vicinity (1100 m) of the shale gas wells area in Wysin (Pomeranian voivodeship, north of Poland), covering the stages of preparation, drilling, hydrofracturing and closing of wells. The results of analysis of air pollution data from Stary Wiec and nearby urban and rural stations, over the period 2012–2017 (starting three years before preparations for hydraulic fracturing) indicated that Stary Wiec represents a clean rural environment with an average concentration of nitrogen oxides, carbon monoxide and particulate matter that is one of the lowest in the Pomeranian region. The aim of this study was to explore the range of potential impact of shale gas exploration on local ambient air quality. Analysis of dependence of the concentration level of pollutants on the wind direction indicated that during the drilling period, when the air was coming directly from the area of the wells, nitrogen oxide concentration increased by 13%. Increases of concentration during the hydro-fracturing period, recorded at the Stary Wiec station, were equal to 108%, 21%, 18%, 12%, 7%, 4%, 1% for nitrogen oxide, non-methane hydrocarbons, carbon monoxide, nitrogen dioxide, particulate matter, carbon dioxide and methane. The results of one-minute concentration values for the period 1–4 September 2016 showed a series of short peaks up to 7.45 ppm for methane and up to 3.03 ppm for non-methane hydrocarbons, being probably the result of operations carried out at the area of the wells.

**Keywords:** air pollution; shale gas; atmospheric methane; atmospheric nonmethane hydrocarbons



**Citation:** Jarosławski, J.; Pawlak, I.; Guzikowski, J.; Pietruczuk, A. Impact of Shale Gas Exploration and Exploitation Activities on the Quality of Ambient Air—The Case Study of Wysin, Poland. *Atmosphere* **2022**, *13*, 1228. <https://doi.org/10.3390/atmos13081228>

Academic Editors: Magdalena Reizer, Zbigniew Nahorski and Jerzy Sowa

Received: 21 June 2022

Accepted: 25 July 2022

Published: 2 August 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Natural gas development and exploration can have a significant impact on air quality [1,2]. Shale gas production contributes to emissions into the air, including methane (CH<sub>4</sub>), nonmethane hydrocarbons (NMHC), carbon dioxide (CO<sub>2</sub>), and hydrogen sulfide (H<sub>2</sub>S) [3], although individual stages of gas production generate different sets of air pollutants [4,5]. The first stage of shale gas production (preparation and drilling) engages a large number of diesel-powered engines (e.g., trucks and pumps) that emit mainly nitrogen oxides (NO<sub>x</sub>), volatile organic compounds (VOCs), CO<sub>2</sub>, and fine particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>) [6,7]. Research by Purvis et al. [8] highlights that the preoperational phase, covering 20 weeks of intensive activities related to preparation for hydraulic fracturing, leads to a significant increase of the annual mean of nitrogen oxide (NO) (3-fold) and NO<sub>x</sub> (2-fold) equal to 5.3 ppb and 10.9 ppb, respectively. Hydraulic fracturing processes often result in uncontrolled CH<sub>4</sub> leaks [9]. At the stage of completion of the well, the process of venting shale gas above the surface can result in increased emission of VOCs [10]. The problem of air pollution is linked to direct emission because drilling and hydraulic fracturing may concern not only the borehole area, but the transport of these pollutants to downwind regions, especially those inhabited by people, which is equally important [11]. The measurements performed by Orak et al. [12] in Marcellus Shale (USA) at the turn of 2015 and

2016 indicated significant increase of  $\text{NO}_x$  during the fracturing phase and of  $\text{C}_2\text{H}_6$  (ethane) and  $\text{CH}_4$  for the flowback phase. The values of the 25th percentile equaled 12.5 ppb, 24 ppb and 2.5 ppm for  $\text{NO}_x$ ,  $\text{C}_2\text{H}_6$  and  $\text{CH}_4$ , respectively, which were significantly higher than for other phases.

Shale gas is regarded as a low-carbon fuel, and its exploitation results in lower  $\text{CO}_2$ ,  $\text{NO}_x$ , and sulfur oxides ( $\text{SO}_x$ ) emissions than conventional coal or oil combustion [12–14]. According to Song et al. [15] replacing coal with shale gas in China reduced the mean monthly concentration of  $\text{PM}_{2.5}$  during the time period 2013–2014 by 70%, compared with the same period in 2011–2012. Similar results were obtained by Zhang et al. [16]. Substitution of coal and biomass fuel by shale gas reduced PM emissions in 2017 in China by 123 kt/y. According to Wang et al. [17], the full environmental impact (mainly associated with water consumption and combustion of diesel fuel during the development of the well) of shale gas is 12.5% greater than natural gas. In contrast, replacing fossil fuels (especially coal and biomass) by shale gas could efficiently reduce emission of PM. Among the conventional/unconventional methods of generating electricity and related to the reduction of harmful effects of pollutant emissions on the environment, the investigations with thermal power plants are worth mentioning. Investigations of Abadan Combined Cycle Power Plant (CCPP) in Iran aimed at improving its performance and reducing emissions, indicated that introducing parabolic solar collectors and a multi-effect desalination cycle decreased  $\text{CO}_2$  emissions from 58.05 to 54.97 kg/MWh of electricity [18]. Further investigations of Abadan (CCPP) showed that the implementation of a multigeneration system caused environmental effects of the modeled system on air pollution production ( $\text{NO}_x$ , CO,  $\text{CO}_2$ ) equal to 8.975 kg/s, which indicated a decrease of pollutant emissions equal to 9.8% [19]. Specific configurations of Abadan CCPP with a solar power tower [20] could reduce emissions into the atmosphere up to 345 63 tons/year. There are numerous objections to uncontrolled methane leakages from natural gas systems during exploration activities [13,21].  $\text{CH}_4$  is the main component of shale gas and the second main greenhouse gas (GHG), and it is much more efficient in absorbing radiation than  $\text{CO}_2$  [21]. Many studies have focused on the contribution of natural gas production to  $\text{CH}_4$  emissions and GHG budget, although their results differ from each other. According to observations by Karion et al. [22], the total fugitive emission of  $\text{CH}_4$  from shale gas operations can be up to 17.3% of total gas production. Research by Shaw et al. [23] showed the results of air quality measurements performed near the UK's first shale gas hydraulic fracturing area (Lancashire, England) at the turn of 2018 and 2019. Observations during flowback operations revealed elevated  $\text{CH}_4$  mixing ratios (30-min mean of up to 5 ppm) after artificial nitrogen lifting of a well. The estimate of mean  $\text{CH}_4$  flux using modeling approaches was approximately  $16 \text{ g s}^{-1}$ .

In the vicinity of the borehole area, a significant emission of  $\text{NO}_x$  and VOC contributes to surface  $\text{O}_3$  formation and results in high  $\text{O}_3$  episodes [24,25]. Schnell et al. [1] analyzed surface  $\text{O}_3$  concentrations in the rural area of the Upper Green River Basin, Wyoming (UGRBWY), during February 2008. Under conditions of a high-pressure system, freezing temperature, snow cover, low wind, and clear sky, the hourly average  $\text{O}_3$  concentration rose from 30 ppb (night) to above 140 ppb (afternoon). Measurements performed by Rappenglück et al. [26] in UGRBWY from January to March 2011 also indicated repeated episodes of the air quality standard (>75 ppb) being exceeded during the winter season. Large amounts of  $\text{NO}_x$  and NMHC emitted during fossil fuel activities accumulate below the shallow stable ( $\approx 50 \text{ m.a.g.l.}$ ) boundary layer, thereby affecting surface  $\text{O}_3$  formation, which reaches maximum 1-h average values above 160 ppb. Cheadle et al. [27] reported that in the North Colorado Range Front (NCRF) in Colorado, on individual summer days with high  $\text{O}_3$  levels, emission related to oil and natural gas extraction activities could contribute up to 30 ppb to  $\text{O}_3$  formation locally. According to Benedict et al. [28] elevated VOC concentrations associated with greater oil and natural gas determine episodes of high  $\text{O}_3$  concentration in NCRF in Colorado. Emission connected with gas exploitation contributed up to 20 ppbv excess  $\text{O}_3$ . Similar results were obtained by Lindaas et al. [29]. They found that amounts of long-lived VOCs, related to oil and natural gas exploitation, are

so high that processes of their oxidation during the day produce sufficient peroxy radicals to significantly impact on O<sub>3</sub> production in NCFR in Colorado during high O<sub>3</sub> days in this region.

The success of shale gas in the United States at the beginning of the 21st century has become a phenomenon on a global scale, mainly due to advantages of geological and natural physical conditions (shallow shale gas formation with a high index of total organic carbon and relatively big thickness) [30]. Additionally, favorable government policies and regulation systems supported development of local shale infrastructure, advanced techniques and technology, especially hydraulic fracturing. In comparison with the United States, the natural conditions in Europe, in terms of geology, topography and water abundance, are more complicated [31]. Additionally, local hostility and lack of acceptance by the population, due to concerns about extreme environmental and health risks, has appeared in many places where attempts have been made to exploit shale gas [32]. Selected shale gas research approaches, with regards air pollutant measurements are listed in Table 1.

**Table 1.** List of shale gas research approaches.

	Author, Date of Publication	Country	Period of Measurements	Measured Pollutants	Measurement Method	Main Goal of the Study
1	Oltmans et al., 2014 [25]	US	2011	O <sub>3</sub> , NO <sub>x</sub> , NMHC	-measurements from tethered balloon -observations from a tall tower	Study of the genesis of episodes of high O <sub>3</sub> concentration values.
2	Karion et al., 2013 [22]	US	2012	CH <sub>4</sub>	-measurements from instrumented single-engine turboprop aircraft	Using mass balance method to estimate CH <sub>4</sub> average natural leak rate.
3	Shah et al., 2020 [33]	UK	2018–2019	CH <sub>4</sub>	-measurements using unmanned aerial vehicle (UAV)	Detection of CH <sub>4</sub> leak rate during first exploratory fracking process.
4	Purvis et al., 2019 [8]	UK	2016–2018	NO <sub>x</sub> , O <sub>3</sub> , SO <sub>2</sub> , NMHC, PM	-in situ monitoring station	Observation of air pollutants before, during and after preparations for hydrofracturing.
5	This study Jarosławski et al., 2022	Poland	2015–2017	NO, NO <sub>2</sub> , NO <sub>x</sub> , O <sub>3</sub> , CO, PM <sub>10</sub> , CO <sub>2</sub> , CH <sub>4</sub> , NMHC	-in situ measuring station	Comparative analysis of pollutants during different stages of well's work cycle and checking meeting of National Air Quality Criteria


The shale rock formation located in the Wysin area belongs to the Baltic-Podlasie-Lublin Basin. This is the largest unconventional gas reservoir in Poland. According to the most reliable estimations, it contains  $346\text{--}768 \times 10^9 \text{ m}^3$  of recoverable shale gas resources (on average four times more than the documented conventional gas resources in Poland) with a maximum of  $1.92 \times 10^{12} \text{ m}^3$  [34]. The shale gas exploratory activities in Poland began a few years ago. The first horizontal drilling was performed in 2011 near the Łebień village in the Pomeranian voivodeship (about 90 km from Wysin). According to a series of measurements of potential air pollution indicators (SO<sub>2</sub>, NO<sub>x</sub>, CO, C<sub>6</sub>H<sub>6</sub>, and H<sub>2</sub>S) made in Łebień by the Regional Inspectorate for Environmental Protection in Gdańsk, no exceeding of the acceptable level of air pollution were observed [35].

This study covers two- years (from August 2015 to July 2017) of continuous observation of the main atmospheric air pollutants (CO, O<sub>3</sub>, PM<sub>10</sub>, NO, NO<sub>2</sub>, and NO<sub>x</sub>), as well as pollutants directly associated with shale gas exploration activities, such as drilling and hydraulic fracturing (CH<sub>4</sub>, NMHC, and CO<sub>2</sub>), from the Stary Wiec station located in the village of Wysin, about 1 km from the borehole area. The measurement campaign was

performed within a framework of the SHEER (Shale gas exploration and exploitation induced risk) project realized under the Horizon 2020 program. The main goal of this work was to present the variation in air pollutant levels at different stages of the natural gas extraction activities and to indicate the potential impact of shale gas exploration activities on the air quality in relation to background levels. For this purpose, a number of statistical analyses were performed: indication of average pollution concentration in different time scales, estimation of high pollutant concentrations, determination of episodes with a particular emphasis on pollutants directly linked to shale gas exploration, and comparison of results obtained at the Stary Wiec station with those from neighboring stations.

## 2. Materials and Methods

An air pollution monitoring station was set up in the small village of Stary Wiec (54.1° N, 18.3° E, 170 m a.s.l.), the closest inhabited place to the boreholes. The distance between the wells area and the station was approximately 1100 m. (Figure 1).



**Figure 1.** Map of Poland with marked Pomeranian region (red square) and locations of the air quality monitoring network sites and Stary Wiec station. (Available online: <https://maps.google.com> (accessed on 19 July 2022)).

Location of the station relative to the wells considering the prevailing wind directions in this region. Approximately 55% of the winds originated from the western directions during the measurement campaign. Moreover, only for 0.8% of the measurement time was the wind speed less than 1 m/s. In this situation, the possible emissions of air pollutants from the drilling site had a good chance of being recorded by the measuring equipment installed at the station. The closest vicinity of the station is dominated by small villages with a population of up to 600 inhabitants. The nearest small town (Skarszewy), with a population of more than 7000, is located 8 km east of the station. Local anthropogenic sources of air pollution were small and typical for rural conditions: several neighboring houses and a small local road located 50 m from the station. Residential buildings mainly included single family houses heated by coal- and wood-burning stoves and several multifamily houses connected to the heat distribution networks. The measuring station


was located in the northwest part of the village. In the immediate vicinity of the station, there was a closed pig farm. Since the pig farm was nonfunctional, it had no effect on the level of pollution in the vicinity of the station. During the westerly winds (which prevail in this region), when there was an inflow of air from the direction of the borehole area, the combustion processes in these houses had no significant effect as they are on the leeward side of the station. The road was occasionally used, with traffic of a dozen cars per hour. Additional heavy traffic, related to activities at the wells area, occurred during the period of exploitation of the wells. In addition, a park with a small pond located in the vicinity of the station was a potential natural source of gases, such as CO<sub>2</sub> and CH<sub>4</sub>. The surrounding landscape was dominated by farmlands and small forests. The nearest significant urban sources of air pollution were located at a distance of 25–50 km from the station (see Figure 1).

The station's container was equipped with a standard set of analyzers measuring concentrations of the air pollutants: NO, NO<sub>2</sub>, NO<sub>x</sub> (sum of NO and NO<sub>2</sub>), CO, O<sub>3</sub>, and PM10. Gaseous pollutants were measured by the API analyzer, models 200, 300, and 400, while PM10 concentration was measured by a TEOM analyzer, model 1400. This standard air pollution monitoring measurement program was extended by measurements of the concentration of CO<sub>2</sub> using a Thermo 410i analyzer and concentrations of CH<sub>4</sub> and NMHC (sum of NMHCs) with a Horiba APHA370 analyzer. Technical specifications of the analyzers employed during the measuring campaign are listed in Table A1 in the Appendix A. Meteorological parameters were recorded by a collocated automatic weather station.


To ensure adequate data quality, i.e., total measurement uncertainty less than 15% in accordance with the requirements used in air quality monitoring networks in Poland, standard quality assurance procedures analogous to those routinely used in air pollution monitoring networks were applied during measurements, including regular calibrations with certified gas mixtures and a certified photometer (in the case of ozone), data verification and validation, and regular control of, and servicing of, analyzers and container systems. Instantaneous values of all pollutants' concentrations were converted into 1-h averages, and these averages were used for further analysis.

The measuring campaign in Sary Wiek covered a wide range of air pollutants, both main gaseous pollutants measured as standard at state air quality monitoring stations (NO, NO<sub>2</sub>, NO<sub>x</sub>, CO, O<sub>3</sub>, PM10) and, additionally, those directly related with shale gas related activities (CO<sub>2</sub>, CH<sub>4</sub>, NMHC). The measurements were performed continuously for two years, from August 2015 to July 2017. This long period of uninterrupted measurements ensured detection of possible air pollution episodes related to all stages of the life cycle of wells: preparatory phase, drilling, hydrofracturing, and closing activities. Additionally, 24 months of monitoring covered two full seasonal cycles, which was important for identifying the natural local variability of each pollutant and possible local sources of pollution not related to the wells' activities, thus allowing the determination of the background values of each analyzed pollutant. In this work we investigated the shale gas industrial development in a rural area in the North of Poland in terms of accompanying air pollutants and GHGs emissions. In this connection, the analysis of the potential impact of activities related to shale gas exploration and exploitation processes on ambient air quality, in the context of environmental protection and the local community, was performed. The comparative analysis of pollutant concentrations during an entire work cycle of the wells (as well as in the period before and after), taking into account meteorological conditions, was considered in terms of the following: (1) whether the National ambient Air Quality Criteria were exceeded (2) occurrence of peaks and deviations from the background values (3) comparison concentrations of all pollutants during different shale gas operation life cycles (drilling and hydrofracturing) relative to the reference period (in this article defined as a remaining period). In order to perform this, we first used the data on the levels of air pollution and meteorological conditions from several of the nearest (7 to 30 km) stations to the regional air pollution monitoring network to assess air quality in the region before


and during the measurement campaign. The analysis of the measurement data (from 2012 to 2017), presented in Figure 2, indicated the lack of significant difference in ambient air pollutants between the baseline period (2012–2014) and the period of intensive activity in the shale gas area (2015–2017). To enable the estimation of the potential impact of shale gas exploration on air quality, an examination on the influence of wind direction on the measured mixing ratios of each pollutant was performed (Figures A1–A5 in Appendix A). Air pollution data were assigned to one of eight wind direction compartments (N, NE, E, SE, S, SW, W, NW) depending on the phase of activity at the area of the wells. Furthermore, an additional compartment for wind coming directly ( $\pm 5^\circ$ ) from the borehole area was determined. Wind rose graphs were used to present the amounts of pollutants averaged for each sector of the wind and to compare these with the amounts estimated for the wind coming directly from the shale gas site. To determine whether or not the air quality standards for human health and plant protection were met the compliance of NO, NO<sub>2</sub>, CO, O<sub>3</sub> and PM<sub>10</sub> with the Air Quality Directive was evaluated. Finally, diurnal and seasonal analyses of CH<sub>4</sub>, NMHC and CO<sub>2</sub> as the gases directly related to shale gas operations were carried out (Figure 3). The detailed analysis of the high temporal resolution data revealed the existence of episodes of elevated concentrations of CH<sub>4</sub> and NMHC registered in the period 1–4 September 2016 (Figure 4). To determine the origin (artificial or biogenic) of high GHGs concentration calculations of CO<sub>2</sub>/CH<sub>4</sub> ratio were performed (Figure 5). Additionally, statistical analysis of episodes recorded for the period 1–4 September 2016, based on correlation coefficient, *p*-value and CH<sub>4</sub>/NMHC ratio values, was performed to confirm whether or not the shale gas related emissions were a potential source of episodes of elevated concentrations of CH<sub>4</sub>.




**Figure 2.** Monthly means of nitrogen oxide (a) and surface O<sub>3</sub> (b) concentrations at Stary Wiek and the surrounding stations for the period 2012–2017.



**Figure 3.** Diurnal variation (mean value together with 1 standard deviation) of CO<sub>2</sub> (a) and CH<sub>4</sub> (b) at the Stary Wiecek station, August 2015–July 2017.



**Figure 4.** Concentration of CH<sub>4</sub> and NMHC over 1st (a), 2nd (b) and 3rd (c) September 2016, measured at the Stary Wiecek station.



**Figure 5.** Values of the  $\text{CO}_2/\text{CH}_4$  ratio over 1–4 September 2016, measured at the Stary Wiec station.

### 3. Results

#### 3.1. Analysis of Measurement Results from the Regional Network of Air Pollution Monitoring

Measurement results of pollutants, namely  $\text{NO}$ ,  $\text{NO}_2$ ,  $\text{CO}$ ,  $\text{O}_3$ , and  $\text{PM}_{10}$ , from the five nearest air pollution monitoring stations, namely, Gdańsk, Starogard Gdański, Kościerzyna, Tczew (reflecting urban background conditions), and Liniewko Kościerskie (reflecting rural background conditions), were used for the general assessment of air quality level in the region before starting any activities related to shale gas exploration and during the measurement campaign in Stary Wiec. Data from the period 2012–2017 were used for the analysis. Results of the available measurements (monthly means) for primary ( $\text{NO}$ ) and secondary ( $\text{O}_3$ ) pollutants are shown in Figure 2.

The analysis of data revealed distinct natural seasonal and year-to-year variations in concentrations of all measured air pollutants in the region. This variation resulted especially from changes in meteorological conditions during the year and changes in emissions from local anthropogenic sources (e.g., traffic and heating systems of buildings in the winter). The period 2015–2017 did not differ from previous years in terms of concentrations of individual pollutants (see Table A2 in the Appendix A). Statistics of pollutant levels measured at Stary Wiec and at nearby rural and urban stations during the measurement campaign are presented in Table 2.

**Table 2.** Statistics for air pollution data from Stary Wiec and nearby rural and urban stations, based on hourly means, collected from August 2015 to July 2017. Mean values for this period, together with standard deviations ( $1\sigma$ ), are presented.

	Stary Wiec	Liniewko	Kościerzyna	Tczew	Starogard	Gdańsk
<b>NO (ppb)</b>	$0.87 \pm 1.59$	$1.03 \pm 0.75$	$6.05 \pm 13.28$	$4.73 \pm 13.60$	$7.86 \pm 13.19$	$3.72 \pm 9.75$
<b>NO<sub>2</sub> (ppb)</b>	$4.20 \pm 2.94$	$3.15 \pm 2.35$	$8.57 \pm 5.41$	$7.21 \pm 6.55$	$7.76 \pm 8.04$	$8.74 \pm 6.79$
<b>O<sub>3</sub> (ppb)</b>	$27.37 \pm 13.18$	$26.68 \pm 12.70$	$22.01 \pm 12.21$			$22.93 \pm 12.04$
<b>CO (ppb)</b>	$242.5 \pm 159.5$	$316.4 \pm 96.6$	$484.9 \pm 400.0$	$285.7 \pm 168.4$	$361.4 \pm 297.6$	$323.8 \pm 146.6$
<b>PM<sub>10</sub> (<math>\mu\text{g}/\text{m}^3</math>)</b>	$16.18 \pm 13.65$		$31.02 \pm 28.45$	$20.35 \pm 17.31$	$42.13 \pm 44.64$	$17.60 \pm 15.45$

Air pollutant levels measured in Stary Wiec were representative of a clean, rural environment. Very low, close to zero, levels of  $\text{NO}$ , a pollutant with a lifetime of approximately 1 h, indicated the lack of significant engine exhaust type sources of air pollution in the vicinity of the station. Pollutants such as  $\text{CO}$ ,  $\text{NO}_2$ , and  $\text{PM}_{10}$  were the lowest, or one of the lowest, in the region during the measurement period. However, the levels of the secondary pollutant  $\text{O}_3$  were at a relatively high level, which is a known typical phenomenon for rural areas [36]. Higher  $\text{O}_3$  concentrations at rural stations than at urban stations are

due to less efficient mechanisms for O<sub>3</sub> destruction [37], the most important of which is the lack of efficient NO emission sources that can reduce the reactions of O<sub>3</sub> destruction (NO+O<sub>3</sub>→NO<sub>2</sub>+O<sub>2</sub>). Most episodes of elevated pollutant concentrations registered at the Stary Wiec station during the 2015–2017 period resulted from regional scale phenomena, such as favorable meteorological conditions or transport of pollutants from regional sources, e.g., large cities of the Pomeranian Voivodeship.

### 3.2. Analysis of Air Pollutant Measurement Results from Stary Wiec Station

As described in Section 1, there were no significant sources of air pollution in the immediate vicinity of the station; hence, it can be assumed that the levels of pollutants in the area of the station were largely due to their transportation to the place of measurement from more distant source areas. Thus, pollutants emitted in the Wysin wells area could have been registered by the station's equipment if they were transported to the sampling point when favorable wind conditions occurred.

To assess the possible impact of pollutant emissions from the wells area on the ambient air quality in the vicinity of the station, an analysis of the distribution of air pollutant concentrations recorded at the station, depending on wind direction, was performed. For this purpose, the average concentrations of individual air pollutants were calculated for eight wind direction compartments. In addition, the average concentrations of pollutants for wind blowing directly ( $\pm 5^\circ$ ) from the wells were determined. The measurement time was divided into three periods in accordance with the actions taken at the boreholes: the drilling period (September–October 2015), the hydrofracturing period (June–July 2016), and the remaining period of measurements (August 2015–July 2017, excluding drilling and hydrofracturing periods).

Significance of differences between the average values of concentrations of pollutants from the NW (north-west) sector and for air originating directly from the well was confirmed by the F-test (to determine whether the analyzed samples were characterized by equal/unequal variance) and the Student's *t*-test with two samples assuming equal or unequal variances (to determine whether the differences of mean values in two analyzed samples were statistically significant at the significance level of 0.05). Results are shown in Table 3.

**Table 3.** Statistical significance according to the *t*-test results (at  $\alpha = 0.05$ ) of differences between concentrations of individual air pollutants for the air coming to the measurement point directly from the wells area and from the NW sector during drilling, hydrofracturing, and the remaining period. Note: Statistically significant cases (*p*-value < 0.05) are bolded. Units as shown in Table 2, except CH<sub>4</sub> (ppm) and NMHC (ppmc).

	Drilling			Hydrofracturing			Remaining Period		
	NW	Well	<i>p</i> -Value	NW	Well	<i>p</i> -Value	NW	Well	<i>p</i> -Value
	Mean	Mean		Mean	Mean		Mean	Mean	
NO	1.41	1.72	0.217	<b>0.58</b>	<b>1.21</b>	<b>0.044</b>	0.95	0.83	0.316
NO <sub>2</sub>	4.18	5.04	0.065	2.74	3.06	0.170	3.51	3.63	0.256
CO	215.5	187.7	0.085	<b>132.96</b>	<b>157.03</b>	<b>0.005</b>	229.08	218.05	0.114
PM <sub>10</sub>	<b>16.02</b>	<b>13.02</b>	<b>0.046</b>	12.33	13.18	0.599	12.91	13.07	0.728
O <sub>3</sub>	20.17	20.89	0.647	27.29	24.24	0.281	<b>29.66</b>	<b>28.42</b>	<b>0.019</b>
CO <sub>2</sub>	426.78	420.68	0.265	406.49	423.24	0.135	412.23	410.57	0.181
CH <sub>4</sub>	<b>1.99</b>	<b>1.95</b>	<b>0.023</b>	1.93	1.95	0.197	1.93	1.93	0.521
NMHC	0.06	0.06	0.85	0.02	0.02	0.147	0.03	0.03	0.216

Measurement results are shown in Figures A1–A5 in the Appendix A. These figures show the dependence of the concentrations of pollutants on the wind direction for a previously defined period. It can be observed that the cleanest air conditions occurred during advection of air from the western directions, while the highest levels of pollutants

appeared while air was coming to the measuring point from the east or south. This situation roughly corresponded to the distribution of sources of air pollution around the measurement site. Cleaner air coming from the West was often of North or West Atlantic origin, while the air from the eastern sectors, mainly of continental origin, passed over relatively close significant sources of air pollution (several larger cities and the motorway located nearby) on its way to the measuring point.

During the low activity period in the wells area, the concentrations of air pollutants registered at the Stary Wiec station did not increase significantly when direct transport from the area of the wells occurred, in comparison to that when the air was coming from the entire NW sector. The relative differences in concentrations were small, in the order of a few percent (up to 5%), except for NO (−13%). For most air pollutants, nonsignificant changes in concentrations were recorded in the air coming directly from the well region as compared to their average values for the whole sector. Significantly lower values were only recorded for O<sub>3</sub>; however, the absolute differences remained very small (approximately 1 ppb).

A different situation occurred during the periods of increased activity at the wells, especially during hydrofracturing. Most of the pollutants occurred in lower concentrations during this period, because of natural seasonal variation. Hydrofracturing was performed in the summer, when the levels of air pollutants (except for O<sub>3</sub>) were usually at the lowest of the year. However, the levels of some of the pollutants in the air arriving at the measurement point directly from the wells area increased significantly according to the *t*-test results, compared to the average values registered at the same time for the air coming from the NW sector during the hydrofracturing period. A significant relative increase (108% and 18%) occurred for NO and CO, respectively. Increase in the levels of NO<sub>2</sub> (12%), PM<sub>10</sub> (7%), and NMHC (21%), although not statistically significant, were also detected. All the aforementioned pollutants were the components of engine exhausts. Therefore, an observed increase in air pollution levels could be the result of intense work of, for example, the pump systems. Moreover, smaller (by 11%) levels of O<sub>3</sub> observed simultaneously with higher NO concentrations confirmed this hypothesis, as O<sub>3</sub> levels are usually anti-correlated with NO levels, especially when close to the source of emission, because of chemical reactions occurring between them. Slight increases in the concentrations of CH<sub>4</sub> and CO<sub>2</sub> were also detected.

For the drilling period, the results were ambiguous. A statistically nonsignificant increase in NO<sub>x</sub> (by approximately 20%) occurred. However, for the other pollutants, the concentrations were usually lower than the average for the NW sector, up to 19% lower for PM<sub>10</sub>. In two cases, those relating to PM<sub>10</sub> and CH<sub>4</sub>, the lower values were statistically significant.

The analysis of the above results showed that, despite a relatively high increase in the concentrations of some pollutants during the hydrofracturing period, their absolute concentrations remained at a very low level; for example, 1.2 ppb for NO or 5.0 ppb for NO<sub>2</sub>, during the hydrofracturing and drilling periods, respectively. Thus, even in the periods of intense activity at the boreholes, the air in the vicinity of the Stary Wiec station remained relatively clean. The analysis of compliance with the Air Quality Directive applicable to the pollutants (Table 4) also showed that the relevant air quality criteria were met for most pollutants.

**Table 4.** Air quality criteria applicable to the pollutants measured at the Stary Wiec station <sup>(1)</sup> Information threshold, <sup>(2)</sup> Alarm threshold, <sup>(3)</sup> Accumulated O<sub>3</sub> exposure indicator (AOT40), <sup>(4)</sup> Protection of vegetation. Normal font-standards for human health protection, italics-standards for plant protection.

Pollutant	Averaging Period	Acceptable Level [ppb]	Acceptable Number of Occurrences during Calendar Year	Number of Exceedances in Stary Wiec	Average or Maximum Concentration Value in Stary Wiec
NO <sub>2</sub>	1 h	104.6 ppb	18	0 (2015–2017)	
NO <sub>2</sub>	1 year	20.9 ppb			4.3 ppb (2016)
PM <sub>10</sub>	24 h	50 µg/m <sup>3</sup>	35	11 (2015–2017)	
PM <sub>10</sub>	1 year	40 µg/m <sup>3</sup>			15.3 µg/m <sup>3</sup> (2016)
CO	8 h running mean	8620 ppb			max 1385 ppb
O <sub>3</sub>	8 h running mean	60 ppb	25 days	8 (2015) 13 (2016) 2 (2017)	
O <sub>3</sub>	1 h	90 ppb <sup>(1)</sup>		2 (2015)	
O <sub>3</sub>	1 h	120 ppb <sup>(2)</sup>		0 (2015–2017)	
O <sub>3</sub>	May–July	9000 ppb*h <sup>(3)</sup>			7696 ppb*h (2016) 2817 ppb*h (2017)
NO <sub>x</sub>	1 year	15.7 ppb <sup>(4)</sup>			5.0 ppb (2016)

By analyzing the air quality standards for human health protection, it can be stated that during the two-year measuring period, most noticeable were the relatively high values in ambient O<sub>3</sub> concentrations. The information target value was exceeded twice in August 2015. Although the air quality limit values for O<sub>3</sub> were frequently exceeded during summer months, it should be noted that exceedances registered in the period 2015–2017 originated from episodes of a larger, regional scale, which were not related to works at the area of the wells. Permissible levels of the remaining pollutants (NO<sub>2</sub>, PM<sub>10</sub>, and CO) for the protection of human health were not exceeded.

### 3.3. Methane, NMHCs, and CO<sub>2</sub> Concentration Measurement Results

The monitoring of CH<sub>4</sub>, NMHC, and CO<sub>2</sub> levels was included into Stary Wiec station's measurement program to track the possible emissions of these gases accompanying the exploration, and subsequent exploitation, of shale gas wells in Stary Wiec. The analysis of the data collected during the two years of measurements showed the existence of well-known seasonal (summer minimum and winter maximum) and diurnal (minimum during the day and maximum at night) cycles of CO<sub>2</sub> and CH<sub>4</sub> related to the activity of natural sources and sinks, namely photosynthesis and respiration of the biosphere for CO<sub>2</sub> and methanogenesis for CH<sub>4</sub>. CO<sub>2</sub> is the basic substrate of the photosynthesis process occurring with the participation of light energy, most often solar energy. Photosynthesis processes are an important sink of CO<sub>2</sub> in the natural ecosystem during the daytime. In contrast, the processes of respiration of plants and soil are significant natural sources of CO<sub>2</sub> and contribute to its increase during the night (in the absence of photosynthesis). Methanogenesis is the process of biological formation of CH<sub>4</sub> during decomposition of organic compounds (e.g., acetate, formate, CO<sub>2</sub>, hydrogen) by anaerobic microorganisms (mainly from the Archaea domain). Figure 3 shows the results of measurements of diurnal variations of CO<sub>2</sub> and CH<sub>4</sub> performed over the entire experimental period.

Both gas species had similar, distinct diurnal distribution, characterized by maximum values during late night–early morning (02:00–04:00 for CO<sub>2</sub> and 03:00–05:00 for CH<sub>4</sub>),

followed by a gradual decrease, reaching a minimum around 12:00–16:00, and then a gradual increase through the evening and night until the maximum peak the next day. Thus, it can be stated that the concentration values of both gases ( $\text{CO}_2$  and  $\text{CH}_4$ ) were cumulated and constantly increasing in the afternoon through evening and night until an early morning maximum on the next day. The maximum values of  $\text{CO}_2$  varied between 431.5 ppm and 433.6 ppm, while the lowest value was around 399 ppm. Daily concentration values of  $\text{CO}_2$  were characterized by a very high amplitude in the summer months, with the maximum reaching around 100 ppm. Hence, high values probably resulted from the presence of important natural sources and sinks in the vicinity of the station (pond and park with deciduous trees). During winter, the amplitude values were almost negligible, oscillating around several ppm. The maximum values of  $\text{CH}_4$  increased to around 1.962 ppm, while the minimum dropped to 1.916 ppm. Stronger amplitudes of diurnal variation were noted during summer, with the maximum values in July 2016, reaching 0.125 ppm.

The mean values at Stary Wiec for the recording period of August 2015–July 2017 were  $1.936 \pm 0.064$  ppm for  $\text{CH}_4$ ,  $414 \pm 29$  ppm for  $\text{CO}_2$ , and  $0.039 \pm 0.034$  ppmc for NMHC. These levels were equal to, or slightly higher than, those reported for the Poland remote mountain region [38,39].

Concentrations of all three analyzed components in the air coming directly from the wells were higher than the average values for the NW sector in the hydrofracturing period, ranging from 1% for  $\text{CH}_4$  up to 21% for NMHC. However, this increase was not statistically significant. For the remaining period, the corresponding differences were below 1% for  $\text{CH}_4$  and  $\text{CO}_2$  and below 5% for NMHC.

Variability in  $\text{CH}_4$  concentrations recorded at Stary Wiec was relatively small (standard deviation 0.06 ppm); hence, more attention was paid to the analysis of episodes of elevated  $\text{CH}_4$  concentrations, as potentially coming from the wells. The highest concentration of  $\text{CH}_4$  (1-h average), 2.81 ppm with more than 10 standard deviations from the average value, was registered on 1 September 2016, outside the drilling and hydrofracturing period. A more detailed analysis of the one-minute concentrations of  $\text{CH}_4$ ,  $\text{CO}_2$ , and NMHC for this, and for several subsequent days, revealed a series of short  $\text{CH}_4$  and NMHC concentration peaks, up to 7.45 ppm for  $\text{CH}_4$  and up to 3.03 ppm for NMHC.  $\text{CO}_2$  concentrations remained at mean levels. The results of the measurements are shown in Figure 4.

The configuration and times of occurrence of the concentration peaks were similar for three consecutive days, with decreasing concentration values, indicating their artificial origin. The wind direction analysis for this period indicated the wells area as a possible methane source.

Analysis of  $\text{CO}_2$  and  $\text{CH}_4$  concentration ratio (ppm/ppm) is sometimes used as a tracer in investigating the origin of GHGs [40]. A low  $\text{CO}_2/\text{CH}_4$  ratio is often interpreted as an indicator of significant anthropogenic impact [41]. This analysis was performed for the time of occurrence of the abovementioned  $\text{CH}_4$  and NMHC episodes. The graph of these ratios is shown in Figure 5.

Usually, its values remained relatively stable during the daytime, and the mean value for the entire measurement period was  $219 \pm 12$ , with an increase during nights, because of strong  $\text{CO}_2$  production through vegetation in a park near the station. During the duration of the episodes, the ratio systematically dropped down to 53, indicating a non-biogenic source of  $\text{CH}_4$ , probably due to some technical procedures at the wells, which was associated with the release of certain amounts of  $\text{CH}_4$  directly into the atmosphere. Unfortunately, the site operator did not provide information about the type of activity being conducted at this time at the Wysin site.

Additionally, to investigate the predominance of the thermogenic origin of increased  $\text{CH}_4$  concentration during the episodes, an analysis, based on the relationship between  $\text{CH}_4$  and NMHC, was performed. Characteristics of the total number of episodes observed for the period 1–4 September 2016 are presented in Table 5.

**Table 5.** Characteristic of CH<sub>4</sub> episode events observed for 1–4 September 2016.

Episode Number	Date	Max CH <sub>4</sub> Concentration [ppm]	Max NMHC Concentration [ppm]	Correlation Coefficient CH <sub>4</sub> -NMHC	<i>p</i> -Value	CH <sub>4</sub> /NMHC Ratio
1	1 September 2016	7.45	3.03	0.998	0.000	6.95
2	2 September 2016	2.7	0.39	0.980	0.020	7.52
3	2 September 2016	2.81	0.45	0.997	0.003	7.01
4	2 September 2016	2.32	0.23	0.995	0.000	10.55
5	2 September 2016	5.36	1.62	0.999	0.000	4.50
6	3 September 2016	2.36	0.20	0.906	0.000	13.14
7	3 September 2016	2.75	0.36	0.958	0.000	10.67
8	3 September 2016	2.30	0.15	0.680	0.064	15.46
9	3 September 2016	2.35	0.11	0.859	0.062	23.01
10	3 September 2016	2.56	0.31	0.997	0.000	11.46
11	3 September 2016	3.42	0.69	0.998	0.000	7.80
12	3 September 2016	3.47	0.73	0.999	0.000	7.54
13	3 September 2016	2.74	0.39	0.999	0.000	8.11
14	3 September 2016	2.68	0.35	0.992	0.001	9.36
15	3 September 2016	3.35	0.66	0.994	0.000	6.42

The concentration values of CH<sub>4</sub> (1-min average) qualified as episodes were greater than the average value by at least 5 deviations, reaching the maximum value of 7.45 ppm (mean  $2.91 \pm 0.92$  ppm), whereas almost all the values of NMHC were greater than the average value by at least 3 standard deviations. The maximum value of NMHC was 3.03 ppm (mean  $0.43 \pm 0.49$  ppm). A significant correlation was observed between CH<sub>4</sub> and NMHC levels ( $p < 0.05$ ) with correlation coefficient above 0.6, and a low value of the CH<sub>4</sub>/NMHC ratio was recorded in almost all episode cases (except episode numbers 8 and 9). Correlation coefficient values, in most cases, were above 0.9, thus indicating a strong association between species. The CH<sub>4</sub>/NMHC ratio values were significantly lower than the mean value for the entire experiment (90) and dropped down to approximately 5. These calculations indicated the possible influence of a thermogenic source of the CH<sub>4</sub> episodes. The origin of episode numbers 8 and 9 is not clear. Correlation coefficient values were slightly smaller than the others (0.680 and 0.859, respectively), and *p*-values (0.064 and 0.062, respectively) were above the adopted significance ( $p < 0.05$ ). The CH<sub>4</sub>/NMHC ratio values were the highest of all (15.46 and 23.01, respectively), but remained significantly lower than the average value for the entire measurement period. The potential sources of elevated CH<sub>4</sub> levels in episodes 8 and 9 may indicate that the CH<sub>4</sub>/NMHC ratio included both thermogenic and biogenic contributions, but these inferences are not unequivocal.

For episode number 1 the methane plume possibly emitted from the well was modeled by the HYSPLIT model. The results are shown on Figure A6 in the Appendix A.

#### 4. Conclusions

This work presents the first long-term (two year) measurement assessment of the air quality impact of activities performed at a shale gas site in Poland.

The development of a shale gas exploration and exploitation site in Wysin (over the period 2015–2017) consisted of a full cycle of technical works, including the following phases, each of which has potential for pollution emission to the atmosphere: pre-operational phase, well drilling, hydro-fracturing, closing the well and site restoration. Continuous monitoring of levels of air pollutants in the vicinity of shale gas wells supported the hypothesis that emission from shale gas extraction activities in the Wysin area has a negligible impact on ambient air quality at a distance of about 1100 m from the wells on their leeward side.

Concentrations of pollutants measured at the station strongly depend on the kind of activities performed at the borehole area. The highest, partly statistically significant increases in level of selected pollutants were noted during increased activity at the wells (during drilling and hydro-fracturing periods) in the conditions of direct transport of air

from the area of the wells to the measurement point. The works of drilling and hydro-fracturing involved intensified vehicle activity, due to delivery of equipment to the site and its operation, and led to significant increases, especially in NO<sub>x</sub> emissions. Concentration of NO during intensive activity at the borehole area increased to an average value of 1.72 ppb (during drilling) and 1.21 ppb (during hydro-fracturing), compared to 0.83 ppb measured during the remaining period when no activities related to the borehole were performed.

Based on the observations of CH<sub>4</sub>, CO<sub>2</sub> and NMHC, the gases directly related to shale gas exploration activities, the average concentrations registered for the period 2015–2017 were equal to  $1.936 \pm 0.064$  ppm for CH<sub>4</sub>,  $414 \pm 29$  ppm for CO<sub>2</sub>, and  $0.039 \pm 0.034$  ppm for NMHC. During the measurement period, several episodes of elevated CH<sub>4</sub> concentration values, with maximum one-hour average equal to 2.81 ppm (more than 10 standard deviations from the average value), occurred in September 2016. Detailed analysis of the one-minute concentrations of CH<sub>4</sub>, CO<sub>2</sub> and NMHC revealed a series of short CH<sub>4</sub> and NMHC concentration peaks, up to 7.45 ppm for CH<sub>4</sub> and up to 3.03 ppm for NMHC. The time of appearance of these peaks was similar for three consecutive days, with decreasing concentration values, which could indicate their artificial origins. CH<sub>4</sub>/CO<sub>2</sub> ratio values during the measurement period were characterized by relatively stable values  $\approx 219 \pm 12$ . Over 1–4 September CH<sub>4</sub>/CO<sub>2</sub> ratio values dropped down to 53. Due to lack of information from the site's operator, we cannot unambiguously state what kind of operations were performed during these days at the borehole area and what exactly caused the high emission of CH<sub>4</sub> and NMHC pollutants. However, analysis of wind direction during these days indicated the borehole area as a potential CH<sub>4</sub>/NMHC source.

Results of the analysis of measurements taken during shale gas exploration activities indicated that significant increases of concentrations of air pollutants (except of NO<sub>x</sub>, CH<sub>4</sub> and NMHC, mentioned before) were not observed. Taking into account the Air Quality Criteria, applicable to individual pollutants, no exceedances of threshold values (with the exception of O<sub>3</sub>) were observed. Relatively low levels of concentrations of air pollutants measured during drilling, hydraulic fracturing and shale gas extraction at Stary Wiec could be determined, especially by the scale of operations across the time and space, which were relatively small. Additionally, exploration activities concerned only two wells. It can be assumed that the use of shale gas extraction technology, taking into account: risk-mitigation measures and technical quality control, could ensure environmentally safe exploration and exploitation of shale gas.

**Author Contributions:** Conceptualization, J.J. and I.P.; methodology, J.J.; formal analysis, I.P. and J.G.; writing—original draft preparation, I.P. and J.J.; writing—review and editing, J.J.; visualization, I.P.; corrections and consultations, A.P.; supervision, J.J. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by SHEER: “Shale Gas Exploration and Exploitation Induced Risks” project funded from Horizon 2020–R&I Framework Programme, call H2020-LCE-2014-1, Grant Agreement No. 640896 and partially supported by the Polish Ministry of Education and Science (Decision no. 13/E-41/SPUB/SP/2020) and the Chief Inspectorate of Environmental Protection, Poland under Grant number GIOŚ/19/2021/DMS/NFOŚ.

**Institutional Review Board Statement:** Not applicable.


**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.


**Conflicts of Interest:** The authors declare no conflict of interest.

### Appendix A


Air pollutant concentrations measured at the Stary Wiec station vs. wind direction





**Figure A1.** Average levels of NO and NO<sub>2</sub> vs. wind direction measured at the Stary Wiec station during drilling (DR), hydrofracturing (HF), and for the remaining period of measurements. All concentration units are in parts per billion (ppb). The red bars indicate the air mass coming directly from the area of the wells.




**Figure A2.** Average levels of CO and PM<sub>10</sub> vs. wind direction measured at the Stary Wiec station during drilling (DR), hydrofracturing (HF), and for the remaining period of measurements. Concentration units are in parts per billion (ppb), except for PM<sub>10</sub> (µg/m<sup>3</sup>). The red bars indicate the air mass coming directly from the area of the wells.



**Figure A3.** Average levels of O<sub>3</sub> vs. wind direction measured at the Stary Wiec station during drilling (DR), hydrofracturing (HF), and for the remaining period of measurements. All concentration units are in parts per billion (ppb). The red bars indicate the air mass coming directly from the area of the wells.



**Figure A4.** Average levels of CO<sub>2</sub> and CH<sub>4</sub> vs. wind direction measured at the Stary Wiec station during drilling (DR), hydrofracturing (HF), and for the remaining period of measurements. Concentrations units for CO<sub>2</sub> and CH<sub>4</sub> are parts per million (ppm).



**Figure A5.** Average levels of NMHC vs. wind direction measured at the Stary Wiec station during drilling (DR), hydrofracturing (HF), and for the remaining period of measurements. Concentration units for NMHC are ppmc.

Methane was the pollutant for which there was the most distinct influence of shale gas extraction activities on ambient air quality at the Stary Wiec station. From 1 to 3 September 2016, a series of short (registered on one-min average data) methane concentration peaks, with a maximum value equal to 7.45 ppm, were noticed. The way in which the repeatability of these peaks occurred indicated a probable artificial origin of the

episode, whereas analysis of wind direction revealed the borehole area as the potential source. To confirm these assumptions, a 3D numerical simulation of dispersion of CH<sub>4</sub> in the area surrounding a station was performed. The borehole area located to the north-west of the station was selected as an emission point. The results of modeling for 01.09.2016 7 pm are shown in Figure A6 A pollutant plume emitted at the well location extended from the NW and passed through the measurement site. The calculation of the pollutant emission rate on the basis of hourly mean pollutant concentration had height 4 m a.g.l. points to the value of 120 kg/h.

The numerical study used the Weather Research and Forecasting (WRF) Model [42] as a source of meteorological data and the HYSPLIT dispersion model [43] for the CH<sub>4</sub> modeling.

The WRF Model is a mesoscale numerical weather prediction system, dedicated to atmospheric research and forecasting of applications. It contains two dynamical cores and serves as a data assimilation system and software architecture, enabling parallel calculations and system extensibility. The Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT), developed by NOAA's Air Resources Laboratory, is one of the most widely used models for atmospheric trajectory and dispersion calculations. This model is widely used for both determining atmospheric trajectory and dispersion calculations.

The initial and boundary meteorology conditions for the WRF Model were used from the NCAR GFS [44] system.

WRF settings:

Domain solution

d1: 14 km × 14 km 51 vertical level 1 h

d2: 4.5 km × 4.5 km 51 vertical level 1 h

d3: 1.2 km × 1.2 km 51 vertical level 15 min

d4: 0.4 km × 0.4 km 51 vertical level 1 min

Simulation start: 01.09.2016 12 UTC end 02.09.2016 00 UTC

The Physics and microphysics schema configuration for WRF Model was performed based on the study by Werner et al. [45].


HYSPLIT settings:

Domain solution:

Setting for the domain was 0.4 km × 0.4 km. Vertically it was 40 levels between 0 m to 224 m above ground level.

**Table A1.** Technical specifications of instruments employed during measuring campaign.

Measured Pollutant	NO/NO <sub>2</sub> /NO <sub>x</sub>	CO	O <sub>3</sub>	PM <sub>10</sub>	CO <sub>2</sub>	CH <sub>4</sub> /NMHC
Instrument	API 200	API 300	API400	TEOM 1400	Thermo 410i	Horiba APHA370
Measurement technique	Chemiluminescence	Infrared absorption	Absorption of UV light	Tapered Element Oscillating Microbalance (TEOM)	NDIR technology	Selective combustion+cross modulation+flame ionization detection
Zero drift	<0.5 ppb/24 h	<20 ppb/24 h	<1.0 ppb/24 h		±1 ppm	±0.1 ppmC/day
Lower detectable limit	0.4 ppb	<20 ppb	<0.6 ppb	<5 µg/m <sup>3</sup>	1 ppm	0.02 ppmC (3σ)
Precision	0.5%	0.5%	0.5%	0.75%	1%	1%
Range	0–50 ppb to 0–20 ppm	0–100 ppb to 0–100 ppm	100 ppb to 10 ppm	0–5 mg/m <sup>3</sup>		0 ppmC–50 ppmC



**Figure A6.** Methane plume modeled by the HYSPLIT model for 1 September 2016, revealed the following values: CH<sub>4</sub> = 2.81 ppm (hourly mean), NMHC = 0.2 ppm, estimated CH<sub>4</sub> emission rate = 120 kg/h. Left and horizontal axis-geographical coordinates, right axis-additional methane concentration above the Stry Wiec background level, i.e., approximately 1.9 ppm. Red circle-well location, black circle-monitoring site.

**Table A2.** Statistics for air pollution data from air quality monitoring stations based on hourly means, collected from January 2012 to July 2015. Mean values for this period, together with standard deviations ( $1\sigma$ ), are presented.

	Liniewko	Kościerzyna	Tczew	Starogard	Gdańsk
NO (ppb)	0.86 ± 0.90	5.72 ± 14.33	4.30 ± 10.18	7.37 ± 11.56	2.93 ± 6.64
NO <sub>2</sub> (ppb)	3.38 ± 2.44	8.30 ± 6.00	7.30 ± 5.60	8.34 ± 5.97	7.67 ± 6.03
O <sub>3</sub> (ppb)	30.14 ± 11.89	22.20 ± 12.81			23.97 ± 12.13
CO (ppb)	255.36 ± 107.12	437.45 ± 307.92	309.61 ± 189.51	386.57 ± 278.28	266.34 ± 136.86
PM10 (µg/m <sup>3</sup> )		39.73 ± 36.80	22.41 ± 17.37	39.65 ± 39.31	17.06 ± 12.81

## References

- Schnell, R.C.; Oltmans, S.J.; Neely, R.R.; Endres, M.S.; Molenaar, J.V.; White, A.B. Rapid photochemical production of ozone at high concentrations in a rural site during winter. *Nat. Geosci.* **2009**, *2*, 120–122. [[CrossRef](#)]
- Katzenstein, A.S.; Doezema, L.A.; Simpson, I.J.; Blake, D.R.; Rowland, F.S. Extensive regional atmospheric hydrocarbon pollution in the southwestern United States. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 11975–11979. [[CrossRef](#)] [[PubMed](#)]
- Skone, T.J. *Life Cycle Greenhouse Gas Inventory of Natural Gas Extraction, Delivery and Electricity Production*; Final Report; National Energy Technology Laboratory: Pittsburgh, PA, USA, 2011. Available online: <http://www.canadiancleanpowercoalition.com/files/4713/2630/3388/LA7%20-%20NG-GHG-LCI.pdf> (accessed on 27 March 2018).
- Sun, R.; Wang, Z. A comprehensive environmental impact assessment method for shale gas development. *Nat. Gas Ind. B* **2015**, *2*, 203–210. [[CrossRef](#)]
- Roy, A.A.; Adams, P.J.; Robinson, A.L. Air pollutant emissions from the development, production, and processing of Marcellus Shale natural gas. *J. Air Waste Manag. Assoc.* **2014**, *64*, 19–37. [[CrossRef](#)]

6. U.S. Environmental Protection Agency (EPA). Nonroad Engines, Equipment and Vehicles: Locomotives. 2013. Available online: <http://www.epa.gov/otaq/locomotives.htm> (accessed on 22 March 2018).
7. U.S. Environmental Protection Agency (EPA). Nonroad Engines, Equipment and Vehicles: Nonroad Diesel Engines. 2013. Available online: <http://www.epa.gov/otaq/nonroad-diesel.htm> (accessed on 22 March 2018).
8. Purvis, R.M.; Lewis, A.C.; Hopkins, J.R.; Wilde, S.E.; Dunmore, R.E.; Allen, G.; Pitt, J.; Ward, R.S. Effects of ‘pre-fracking’ operations on ambient air quality at a shale gas exploration site in rural North Yorkshire, England. *Sci. Total Environ.* **2019**, *673*, 445–454. [[CrossRef](#)] [[PubMed](#)]
9. Brandt, A.R.; Heath, G.A.; Kort, E.A.; O’Sullivan, F.; Pétron, G.; Jordaan, S.M.; Tans, P.; Wilcox, J.; Gopstein, A.M.; Arent, D.; et al. Methane Leaks from North American Natural Gas Systems. *Science* **2014**, *343*, 733–735. [[CrossRef](#)] [[PubMed](#)]
10. Armendariz, A. Emissions from Natural Gas Production in the Barnett Shale Area and Opportunities for Cost-Effective Improvements. Ph.D. Thesis, Southern Methodist University, Dallas, TX, USA, 2009.
11. Kembal-Cook, S.; Bar-Ilan, A.; Grant, J.; Parker, L.; Jung, J.; Santamaria, W.; Mathews, J.; Yarwood, G. Ozone impacts of Natural gas development in the Haynesville shale. *Environ. Sci. Technol.* **2010**, *44*, 9357–9363. [[CrossRef](#)] [[PubMed](#)]
12. Orak, N.H.; Reeder, M.; Pekney, N.J. Identifying and quantifying source contributions of air quality contaminants during unconventional shale gas extraction. *Atmos. Chem. Phys.* **2021**, *21*, 4729–4739. [[CrossRef](#)]
13. Burnham, A.; Clark, C. *Examining the Impacts of Methane Leakage on Life-Cycle Greenhouse Gas Emissions of Shale and Conventional Natural Gas*; Air & Waste Management Association: Pittsburgh, PA, USA, 2012; pp. 8–12.
14. De Gouw, J.A.; Parrish, D.D.; Frost, G.J.; Trainer, M. Reduced emissions of CO<sub>2</sub>, NO<sub>x</sub>, and SO<sub>2</sub> from US power plants owing to switch from coal to natural gas with combined cycle technology. *Earth’s Future* **2014**, *2*, 75–82. [[CrossRef](#)]
15. Song, W.; Chang, Y.; Liu, X.; Li, K.; Gong, Y.; He, G.; Wang, X.; Christie, P.; Zheng, M.; Dore, A.J.; et al. A Multiyear Assessment of Air Quality Benefits from China’s Emerging Shale Gas Revolution: Urumqi as a Case Study. *Environ. Sci. Technol.* **2015**, *49*, 2066–2072. [[CrossRef](#)] [[PubMed](#)]
16. Zhang, Y.; Wang, H.; Han, Y.; Wang, D.; Zhu, G.; Lu, X. Shale gas development in China: Implications for indoor and outdoor air quality and greenhouse gas emissions. *Environ. Int.* **2020**, *141*, 105727. [[CrossRef](#)]
17. Wang, S.; Tang, X.; Wang, J.; Zhang, B.; Sun, W.; Höök, M. Environmental impacts from conventional and shale gas and oil development in China considering regional differences and well depth. *Resour. Conserv. Recycl.* **2021**, *167*, 105368. [[CrossRef](#)]
18. Javadi, M.A.; Khalaji, M.; Ghasemiasl, R. Exergoeconomic and environmental analysis of a combined Power and water desalination plant with parabolic solar collector. *Desalination Water Treat.* **2020**, *193*, 212–223. [[CrossRef](#)]
19. Ghasemiasl, R.; Abhari, M.K.; Javadi, M.A.; Ghomashi, H. 4E investigating of a combined power plant and converting it to a multigeneration system to reduce environmental pollutant production and sustainable development. *Energy Convers. Manag.* **2021**, *245*, 114468. [[CrossRef](#)]
20. Javadi, M.A.; Najafi, N.J.; Abhari, M.K.; Jabery, R.; Pourtaba, H. 4E analysis of three different configurations of a combined cycle power plant integrated with a solar power tower system. *Sustain. Energy Technol. Assess.* **2021**, *48*, 101599. [[CrossRef](#)]
21. Meng, Q. The impacts of fracking on the environment: A total environmental study paradigm. *Sci. Total Environ.* **2017**, *580*, 953–957. [[CrossRef](#)]
22. Karion, A.; Sweeney, C.; Pétron, G.; Frost, G.; Hardesty, R.M.; Kofler, J.; Miller, B.R.; Newberger, T.; Wolter, S.; Banta, R.; et al. Methane emissions estimate from airborne measurements over a western United States natural gas field. *Geophys. Res. Lett.* **2013**, *40*, 4393–4397. [[CrossRef](#)]
23. Shaw, J.T.; Allen, G.; Pitt, J.; Shah, A.; Wilde, S.; Stamford, L.; Fan, Z.; Ricketts, H.; Williams, P.I.; Bateson, P.; et al. Methane flux from flowback operations at a shale gas site. *J. Air Waste Manag. Assoc.* **2020**, *70*, 1324–1339. [[CrossRef](#)] [[PubMed](#)]
24. Ahmadov, R.; McKeen, S.; Trainer, M.; Banta, R.; Brewer, A.; Brown, S.; Edwards, P.M.; de Gouw, J.A.; Frost, G.J.; Gilman, J.; et al. Understanding high wintertime ozone pollution events in an oil—And natural gas—Producing region of the western US. *Atmos. Chem. Phys.* **2015**, *15*, 411–429. [[CrossRef](#)]
25. Oltmans, S.; Schnell, R.; Johnson, B.; Pétron, G.; Mefford, T.; Neely III, R. Anatomy of wintertime ozone associated with oil and natural gas extraction activity in Wyoming and Utah. *Elem. Sci. Anthr.* **2014**, *2*, 24. [[CrossRef](#)]
26. Rappenglück, B.; Ackermann, L.; Alvarez, S.; Golovko, J.; Buhr, M.; Field, R.A.; Soltis, J.; Montague, D.C.; Hauze, B.; Adamson, S.; et al. Strong wintertime ozone events in the Upper Green River basin Wyoming. *Atmos. Chem. Phys.* **2014**, *14*, 4909–4934. [[CrossRef](#)]
27. Cheadle, L.C.; Oltmans, S.J.; Pétron, G.; Schnell, R.C.; Mattson, E.J.; Herndon, S.C.; Thompson, A.M.; Blake, D.R.; McClure-Begley, A. Surface ozone in the Colorado northern Front Range and the influence of oil and gas development during FRAPPE/DISCOVER-AQ in summer 2014. *Elem. Sci. Anthr.* **2017**, *5*, 61. [[CrossRef](#)]
28. Benedict, K.B.; Zhou, Y.; Sive, B.C.; Prenni, A.J.; Gebhart, K.A.; Fisher, E.V.; Evansoski-Cole, A.; Sullivan, A.P.; Callahan, S.; Schichtel, B.A.; et al. Volatile organic compounds and ozone in Rocky Mountain National Park during FRAPPÉ. *Atmos. Chem. Phys.* **2019**, *19*, 499–521. [[CrossRef](#)]
29. Lindaas, J.; Farmer, D.K.; Pollack, I.B.; Abeleira, A.; Flocke, F.; Fisher, E.V. Acyl Peroxy Nitrates Link Oil and Natural Gas Emissions to High Ozone Abundances in the Colorado Front Range During Summer 2015. *J. Geophys. Res. Atmos.* **2019**, *124*, 2336–2350. [[CrossRef](#)]
30. Arthur, J.D.; Langhus, B.; Alleman, D. An overview of modern shale gas development in the United States. *All Consult.* **2008**, *3*, 14–17.

31. Le, M.T. An assessment of the potential for the development of the shale gas industry in countries outside of North America. *Heliyon* **2018**, *4*, 2. [[CrossRef](#)]
32. Patel, S. Europe Embraces Shale Gas. *Power Magazine*, 1 April 2013. Available online: <https://www.powermag.com/europe-embraces-shale-gas/> (accessed on 26 May 2022).
33. Shah, A.; Ricketts, H.; Pitt, J.R.; Shaw, J.T.; Kabbabe, K.; Leen, J.B.; Allen, G. Unmanned aerial vehicle observations of cold venting from exploratory hydraulic fracturing in the United Kingdom. *Environ. Res. Commun.* **2020**, *2*, 021003. [[CrossRef](#)]
34. Państwowy Instytut Badawczy (PIG). Ocena Zasobów Wydobywalnych Gazu Ziarnego i Ropy Naftowej w Formacjach Łupkowych Dolnego Paleozoiku w Polsce (Basen Bałtycko-Podlasko-Lubelski). Raport Pierwszy; 2012. Available online: <https://www.pgi.gov.pl/docman-tree-all/aktualnosci-2012/zasoby-gazu/771-raport-pl/file.html> (accessed on 23 March 2018).
35. Polish Geological Institute. Environmental Aspects of Hydraulic Fracturing Treatment Performed on the Łebień LE-2H Well. Final Report; 2011. Available online: <https://www.pgi.gov.pl/en/dokumenty-pig-pib-all/kopalnia-wiedzy/gaz-lupkowy/786-the-lebien-report/file.html> (accessed on 10 September 2021).
36. Pawlak, I.; Jarosławski, J. The influence of selected meteorological parameters on the concentration of surface ozone in the central region of Poland. *Atmos. Ocean.* **2015**, *53*, 126–139. [[CrossRef](#)]
37. Saitanis, C.J. Background ozone monitoring and phytodetection in the greater rural area of Corinth–Greece. *Chemosphere* **2003**, *51*, 913–923. [[CrossRef](#)]
38. Nęcki, J.M.; Gałkowski, M.; Chmura, Ł.; Gerbig, C.; Zimnoch, M.; Zięba, D.; Bartyzel, J.; Wołkowicz, W.; Różański, K. Regional Representativeness of CH<sub>4</sub> and NO<sub>2</sub> Mixing Ratio Measurements at High-Altitude Mountain Station Kasprowy Wierch, Southern Poland. *Aerosol Air Qual. Res.* **2016**, *16*, 568–580. [[CrossRef](#)]
39. Różański, K.; Chmura, Ł.; Gałkowski, M.; Nęcki, J.; Zimnoch, M.; Bartyzel, J. Monitoring of greenhouse gases in the atmosphere—A Polish perspective. *Pap. Glob. Chang.* **2016**, *23*, 111–126. [[CrossRef](#)]
40. Popa, M.E.; Vollmer, M.K.; Jordan, A.; Brand, W.A.; Pathirana, S.L.; Rothe, M.; Röckmann, T. Vehicle emissions of greenhouse gases and related tracers from a tunnel study: CO:CO<sub>2</sub>, N<sub>2</sub>O:CO<sub>2</sub>, CH<sub>4</sub>:CO<sub>2</sub>, O<sub>2</sub>:CO<sub>2</sub> ratios, and the stable isotopes <sup>13</sup>C and <sup>18</sup>O in CO<sub>2</sub> and CO. *Atmos. Chem. Phys.* **2013**, *14*, 2105–2123. [[CrossRef](#)]
41. Stanley, E.H.; Casson, N.J.; Christel, S.T.; Crawford, J.T.; Loken, L.C.; Oliver, S.K. The ecology of methane in streams and rivers: Patterns, controls, and global significance. *Ecol. Monogr.* **2016**, *86*, 146–171. [[CrossRef](#)]
42. Skamarock, W.C.; Klemp, J.B.; Dudhia, J.; Gill, D.O.; Liu, Z.; Berner, J.; Wang, W.; Powers, J.G.; Duda, M.G.; Barker, D.M.; et al. *A Description of the Advanced Research WRF Model Version 4.1*; National Center for Atmospheric Research: Boulder, CO, USA, 2019. [[CrossRef](#)]
43. Stein, A.F.; Draxler, R.R.; Rolph, G.D.; Stunder, B.J.B.; Cohen, M.D.; Ngan, F. NOAA’s HYSPLIT atmospheric transport and dispersion modeling system. *Bull. Amer. Meteor. Soc.* **2015**, *96*, 2059–2077. [[CrossRef](#)]
44. NCEP/National Weather Service/NOAA/U.S. Department of Commerce, NCEP GFS 0.25 Degree Global Forecast Auxiliary Grids Historical Archive; Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory: Boulder, CO, USA, 2015. [[CrossRef](#)]
45. Werner, M.; Kryza, M.; Guzikowski, J. Can Data Assimilation of Surface PM<sub>2.5</sub> and Satellite AOD Improve WRF-Chem Forecasting? A Case Study for Two Scenarios of Particulate Air Pollution Episodes in Poland. *Remote Sens.* **2019**, *11*, 2364. [[CrossRef](#)]