

Erratum



Erratum: Feng *et al.* A Method for Deriving the Boundary Layer Mixing Height from MODIS Atmospheric Profile Data. *Atmosphere*, 2015, 6, 1346-1361

Xueliang Feng, Bingfang Wu * and Nana Yan

Key Laboratory of Digital Earth Science, Institute of Remote Sensing and Digital Earth (RADI), Chinese Academy of Sciences, Beijing 100094, China; fengxl@radi.ac.cn (X.F.); yannn@radi.ac.cn (N.Y.)

* Correspondence: wubf@radi.ac.cn; Tel.: +86-10-6485-5689; Fax: +86-10-6485-8721

Academic Editors: John Boland and Robert W. Talbot Received: 23 February 2016; Accepted: 3 March 2016; Published: 10 March 2016

The authors would like to correct the acknowledgements of this article [1] as follows:

Acknowledgments: This research was supported by the National Natural Sciences Foundation of China (Number: 41271424 and 91025007), and Special Fund for Grains-Scientific Research in the Public Interest (Grant No. 201313009-2). We thank the HiWATER project for providing radiosonde data, and the MODIS data-processing group for atmospheric moisture profile data. We would also like to thank the anonymous reviewers whose useful comments improved the paper.

The authors would like to apologize for any inconvenience this may have caused to the readers.

References

1. Feng, X.; Wu, B.; Yan, N. A method for deriving the boundary layer mixing height from MODIS atmospheric profile data. *Atmosphere* **2015**, *6*, 1346–1361. [CrossRef]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (http://creativecommons.org/licenses/by/4.0/).