

Supplementary

Table 1. Environmental data of lakes extracted from the literature and considered in the analyses.

	Lat.	Lon.	Altitude (m)	Area (km ²)	Max. depth (m)	EC. (μS/cm)	pH	Annual P. (mm)	Average T. (°C)	Min. T. (°C)	Max. T. (°C)
Amvrakia	38.76	21.18	28	14.53	50	950	7.5	815.37	17.77	5.94	29.1
Kastoria	40.52	21.29	629	28.85	9.1	325	8	894.94	12.34	-3.15	25.04
Lysimachia	38.56	21.38	16	13.05	8.1	332	7.8	713.34	17.98	6.29	29.25
Megali Prespa	40.90	21.00	852	281.68	55	220	8.5	827.98	12.02	-3.53	24.8
Mikri Prespa	40.75	21.08	853	47.38	8.4	293	7.8	842.96	12.05	-3.46	24.81
Ozeros	38.65	21.22	22	9.39	5.6	450	8	765.98	18.15	6.5	29.39
Pamvotis	39.66	20.88	470	19.24	7.5	353	8.21	1254.15	13.93	0.35	26.35
Petron	40.72	21.69	527	12.36	5	804	6.9	964.34	12.71	-3.24	25.08
Trichonis	38.56	21.55	18	96.52	57	320	8.1	707.01	17.08	4.97	28.56
Vegoritisi	40.74	21.78	524	53.97	48	718	7.6	982.81	12.78	-3.25	25.1
Volvi	40.67	23.48	37	72.02	27.3	1040	8.5	712.97	16.38	2.23	28.29
Zazari	40.62	21.55	602	1.7	5.5	177	7.5	954.54	12.63	-3.11	25.09
Zirou	39.24	20.85	49	0.42	70	333	8.18	1045.04	16.43	4.27	27.88
Skadar	42.18	19.30	6	365	8.3	225	8.15	1393.41	15.06	0.98	27.33
Ohrid	41.04	20.72	693	358	293	208	8.48	898.85	11.84	-3.38	24.76
Srebarna	44.11	27.07	11	1.9	3.3	567	9.03	743.37	13.36	-5.13	26.19




Figure S1. Partial dependence plots showing the responses of fish (left column of plots) and macrophyte richness (right column) to each predictor.