

Supplementary Information

Enhanced Adsorptive Removal of Dyes using Mandarin Peel Biochars via Chemical Activation with NH₄Cl and ZnCl₂

Hyunjun Park ¹, Jiseok Kim ¹, Yong-Gu Lee ^{1,*}, and Kangmin Chon ^{1,2,*}

¹ Department of Environmental Engineering, College of Engineering, Kangwon National University, Kangwondaehak-gil, 1, Chuncheon-si, Gangwon-do 24341, Republic of Korea; hjpark1597@naver.com (H.P.); skyjskim95@gmail.com (J.K.)

² Department of Integrated Energy and Infra system, Kangwon National University, Kangwondaehak-gil, 1, Chuncheon-si, Gangwon-do 24341, Republic of Korea

* Correspondence: yglee19@kangwon.ac.kr (Y.-G.L.); kmchon@kangwon.ac.kr (K.C.); Tel.: +82-33-250-6352 (K.C.)

Table S1. The physicochemical characteristics of the dyes.

Compound (Abbreviation)	Formula ^a	MW ^a (g/mol)	Charge ^a				log D ^a				log P ^a	pKa ^a
			pH 3.0	pH 5.0	pH 7.0	pH 9.0	pH 3.0	pH 5.0	pH 7.0	pH 9.0		
Methyl Orange (MO)	C ₁₄ H ₁₄ N ₃ NaO ₃ S	327.3	-0.21	-0.96	-1.00	-1.00	2.53	1.54	1.30	1.29	2.61	3.58
Fast Green FCF (FG)	C ₃₇ H ₃₄ N ₂ Na ₂ O ₁₀ S ₃	808.8	-1.24	-1.97	-2.01	-2.53	1.18	0.17	-	-	2.31	3.11

^a ChemAxon (<http://www.chemicalize.org>, 2nd January 2021); log D = distribution coefficient; log P = partition coefficient.