

Article

Differential response of nutrients to seasonal hydrological changes and a rain event in a subtropical watershed, southeastern China

Xiaolan Guo ¹, Xiuli Yan ^{1,2*}, Hongyan Bao ³, Junwen Wu ^{1,2} and Shuh-Ji Kao ^{3,4}

¹ Guangdong Provincial Key Laboratory of Marine Disaster Prediction and Prevention, Shantou University, Shantou 515063, China; 20xlguo@stu.edu.cn (X.G.); wujw@stu.edu.cn (J.W.)

² Southern Marine Science and Engineering Guangdong Laboratory, Guangzhou 511458, China

³ State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen 361002, China; baohy@xmu.edu.cn (H.B.); sjkao@xmu.edu.cn (S.-J.K.)

⁴ State Key Laboratory of Marine Resource Utilization in South China Sea, Hainan University, Haikou 570228, China

* Correspondence: yanxl@stu.edu.cn




Figure S1. Relationships between nutrient fluxes and river discharge in (a) the NS and (b) the WS.




Figure S2. Time series of nutrient fluxes in the NS (a-c) and the WS (e-f). (a) and (d) are SRP fluxes, (b) and (e) are DSi fluxes, (c) and (f) are DIN fluxes. Gray shading indicates the runoff.