

Supplementary Material

Reuse of Pretreated Agro-Industrial Wastewaters for Hydroponic Production of Lettuce

Alexandra Afonso ^{1,2,*}, Mariana Regato ³, Mariana Patanita ⁴, Silvana Luz ^{1,5}, Maria João Carvalho ¹, Annabel Fernandes ², Ana Lopes ², Adelaide Almeida ^{1,2}, Idália Costa ³ and Fátima Carvalho ^{1,2}

¹ Department of Applied Sciences and Technologies, Escola Superior Agrária de Beja, Instituto Politécnico de Beja, 7800-309 Beja, Portugal; silvana.g.luz@gmail.com (S.L.); joaobcarvalho@ipbeja.pt (M.J.C.); maalmeida@ipbeja.pt (A.A.); mfcarvalho@ipbeja.pt (F.C.)



² Fiber Materials and Environmental Technologies (FibEnTech-UBI), Universidade da Beira Interior, R. Marquês de Ávila e Bolama, 6201-001 Covilhã, Portugal; annabelf@ubi.pt (A.F.); analopes@ubi.pt (A.L.)

³ Department of Bioscience, Escola Superior Agrária de Beja, Instituto Politécnico de Beja, 7800-295 Beja, Portugal; mare@ipbeja.pt (M.R.); idalia.costa@ipbeja.pt (I.C.)

⁴ MED—Mediterranean Institute for Agriculture, Environment and Development & CHANGE—Global Change and Sustainability Institute, Institute for Advanced Studies and Research, Universidade de Évora, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal; mpatanita@uevora.pt

⁵ Departamento de Ingeniería Química y Química Física, Instituto Universitario de Investigación del Agua, Cambio Climático y Sostenibilidad (IACYS), Universidad de Extremadura, 06006 Badajoz, Spain

* Correspondence: alexandra.afonso@ipbeja.pt






Figure S1. Variation of (a) temperature, (b) ORP, (c) salinity, and (d) TDS in the different nutrient solutions, OMW-T, WW-T, CWW-T, and CNS, during the experimental period of the lettuce growth in the hydroponic system.