


Article

Space–Time Spectral Collocation Method for Solving Burgers Equations with the Convergence Analysis

Yu Huang ¹, Mohammad Hadi Noori Skandari ², Fatemeh Mohammadizadeh ²,
Hojjat Ahsani Tehrani ², Svetlin Georgiev Georgiev ³, Emran Tohidi ⁴ and Stanford Shateyi ^{5,*}

¹ College of Mathematics and Statistics, Nanjing University of Information Science and Technology, Nanjing 210044, China

² Faculty of Mathematical Sciences, Shahrood University of Technology, Shahrood 3619995161, Iran

³ Department of Applied Mathematics, Sorbonne University, 75005 Paris, France

⁴ Department of Mathematics, Kosar University of Bojnord, P. O. Box 9415615458, Bojnord, Iran

⁵ Department of Mathematics and Applied Mathematics, University of Venda, P. Bag X5050, Thohoyandu 0950, South Africa

* Correspondence: stanford.shateyi@univen.ac.za; Tel.: +27-15962-8163

Received: 27 August 2019; Accepted: 4 October 2019; Published: 22 November 2019



Abstract: This article deals with a numerical approach based on the symmetric space-time Chebyshev spectral collocation method for solving different types of Burgers equations with Dirichlet boundary conditions. In this method, the variables of the equation are first approximated by interpolating polynomials and then discretized at the Chebyshev–Gauss–Lobatto points. Thus, we get a system of algebraic equations whose solution is the set of unknown coefficients of the approximate solution of the main problem. We investigate the convergence of the suggested numerical scheme and compare the proposed method with several recent approaches through examining some test problems.

Keywords: Burgers equation; Chebyshev spectral collocation method; Chebyshev–Gauss–Lobatto nodes; convergence analysis

1. Introduction

Many phenomena in physics, biology and engineering can be modelled mathematically by partial differential equations (PDEs). The Burgers equation is one of the most important PDEs to be surveyed in the recent years by many researchers [1,2]. This equation describes various kinds of phenomena in plasma physics, solid state physics, optical fibers, fluid dynamics, chemical kinetics, non-linear acoustics, gas dynamics, traffic flow, etc.

Also, the generalized Burgers–Fisher equation is one of the most important classes of non-linear PDEs which has appeared in several categories of applications, such as shockwave formation, turbulence, heat conduction, sound waves in viscous medium, and some other fields of applied branches of science and engineering [3]. Moreover, The Burgers–Huxley equation has been considered to be an evolution equation that describes nerve pulse propagation in biology from which molecular CB properties shall be computed. The generalized Burgers–Huxley equation was investigated to describe the interaction between reaction mechanisms, convection effects, and diffusion transport [4].

Since an analytical in a closed-form solution is generally unavailable for non-linear PDEs, numerical methods are widely used for solving them. There are some effective numerical methods to solve PDEs, especially for the Burgers equation. In [5], a comprehensive review of some techniques is presented. Berger and Kohn in [6] used the mesh refinement method. Budd et al. in [7] applied mesh movement. Soheili et al. used a moving-mesh PDE (MMPDE) approach [8]. In [9], Ramadan et al. suggested a method based on collocation of septic B-splines over finite elements for numerical

solutions of the non-linear Burgers equation. Ramadan and El-Danaf considered the solution of the modified Burgers equation using the collocation method with quintic splines [10]. Haq et al. in [11] formulated simple classical radial basis functions (RBFs) collocation method for the numerical solution of the non-linear dispersive and dissipative Burgers equation. Both Orac et al. in [12] and Lepik in [13] investigated the numerical solutions using Haar wavelet. Inan and Bahadir described implicit exponential difference method in two cases: finite and fully finite [14]. Saka and Dog used quintic B-spline collocation procedure [15]. Irk in [16] used sextic B-spline collocation technique. In [17], Demiray suggested the hyperbolic tangent method and presented travelling wave solution for the perturbed Burgers equation. Hon and Mao in [18] applied the multiquadric (MQ) as a spatial approximation scheme. Schulze-Halberg discussed a linearization method for solving Burgers equation with time dependent coefficients and a non-linear forcing term [19]. In [20], Seydaoglu presented the high-order splitting methods. Guo et al. proposed a high-order finite-volume compact scheme [21]. In [22], Mukundan and Awasthi used new efficient numerical techniques for solving one-dimensional quasi-linear Burgers equation. Hammad and El-Azab solved Burgers–Huxley and Burgers–Fisher equations with discretization in time by a new linear approximation scheme and in space by a high order compact finite difference method in [23]. In [24], Arora and Kumar used a new numerical method entitled “modified cubic B-spline differential quadrature method (MCB-DQM)” to find the approximate solution of the Burgers equations. El-Wakil et al. presented the Burgers equation and some other PDEs with self-similar solutions [25]. Singh et al. in [26] approximated numerical solutions for the generalized Burgers–Huxley (gBH) equation using modified cubic B-spline differential quadrature method (MCB-DQM). The scheme was based on the differential quadrature method in which the weighted coefficients were computed using modified cubic B-splines as a set of basis functions.

The spectral collocation (SC) method is one of the most important methods to solve continuous-time problems including ODEs and PDEs systems in various fields of science and engineering. In this method, an interpolating polynomial is applied to approximate the unknown function. In fact, unknown function in the problem can be expressed in the term of the approximate values at the special nodes. This method shows suitable results in comparison with those of other methods, since it has used the orthogonal polynomials for instance, Legendre and Chebyshev polynomials.

Weinan in [27] analyzed numerical methods for some evolutionary equations which admit semigroup formulations and the author shows the spectral accuracy of the spectral and pseudospectral methods for the Burgers equation. Xiao et al. used the non-linear Petrov–Galerkin method to reduce the order of Navier–Stokes equations and improved the stability of ROM results without tuning parameters. Also, the obtained numerical results show that the proposed POD Petrov–Galerkin method gives more accurate and stable results than the corresponding results obtained by using the POD Bubnov–Galerkin method [28]. In [29], a generalized Langevin equation is investigated and shown that the form of its coefficients depends critically on the assumption of continuity of the reconstructed trajectory. In [30], the two-dimensional unsteady Burgers equation is presented and the authors use the 4-bit lattice Boltzmann model to solve the 2D unsteady Burgers equation. In [31], numerical solutions for the 2D Burgers equation are computed using higher-order accurate finite difference schemes. More precisely, the author used the fourth-order accurate Du Fort Frankel scheme for solving the two-dimensional Burgers equation.

In this paper, we apply the symmetric space-time Chebyshev SC (CSC) method for solving Burgers equation and compare the associated results with those of some other aforementioned well-known methods. Through the numerical examples, we show that CSC method is more effective than other methods and we can achieve to more precise results for the solution of Burgers equations. In fact, we see that the number of discretization points (i.e., the CGL points in CSC method) and also the error of CSC method are less than the others used to solve Burgers equations.

This paper is organized as follows: In Section 2, Burgers equation and its different types are introduced. The CSC method for Burgers equation is implemented in Section 3. In Section 4,

the convergence of the CSC method is analyzed. Next, the two-dimensional Burgers equation is introduced in Section 5 and the CSC method is applied to solve the equation. Then, Section 6 contains numerical examples to solve the Burgers equations in both cases of one and two-dimensional. Also, the figures of errors (associated with the proposed method) are depicted in some cases which confirm the efficiency of our suggested numerical scheme. Finally, the paper is concluded with a reasonable conclusion.

2. Burgers Equation

Three important types of Burgers equations are as follows:

1. For a field $V(.,.)$ and diffusion coefficient (or viscosity, as in the original fluid mechanical context) ε , the general form of viscous Burgers equation is as follows

$$V_t + VV_x - \varepsilon V_{xx} = 0, \quad a \leq x \leq b, \quad t > 0. \quad (1)$$

2. The Burgers–Fisher equation is a non-linear PDE of second order of the form

$$V_t - V_{xx} + \alpha(t)VV_x = \beta(t)V(1 - V), \quad a \leq x \leq b, \quad (2)$$

where $\alpha(.)$ and $\beta(.)$ are given functions. It plays an important role in various fields of gas dynamics, traffic flow, physics applications, financial and applied mathematics [23].

3. The Burgers–Huxley equation is as follows

$$V_t + \alpha V^\delta V_x - \beta V_{xx} = \gamma V(1 - V^\delta)(V^\delta - \eta), \quad 0 \leq x \leq 1, \quad (3)$$

where $\alpha, \beta, \delta, \gamma$ and η are given constants. The Burgers–Huxley as a non-linear PDE describes the interaction between reaction mechanisms, convection effects, and diffusion transports [23].

In this paper, we represent the Burgers Equations (1)–(3) as the following general form

$$V_t = \Lambda(t, V, V_x, V_{xx}). \quad (4)$$

The time initial and space boundary conditions (in Dirichlet form) for Burgers Equation (4) are usually given as follows

$$V(0, x) = f(x), \quad a \leq x \leq b, \quad (5)$$

$$V(t, a) = g_1(t), \quad t \in [T_0, T_1], \quad (6)$$

$$V(t, b) = g_2(t), \quad t \in [T_0, T_1]. \quad (7)$$

3. Implementing the CSC Method

Here, the CSC method for Burgers Equation (4) with conditions (5)–(7) is presented. The CSC method [32–34] is one of the most efficient numerical methods to solve continuous-time problems. Recently, some researchers have applied it to solve special problems (see [35–37]). One of the most important advantages of CSC method in comparison with other approximate methods is the high degree of accuracy that CSC approximations offer. Also, the CSC underlying polynomial space spanned by Chebyshev orthogonal polynomials on the interval $[-1, 1]$ with respect to a weight function $w(t) = \frac{1}{\sqrt{1-t^2}}$. In the CSC method, we use some points on the interval $[-1, 1]$ to discretize

the problem. Hence, we first transform the variables of Equation (4) to this interval by the following relations

$$\begin{cases} t = \frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2}, & t \in [T_0, T_1], \bar{t} \in [-1, 1], \\ x = \frac{b-a}{2} \bar{x} + \frac{b+a}{2}, & x \in [a, b], \bar{x} \in [-1, 1]. \end{cases} \quad (8)$$

Therefore, Burgers Equation (4)–(7) converted into the following form

$$\begin{cases} U_{\bar{t}} = \psi(\bar{t}, U(\bar{t}, \bar{x}), U_{\bar{x}}(\bar{t}, \bar{x}), U_{\bar{x}\bar{x}}(\bar{t}, \bar{x})), \\ U(-1, \bar{x}) = F(\bar{x}), \\ U(\bar{t}, -1) = G_1(\bar{t}), \\ U(\bar{t}, 1) = G_2(\bar{t}), \end{cases} \quad (9)$$

where

$$\begin{aligned} \psi(\bar{t}, U, U_{\bar{x}}, U_{\bar{x}\bar{x}}) &= \frac{T_1 - T_0}{2} \Lambda \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2}, U, U_{\bar{x}}, U_{\bar{x}\bar{x}} \right) \\ U(\bar{t}, \bar{x}) &= V \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2}, \frac{b-a}{2} \bar{x} + \frac{b+a}{2} \right), \\ U_{\bar{x}}(\bar{t}, \bar{x}) &= \frac{2}{b-a} V_x \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2}, \frac{b-a}{2} \bar{x} + \frac{b+a}{2} \right), \\ U_{\bar{x}\bar{x}}(\bar{t}, \bar{x}) &= \left(\frac{2}{b-a} \right)^2 V_{xx} \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2}, \frac{b-a}{2} \bar{x} + \frac{b+a}{2} \right), \\ F(\bar{x}) &= f \left(\frac{b-a}{2} \bar{x} + \frac{b+a}{2} \right), \\ G_1(\bar{t}) &= g_1 \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2} \right), \\ G_2(\bar{t}) &= g_2 \left(\frac{T_1 - T_0}{2} \bar{t} + \frac{T_1 + T_0}{2} \right). \end{aligned}$$

To discretize system (9), we use the CGL points on $[-1, 1]$ which are defined by the following relations

$$\bar{x}_k = \bar{t}_k = \cos\left(\frac{N-k}{N}\pi\right), \quad k = 0, 1, \dots, N, \quad (10)$$

where they are the roots of $(1 - \bar{t}^2) \frac{dT_N}{d\bar{t}}$ and $T_N(\cdot)$ is the Chebyshev polynomial of order N . It should be noted that the Chebyshev polynomials are expressed by

$$T_j(\bar{t}) = \cos(j \cos^{-1}(\bar{t})), \quad \bar{t} \in [-1, 1], \quad j = 0, 1, \dots, N, \quad (11)$$

and it is easy to show that

$$\begin{cases} T_0(\bar{t}) = 1, \quad T_1(\bar{t}) = \bar{t} \\ T_{j+1}(\bar{t}) = 2\bar{t}T_j(\bar{t}) - T_{j-1}(\bar{t}), \quad j = 1, 2, \dots \end{cases}$$

For interpolating in the CSC method, the following Lagrange polynomials are used

$$L_k(\bar{t}) = \prod_{\substack{j=0 \\ j \neq k}}^N \frac{\bar{t} - \bar{t}_j}{\bar{t}_k - \bar{t}_j} = \frac{2}{N\mu_k} \sum_{j=0}^N \frac{1}{\mu_j} T_j(\bar{t}_k) T_j(\bar{t}), \quad k = 0, 1, \dots, N, \quad \bar{t} \in [-1, 1],$$

where

$$\mu_j = \begin{cases} 2, & \text{if } j = 0, N, \\ 1, & \text{if } 1 \leq j \leq N-1, \end{cases}$$

and we have

$$L_j(\bar{t}_k) = \delta_{jk} = \begin{cases} 1, & \text{if } j = k, \\ 0, & \text{if } j \neq k. \end{cases} \quad (12)$$

In the CSC method, to approximate the solution of Burgers Equation (9), we use the following polynomial interpolation

$$U(\bar{t}, \bar{x}) \approx U^N(\bar{t}, \bar{x}) = \sum_{i=0}^N \sum_{j=0}^N \bar{a}_{ij}^N L_i(\bar{t}) L_j(\bar{x}). \quad (13)$$

By (12), we have

$$U^N(\bar{t}_i, \bar{x}_j) = \bar{a}_{ij}^N. \quad (14)$$

To express the derivative $U_{\bar{t}}^N(\cdot, \cdot)$, $U_{\bar{x}}^N(\cdot, \cdot)$ and $U_{\bar{x}\bar{x}}^N(\cdot, \cdot)$ in terms of $U^N(\cdot, \cdot)$ at the node points \bar{t}_k , we can use the matrix multiplication $D = (D_{kj})$ and get

$$\begin{cases} U_{\bar{t}}^N(\bar{t}_p, \bar{x}_k) = \sum_{i=0}^N \bar{a}_{ik}^N D_{pi}, \\ U_{\bar{x}}^N(\bar{t}_p, \bar{x}_k) = \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \\ U_{\bar{x}\bar{x}}^N(\bar{t}_p, \bar{x}_k) = \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj}, \quad k, p = 0, 1, \dots, N, \end{cases} \quad (15)$$

where

$$D_{kj} = L'_j(\bar{t}_k) = \begin{cases} \frac{\mu_k}{\mu_j} (-1)^{k+j} \frac{1}{\bar{t}_k - \bar{t}_j}, & \text{if } j \neq k, \\ -\frac{\bar{t}_k}{2 - 2\bar{t}_k^2}, & \text{if } 0 \leq j = k \leq N-1, \\ -\frac{2N^2 + 1}{6}, & \text{if } j = k = 0, \\ \frac{2N^2 + 1}{6}, & \text{if } j = k = N, \end{cases} \quad (16)$$

$\hat{D} = D \cdot D = (\hat{D}_{kj})$, $\hat{D}_{kj} = \sum_{l=0}^N D_{kl} D_{lj}$, $k, j = 0, 1, \dots, N$. In fact, multiplication by matrix D transforms a vector of the state variables at the CGL points to the vector of approximate derivatives at these points. Now, by relations (14) and (15), the system (9) can be converted into the following system of algebraic equations

$$\begin{cases} \sum_{i=0}^N \bar{a}_{ik}^N D_{pi} - \psi(\bar{t}_p, \bar{a}_{pk}^N, \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj}) = 0, \\ \bar{a}_{0k}^N = F(\bar{x}_k); \quad k = 0, 1, \dots, N, \\ \bar{a}_{p0}^N = G_1(\bar{t}_p), \quad \bar{a}_{pN}^N = G_2(\bar{t}_p); \quad p = 1, \dots, N. \end{cases} \quad (17)$$

Here, by solving above system with respect to $(\bar{a}_{pk}^N; p, k = 0, 1, \dots, N)$, we can obtain continuous and pointwise approximate solutions (13) and (14), respectively.

4. The Convergence of the Method

In this section, first we give definition of the modulus of continuity and then analyze the convergence of the presented method.

Assume that $\bar{\Omega} = [-1, 1] \times [-1, 1]$. With $C^r(\bar{\Omega})$ we denote the space of the continuous functions with continuous derivatives of r th order.

Definition 1. Function $W : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the following properties is called a modulus of continuity [38]

1. W is increasing,
2. $\lim_{z \rightarrow 0} W(z) = 0$,
3. for any z_1 and $z_2 \in \mathbb{R}^+$, $W(z_1 + z_2) \leq W(z_1) + W(z_2)$,
4. there exists a constant c such that for all $0 < z \leq 2$, $cW(z) \geq z$.

Some important modulus of continuity can be defined as

$$W(z) = z^\alpha, \quad 0 < \alpha \leq 1. \quad (18)$$

Now, assume that B^2 is a unit circle in \mathbb{R}^2 . We say that a continuous function $f(\cdot, \cdot)$ on $\bar{\Omega}$ admits $W(\cdot)$ as a modulus of continuity, if the following value is finite

$$|f(\cdot, \cdot)|_W = \sup \left\{ \frac{|f(\bar{t}, \bar{x}) - f(\tilde{t}, \tilde{x})|}{W(\|(\bar{t}, \bar{x}) - (\tilde{t}, \tilde{x})\|_\infty)} : (\bar{t}, \bar{x}), (\tilde{t}, \tilde{x}) \in \bar{\Omega}, (\bar{t}, \bar{x}) \neq (\tilde{t}, \tilde{x}) \right\}, \quad (19)$$

where

$$\|(\bar{t}, \bar{x}) - (\tilde{t}, \tilde{x})\|_\infty = \max\{|\bar{t} - \tilde{t}|, |\bar{x} - \tilde{x}|\}; \quad (\bar{t}, \bar{x}), (\tilde{t}, \tilde{x}) \in \bar{\Omega}, (\bar{t}, \bar{x}) \neq (\tilde{t}, \tilde{x}). \quad (20)$$

With $C_W^1(B^2)$ we denote the space of all functions $f(\cdot, \cdot)$ on B^2 with continuous first-order partial derivatives, and let it be endowed with the following norm

$$\|f(\cdot, \cdot)\|_{1,W} = \|f(\cdot, \cdot)\|_\infty + \|f_{\bar{t}}(\cdot, \cdot)\|_\infty + \|f_{\bar{x}}(\cdot, \cdot)\|_\infty + |f_{\bar{t}}(\cdot, \cdot)|_W + |f_{\bar{x}}(\cdot, \cdot)|_W. \quad (21)$$

Next, define

$$C_W^1(\bar{\Omega}) = \left\{ f(\cdot, \cdot) \in C^1(\bar{\Omega}) : \forall (\bar{t}, \bar{x}) \in \bar{\Omega}, \exists \text{ map } \phi : B^2 \rightarrow \bar{\Omega}, \text{ s.t. } \right. \\ \left. (\tilde{t}, \tilde{x}) \in \text{int}(\phi(B^2)) \text{ and } f \circ \phi(\cdot, \cdot) \in C_W^1(B^2) \right\}. \quad (22)$$

It can be proved that if $\bar{\Omega} = \bigcup_{i=1}^l \text{int}(\phi_i(B^2))$ for some ϕ_1, \dots, ϕ_l , then $f(\cdot, \cdot) \in C_W^1(\bar{\Omega})$ if and only if $f \circ \phi_i(\cdot, \cdot) \in C_W^1(B^2)$ for each $i = 1, \dots, l$. Moreover, $C_W^1(\bar{\Omega})$ with norm

$$\|f(\cdot, \cdot)\|_{1,W} = \sum_{i=1}^l \|f \circ \phi_i(\cdot, \cdot)\|_{1,W}, \quad (23)$$

is a Banach space (for more details see [38]). At follows, we show the space of all polynomials of total degree at most $2N$ on $\bar{\Omega}$ by $\text{Pol}(N, N, \bar{\Omega})$, i.e.,

$$\text{Pol}(N, N, \bar{\Omega}) = \left\{ \eta(\bar{t}, \bar{x}) = \sum_{i=0}^N \sum_{j=0}^N \gamma_{ij} \bar{t}^i \bar{x}^j : (\bar{t}, \bar{x}) \in \bar{\Omega}, \gamma_{ij} \in \mathbb{R} \right\}.$$

Theorem 1. For any $f(\cdot, \cdot) \in C_W^1(\bar{\Omega})$, there is a polynomial $\eta(\cdot, \cdot) \in \text{Pol}(N, N, \bar{\Omega})$ such that

$$\|f(\cdot, \cdot) - \eta(\cdot, \cdot)\|_\infty \leq \frac{c_0 c_1}{2N} W\left(\frac{1}{2N}\right), \quad (24)$$

where $c_1 = \|f(\cdot, \cdot)\|_{1,W}$ and c_0 is a constant that depends on $W(\cdot)$, but independent of N .

Proof. The proof is a result of Theorem 2.1 in [38]. \square

To prove the existence of solutions of the system (17), we convert it in the following form

$$\begin{cases} |\sum_{i=0}^N \bar{a}_{ik}^N D_{pi} - \psi(\bar{t}_p, \bar{a}_{pk}^N, \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj})| \\ \leq \frac{\sqrt{N}}{2N-1} W(\frac{1}{2N-1}), \quad p, k = 1, 2, \dots, N-1, \\ |\bar{a}_{p0}^N - G_1(\bar{t}_p)| \leq \frac{\sqrt{N}}{2N-1} W(\frac{1}{2N-1}), |\bar{a}_{pN}^N - G_2(\bar{t}_p)| \leq \frac{\sqrt{N}}{2N-1} W(\frac{1}{2N-1}), \quad p = 0, 1, \dots, N, \\ |\bar{a}_{0k}^N - F(\bar{x}_k)| \leq \frac{\sqrt{N}}{2N-1} W(\frac{1}{2N-1}), \quad k = 0, 1, \dots, N, \end{cases} \quad (25)$$

where N is sufficiently big and $W(\cdot)$ is a given modulus of continuity. Since $\lim_{N \rightarrow \infty} \frac{\sqrt{N}}{2N-1} W(\frac{1}{2N-1}) = 0$, any solution $\bar{a}_N = (\bar{a}_{pk}^N; p, k = 0, 1, \dots, N)$ for system of algebraic inequalities (25) is a solution for system of algebraic Equation (17) when N tends to infinity.

We assume that ψ has bounded and continuous derivatives with respect to its arguments. Hence, there exists a constant M such that

$$|\psi(\bar{t}, U, U_{\bar{x}}, U_{\bar{x}\bar{x}}) - \psi(\bar{t}, \tilde{U}, \tilde{U}_{\bar{x}}, \tilde{U}_{\bar{x}\bar{x}})| \leq M|U - \tilde{U}|. \quad (26)$$

Now we will show that the system (25) has at least one solution \bar{a}_N .

Theorem 2. Let $U(\cdot, \cdot)$ be a solution for system (9) where $U(\cdot, \cdot)$ is in $C_W^1(\bar{\Omega})$. Then there is a positive integer K such that for any $N \geq K$, system (25) has a solution as

$$\bar{a}_N = (\bar{a}_{pk}; p, k = 0, 1, \dots, N), \quad (27)$$

which satisfies

$$|U(\bar{t}_p, \bar{x}_k) - \bar{a}_{pk}^N| \leq \frac{L}{2N-1} W(\frac{1}{2N-1}), \quad p, k = 0, 1, \dots, N, \quad (28)$$

where L is a positive constant independent of N .

Proof of Theorem 2. Assume that $\eta(\cdot, \cdot)$ in $Pol(N-1, N, \bar{\Omega})$ is the best polynomial approximation of $U_{\bar{t}}(\cdot, \cdot)$. By Theorem 1, we get

$$\|U_{\bar{t}}(\bar{t}, \bar{x}) - \eta(\bar{t}, \bar{x})\|_{\infty} \leq \frac{\gamma}{2N-1} W(\frac{1}{2N-1}), \quad (\bar{t}, \bar{x}) \in \bar{\Omega}, \quad (29)$$

where γ is a constant independent of N . We define

$$\tilde{U}(\bar{t}, \bar{x}) = U(-1, \bar{x}) + \int_{-1}^{\bar{t}} \eta(\tau, \bar{x}) d\tau, \quad (\bar{t}, \bar{x}) \in \bar{\Omega}, \quad (30)$$

and

$$\bar{a}_{pk}^N = \tilde{U}(\bar{t}_p, \bar{x}_k); \quad p, k = 0, 1, \dots, N. \quad (31)$$

We will see that $\bar{a}_N = (\bar{a}_{pk}^N; p, k = 0, 1, \dots, N)$ satisfies system (25). By (29)–(31), for $(\bar{t}, \bar{x}) \in \bar{\Omega}$, we get

$$\begin{aligned} |U(\bar{t}, \bar{x}) - \tilde{U}(\bar{t}, \bar{x})| &= \left| \int_{-1}^{\bar{t}} (U_{\bar{t}}(\tau, \bar{x}) - \eta(\tau, \bar{x})) d\tau \right| \leq \int_{-1}^{\bar{t}} |U_{\bar{t}}(\tau, \bar{x}) - \eta(\tau, \bar{x})| d\tau \\ &\leq \frac{\gamma}{2N-1} W(\frac{1}{2N-1}) \int_{-1}^{\bar{t}} d\tau \leq \frac{2\gamma}{2N-1} W(\frac{1}{2N-1}). \end{aligned} \quad (32)$$

Now, by the relation (30), the function $\tilde{U}(\cdot, \bar{x})$, $\bar{x} \in [-1, 1]$, is a polynomial of total degree at most $2N$. Hence, its derivatives at CGL nodes $\bar{t}_0, \bar{t}_1, \dots, \bar{t}_N$ are exactly equal to the value of the polynomial at the nodes multiplied by the differential matrix D , defined by (16). Thus, we have

$$\sum_{i=0}^N \bar{a}_{ik}^N D_{pi} = \tilde{U}_{\bar{t}}(\bar{t}_p, \bar{x}_k); \quad p, k = 0, 1, \dots, N. \quad (33)$$

Therefore, by the relations (26) and (32), we get

$$\begin{aligned} & \left| \sum_{i=0}^N \bar{a}_{ik}^N D_{pi} - \psi \left(\bar{t}_p, \bar{a}_{pk}^N, \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj} \right) \right| \\ & \leq |\tilde{U}_{\bar{t}}(\bar{t}_p, \bar{x}_k) - U_{\bar{t}}(\bar{t}_p, \bar{x}_k)| + \\ & \left| U_{\bar{t}}(\bar{t}_p, \bar{x}_k) - \psi \left(\bar{t}_p, \bar{a}_{pk}^N, \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj} \right) \right| \\ & = |\eta(\bar{t}_p, \bar{x}_k) - U_{\bar{t}}(\bar{t}_p, \bar{x}_k)| + \\ & \left| \psi \left(\bar{t}_p, U(\bar{t}_p, \bar{x}_k), U_{\bar{x}}(\bar{t}_p, \bar{x}_k), U_{\bar{x}\bar{x}}(\bar{t}_p, \bar{x}_k) \right) - \psi \left(\bar{t}_p, \bar{a}_{pk}^N, \sum_{j=0}^N \bar{a}_{pj}^N D_{kj}, \sum_{j=0}^N \bar{a}_{pj}^N \hat{D}_{kj} \right) \right| \\ & \leq |\eta(\bar{t}_p, \bar{x}_k) - U_{\bar{t}}(\bar{t}_p, \bar{x}_k)| + M \left| U(\bar{t}_p, \bar{x}_k) - \bar{a}_{pk}^N \right| \\ & \leq \frac{\gamma}{2N-1} W\left(\frac{1}{2N-1}\right) + M \frac{2\gamma}{2N-1} W\left(\frac{1}{2N-1}\right) \\ & = \frac{\gamma(2M+1)}{2N-1} W\left(\frac{1}{2N-1}\right), \quad p, k = 1, \dots, N-1, \end{aligned} \quad (34)$$

where M is a Lipschitz constant which satisfies the relation (26). Furthermore, for boundary conditions, we get

$$\begin{aligned} |\tilde{U}(-1, \bar{x}_k) - F(\bar{x}_k)| & \leq |\tilde{U}(-1, \bar{x}_k) - U(-1, \bar{x}_k)| + |U(-1, \bar{x}_k) - F(\bar{x}_k)| \\ & \leq \frac{2\gamma}{2N-1} W\left(\frac{1}{2N-1}\right), \quad k = 0, 1, \dots, N. \end{aligned} \quad (35)$$

Also, for all $p = 0, 1, \dots, N$,

$$|\tilde{U}(\bar{t}_p, -1) - G_1(\bar{t}_p)| = |\tilde{U}(\bar{t}_p, -1) - U(\bar{t}_p, -1)| \leq \frac{2\gamma}{2N-1} W\left(\frac{1}{2N-1}\right), \quad (36)$$

$$|\tilde{U}(\bar{t}_p, 1) - G_2(\bar{t}_p)| = |\tilde{U}(\bar{t}_p, 1) - U(\bar{t}_p, 1)| \leq \frac{2\gamma}{2N-1} W\left(\frac{1}{2N-1}\right). \quad (37)$$

Hence, if we select K such that

$$\max\{\gamma(2M+1), 2\gamma\} \leq \sqrt{N},$$

for $N \geq K$, then by (34)–(37), \bar{a}_N satisfies the system (25). This completes the proof. \square

Now, we will show that the sequence of the solutions for the system (25) and the sequence of their interpolating polynomials converge to the solution of the system (9).

Theorem 3. Let $\{\bar{a}_N = \bar{a}_{pk}^N; p, k = 0, 1, \dots, N\}_{N=K}^\infty$ be a sequence of solutions for the system (25) and $\{U^N(\cdot, \cdot)\}_{N=K}^\infty$ be their interpolating polynomials sequence defined by (13). Also, we assume that for any \bar{x} in $[-1, 1]$, the sequence $\{(U^N(-1, \bar{x}), U_{\bar{t}}^N(\cdot, \cdot))\}_{N=K}^\infty$ has a subsequence $\{(U^{N_i}(-1, \bar{x}), U_{\bar{t}}^{N_i}(\cdot, \cdot))\}_{i=0}^\infty$ that

uniformly converges to $(\phi^\infty(\bar{x}), \lambda(\cdot, \cdot))$, where $\lambda(\cdot, \cdot) \in C^2(\bar{\Omega})$, $\phi^\infty(\cdot) \in C^2([-1, 1])$ and $\lim_{i \rightarrow \infty} N_i = \infty$. Then the pair

$$\tilde{U}(\bar{t}, \bar{x}) = \lim_{i \rightarrow \infty} U^{N_i}(\bar{t}, \bar{x}), \quad (38)$$

for $(\bar{t}, \bar{x}) \in \bar{\Omega}$, is a solution of the system (9).

Proof of Theorem 3. By our assumptions, we have

$$\tilde{U}(\bar{t}, \bar{x}) = \phi^\infty(\bar{x}) + \int_{-1}^{\bar{t}} \lambda(\tau, \bar{x}) d\tau. \quad (39)$$

We first show that $\tilde{U}(\bar{t}, \bar{x})$ for $\bar{t} \in [-1, 1]$ and $\bar{x} = x_k$, $k = 0, 1, \dots, N$ satisfy the system (9). Assume that $\tilde{U}(\cdot, \bar{x}_k)$ for some $k = 1, \dots, N-1$ does not satisfy the first equation of (9). Then, there is a τ in $(-1, 1)$ such that

$$\tilde{U}_t(\tau, \bar{x}_k) - \psi(\bar{t}_p, \tilde{U}(\tau, \bar{x}_k), \tilde{U}_{\bar{x}}(\tau, \bar{x}_k), \tilde{U}_{\bar{x}\bar{x}}(\tau, \bar{x}_k)) \neq 0.$$

Since the CGL nodes $\{\bar{t}_p\}_{p=0}^N$ are dense in $[-1, 1]$ when $N \rightarrow \infty$, there is a sequence $\{\bar{t}_{l_{N_i}}\}_{i=1}^\infty$ such that $0 < l_{N_i} < N_i$ and $\lim_{i \rightarrow \infty} \bar{t}_{l_{N_i}} = \tau$. Thus,

$$\begin{aligned} & \lim_{i \rightarrow \infty} \left(\tilde{U}_{\bar{t}}(\bar{t}_{l_{N_i}}, \bar{x}_k) - \psi(\bar{t}_p, \tilde{U}(\bar{t}_{l_{N_i}}, \bar{x}_k), \tilde{U}_{\bar{x}}(\bar{t}_{l_{N_i}}, \bar{x}_k), \tilde{U}_{\bar{x}\bar{x}}(\bar{t}_{l_{N_i}}, \bar{x}_k)) \right) \\ &= \tilde{U}_t(\tau, \bar{x}_k) - \psi(\bar{t}_p, \tilde{U}(\tau, \bar{x}_k), \tilde{U}_{\bar{x}}(\tau, \bar{x}_k), \tilde{U}_{\bar{x}\bar{x}}(\tau, \bar{x}_k)) \neq 0. \end{aligned} \quad (40)$$

On the other hand, since $\lim_{i \rightarrow \infty} \frac{\sqrt{N_i}}{2N_i-1} W(\frac{1}{2N_i-1}) = 0$, by (25) we obtain

$$\lim_{i \rightarrow \infty} \left(\tilde{U}_{\bar{t}}(\bar{t}_{l_{N_i}}, \bar{x}_k) - \psi(\bar{t}_p, \tilde{U}(\bar{t}_{l_{N_i}}, \bar{x}_k), \tilde{U}_{\bar{x}}(\bar{t}_{l_{N_i}}, \bar{x}_k), \tilde{U}_{\bar{x}\bar{x}}(\bar{t}_{l_{N_i}}, \bar{x}_k)) \right) = 0,$$

which contradicts with (40). Thus, $\tilde{U}(\bar{t}, \bar{x})$ (for all $\bar{t} \in [-1, 1]$ and $\bar{x} = \bar{x}_k$, $k = 1, \dots, N-1$) satisfies the first equation of (9). Also, it can be easily proven that $\tilde{U}(\cdot, \bar{x}_k)$, $k = 0, 1, \dots, N$, satisfies the boundary conditions. For example, we show that $\tilde{U}(-1, \bar{x}_k) = F(\bar{x}_k)$ for $k = 0, 1, \dots, N$. We have

$$\begin{aligned} 0 &\leq |\tilde{U}(-1, \bar{x}_k) - F(\bar{x}_k)| = \left| \lim_{i \rightarrow \infty} U^{N_i}(-1, \bar{x}_k) - F(\bar{x}_k) \right| = \lim_{i \rightarrow \infty} |U^{N_i}(-1, \bar{x}_k) - F(\bar{x}_k)| \\ &= \lim_{i \rightarrow \infty} |\bar{a}_{0k}^{N_i} - F(\bar{x}_k)| \leq \lim_{i \rightarrow \infty} \frac{\sqrt{N_i}}{2N_i-1} W(\frac{1}{2N_i-1}) = 0. \end{aligned}$$

Hence, $\tilde{U}(-1, \bar{x}_k) = F(\bar{x}_k)$ for all $k = 0, 1, \dots, N$. Now, we know that the nodes $\{\bar{x}_k\}_{k=0}^N$ are dense in $[-1, 1]$ when $N \rightarrow \infty$. Therefore the pair $\tilde{U}(\cdot, \cdot)$, defined by (38), is a solution of (9) on $\bar{\Omega} = [-1, 1] \times [-1, 1]$. This completes the proof. \square

5. The Generalization of the CSC Method for Two-Dimensional Burgers Equation

In this section, we introduce the two-dimensional Burgers equation and solve it by using the CSC method. The two-dimensional Burgers equation is as follows

$$\begin{aligned} U_t(x, y, t) + U(x, y, t)U_x(x, y, t) + U(x, y, t)U_y(x, y, t) &= \varepsilon(U_{xx}(x, y, t) + U_{yy}(x, y, t)), \\ (x, y, t) &\in [0, 1] \times [0, 1] \times [0, T], \end{aligned} \quad (41)$$

and the time initial and space boundary conditions are as follows

$$U(x, y, 0) = f(x, y), \quad (42)$$

$$U(0, y, t) = h_1(y, t), \quad (43)$$

$$U(1, y, t) = h_2(y, t), \quad (44)$$

$$U(x, 0, t) = h_3(x, t), \quad (45)$$

$$U(x, 1, t) = h_4(x, t). \quad (46)$$

As we explained \bar{x}_k, \bar{y}_n and \bar{t}_p are the CGL points which were defined in the Section 3.

Now, for obtaining the numerical solution of the two-dimensional Burgers Equation (41) by applying the CSC method, the interpolation polynomial is following

$$U(\bar{x}, \bar{y}, \bar{t}) \approx U^N(\bar{x}, \bar{y}, \bar{t}) = \sum_{i=0}^N \sum_{j=0}^N \sum_{l=0}^N \bar{a}_{ijl}^N L_i(\bar{x}) L_j(\bar{y}) L_l(\bar{t}). \quad (47)$$

By (12), we have

$$U^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \bar{a}_{knp}^N \quad (48)$$

To represent the derivatives $U_{\bar{t}}^N(\cdot, \cdot, \cdot), U_{\bar{x}}^N(\cdot, \cdot, \cdot), U_{\bar{y}}^N(\cdot, \cdot, \cdot), U_{\bar{x}\bar{x}}^N(\cdot, \cdot, \cdot)$ and $U_{\bar{y}\bar{y}}^N(\cdot, \cdot, \cdot)$ in terms of $U^N(\cdot, \cdot, \cdot)$ at the node points $(\bar{x}_k, \bar{y}_n, \bar{t}_p)$, by using the matrix multiplication $D = (D_{kj})$, we have

$$\begin{cases} U_{\bar{t}}^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \sum_{l=0}^N \bar{a}_{knl}^N D_{pl}, \\ U_{\bar{x}}^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \sum_{i=0}^N \bar{a}_{inp}^N D_{ki}, \\ U_{\bar{y}}^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \sum_{j=0}^N \bar{a}_{kjp}^N D_{nj}, \\ U_{\bar{x}\bar{x}}^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \sum_{i=0}^N \bar{a}_{inp}^N \hat{D}_{kj}, \\ U_{\bar{y}\bar{y}}^N(\bar{x}_k, \bar{y}_n, \bar{t}_p) = \sum_{j=0}^N \bar{a}_{kjp}^N \hat{D}_{nj}, \quad k, n, p = 0, 1, \dots, N, \end{cases} \quad (49)$$

where D_{kj} and \hat{D}_{kj} were defined in the Section 3.

Now, by applying relations (48) and (49) Equation (41) can be rewritten as

$$\begin{cases} \sum_{l=0}^N \bar{a}_{knl}^N D_{pl} + \bar{a}_{knp}^N \left(\sum_{i=0}^N \bar{a}_{inp}^N D_{ki} + \sum_{j=0}^N \bar{a}_{kjp}^N D_{nj} \right) - \varepsilon \left(\sum_{i=0}^N \bar{a}_{inp}^N \hat{D}_{kj} + \sum_{j=0}^N \bar{a}_{kjp}^N \hat{D}_{nj} \right) = 0 \\ \bar{a}_{kn0} = f(\bar{x}_k, \bar{y}_n), \quad k, n = 0, 1, \dots, N, \\ \bar{a}_{0np} = h_1(\bar{y}_n, \bar{t}_p), \quad n = 0, 1, \dots, N, \quad p = 1, \dots, N, \\ \bar{a}_{Nnp} = h_2(\bar{y}_n, \bar{t}_p), \quad n = 0, 1, \dots, N, \quad p = 1, \dots, N, \\ \bar{a}_{k0p} = h_3(\bar{x}_k, \bar{t}_p), \quad k = 1, \dots, N-1, \quad p = 1, \dots, N, \\ \bar{a}_{kNp} = h_4(\bar{x}_k, \bar{t}_p), \quad k = 1, \dots, N-1, \quad p = 1, \dots, N, \end{cases} \quad (50)$$

Now, we can obtain the numerical solution of the above system with respect to $(\bar{a}_{knp}^N; k, n, p = 0, 1, \dots, N)$.

6. Numerical Examples

In the following examples, we use the Levenberg-Marquardt method (a quasi-Newton method) for FSOLVE command in MATLAB software to solve the algebraic system (17). We calculate L_2 and L_∞ errors as follows

$$\begin{aligned} L_2 &= \| U(\bar{t}, \cdot) - U^N(\bar{t}, \cdot) \|_2 = \left(\sum_{j=0}^N | U(\bar{t}, x_j) - U^N(\bar{t}, x_j) |^2 \right)^{\frac{1}{2}}, \\ L_\infty &= \| U(\bar{t}, \cdot) - U^N(\bar{t}, \cdot) \|_\infty = \max_{0 \leq j \leq N} | U(\bar{t}, x_j) - U^N(\bar{t}, x_j) |, \end{aligned} \quad (51)$$

where x_j , $j = 0, 1, \dots, N$, are the collocation points, $\bar{t} \in [-1, 1]$ is a given point and $U(\cdot, \cdot)$ and $U^N(\cdot, \cdot)$ are the analytical and approximate solutions, respectively. Also, the absolute error can be obtained by

$$E(\bar{t}, \bar{x}) = |U(\bar{t}, \bar{x}) - U^N(\bar{t}, \bar{x})|, \quad (\bar{t}, \bar{x}) \in [-1, 1] \times [-1, 1].$$

Example 1. Consider the Burgers–Fisher Equation (2), where $T_0 = -0.2$, $T_1 = 0$, $a = -1$, $b = 0$, $\alpha(t) = 24$ and $\beta(t) = -48$ for $t \in [T_0, T_1]$. Also, we assume that the boundary conditions are given by

$$\begin{aligned} U(t, -1) &= \frac{1}{2} - \frac{1}{2} \tanh[6(-1 - 8t)], \\ U(t, 0) &= \frac{1}{2} - \frac{1}{2} \tanh[6(-8t)], \end{aligned}$$

and the initial condition is as follows

$$U(-0.2, x) = \frac{1}{2} - \frac{1}{2} \tanh[6(x + 1.6)].$$

Then the exact solution is

$$U(t, x) = \frac{1}{2} - \frac{1}{2} \tanh[6(x - 8t)].$$

We gain the numerical results by the CSC method at $t = -0.1, -0.05, -0.04, -0.035, -0.03$ for $N = 30 \times 30$ which are shown in Table 1. We observe that our numerical results are better than the results of MMPDE methods [8], which they are obtained for $\Delta t = 10^{-6}$ and $\Delta x = \frac{1}{60}$ (or $N = 200,000 \times 60$). In Table 1, L_2 errors are presented. In Figures 1 and 2, we show the approximate solution and absolute error, respectively. In Figure 3, we represent the exact and approximate solutions for $t = -0.1, -0.05$ and -0.03 . We also illustrate L_2 errors in Figure 4.

Table 1. Comparison of the L_2 error for Example 1.

Method	N	$t = -0.1$	$t = -0.05$	$t = -0.04$	$t = -0.035$	$t = -0.03$
Presented method	30×30	1.8293×10^{-4}	1.1920×10^{-4}	1.2691×10^{-4}	1.4053×10^{-4}	1.4187×10^{-4}
Mesh for optimal M [8]	$200,000 \times 60$	2.1×10^{-3}	2.9×10^{-3}	3.4×10^{-3}	3.7×10^{-3}	4.2×10^{-3}
Mesh for arc-length M [8]	$200,000 \times 60$	2.7×10^{-3}	8×10^{-3}	7.9×10^{-3}	7.6×10^{-3}	7.1×10^{-3}
Mesh for curvature M [8]	$200,000 \times 60$	2.1×10^{-3}	2.4×10^{-3}	2.4×10^{-3}	2.4×10^{-3}	2.5×10^{-3}




Figure 1. The approximate solution for Example 1.




Figure 2. The absolute error $E(.,.)$ for example 1




Figure 3. Exact and approximate solutions for $t = -0.1, -0.05$ and -0.03 .




Figure 4. L_2 errors for Example 1.

Example 2. Consider the Burgers–Fisher Equation (2), where $T_0 = -0.05$, $T_1 = 0.05$, $a = -1$, $b = 0$, $\alpha(t) = 20$ and $\beta(t) = -1 + 3\sin t$ for $t \in [T_0, T_1]$. Also, consider the boundary conditions

$$U(t, -1) = \frac{\cosh[-1 - 3\cos t] + \sinh[-1 - 3\cos t] - \sqrt{5}}{-4\cosh[-1 - 3\cos t] + 6\sinh[-1 - 3\cos t]},$$

$$U(t, 0) = \frac{\cosh[-3\cos t] + \sinh[-3\cos t] - \sqrt{5}}{-4\cosh[-3\cos t] + 6\sinh[-3\cos t]},$$

and the initial condition

$$U(-0.05, x) = \frac{\cosh[x - 3\cos(-0.05)] + \sinh[x - 3\cos(-0.05)] - \sqrt{5}}{-4\cosh[x - 3\cos(-0.05)] + 6\sinh[x - 3\cos(-0.05)]}.$$

By these, the exact solution is as follows

$$U(t, x) = \frac{\cosh[x - 3\cos t] + \sinh[x - 3\cos t] - \sqrt{5}}{-4\cosh[x - 3\cos t] + 6\sinh[x - 3\cos t]}.$$

We solve the system (17) according to this example. Table 2 shows the L_2 errors at $t = -0.05, -0.025, 0, 0.025$ and 0.05 for CSC method and MMPDE methods [8]. Our numerical results are satisfied $N = 10 \times 10$ (or equivalently $\Delta x = 0.1$ and $\Delta t = 0.01$) and results of the MMPDE methods are with $\Delta x = \frac{1}{40}$ and $\Delta t = 10^{-4}$ (or equivalently $N = 1000 \times 40$). In Figures 5, 6 and 7, we illustrate the approximate solution, absolute error and the L_2 error, respectively.

Table 2. Comparison of the L_2 error for Example 2.

Method	N	$t = -0.05$	$t = -0.025$	$t = 0$	$t = 0.025$	$t = 0.05$
Presented method	10×10	4.8044×10^{-6}	2.4602×10^{-4}	4.1079×10^{-4}	5.3422×10^{-4}	6.2951×10^{-4}
Mesh for optimal M [8]	1000×40	2.2×10^{-3}	4×10^{-3}	5.3×10^{-3}	3.5×10^{-3}	1.2×10^{-3}
Mesh for arc-length M [8]	1000×40	1.8×10^{-3}	4.1×10^{-3}	6.6×10^{-3}	6.8×10^{-3}	2.3×10^{-3}
Mesh for curvature M [8]	1000×40	2.2×10^{-3}	4×10^{-3}	5.2×10^{-3}	3.6×10^{-3}	9.6×10^{-4}




Figure 5. The approximate solution for Example 2.





Figure 6. The absolute error $E(.,.)$ for Example 2.

Figure 7. L_2 errors for Example 2.

Example 3. Consider the Burgers–Huxley Equation (3) with $\alpha = 1$, $\delta = 2$ and $\gamma = 0$. Therefore, it can be written as follows

$$U_t + U^2 U_x - \beta U_{xx} = 0, \quad (52)$$

where $T_0 = 1$, $T_1 = 10$, $a = 0$ and $b = 1$. Also, consider the boundary conditions

$$\begin{aligned} U(t, 0) &= 0, \\ U(t, 1) &= \frac{1}{t + \frac{t\sqrt{t}}{c_0} \exp(\frac{1}{4\beta t})}, \end{aligned} \quad (53)$$

and the initial condition

$$U(1, x) = \frac{x}{1 + \frac{1}{c_0} \exp(\frac{x^2}{4\beta})}, \quad (54)$$

where $0 < c_0 < 1$. Hence, the exact solution is given by

$$U(t, x) = \frac{\frac{x}{t}}{1 + \frac{\sqrt{t}}{c_0} \exp(\frac{x^2}{4\beta t})}.$$

We take $c_0 = 0.5$ and $\beta = 0.01$. Table 3 shows the L_∞ errors by using CSC method and given methods in [9,12,15,16] for $t = 2, 4$ and 6 and $N = 9 \times 9$. In Figures 8 and 9, we show the approximate solution and absolute error, respectively. Also, in Figure 10, we represent the exact and approximate solutions for $t = 2, 6$ and 10. Moreover, Figure 11 shows the L_∞ error.

Table 3. Comparison of the L_∞ error for Example 3.

Method	N	$T = 2$	$T = 6$	$T = 10$
Presented method	9×9	5.5673×10^{-4}	4.4466×10^{-4}	3.0034×10^{-4}
Haar wavelet method [12]	16×900	7.5978×10^{-4}	4.6335×10^{-4}	1.16480×10^{-3}
QBC method [10]	200×900	1.21698×10^{-3}	7.2249×10^{-4}	1.28124×10^{-3}
SBC method [9]	50×900	1.70309×10^{-3}	7.6105×10^{-4}	1.80239×10^{-3}
QBCA1 method [15]	200×900	8.1680×10^{-4}	5.2579×10^{-4}	1.28125×10^{-3}
QBCA2 method [15]	200×900	8.2212×10^{-4}	5.2579×10^{-4}	1.28125×10^{-3}
SBC1 method [16]	200×900	8.2934×10^{-4}	...	1.28127×10^{-3}
SBC2 method [16]	200×900	8.2734×10^{-4}	...	1.28127×10^{-3}




Figure 8. The approximate solution for Example 3.




Figure 9. The absolute error $E(.,.)$ for Example 3.





Figure 10. Exact and approximate solutions for $t = 2, 6$ and 10 .

Figure 11. L_∞ errors for Example 3.

Example 4. Consider the Burgers Equation (52) for $x \in [0, 1.3]$ and

$$\begin{cases} U(t, 0) = 0, \\ U(t, 1) = \frac{1.3}{t + \frac{t\sqrt{t}}{c_0} \exp(\frac{1.69}{4\beta t})}, \\ U(1, x) = \frac{x}{1 + \frac{1}{c_0} \exp(\frac{x^2}{4\beta})}. \end{cases}$$

The numerical results of our method and other methods [12,16] are displayed in Table 4 for different values of t . The L_∞ error results for this example is depicted in Table 4 along with the comparison of the error computed by the present method and other methods. In Figures 12 and 13, the approximate solution and absolute error are shown, respectively. Also, in Figure 14, we represent the exact and approximate solutions for $t = 2, 6$ and 10 .

Table 4. Comparison of the L_∞ error for Example 4.

Method	N	$T = 2$	$T = 6$	$T = 10$
Presented method	10×10	5.306×10^{-4}	4.294×10^{-4}	3.166×10^{-4}
Haar wavelet method [12]	16×900	7.2890×10^{-4}	4.5606×10^{-4}	3.2374×10^{-4}
SBC1 method [16]	260×900	8.2934×10^{-4}	...	3.2723×10^{-4}
SBC2 method [16]	260×900	8.2734×10^{-4}	...	3.2337×10^{-4}




Figure 12. The approximate solution for Example 4.




Figure 13. The absolute error $E(.,.)$ for Example 4.




Figure 14. Exact and approximate solutions for $t = 2, 6$ and 10 .

Example 5. Consider the Burgers Equation (1), where $T_0 = 1$, $T_1 = 5$, $a = 0$, $b = 8$ and $\varepsilon = 0.5$. The initial condition for the current problem is

$$U(1, x) = \frac{x}{1 + \exp(\frac{1}{4\varepsilon}(x^2 - \frac{1}{4}))}, \quad (55)$$

and the boundary conditions are

$$\begin{aligned} U(t, 0) &= 0, \\ U(t, 8) &= \frac{8}{t + t(\frac{t}{\exp(\frac{1}{8\varepsilon})})^{\frac{1}{2}} \exp(\frac{16}{\varepsilon t})}. \end{aligned} \quad (56)$$



Here, we have the following analytical solution

$$U(t, x) = \frac{\frac{x}{t}}{1 + (\frac{t}{\exp(\frac{1}{8\varepsilon})})^{\frac{1}{2}} \exp(\frac{x^2}{4\varepsilon t})}.$$

In Figures 15 and 16, we show the approximate solution and absolute error, respectively. Also, Figure 17 illustrates the comparison between the exact solution and numerical solution given by the proposed method. Also, we compare the L_∞ and L_2 errors which are computed by the present method and other methods [14] in Table 5. We can observe that the results of CSC for $N = 30 \times 30$ are better than the results of method of Inan and Bahadır [14] for $N = 320 \times 40,000$.

Table 5. Comparison of the L_∞ and L_2 errors for Example 5

Method	T	N	L_2	L_∞
Presented method	$T = 1.5$	30×30	3.2025×10^{-8}	1.2611×10^{-7}
$I - EFD$ method [14]	$T = 1.5$	$320 \times 40,000$	2.1×10^{-5}	1.8×10^{-5}
$FI - EFD$ method [14]	$T = 1.5$	$320 \times 40,000$	2.2×10^{-5}	1.9×10^{-5}
Presented method	$T = 3$	30×30	3.2026×10^{-8}	6.9546×10^{-9}
$I - EFD$ method [14]	$T = 3$	$320 \times 40,000$	2.2×10^{-5}	3.8×10^{-5}
$FI - EFD$ method [14]	$T = 3$	$320 \times 40,000$	2.3×10^{-5}	1.8×10^{-5}
Presented method	$T = 4.5$	30×30	3.2028×10^{-8}	1.6022×10^{-9}
$I - EFD$ method [14]	$T = 4.5$	$320 \times 40,000$	4.08×10^{-4}	7.43×10^{-4}
$FI - EFD$ method [14]	$T = 4.5$	$320 \times 40,000$	4.08×10^{-4}	7.43×10^{-4}

**Figure 15.** The approximate solution for Example 5.**Figure 16.** The absolute error $E(.,.)$ for Example 5.




Figure 17. Exact and approximate solutions for Example 5.

Example 6. By considering the two-dimensional Burgers Equation (41) with $T = 1$ and $\varepsilon = 0.1, 0.2, 0.5$ and 1 , the initial condition is as follow [30,31]

$$U(x, y, 0) = \frac{1}{1 + \exp\left(\frac{(x+y)}{2\varepsilon}\right)}, \quad (57)$$

and the boundary conditions are

$$U(0, y, t) = \frac{1}{1 + \exp\left(\frac{(y-t)}{2\varepsilon}\right)},$$

$$U(1, y, t) = \frac{1}{1 + \exp\left(\frac{(1+y-t)}{2\varepsilon}\right)}, \quad (58)$$

$$U(x, 0, t) = \frac{1}{1 + \exp\left(\frac{(x-t)}{2\varepsilon}\right)}, \quad (59)$$

$$U(x, 1, t) = \frac{1}{1 + \exp\left(\frac{(x+1-t)}{2\varepsilon}\right)}. \quad (60)$$

By these considerations, the exact solution is

$$U(x, y, t) = \frac{1}{1 + \exp\left(\frac{(x+y-t)}{2\varepsilon}\right)}.$$

We calculate approximate solutions and absolute errors in different $\varepsilon = 0.1, 0.2, 0.5$ and 1 with $N = 10$. In Figures 18 and 19, we observe the approximate solutions and absolute errors with $\varepsilon = 0.1$, respectively. Also, Figures 20 and 21 illustrate the numerical solution and absolute error given by the presented method for $\varepsilon = 0.1$. The approximate solutions and absolute errors with $\varepsilon = 0.5$ and 1 are expressed in Figures 22–25, respectively. Moreover, we computed the L_∞ errors for various ε at $t = t_0$ which are shown in Table 6.

Table 6. The L_∞ error for various ε Example 6.

The CSC Method	$\varepsilon = 0.1$	$\varepsilon = 0.2$	$\varepsilon = 0.5$	$\varepsilon = 1$
L_∞	1.076×10^{-4}	3.421×10^{-5}	6.073×10^{-5}	7.333×10^{-6}




Figure 18. The approximate solution for $\varepsilon = 0.1$ Example 6.




Figure 19. The absolute error for $\varepsilon = 0.1$ Example 6.




Figure 20. The approximate solution for $\varepsilon = 0.2$ Example 6.




Figure 21. The absolute error for $\varepsilon = 0.2$ Example 6.




Figure 22. The approximate solution for $\varepsilon = 0.5$ Example 6.




Figure 23. The absolute error for $\varepsilon = 0.5$ Example 6.




Figure 24. The approximate solution for $\varepsilon = 1$ Example 6.




Figure 25. The absolute error for $\varepsilon = 1$ Example 6.

7. Conclusions

In this article, we used the Chebyshev spectral collocation method to get numerical solutions for different types of one and two-dimensional Burgers equation. We analyzed the convergence of the CSC method by using the concept of module of continuity and compared the obtained approximate solutions with those of other methods. We showed that the CSC method has very high accuracy and it is more precise with respect to the other numerical methods. Our investigations can be used in the three-dimensional case and we prepare these investigations as a future article.

Author Contributions: The authors contributed equally to this work.

Funding: This research was funded by the National Natural Science Foundation of China Projects grant number [11601240, 11771214].

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

CSC Chebyshev Spectral Collocation
CGL Chebyshev–Gauss–Lobatto

References

1. Eltayeb, H.; Bachar, I.; Kilicman, A. On conformable double laplace transform and one dimensional fractional coupled Burgers equation. *Symmetry* **2019**, *11*, 417. [\[CrossRef\]](#)
2. Zhang, X.; Zhang, Y. Some similarity solutions and numerical solutions to the Time-Fractional Burgers system. *Symmetry* **2019**, *11*, 37–50. [\[CrossRef\]](#)
3. Malik, S.A.; Qureshi, I.M.; Amir, M.; Malik, A.N.; Haq, I. Numerical solution to generalized Burgers-Fisher equation using Exp-Function method hybridized with heuristic computation. *PLoS ONE* **2015**, *10*, e0121728. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Ray, S.S.; Gupta, A.K. On the Solution of Burgers-Huxley and Huxley equation using wavelet collocation method. *Cmes-Comp. Model. Eng.* **2013**, *91*, 409–424. [\[CrossRef\]](#)
5. Dhawan, S.; Kapoor, S.; Kumar, S.; Rawat, S. Contemporary review of techniques for the solution of nonlinear Burgers equation. *J. Comput. Sci.* **2012**, *3*, 405–419. [\[CrossRef\]](#)
6. Berger, M.; Kohn, R.V. A rescaling algorithm for the numerical calculation of blowing-up solutions. *Commun. Pur. Appl. Anal.* **1988**, *41*, 841–863. [\[CrossRef\]](#)
7. Budd, C.J.; Huang, W.; Russell, R.D. Moving mesh methods for problems with blow-up. *SIAM J. Sci. Comput.* **1996**, *17*, 305–327. [\[CrossRef\]](#)
8. Soheili, A.R.; Kerayechian, A.; Davoodi, N. Adaptive numerical method for Burgers'-type nonlinear equations. *Appl. Math. Comput.* **2012**, *219*, 3486–3495. [\[CrossRef\]](#)
9. Ramadan, M.A.; El-Danaf, T.S.; Abd Alaal, F.E.I. A numerical solution of the Burgers' equation using septic B-splines. *Chaos. Soliton. Fract.* **2005**, *26*, 795–804. [\[CrossRef\]](#)
10. Ramadan, M.A.; El-Danaf, T.S. Numerical treatment for the modified Burgers equation. *Math. Comput. Simul.* **2005**, *70*, 90–98. [\[CrossRef\]](#)
11. Haq, S.; Ul-Islam, S.; Uddin, M. A mesh-free method for the numerical solution of the KdV-Burgers' equation. *Appl. Math. Model.* **2009**, *33*, 3442–3449. [\[CrossRef\]](#)
12. Oruc, O.; Bulut, F.; Esen, A.; A Haar wavelet-finite difference hybrid method for the numerical solution of the modified Burgers' equation. *J. Math. Chem.* **2015**, *53*, 1592–1607. [\[CrossRef\]](#)
13. Lepik, U. Numerical solution of evolution equations by the Haar wavelet method. *Appl. Math. Comput.* **2007**, *185*, 695–704. [\[CrossRef\]](#)
14. Inan, B.; Bahadir, A.R. Numerical solution of the one-dimensional Burgers' equation: Implicit and fully implicit exponential finite difference methods. *Pramana J. Phys.* **2013**, *81*, 547–556. [\[CrossRef\]](#)
15. Saka, B.; Dag, I. A numerical study of the Burgers' equation. *J. Franklin. I.* **2008**, *345*, 328–348. [\[CrossRef\]](#)
16. Irk, D. Sextic B-spline collocation method for the modified Burgers' equation. *Kybernetes* **2009**, *38*, 1599–1620. [\[CrossRef\]](#)
17. Demiray, H. A note on the travelling wave solution to the perturbed Burgers equation. *Appl. Math. Model.* **2002**, *26*, 37–40. [\[CrossRef\]](#)
18. Hon, Y.C.; Mao, X.Z. An efficient numerical scheme for Burgers' equation. *Appl. Math. Comput.* **1998**, *95*, 37–50. [\[CrossRef\]](#)
19. Schulze-Halberg, A. Burgers' equation with time-dependent coefficients and nonlinear forcing term: Linearization and exact solvability. *Commun. Nonlinear. Sci.* **2015**, *22*, 1068–1083. [\[CrossRef\]](#)
20. Seydaoglu, M.; Erdogan, U.; Ozis, T. Numerical solution of Burgers' equation with high order splitting methods. *J. Comput. Appl. Math.* **2016**, *291*, 410–421. [\[CrossRef\]](#)
21. Guo, Y.; Shi, Y.; Li, Y. A fifth order finite volume weighted compact scheme for solving one-dimensional Burgers equation. *Appl. Math. Comput.* **2016**, *281*, 172–185. [\[CrossRef\]](#)
22. Mukundan, V.; Awasthi, A. Efficient numerical techniques for Burgers' equation. *Appl. Math. Comput.* **2015**, *262*, 282–297. [\[CrossRef\]](#)
23. Hammad, D.A.; El-Azab, M.S. 2N order compact finite difference scheme with collocation method for solving the generalized Burgers'-Huxley and Burgers'-Fisher equations. *Appl. Math. Comput.* **2015**, *258*, 296–311. [\[CrossRef\]](#)
24. Arora, G.; Kumar Singh, B. Numerical solution of Burgers equation with modified cubic B-spline differential quadrature method. *Appl. Math. Comput.* **2013**, *224*, 166–177. [\[CrossRef\]](#)

25. El-Wakil, S.A.; Abulwafa, E.M.; El-hanbaly, A.M.; El-Shewy, E.K.; Abd-El-Hamid, H.M. Self-similar solutions for some nonlinear evolution equations: KdV, mKdV and Burgers equations. *J. Assoc. Arab Univ. Basic Appl.* **2016**, *19*, 44–51. [\[CrossRef\]](#)
26. Singh, B.K.; Arora, G.; Singh, M.K. A numerical scheme for the generalized Burgers'-Huxley equation. *J. Egypt. Math. Soc.* **2016**, *24*, 629–637. [\[CrossRef\]](#)
27. Weinan, E. Convergence of spectral methods for Burgers' equation, *SIAM J. Numer. Anal.* **1992**, *29*, 1520–1541. [\[CrossRef\]](#)
28. Xiao, D.; Fang, F.; Du, J.; Pain, C.C.; Navon, I.M.; Buchon, A.G.; ElSheikh, A.H.; Hu, G.. Non-Linear Petrov-Galerkin methods for reduced order modelling of the Navier-Stokes equations using a mixed finite element pair. *Comp. Methods Appl. Mech. Engrg.* **2013**, *255*, 147–157. [\[CrossRef\]](#)
29. Prodanov, D. Analytical and numerical treatments of conservative diffusions and the Burgers equation. *Entropy* **2018**, *20*, 492. [\[CrossRef\]](#)
30. Duan Y.; Liu, R. Lattice Boltzmann model for two-dimensional unsteady Burgers' equation. *J. Comput. Appl. Math.* **2007**, *206*, 432–439. [\[CrossRef\]](#)
31. Radwan, S.F.; Comparison of higher-order accurate schemes for solving the two-dimensional unsteady Burgers' equation. *J. Comput. Appl. Math.* **2005**, *174*, 383–397. [\[CrossRef\]](#)
32. Elnegar, G.N.; Kazemi, M.A. Pseudo-spectral Chebyshev optimal control of constrained nonlinear dynamical systems. *Comput. Optim. Appl.* **1998**, *11*, 195–217.:1018694111831. [\[CrossRef\]](#)
33. Gong, Q.; Ross, I.M.; Fahroo, F. A Chebyshev pseudo-spectral method for nonlinear constrained optimal control problems. In Proceedings of the 48th IEEE Conference on Decision and Control (CDC) held jointly with 2009 28th Chinese Control Conference, Shanghai, China, 29 January 2010. [\[CrossRef\]](#)
34. Fahroo, F.; Ross, I.M. Direct trajectory optimization by a Chebyshev pseudo-spectral method. *J. Guid. Control. Dynam.* **2002**, *25*, 160–166. [\[CrossRef\]](#)
35. Noori Skandari, M.H.; Ghaznavi, M. Chebyshev Pseudo-Spectral method for Bratu's problem. *Iran. J. Sci. Technol. Trans. Sci.* **2017**, *41*, 913–921. [\[CrossRef\]](#)
36. Ghaznavi, M.; Noori Skandari, M.H. An efficient Pseudo-Spectral method for nonsmooth dynamical systems, *Iran. J. Sci. Technol. Trans. Sci.* **2016**, *42*, 635–646. [\[CrossRef\]](#)
37. Noori Skandari, M.H.; Kamyad, A.V.; Effati, S. Generalized Euler-Lagrange equation for nonsmooth calculus of variations. *Nonlinear. Dyn.* **2014**, *75*, 85–100. [\[CrossRef\]](#)
38. Ragozin, D.L. Polynomial approximation on compact manifolds and homogeneous spaces. *Trans. Amer. Math. Soc.* **1971**, *162*, 157–170. [\[CrossRef\]](#)



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).