




Article

Mixed Type Nondifferentiable Higher-Order Symmetric Duality over Cones

Izhar Ahmad ^{1,†} , Khushboo Verma ^{2,†}  and Suliman Al-Homidan ^{1,*,†} 

¹ Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia; drizhar@kfupm.edu.sa

² Department of Applied Sciences and Humanities, Faculty of Engineering, University of Lucknow, New Campus, Lucknow 226 021, India; 1986khushi@gmail.com

* Correspondence: homidan@kfupm.edu.sa

† These authors contributed equally to this work.

Received: 29 December 2019; Accepted: 1 February 2020; Published: 11 February 2020



Abstract: A new mixed type nondifferentiable higher-order symmetric dual programs over cones is formulated. As of now, in the literature, either Wolfe-type or Mond–Weir-type nondifferentiable symmetric duals have been studied. However, we present a unified dual model and discuss weak, strong, and converse duality theorems for such programs under higher-order \mathcal{F} -convexity/higher-order \mathcal{F} -pseudoconvexity. Self-duality is also discussed. Our dual programs and results generalize some dual formulations and results appeared in the literature. Two non-trivial examples are given to show the uniqueness of higher-order \mathcal{F} -convex/higher-order \mathcal{F} -pseudoconvex functions and existence of higher-order symmetric dual programs.

Keywords: higher-order nondifferentiable programming; symmetric duality; self duality higher-order \mathcal{F} -convexity/higher-order \mathcal{F} -pseudoconvexity; duality theorems

1. Introduction

The study of higher-order duality has computational advantages over the first-order duality when approximations are used, as it provides tighter bound for the value of the objective function. Mangasarian [1] formulated a class of higher-order duality in nonlinear problems. Later on, Mond and Zhang [2] obtained various higher-order duality results under higher-order invexity assumptions. Chen [3] discussed Mond–Weir-type higher-order symmetric duality involving F -convex functions, whereas Mishra [4] obtained Mond–Weir-type higher-order symmetric duality theorems under generalized invexity. Khurana [5] presented symmetric duality results for a Mond–Weir-type dual programs over arbitrary cones under cone-pseudoinvexity and strongly cone-pseudoinvexity. Later on, a higher-order Mond–Weir-type nondifferentiable multiobjective dual problem is formulated and established duality relations involving higher-order (F, α, ρ, d) type-I functions by Ahmad et al. [6]. In [7], Gupta and Jayswal obtained multiobjective higher-order symmetric duality results under higher-order K -preinvexity/ K -pseudoinvexity.

The theoretical and algorithmic concepts of mixed duality in nonlinear programming problems are interesting and useful. Under K -preinvexity and K -pseudoinvexity assumptions, Ahmad and Husain [8] formulated multiobjective mixed type symmetric dual programs over cones and proved duality results. Chandra et al. [9] studied mixed type symmetric duality results. Xu [10] proved duality theorems for two mixed type duals of a multiobjective programming problem. Yang et al. [11] presented mixed type symmetric duality for nondifferentiable nonlinear programming problems. By ignoring nonnegativity constraints of symmetric dual given in [12], Ahmad [13] presented multiobjective mixed type symmetric duality results. Recently, Verma et al. [14] formulated a

higher-order nondifferentiable mixed symmetric dual model and duality results are studied under higher-order invexity/generalized invexity.

In the present paper, a new mixed type nondifferentiable higher-order symmetric dual programs over cones are formulated. As of now, in the literature, either Wolfe-type or Mond–Weir-type nondifferentiable symmetric dual programs have been discussed. However, our model unifies both dual programs. Under higher-order \mathcal{F} -convexity/higher-order \mathcal{F} -pseudoconvexity, appropriate duality theorems are proved. Self-duality is also discussed. Our study extends and generalizes the existing results appeared in [15–21]. Two non-trivial examples are given to show the uniqueness of higher-order \mathcal{F} -convex/higher-order \mathcal{F} -pseudoconvex functions and existence of higher-order symmetric dual programs.

2. Preliminaries

We consider the following nonlinear programming problem.

$$(P) \quad \begin{aligned} &\text{Minimize } F(x) \\ &x \in X, \end{aligned}$$

where $X \subseteq \mathbb{R}^n$ and $F : X \rightarrow \mathbb{R}$.

2.1. Definitions

(a) The support function $s(x|E)$ of E is given by

$$s(x|E) = \max\{x^T y : y \in E\}.$$

The subdifferential of $s(x|E)$ is defined by

$$\partial s(x|E) = \{z \in E : z^T x = s(x|E)\}.$$

The normal cone for any convex set $S \subset \mathbb{R}^n$ at a point $x \in S$ is defined by

$$N_S(x) = \{y \in \mathbb{R}^n : y^T(z - x) \leq 0 \text{ for all } z \in S\}.$$

It is readily verified that for a compact convex set D , y is in $N_D(x)$ if and only if

$$s(y|D) = x^T y.$$

(b) Let C be a closed convex cone in \mathbb{R}^n with nonempty interior. The positive polar cone C^* of C is given by

$$C^* = \{q \in \mathbb{R}^n : x^T q \geq 0 \text{ for all } x \in C\}.$$

(c) A function $\mathcal{F} : X \times X \times \mathbb{R}^n \mapsto \mathbb{R}$ (where $X \subseteq \mathbb{R}^n$) is sublinear in its third component, if for all $x, y \in X$:

- (i) $\mathcal{F}(x, y; b_1 + b_2) \leq \mathcal{F}(x, y; b_1) + \mathcal{F}(x, y; b_2)$, for all $b_1, b_2 \in \mathbb{R}^n$; and
- (ii) $\mathcal{F}(x, y; \alpha b) = \alpha \mathcal{F}(x, y; b)$, for all $\alpha \in \mathbb{R}$, for all $b \in \mathbb{R}^n$.

(d) [3] A function $\phi : \mathbb{R}^n \mapsto \mathbb{R}$ is called higher-order \mathcal{F} -convex at $u \in \mathbb{R}^n$ with respect to $g : \mathbb{R}^n \times \mathbb{R}^n \mapsto \mathbb{R}$, if for all $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$,

$$\phi(x) - \phi(u) - g(u, p) + p^T \nabla_p g(u, p) \geq \mathcal{F}(x, u; \nabla_x \phi(u) + \nabla_p g(u, p)).$$

(e) [3] A function $\phi : \mathbb{R}^n \mapsto \mathbb{R}$ is called higher-order \mathcal{F} -pseudoconvex at $u \in \mathbb{R}^n$ with respect to $g : \mathbb{R}^n \times \mathbb{R}^n \mapsto \mathbb{R}$, if for all $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$,

$$\mathcal{F}(x, u; \nabla_x \phi(u) + \nabla_p g(u, p)) \geq 0 \Rightarrow \phi(x) - \phi(u) - g(u, p) + p^T \nabla_p g(u, p) \geq 0.$$

The above definitions (d) and (e) are validated by the following example.

2.2. Example

Consider $\phi(x) = e^{-x} + x^2$, $F_{x,u}(a) = |a|(x^2 - u^2)$, $g(u, p) = \frac{p}{u+1}$ and $X = \{x : x \geq 1\}$.

Now, we show that the function $\phi(x)$ is a higher-order \mathcal{F} -pseudoconvex at $u = 1$ and for all $x \geq 1$.

$$\begin{aligned} F(x, u; \nabla_x \phi(u) + \nabla_p g(u, p)) &= |\nabla_x \phi(u) + \nabla_p g(u, p)|(x^2 - u^2) \\ &= |-e^{-u} + 2u + \frac{1}{u+1}|(x^2 - u^2), \end{aligned}$$

For $u = 1$, the above expression reduces to

$$F(x, 1; \nabla_x \phi(1) + \nabla_p g(1, p)) = |-e^{-1} + 2 + \frac{1}{2}|(x^2 - 1) \geq 0, \text{ for } x \geq 1, \quad (1)$$

Now,

$$\phi(x) - \phi(u) - g(u, p) + p^T \nabla_p g(u, p) = e^{-x} + x^2 - e^{-u} - u^2 - \frac{p}{u+1} + \frac{p}{u+1}.$$

At $u = 1$, the above equality becomes

$$\phi(x) - \phi(u) - g(u, p) + p^T \nabla_p g(u, p) = e^{-x} + x^2 - \frac{1}{e} - 1 \geq 0, \text{ for } x \geq 1, \quad (2)$$

The Equations (1) and (2) show that function $\phi(x)$ is a higher-order \mathcal{F} -pseudoconvex at $u = 1$ and for all $x \geq 1$ (see Figures 1–3).

Now, we determine whether $\phi(x)$ is a higher-order \mathcal{F} -convex function.

$$\begin{aligned} \phi(x) - \phi(u) - g(u, p) + p^T \nabla_p g(u, p) - F(x, u; \nabla_x \phi(u) + \nabla_p g(u, p)) \\ = e^{-x} + x^2 - e^{-u} - u^2 - \frac{p}{u+1} + \frac{p}{u+1} - |-e^{-u} + 2u + \frac{1}{u+1}|(x^2 - u^2). \end{aligned}$$

At $u = 1$

$$\begin{aligned} &= e^{-x} + x^2 - \frac{1}{e} - 1 - |-e^{-1} + 2 + \frac{1}{2}|(x^2 - 1) \equiv \chi, \text{ (see Figure 4)} \\ &\not\geq 0, \text{ (for } x \geq 2), \end{aligned}$$

which shows that the function $\phi(x)$ is not a higher-order \mathcal{F} -convex (see Figure 4) at $u = 1$ and $x = 2$. This concludes that every higher-order \mathcal{F} -convex function is a higher-order \mathcal{F} -pseudoconvex function, but the converse is not necessarily true. (See Appendix A for codes of Figures 1–4).




Figure 1. Graph of $F_{x,u}(\cdot)$ against x and u .




Figure 2. Graph of $F_{x,u}(\cdot)$ against x .




Figure 3. Graph of ϕ against x .




Figure 4. Graph of χ against x .

3. Higher-Order Mixed Type Symmetric Duality over Cones

For $N = \{1, 2, 3, \dots, n\}$ and $H = \{1, 2, 3, \dots, h\}$, let us assume $A_1 \subset N, B_1 \subset H$ and $A_2 = N \setminus A_1$ and $B_2 = H \setminus B_1$, where $|A_1|$ denotes the number of elements in the set A_1 , also $|A_2|, |B_1|$ and $|B_2|$ are defined similarly.

Again if $|A_1| = 0$, then $|A_2| = n$ and therefore $\mathbb{R}^{|A_1|}$ is a zero-dimensional Euclidean space and $\mathbb{R}^{|A_2|}$ is n -dimensional Euclidean space.

Now $x \in \mathbb{R}^n$ if and only if $x = \{x^1, x^2\}$, i.e. $x^1 \in \mathbb{R}^{|A_1|}, x^2 \in \mathbb{R}^{|A_2|}$. Similarly, any $y \in \mathbb{R}^h$ if and only if $y = \{y^1, y^2\}$, i.e. $y^1 \in \mathbb{R}^{|B_1|}, y^2 \in \mathbb{R}^{|B_2|}$. Let

- (i) $f^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \rightarrow \mathbb{R}$,
- (ii) $f^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \rightarrow \mathbb{R}$,
- (iii) $h^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \times \mathbb{R}^{|A_1|} \rightarrow \mathbb{R}$,
- (iv) $h^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \times \mathbb{R}^{|A_2|} \rightarrow \mathbb{R}$,
- (v) $g^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \times \mathbb{R}^{|B_1|} \rightarrow \mathbb{R}$ and
- (vi) $g^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \times \mathbb{R}^{|B_2|} \rightarrow \mathbb{R}$
be twice differentiable functions.

Now we formulate the following mixed type nondifferentiable higher-order symmetric dual programs over cones.

Primal Problem (MHP):

$$\begin{aligned} \text{Minimize } G(x, y, z, p) = & f^1(x^1, y^1) + s(x^1|E_1) + f^2(x^2, y^2) + s(x^2|E_2) - (y^1)^T z^1 \\ & + g^1(x^1, y^1, p^1) + g^2(x^2, y^2, p^2) - (p^1)^T \nabla_{p^1} g^1(x^1, y^1, p^1) \\ & - (p^2)^T \nabla_{p^2} g^2(x^2, y^2, p^2) - (y^2)^T [\nabla_{y^2} f^2(x^2, y^2) + \nabla_{p^2} g^2(x^2, y^2, p^2)] \end{aligned}$$

subject to

$$-(\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} g^1(x^1, y^1, p^1)) \in C_3^*, \quad (3)$$

$$-(\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} g^2(x^2, y^2, p^2)) \in C_4^*, \quad (4)$$

$$(y^1)^T [\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} g^1(x^1, y^1, p^1)] \geq 0, \quad (5)$$

$$(p^1)^T [\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} g^1(x^1, y^1, p^1)] \geq 0, \quad (6)$$

$$(p^2)^T [\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} g^2(x^2, y^2, p^2)] \geq 0, \quad (7)$$

$$x^1 \in C_1, x^2 \in C_2, y^2 \geq 0, \quad (8)$$

$$z^1 \in D_1, z^2 \in D_2. \quad (9)$$

Dual Problem (MHD):

$$\begin{aligned} \text{Maximize } H(u, v, w, r) = & f^1(u^1, v^1) - S(v^1|D_1) + f^2(u^2, v^2) - S(v^2|D_2) + (u^1)^T w^1 \\ & + h^1(u^1, v^1, r^1) + h^2(u^2, v^2, r^2) - (r^1)^T \nabla_{r^1} h^1(u^1, v^1, r^1) \\ & - (r^2)^T \nabla_{r^2} h^2(u^2, v^2, r^2) - (u^2)^T [\nabla_{u^2} f^2(u^2, v^2) + \nabla_{r^2} h^2(u^2, v^2, r^2)] \end{aligned}$$

subject to

$$(\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)) \in C_1^*, \quad (10)$$

$$(\nabla_{u^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} h^2(u^2, v^2, r^2)) \in C_2^*, \quad (11)$$

$$(u^1)^T [\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)] \leq 0, \quad (12)$$

$$(r^1)^T [\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)] \leq 0, \quad (13)$$

$$(r^2)^T [\nabla_{u^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} h^2(u^2, v^2, r^2)] \leq 0, \quad (14)$$

$$v^1 \in C_3, v^2 \in C_4, u^2 \geq 0, \quad (15)$$

$$w^1 \in E_1, w^2 \in E_2, \quad (16)$$

where, $p^1 \in \mathbb{R}^{|B_1|}$, $p^2 \in \mathbb{R}^{|B_2|}$, $r^1 \in \mathbb{R}^{|A_1|}$, $r^2 \in \mathbb{R}^{|A_2|}$ and E_1 , E_2 , D_1 , and D_2 are compact convex sets in $\mathbb{R}^{|B_1|}$, $\mathbb{R}^{|B_2|}$, $\mathbb{R}^{|A_1|}$, and $\mathbb{R}^{|A_2|}$, respectively.

4. Remark

The above dual programs generalize several models that appeared in the literature can be seen below.

- (A) If $|B_1| = 0$, $|A_1| = 0$, $C_1 = \mathbb{R}^n_+$ and $C_2 = \mathbb{R}^h_+$, then the above programs reduce to non-differentiable Wolfe-type dual programs (see [18]).
- (B) Let $|B_2| = 0$, $|A_2| = 0$, $C_1 = \mathbb{R}^n_+$ and $C_2 = \mathbb{R}^h_+$. Then the above programs become non-differentiable Mond–Weir-type dual programs (see [18]).
- (C) Let $C_1 = \mathbb{R}^n_+$ and $C_2 = \mathbb{R}^h_+$ in (MHP) and (MHD), respectively. Then we get the programs proposed by Verma et al. [14].
- (D) Let $C_1 = \mathbb{R}^n_+$, $C_2 = \mathbb{R}^h_+$, $E = 0$ and $D = 0$ in (MHP) and (MHD), respectively. Then we get the programs discussed by Verma et al. [22].

5. Duality Theorems

In this section, we establish the relations between primal problem (MHP) and dual problem (MHD).

Theorem 1. (Weak Duality) Let $(x^1, x^2, y^1, y^2, z^1, z^2, p^1, p^2)$ be feasible for (MHP) and $(u^1, u^2, v^1, v^2, w^1, w^2, r^1, r^2)$ be feasible for (MHD). Suppose that

- (i) $f^1(., v^1) + (.)^T w^1$ is higher-order \mathcal{F} -pseudo-convex function at u^1 with respect to $h^1(u^1, v^1, r^1)$,
- (ii) $-f^1(x^1, .) + (.)^T z^1$ is higher-order \mathcal{F} -pseudo-convex function at y^1 with respect to $-g^1(x^1, y^1, p^1)$,
- (iii) $f^2(., v^2) + (.)^T w^2$ is higher-order \mathcal{F} -convex function at u^2 with respect to $h^2(u^2, v^2, r^2)$,
- (iv) $-f^2(x^2, .) + (.)^T z^2$ is higher-order \mathcal{F} -convex function at y^2 with respect to $-g^2(x^2, y^2, p^2)$,
- (v) $\mathcal{F}(x^1, u^1; \psi^1) + (u^1)^T \psi^1 + (r^1)^T \psi^1 \geq 0, \forall x^1, u^1 \in C_1, \psi^1 \in C_1^*$,
- (vi) $\mathcal{F}(v^1, y^1; \psi^2) + (y^1)^T \psi^2 + (p^1)^T \psi^2 \geq 0, \forall v^1, y^1 \in C_3, \psi^2 \in C_3^*$,
- (vii) $\mathcal{F}(x^2, u^2; \psi^3) + (u^2)^T \psi^3 + (r^2)^T \psi^3 \geq 0, \forall x^2, u^2 \in C_2, \psi^3 \in C_2^*$ and
- (viii) $\mathcal{F}(v^2, y^2; \psi^4) + (y^2)^T \psi^4 + (p^2)^T \psi^4 \geq 0, \forall v^2, y^2 \in C_4, \psi^4 \in C_4^*$.

Then

$$G(x^1, x^2, y^1, y^2, z^1, z^2, p^1, p^2) \geq H(u^1, u^2, v^1, v^2, w^1, w^2, r^1, r^2). \quad (17)$$

Proof. From hypotheses (iii) and (iv), we get

$$\begin{aligned} f^2(x^2, v^2) + (x^2)^T w^2 - f^2(u^2, v^2) - (u^2)^T w^2 - h^2(u^2, v^2, p^2) + (p^2)^T \nabla_{p^2} h^2(u^2, v^2, p^2) \\ \geq \mathcal{F}(x^2, u^2; \nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)) \end{aligned} \quad (18)$$

and

$$\begin{aligned} f^2(x^2, y^2) - (y^2)^T z^2 - f^2(x^2, v^2) + (v^2)^T z^2 + g^2(x^2, y^2, p^2) - (p^2)^T \nabla_{p^2} g^2(x^2, y^2, p^2) \\ \geq \mathcal{F}(v^2, y^2; -(\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2))). \end{aligned} \quad (19)$$

Let

$$\psi^3 = (\nabla_{u^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} h^2(u^2, v^2, r^2)) \in C_2^*$$

and

$$\psi^4 = -(\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} g^2(x^2, y^2, p^2)) \in C_4^*.$$

Now from hypotheses (vii) and (viii), we get

$$\begin{aligned} \mathcal{F}(x^2, u^2; \nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)) + u^2 [\nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)] \\ \geq -r^2 [\nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)] \end{aligned}$$

and

$$\begin{aligned} \mathcal{F}(v^2, y^2; -(\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2))) - y^2 [\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2)] \\ \geq p^2 [\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2)]. \end{aligned}$$

The above inequalities along with inequalities (7) and (14) imply that

$$\mathcal{F}(x^2, u^2; \nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)) \geq -u^2 [\nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)]$$

and

$$\mathcal{F}(v^2, y^2; -(\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2))) \geq y^2 [\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2)].$$

The above inequalities along with Equations (18) and (19) give

$$\begin{aligned} f^2(x^2, v^2) + (x^2)^T w^2 - f^2(u^2, v^2) - (u^2)^T w^2 - h^2(u^2, v^2, p^2) + (p^2)^T \nabla_{p^2} h^2(u^2, v^2, p^2) \\ \geq -u^2 [\nabla_{x^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)] \end{aligned}$$

and

$$\begin{aligned} f^2(x^2, y^2) - (y^2)^T z^2 - f^2(x^2, v^2) + (v^2)^T z^2 + g^2(x^2, y^2, p^2) - (p^2)^T \nabla_{p^2} g^2(x^2, y^2, p^2) \\ \geq y^2 [\nabla_{y^2} f^2(x^2, y^2) - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2)]. \end{aligned}$$

Now adding the above two inequalities, we obtain

$$\begin{aligned}
& f^2(x^2, y^2) + (x^2)^T w^2 - (y^2)^T z^2 + g^2(x^2, y^2, p^2) - (p^2)^T \nabla_{p^2} g^2(x^2, y^2, p^2) - y^2 [\nabla_{y^2} f^2(x^2, y^2) \\
& \quad - z^2 + \nabla_{p^2} h^2(x^2, y^2, p^2)] \\
& \geq f^2(u^2, v^2) + (u^2)^T w^2 - (v^2)^T z^2 + h^2(u^2, v^2, p^2) - (p^2)^T \nabla_{p^2} h^2(u^2, v^2, p^2) - u^2 [\nabla_{x^2} f^2(u^2, v^2) \\
& \quad + w^2 + \nabla_{r^2} g^2(u^2, v^2, r^2)].
\end{aligned}$$

Noting that $(x^2)^T w^2 \leq s(x^2 | E_2)$ and $(v^2)^T z^2 \leq s(v^2 | D_2)$, we have

$$\begin{aligned}
& f^2(x^2, y^2) + s(x^2 | E_2) + g^2(x^2, y^2, p^2) - y^2 [\nabla_{y^2} f^2(x^2, y^2) + \nabla_{p^2} h^2(x^2, y^2, p^2)] \\
& \geq f^2(u^2, v^2) - s(v^2 | D_2) + h^2(u^2, v^2, p^2) \\
& \quad - (p^2)^T \nabla_{p^2} h^2(u^2, v^2, p^2) - u^2 [\nabla_{x^2} f^2(u^2, v^2) + \nabla_{r^2} h^2(u^2, v^2, r^2)].
\end{aligned} \tag{20}$$

By taking

$$\psi^1 = (\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)) \in C_1^*$$

and

$$\psi^2 = -(\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} g^1(x^1, y^1, p^1)) \in C_3^*,$$

we have

$$\mathcal{F}(x^1, u^1; \nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} g^1(u^1, v^1, r^1)) \geq -(u^1 + r^1) [\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} g^1(u^1, v^1, r^1)]$$

(by the hypothesis (v))

and

$$\mathcal{F}(v^1, y^1; -(\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} h^1(x^1, y^1, p^1))) \geq (y^1 + p^1) [\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} h^1(x^1, y^1, p^1)]$$

(by the hypothesis (vi)).

The above two inequalities along with inequalities (5), (6), (12), and (13) give

$$\mathcal{F}(x^1, u^1; \nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} g^1(u^1, v^1, r^1)) \geq 0$$

and

$$\mathcal{F}(v^1, y^1; -(\nabla_{y^1} f^1(x^1, y^1) - z^1 + \nabla_{p^1} h^1(x^1, y^1, p^1))) \geq 0,$$

which by hypotheses (i) and (ii) imply

$$f^1(x^1, v^1) + (x^1)^T w^1 - f^1(u^1, v^1) - (v^1)^T w^1 - h^1(u^1, v^1, p^1) + (p^1)^T \nabla_{p^1} h^1(u^1, v^1, p^1) \geq 0$$

and

$$f^1(x^1, y^1) + (v^1)^T z^1 - f^1(x^1, v^1) - (y^1)^T z^1 + g^1(x^1, y^1, p^1) - (p^1)^T \nabla_{p^1} g^1(x^1, y^1, p^1) \geq 0.$$

Adding the above two inequalities, we obtain

$$\begin{aligned}
& f^1(x^1, y^1) - (y^1)^T z^1 + (x^1)^T w^1 + g^1(x^1, y^1, p^1) - (p^1)^T \nabla_{p^1} g^1(x^1, y^1, p^1) \\
& \geq f^1(u^1, v^1) + (v^1)^T w^1 - (v^1)^T z^1 + h^1(u^1, v^1, p^1) - (p^1)^T \nabla_{p^1} h^1(u^1, v^1, p^1).
\end{aligned}$$

Using $(x^1)^T w^1 \leq s(x^1|E_1)$ and $(v^1)^T z^1 \leq s(v^1|D_1)$, we have

$$\begin{aligned} f^1(x^1, y^1) - (y^1)^T z^1 + s(x^1|E_1) + g^1(x^1, y^1, p^1) - (p^1)^T \nabla_{p^1} g^1(x^1, y^1, p^1) \\ \geq f^1(u^1, v^1) + (v^1)^T w - s(v^1|D_1) + h^1(u^1, v^1, p^1) - (p^1)^T \nabla_{p^1} h^1(u^1, v^1, p^1). \end{aligned} \quad (21)$$

Combining inequalities (20) and (21), we get

$$G(x^1, x^2, y^1, y^2, z^1, z^2, p^1, p^2) \geq H(u^1, u^2, v^1, v^2, w^1, w^2, r^1, r^2).$$

□

Theorem 2. (Strong Duality)

Let $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1, \bar{p}^2)$ be an optimal solution of (MHP). Suppose that

- (i) $\nabla_{p^1 p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)$ is positive or negative definite matrix and $\nabla_{p^2 p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)$ is negative definite matrix,
- (ii) $\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) \neq 0$ and $\nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) \neq 0$,
- (iii) $(\bar{p}^1)^T [\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0 \Rightarrow \bar{p}^1 = 0$
and
 $\bar{y}^2 [\nabla_{y^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) - \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) + \nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] = 0 \Rightarrow \bar{p}^2 = 0$,
- (iv) $g^1(\bar{x}^1, \bar{y}^1, 0) = h^1(\bar{x}^1, \bar{y}^1, 0)$, $\nabla_{x^1} g^1(\bar{x}^1, \bar{y}^1, 0) = \nabla_{x^1} h^1(\bar{x}^1, \bar{y}^1, 0)$,
 $\nabla_{y^1} g^1(\bar{x}^1, \bar{y}^1, 0) = \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, 0)$, $g^2(\bar{x}^2, \bar{y}^2, 0) = h^2(\bar{x}^2, \bar{y}^2, 0)$ and
 $\nabla_{x^2} g^2(\bar{x}^2, \bar{y}^2, 0) = \nabla_{x^2} h^2(\bar{x}^2, \bar{y}^2, 0)$.

Then,

- (I) $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1 = 0, \bar{r}^2 = 0)$ is feasible for (MHD), and
- (II) $G(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1, \bar{p}^2) = H(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1, \bar{r}^2)$.

Furthermore, if the hypotheses of Theorem 1 are satisfied for all feasible solutions of (MHP) and (MHD), then $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1 = 0, \bar{r}^2 = 0)$ is an optimal solution for (MHD).

Proof. As $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1, \bar{p}^2)$ is an optimal solution of (MHP), by the Fritz John necessary optimality conditions [23,24], there exist $\alpha, \gamma, \delta^1, \delta^2 \in \mathbb{R}$, $\beta^1 \in \mathbb{R}^{|B_1|}$, $\beta^2, \eta \in \mathbb{R}^{|B_2|}$, $\zeta^1 \in \mathbb{R}^{|A_1|}$, $\zeta^2 \in \mathbb{R}^{|A_2|}$ such that the following conditions are satisfied at $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1, \bar{p}^2)$:

$$\begin{aligned} \{ \alpha [\nabla_{x^1} f^1(\bar{x}^1, \bar{y}^1) + \zeta^1 + \nabla_{x^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) - \nabla_{p^1 x^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) \bar{p}^1] + [\nabla_{y^1 x^1} f^1(\bar{x}^1, \bar{y}^1) \\ + \nabla_{p^1 x^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] (\beta^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1) \} (x^1 - \bar{x}^1) \geq 0, \text{ for all } x^1 \in C_1, \end{aligned} \quad (22)$$

$$\begin{aligned} \{ \alpha [\nabla_{x^2} f^2(\bar{x}^2, \bar{y}^2) + \zeta^2 + \nabla_{x^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] + \{ \nabla_{p^2 x^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) \} (\beta^2 - \alpha \bar{y}^2 - \alpha \bar{p}^2 \\ - \delta^2 \bar{p}^2) + \{ \nabla_{y^2 x^2} f^2(\bar{x}^2, \bar{y}^2) \} (\beta^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) \} (x^2 - \bar{x}^2) \geq 0, \text{ for all } x^2 \in C_2, \end{aligned} \quad (23)$$

$$\alpha[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{y^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) - \nabla_{p^1 y^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) \bar{p}^1] + (\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) + \nabla_{p^1 y^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1))(\beta^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1) - \gamma[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$

for all, $y^1 \in \mathbb{R}^{B_1}$,

(24)

$$\{\nabla_{p^2 y^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)\}(\beta^2 - \alpha \bar{y}^2 - \alpha \bar{p}^2 - \delta^2 \bar{p}^2) + \alpha[\nabla_{y^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) - \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] + \{\nabla_{y^2 y^2} f^2(\bar{x}^2, \bar{y}^2)\}(\beta^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) - \eta = 0 \text{ for all, } y^2 \in \mathbb{R}^{B_2},$$
(25)

$$\{\nabla_{p^1 p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)\}(\beta^1 - \alpha \bar{p}^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1) - \delta^1[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$
(26)

$$\{\nabla_{p^2 p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)\}(\beta^2 - \alpha \bar{y}^2 - \alpha \bar{p}^2 - \delta^2 \bar{p}^2) - \delta^2[\nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] = 0,$$
(27)

$$\beta^1[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$
(28)

$$\beta^2[\nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] = 0,$$
(29)

$$\gamma \bar{y}^1[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$
(30)

$$\delta^1 \bar{p}^1[\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$
(31)

$$\delta^2 \bar{p}^2[\nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] = 0,$$
(32)

$$(\alpha - \gamma) y^1 + \beta^1 - \delta^1 p^1 \in N_{D_1}(\bar{z}^1),$$
(33)

$$\beta^2 - \delta^2 p^2 \in N_{D_2}(\bar{z}^2),$$
(34)

$$(\xi^1)^T \bar{x}^1 = S(x^1 | E_1), \xi^1 \in C_1,$$
(35)

$$(\xi^2)^T \bar{x}^2 = S(x^2 | E_2), \xi^2 \in C_2,$$
(36)

$$\eta \bar{y}^2 = 0,$$
(37)

$$(\alpha, \beta^1, \beta^2, \gamma, \delta^1, \delta^2, \eta) \neq 0,$$
(38)

$$(\alpha, \beta^1, \beta^2, \gamma, \delta^1, \delta^2, \eta) \geq 0.$$
(39)

Premultiplying Equations (26) and (27) by $(\beta^1 - \alpha \bar{p}^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1)$ and $(\beta^2 - \alpha \bar{p}^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2)$, respectively, and then using Equations (28)–(32), we get

$$(\beta^1 - \alpha \bar{p}^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1)^T \nabla_{p^1 p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) (\beta^1 - \alpha \bar{p}^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1) = 0,$$
(40)

and

$$\begin{aligned} & (\beta^2 - \alpha \bar{p}^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) \nabla_{p^2 p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) (\beta^2 - \alpha \bar{p}^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) \\ & = -\alpha \delta^2 \bar{y}^2 [\nabla_{y^2} f^2(\bar{x}^2, \bar{y}^2) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)]. \end{aligned}$$
(41)

Using hypothesis (i) in (40), we get

$$\beta^1 = \alpha \bar{p}^1 + \gamma \bar{y}^1 + \delta^1 \bar{p}^1.$$
(42)

Further, using inequalities (4), (8), and (39) in (41), we obtain

$$(\beta^2 - \alpha \bar{p}^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) \nabla_{p^2 p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) (\beta^2 - \alpha \bar{p}^2 - \alpha \bar{y}^2 - \delta^2 \bar{p}^2) \geq 0,$$

which on using hypothesis (i), we obtain

$$\beta^2 = \alpha \bar{p}^2 + \alpha \bar{y}^2 + \delta^2 \bar{p}^2. \quad (43)$$

From Equations (26), (27), along with hypothesis (ii), we obtain

$$\delta^1 = 0 \quad (44)$$

and

$$\delta^2 = 0. \quad (45)$$

Now suppose, $\alpha = 0$. Then, Equations (43) and (45) give $\beta^2 = 0$. The Equation (24) with hypothesis (ii) implies $\gamma = 0$, and Equation (42) yields $\beta^1 = 0$. Therefore, Equation (25) implies $\eta = 0$. Thus $(\alpha, \beta^1, \beta^2, \gamma, \delta^1, \delta^2, \eta) = 0$, which contradicts the Equation (38). Therefore,

$$\alpha > 0. \quad (46)$$

From Equations (28), (30) and (31), we have

$$(\beta^1 - \gamma \bar{y}^1 - \delta^1 \bar{p}^1)^T [\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0,$$

which along with (42), gives

$$\alpha (\bar{p}^1)^T [\nabla_{y^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] = 0. \quad (47)$$

Using hypothesis (iii), we get

$$\bar{p}^1 = 0. \quad (48)$$

Further, from Equations (25), (37), and (43), we obtain

$$\alpha \bar{y}^2 [\nabla_{y^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) - \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) + \nabla_{y^2 y^2} f^2(\bar{x}^2, \bar{y}^2) \bar{p}^2] = 0, \quad (49)$$

which by hypothesis (iii) and Equation (46) implies

$$\bar{p}^2 = 0. \quad (50)$$

Therefore, (42) and (43) reduce to

$$\beta^1 = \gamma \bar{y}^1 \quad (51)$$

and

$$\beta^2 = \alpha \bar{y}^2. \quad (52)$$

Now from Equations (24), (42), and (48) with hypotheses (ii) and (iv), we get

$$\alpha = \gamma > 0. \quad (53)$$

Therefore, Equation (52) implies

$$\bar{y}^2 = \frac{\beta^2}{\alpha} \geq 0. \quad (54)$$

Now, Equations (22) and (23) along with (42), (43), (50) and hypothesis (iv), we obtain

$$\alpha[\nabla_{x^1} f^1(\bar{x}^1, \bar{y}^1) + \bar{\xi}^1 + \nabla_{r^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)](x^1 - \bar{x}^1) \geq 0, \text{ for all } x^1 \in C_1 \quad (55)$$

and

$$\alpha[\nabla_{x^2} f^2(\bar{x}^2, \bar{y}^2) + \bar{\xi}^2 + \nabla_{r^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)](x^2 - \bar{x}^2) \geq 0, \text{ for all } x^2 \in C_2. \quad (56)$$

Let $x^1 \in C_1$, then $\bar{x}^1 + x^1 \in C_1$. Then above inequality (55) implies

$$(\bar{x}^1)^T [\nabla_{x^1} f^1(\bar{x}^1, \bar{y}^1) - \bar{\xi}^1 + \nabla_{r^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1)] \geq 0, \text{ for all } x^1 \in C_1. \quad (57)$$

Similarly,

$$(\bar{x}^2)^T [\nabla_{x^2} f^2(\bar{x}^2, \bar{y}^2) + \bar{\xi}^2 + \nabla_{r^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2)] \geq 0, \text{ for all } x^2 \in C_2. \quad (58)$$

Therefore,

$$\nabla_{x^1} f^1(\bar{x}^1, \bar{y}^1) + \bar{\xi}^1 + \nabla_{r^1} g^1(\bar{x}^1, \bar{y}^1, \bar{p}^1) \in C_1^* \quad (59)$$

$$\nabla_{x^2} f^2(\bar{x}^2, \bar{y}^2) + \bar{\xi}^2 + \nabla_{r^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) \in C_2^*. \quad (60)$$

If we take $\bar{\xi}^1 = \bar{w}^1$ and $\bar{\xi}^2 = \bar{w}^2$, then $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1 = 0, \bar{r}^2 = 0)$ satisfies the dual constraints (10)–(15). Therefore, it is feasible solution for (MHD).

Now from Equations (34), (46), (50), and (52), we get $\bar{y}^2 \in N_{D_2}(\bar{z}^2)$. As D_2 is a compact convex set in $\mathbb{R}^{|A_2|}$,

$$(\bar{y}^2)^T \bar{z}^2 = S(\bar{y}^2 | D_2). \quad (61)$$

Moreover, since $\beta^1 = \gamma \bar{y}^1$ and $\gamma > 0$ then from Equations (33), (48), and (53), we obtain $\bar{y}^1 \in N_{D_1}(\bar{z}^1)$. As D_1 is a compact convex set in $\mathbb{R}^{|A_1|}$,

$$(\bar{y}^1)^T \bar{z}^1 = S(\bar{y}^1 | D_1). \quad (62)$$

Therefore by Equations (35), (36), (48), (50), (61), (62), and hypothesis (iv), we get the equal objective functions value of (MHP) and (MHD) at $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1 = 0, \bar{p}^2 = 0)$ and $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1 = 0, \bar{r}^2 = 0)$. From Theorem 1, we get that $(\bar{x}^1, \bar{x}^2, \bar{y}^1, \bar{y}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1 = 0, \bar{r}^2 = 0)$ is an optimal solution for (MHD). \square

Theorem 3. (Converse Duality) Let $(\bar{u}^1, \bar{u}^2, \bar{v}^1, \bar{v}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1, \bar{r}^2)$ be an optimal solution of (MHD). Suppose that

(i) $\nabla_{r^1, r^1} h^1(\bar{u}^1, \bar{v}^1, \bar{r}^1)$ is positive or negative definite matrix and $\nabla_{r^2, r^2} h^2(\bar{u}^2, \bar{v}^2, \bar{r}^2)$ is negative definite matrix,

(ii) $\nabla_{v^1} f^1(\bar{u}^1, \bar{v}^1) + \bar{w}^1 + \nabla_{r^1} h^1(\bar{u}^1, \bar{v}^1, \bar{r}^1) \neq 0$ and $\nabla_{v^2} f^2(\bar{u}^2, \bar{v}^2) + \bar{w}^2 + \nabla_{r^2} h^2(\bar{u}^2, \bar{v}^2, \bar{r}^2) \neq 0$,

(iii) $(\bar{r}^1)^T [\nabla_{u^1} f^1(\bar{u}^1, \bar{v}^1) + \bar{w}^1 + \nabla_{r^1} h^1(\bar{u}^1, \bar{v}^1, \bar{r}^1)] = 0 \Rightarrow \bar{r}^1 = 0$

and

$$\bar{u}^2 [\nabla_{u^2} h^2(\bar{u}^2, \bar{v}^2, \bar{r}^2) - \nabla_{r^2} h^2(\bar{u}^2, \bar{v}^2, \bar{r}^2) + \nabla_{u^2 u^2} f^2(\bar{u}^2, \bar{v}^2) \bar{r}^2] = 0 \Rightarrow \bar{r}^2 = 0,$$

(iv) $h^1(\bar{u}^1, \bar{v}^1, 0) = g^1(\bar{u}^1, \bar{v}^1, 0)$, $\nabla_{u^1} h^1(\bar{u}^1, \bar{v}^1, 0) = \nabla_{p^1} g^1(\bar{u}^1, \bar{v}^1, 0)$,

$$\nabla_{v^1} h^1(\bar{u}^1, \bar{v}^1, 0) = \nabla_{r^1} h^1(\bar{u}^1, \bar{v}^1, 0), \quad h^2(\bar{u}^2, \bar{v}^2, 0) = g^2(\bar{u}^2, \bar{v}^2, 0) \text{ and}$$

$$\nabla_{u^2} h^2(\bar{u}^2, \bar{v}^2, 0) = \nabla_{p^2} g^2(\bar{u}^2, \bar{v}^2, 0).$$

Then,

(I) $(\bar{u}^1, \bar{u}^2, \bar{v}^1, \bar{v}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1 = 0, \bar{p}^2 = 0)$ is feasible for (MHP), and

(II) $G(\bar{u}^1, \bar{u}^2, \bar{v}^1, \bar{v}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1, \bar{p}^2) = H(\bar{u}^1, \bar{u}^2, \bar{v}^1, \bar{v}^2, \bar{w}^1, \bar{w}^2, \bar{r}^1, \bar{r}^2)$.

Furthermore, if the hypotheses of Theorem 1 are satisfied for all feasible solutions of (MHP) and (MHD), then $(\bar{u}^1, \bar{u}^2, \bar{v}^1, \bar{v}^2, \bar{z}^1, \bar{z}^2, \bar{p}^1 = 0, \bar{p}^2 = 0)$ is an optimal solution for (MHP).

Proof. The proof follows on the line of Theorem 2. \square

5.1. Self Duality

In general, primal problem (MHP) and dual problem (MHD) are not self-duals without including restrictions on f , h and g . If we assume $D_1 = D_2, E_1 = E_2, f^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \rightarrow \mathbb{R}, f^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \rightarrow \mathbb{R}, h^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \times \mathbb{R}^{|A_1|} \rightarrow \mathbb{R}, h^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \times \mathbb{R}^{|A_2|} \rightarrow \mathbb{R}, g^1 : \mathbb{R}^{|A_1|} \times \mathbb{R}^{|B_1|} \times \mathbb{R}^{|B_1|} \rightarrow \mathbb{R}, g^2 : \mathbb{R}^{|A_2|} \times \mathbb{R}^{|B_2|} \times \mathbb{R}^{|B_2|} \rightarrow \mathbb{R}$ are skew symmetric, that is,

$$f^i(u^1, v^1) = -f^i(v^1, u^1), i = 1, 2,$$

$$h^i(u^1, v^1, r^1) = -h^i(v^1, u^1, r^1), i = 1, 2,$$

$$C_1 = C_3$$

and

$$C_2 = C_4.$$

By rewriting the dual problem (MHD) as a minimization problem, we get

$$\begin{aligned} \text{Minimize } H(u, v, w, r) = & -\{f^1(u^1, v^1) - S(v^1|D_1) + f_2(u^2, v^2) - S(v^2|D_2) + (u^1)^T w^1 \\ & + h^1(u^1, v^1, r^1) + h_2(u^2, v^2, r^2) - (r^1)^T \nabla_{r^1} h^1(u^1, v^1, r^1) - (r^2)^T \nabla_{r^2} h^2(u^2, v^2, r^2) \\ & - (u^2)^T [\nabla_{u^2} f^2(u^2, v^2) + \nabla_{r^2} h^2(u^2, v^2, r^2)]\} \end{aligned}$$

subject to

$$\begin{aligned} \nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1) & \in C_1^*, \\ \nabla_{u^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} h^2(u^2, v^2, r^2) & \in C_2^*, \\ (u^1)^T [\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)] & \leq 0, \\ (r^1)^T [\nabla_{u^1} f^1(u^1, v^1) + w^1 + \nabla_{r^1} h^1(u^1, v^1, r^1)] & \leq 0, \\ (r^2)^T [\nabla_{u^2} f^2(u^2, v^2) + w^2 + \nabla_{r^2} h^2(u^2, v^2, r^2)] & \leq 0, \\ v^1 \in C_3, v^2 \in C_4, u^2 \geq 0, & \end{aligned}$$

As f , h and g are skew symmetric, therefore

$$\nabla_{u^1} f^1(u^1, v^1) = -\nabla_{u^1} f^1(v^1, u^1), \quad \nabla_{u^1} f^2(u^2, v^2) = -\nabla_{u^1} f^2(v^2, u^2),$$

$$\nabla_{u^1} h^1(u^1, v^1, r^1) = -\nabla_{u^1} h^1(v^1, u^1, r^1), \quad \text{and} \quad \nabla_{u^1} h^2(u^2, v^2, r^2) = -\nabla_{u^1} h^2(v^2, u^2, r^2).$$

Now the above problem becomes:

$$\begin{aligned} \text{Minimize } H(u, v, w, r) = & f^1(v^1, u^1) + S(v^1|D_1) + f^2(v^2, u^2) + S(v^2|D_2) - (u^1)^T w^1 \\ & + h^1(v^1, u^1, r^1) + h^2(v^2, u^2, r^2) - (r^1)^T \nabla_{r^1} h^1(v^1, u^1, r^1) \\ & - (r^2)^T \nabla_{r^2} h^2(v^2, u^2, r^2) - (u^2)^T [\nabla_{u^2} f^2(v^2, u^2) + \nabla_{r^2} g^2(v^2, u^2, r^2)] \end{aligned}$$

subject to

$$\begin{aligned} -(\nabla_{u^1} f^1(v^1, u^1) + w^1 + \nabla_{r^1} h^1(v^1, u^1, r^1)) &\in C_3^*, \\ -(\nabla_{u^2} f^2(v^2, u^2) + w^2 + \nabla_{r^2} h^2(v^2, u^2, r^2)) &\in C_4^*, \\ (u^1)^T [\nabla_{u^1} f^1(v^1, u^1) + w^1 + \nabla_{r^1} h^1(v^1, u^1, r^1)] &\geq 0, \\ (r^1)^T [\nabla_{u^1} f^1(v^1, u^1) + w^1 + \nabla_{r^1} h^1(v^1, u^1, r^1)] &\geq 0, \\ (r^2)^T [\nabla_{u^2} f^2(v^2, u^2) + w^2 + \nabla_{r^2} h^2(v^2, u^2, r^2)] &\geq 0, \\ v^1 \in C_1, v^2 \in C_2, u^2 \geq 0, \end{aligned}$$

which is identical to (MHP), that is, the objective function and the constraint functions of (MHP) and (MHD) are identical. Thus, (MHP) is a self-dual.

Thus, if $(x^1, x^2, y^1, y^2, w^1, w^2, p^1, p^2)$ is feasible for (MHP), then $(y^1, y^2, x^1, x^2, w^1, w^2, p^1, p^2)$ is feasible for (MHD) and vice versa.

The weak and strong duality theorems are verified by the following example.

5.2. Example

Let $f(x, y) = (x_1^2 + y_1^2 + x_1 + y_1 + x_2^2 + y_2^2 + x_2 + y_2)$, $s(x|E) = \frac{x+|x|}{2}$, $g(x, y, p) = \frac{1}{2} p^T (\nabla_{yy} f(x, y) p)$, $h(x, y, r) = \frac{1}{2} r^T (\nabla_{yy} f(x, y) r)$ and $C_1 = C_2 = C_3 = C_4 = \mathbb{R}_+^2$. Then, the problems (MHP) and (MHD) are as follows.

Primal Problem (MHP):

Minimize $G(x, y, z, p) =$

$$x_1^2 + x_2^2 + x_1 + x_2 + y_1^2 + y_2^2 + y_1 + y_2 + \frac{x_1+|x_1|}{2} + \frac{x_2+|x_2|}{2} - y_1 z_1 - p_1^2 - p_2^2 - y_2(2y_2 + 1 + 2p_2)$$

subject to

$$\begin{aligned} (2y_1 + 1 - z_1 + 2p_1) &\leq 0, \\ (2y_2 + 1 - z_2 + 2p_2) &\leq 0, \\ (2y_1^2 + y_1 - y_1 z_1 + 2p_1 y_1) &\geq 0, \\ (2p_1 y_1 + p_1 - p_1 z_1 + 2p_1^2) &\geq 0, \\ (2p_2 y_2 + p_2 - p_2 z_2 + 2p_2^2) &\geq 0, \\ x, z, y_2 &\geq 0. \end{aligned}$$

Dual Problem (MHD):

Maximize $H(u, v, w, r) =$

$$u_1^2 + u_2^2 + u_1 + u_2 + v_1^2 + v_2^2 + v_1 + v_2 - \frac{v_1+|v_1|}{2} - \frac{v_2+|v_2|}{2} + u_1 w_1 - r_1^2 - r_2^2 - u_2(2u_2 + 1 + 2r_2)$$

subject to

$$\begin{aligned}(2u_1 + 1 + w_1 + 2r_1) &\geq 0, \\ (2u_2 + 1 + w_2 + 2r_2) &\geq 0, \\ (2u_1^2 + u_1 + u_1w_1 + 2r_1u_1) &\leq 0, \\ (2r_1u_1 + r_1 + r_1w_1 + 2r_1^2) &\leq 0, \\ (2r_2u_2 + r_2 + r_2w_2 + 2r_2^2) &\leq 0, \\ v, w, u_2 &\geq 0,\end{aligned}$$

where

- (i) $f(x, y) : \mathbb{R}^2 \times \mathbb{R}^2 \longrightarrow \mathbb{R}$,
- (ii) $g(x, y, p) : \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \longrightarrow \mathbb{R}$ and
- (iii) $h(x, y, r) : \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \longrightarrow \mathbb{R}$.

$$\text{Let } \bar{x} = \bar{y} = \bar{u} = \bar{v} = \bar{w} = \bar{z} = \bar{p} = \bar{r} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Then, (x, y, z, p) is feasible for (MHP) and (u, v, w, r) is feasible for (MHD).

Now, $f^1(., v) + (.)^T w^1 = (u_1^2 + u_1 + u_1 w_1)$ is a higher-order \mathcal{F} -pseudo-convex function at $\bar{u}^1 = 0$ with respect to h^1 and $-f^1(x, .) + (.)^T z^1 = (-y_1^2 - y_1 + y_1 z_1)$ is a higher-order \mathcal{F} -pseudo-convex function at $\bar{y}^1 = 0$ with respect to $-g^1$. Now, $f^2(., v) + (.)^T w^2 = (u_2^2 + u_2 + u_2 w_2)$ is a higher-order \mathcal{F} -convex function at $\bar{u}^2 = 0$ with respect to h^2 . Similarly, $-f^2(x, .) + (.)^T z^2 = (-y_2^2 - y_2 + y_2 z_2)$ is a higher-order \mathcal{F} -convex function at $\bar{y}^2 = 0$ with respect to $-g^2$.

Also conditions (v) to (viii) of weak duality theorem (Theorem 1) are satisfied for $\bar{x} = \bar{y} = \bar{u} = \bar{v} = \bar{w} = \bar{z} = \bar{p} = \bar{r} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$. Therefore, weak duality theorem is satisfied.

Furthermore,

- (i) $\nabla_{p^1 p^1} g^1(\bar{x}, \bar{y}, \bar{p}) = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$ is positive definite matrix

and

$$\nabla_{p^2 p^2} g^2(\bar{x}, \bar{y}, \bar{p}) = \begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix} \text{ is negative definite matrix,}$$

- (ii) $\{\nabla_{y^1} f^1(\bar{x}, \bar{y}) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}, \bar{y}, \bar{p}), \nabla_{y^2} f^2(\bar{x}, \bar{y}) - \bar{z}^2 + \nabla_{p^2} g^2(\bar{x}, \bar{y}, \bar{p})\} = \begin{pmatrix} 1 \\ 1 \end{pmatrix},$

- (iii) $(\bar{p}^1)^T \{\nabla_{y^1} f^1(\bar{x}, \bar{y}) - \bar{z}^1 + \nabla_{p^1} g^1(\bar{x}, \bar{y}, \bar{p})\} = 0 \implies \bar{p}^1 = 0$ and $\bar{y}^2 [\nabla_{y^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) - \nabla_{p^2} g^2(\bar{x}^2, \bar{y}^2, \bar{p}^2) + \nabla_{y^2 y^2} f^2(\bar{x}^2, \bar{y}^2) \bar{p}^2] = 0 \implies \bar{p}^2 = 0.$

(As \bar{p} initially taken as zero).

- (iv) $h^i(\bar{x}, \bar{y}, 0) = g^i(\bar{x}, \bar{y}, 0)$, $\nabla_x h^i(\bar{x}, \bar{y}, 0) = \nabla_x g^i(\bar{x}, \bar{y}, 0)$, $\nabla_{y^1} g^1(\bar{x}^1, \bar{y}^1, 0) = \nabla_{p^1} g^1(\bar{x}^1, \bar{y}^1, 0)$, for $i = 1, 2$.

Therefore, all the assumptions of strong duality theorem (Theorem 2) are also satisfied. Now objective function value is equal to 0.

6. Special Cases

- (i) Let $C_1 = \mathbb{R}^n_+$, $C_2 = \mathbb{R}^h_+$, $|B_1| = 0$, $|A_1| = 0$, $E_1 = 0$ and $D_1 = 0$ in (MHP) and (MHD). Then, we get the programs proposed by Verma and Gulati [21].
- (ii) If $C_1 = \mathbb{R}^n_+$, $C_3 = \mathbb{R}^h_+$, removed the higher-order and non-differentiable terms and omission of inequalities (6), (7), (13), and (14), then we get the model presented by Chandra et al. [9].
- (iii) Let $C_1 = \mathbb{R}^n_+$ and $C_3 = \mathbb{R}^h_+$, $|A_2| = 0$, $|B_2| = 0$, $p = 0$ and $r = 0$. Then, our programs are reduced to the programs presented in [25].
- (iv) If $C_1 = -C_1$, $C_3 = -C_3$, $|B_2| = 0$, $|A_2| = 0$, $E_1 = 0$, $D_1 = 0$ and omission of inequalities (7) and (14), then we get the dual programs discussed in [26].
- (v) If $C_2 = -C_2$, $C_4 = -C_4$, $|B_1| = 0$, $|A_1| = 0$, $E_2 = 0$, $D_2 = 0$ and omission of inequalities (6) and (13), then we get the dual programs studied in [26].
- (vi) If $C_2 = -C_2$, $C_4 = -C_4$, $|B_1| = 0$, $|A_1| = 0$, $E_2 = 0$, $D_2 = 0$, $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$ and $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$, and omission of inequalities (6) and (13), then we get the dual programs is equal to obtained in [27].

For $C_2 = \mathbb{R}^n_+$, $C_4 = \mathbb{R}^h_+$, and omission of inequalities (6) and (13), we get the following special cases (vii) to (viii):

- (vii) If $|B_1| = 0$ and $|A_1| = 0$, then we get the programs considered in [17].
- (viii) Let $|B_1| = 0$, $|A_1| = 0$, $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$ and $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$. Then, our programs become the programs presented in [16].

For $C_1 = \mathbb{R}^n_+$, $C_3 = \mathbb{R}^h_+$, and omission of inequalities (7) and (14), we get the following special cases (ix) to (xiii):

- (ix) If $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$ and $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$, then we get the programs obtained in [15].
- (x) If $|B_2| = 0$, $|A_2| = 0$, $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$ and $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$, then we get the dual programs derived in [19].
- (xi) If $|B_2| = 0$, $|A_2| = 0$, $E_1 = 0$, $D_1 = 0$, $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$ and $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$, then we get the dual programs studied in [4].
- (xii) If $|B_2| = 0$, $|A_2| = 0$, $g(x, y, p) = (1/2)p^T \nabla_{yy} f(x, y)p$, $h(u, v, r) = (1/2)r^T \nabla_{xx} f(u, v)r$, $E_1 = 0$ and $D_1 = 0$, then we get the dual programs presented in [28].
- (xiii) Let $C_1 = -C_1$, $C_3 = -C_3$, $|A_2| = 0$, $|B_2| = 0$, $p = 0$, $r = 0$, $E_1 = 0$ and $D_1 = 0$. Then, we obtain symmetric dual programs [29,30].

7. Conclusions

A new mixed type nondifferentiable higher-order symmetric dual programs over cones has been formulated. It is also shown the uniqueness of higher-order \mathcal{F} -convexity/higher-order \mathcal{F} -pseudoconvexity by a non-trivial example. Under these assumptions, the weak, strong, and converse duality theorems have been derived, and verified by an example. Self-duality has been also discussed. The presented programs and results generalize some existing duals and their corresponding theorems appeared in the literature. The symmetric duality between (MHP) and (MHD) can be extended for fractional and multiobjective programming problems over cones.

Author Contributions: All authors contributed equally and significantly in writing this article. All authors have read and agreed to the published version of the manuscript.

Funding: Deanship of Research, King Fahd University of Petroleum and Minerals, Saudi Arabia, Project No. IN171012.

Acknowledgments: The first and third authors would like to thank the King Fahd University of Petroleum and Minerals, Saudi Arabia to provide the financial support under the Internal Research Project no. IN171012. The authors are thankful to referees for their valuable suggestions which improved the results and presentation of this article.

Conflicts of Interest: The authors declare no conflicts of interest.

Appendix A

The Figures 1–4 are plotted in Wolfram Mathematica. The codes are given below:

Figure 1: `Plot3D[(-u^2 + x^2)(-e^(-u) + 2u + 1/(1 + u)), x, 0, 5, u, 0, 5]`

Figure 2: `Plot[(-1 + x^2)(-e^(-1) + 2 + 1/2), x, 0, 5, Frame -> True, FrameLabel -> x]`

Figure 3: `Plot[(e^(-x) + x^2 - 1/e - 1), x, 0, 5, Frame -> True, FrameLabel -> x]`

Figure 4: `Plot[e^(-x) + x^2 - 1/e - 1 - 2.1321205588(x^2 - 1), x, 0, 4, Frame -> True, FrameLabel -> x]`.

References

1. Mangasarian, O.L. Second and higher-order duality in nonlinear programming. *J. Math. Anal. Appl.* **1975**, *51*, 607–620. [\[CrossRef\]](#)
2. Mond, B.; Zhang, J. Higher-order invexity and duality in mathematical programming. In *Generalized Convexity, Generalized Monotonicity: Recent Results*; Crouzeix, J.P., Legaz, J.E.M., Volle, M., Eds.; Kluwer Academic: Dordrecht, The Netherlands, 1998; pp. 357–372.
3. Chen, X. Higher-order symmetric duality in nondifferentiable multiobjective programming problems. *J. Math. Anal. Appl.* **2004**, *290*, 423–435. [\[CrossRef\]](#)
4. Mishra, S.K. Nondifferentiable higher-order symmetric duality in mathematical programming with generalized invexity. *Eur. Oper. Res.* **2005**, *167*, 28–34. [\[CrossRef\]](#)
5. Khurana, S. Symmetric duality in multiobjective programming involving generalized cone-invex functions. *Eur. Oper. Res.* **2005**, *165*, 592–597. [\[CrossRef\]](#)
6. Ahmad, I.; Husain, Z.; Sharma, S. Higher-order duality in non-differentiable multiobjective programming. *Numer. Funct. Anal. Optim.* **2007**, *28*, 989–1002. [\[CrossRef\]](#)
7. Gupta, S.K.; Jayswal, A. Multiobjective higher-order symmetric duality involving generalized cone-invex functions. *Comput. Math. Appl.* **2010**, *60*, 3187–3192. [\[CrossRef\]](#)
8. Ahmad, I.; Husain, Z. Multiobjective mixed symmetric duality involving cones. *Comput. Math. Appl.* **2010**, *59*, 319–326. [\[CrossRef\]](#)
9. Chandra, S.; Husain, I. On mixed symmetric duality in mathematical programming. *Opsearch* **1999**, *36*, 165–171. [\[CrossRef\]](#)
10. Xu, Z. Mixed type duality in multiobjective programming problems. *J. Math. Anal. Appl.* **1996**, *198*, 621–635. [\[CrossRef\]](#)
11. Yang, X.M.; Teo, K.L.; Yang, X.Q. Mixed symmetric duality in nondifferentiable mathematical programming. *Indian J. Pure Appl. Math.* **2003**, *34*, 805–815.

12. Bector, C.R.; Chandra, S. On mixed symmetric duality in multiobjective programming. *Opsearch* **1999**, *36*, 399–407. [\[CrossRef\]](#)
13. Ahmad, I. Multiobjective mixed symmetric duality with invexity. *N. Z. J. Math.* **2005**, *34*, 1–9.
14. Verma, K.; Mathur, P.; Gulati, T.R. A new approach on mixed type nondifferentiable higher-order symmetric duality. *J. Oper. Res. Soc. China* **2019**, *7*, 321–335. [\[CrossRef\]](#)
15. Agarwal, R.P.; Ahmad, I.; Gupta, S.K.; Kailey, N. Generalized Second-order mixed symmetric duality in nondifferentiable mathematical programming. In *Abstract and Applied Analysis*; Hindawi: London, UK; New York, NY, USA, 2011; Volume 103597, 14p.
16. Gulati, T.R.; Gupta, S.K. Wolfe-type second order symmetric duality in nondifferentiable programming. *J. Math. Anal. Appl.* **2005**, *310*, 247–253. [\[CrossRef\]](#)
17. Gulati, T.R.; Gupta, S.K. Higher order nondifferentiable symmetric duality with generalized F-convexity. *J. Math. Anal. Appl.* **2007**, *329*, 229–237. [\[CrossRef\]](#)
18. Gulati, T.R.; Verma, K. Nondifferentiable higher-order symmetric duality under invexity/generalized invexity. *Filomat* **2014**, *28*, 1661–1674. [\[CrossRef\]](#)
19. Hou, S.H.; Yang, X.M. On second-order symmetric duality in nondifferentiable programming. *J. Math. Anal. Appl.* **2001**, *255*, 488–491. [\[CrossRef\]](#)
20. Verma, K.; Gulati, T.R. Higher order symmetric duality using generalized invexity. In *Proceeding of the 3rd International Conference on Operations Research and Statistics (ORS 2013)*, Singapore, 22–23 April 2013. [\[CrossRef\]](#)
21. Verma, K.; Gulati, T.R. Wolfe-type higher-order symmetric duality under invexity. *J. Appl. Math. Inform.* **2014**, *32*, 153–159. [\[CrossRef\]](#)
22. Verma, K.; Mathur, P.; Gulati, T.R. Mixed type higher order symmetric duality over cones. *J. Model. Optim.* **2017**, *7*, 1–7. [\[CrossRef\]](#)
23. Schechter, M. More on subgradient duality. *J. Math. Anal. Appl.* **1979**, *71*, 251–262. [\[CrossRef\]](#)
24. Suneja, S.K.; Agarwal, S.; Dawar, S. Multiobjective symmetric duality involving cones. *Eur. Oper. Res.* **2002**, *141*, 471–479. [\[CrossRef\]](#)
25. Mond, B.; Schechter, M. Nondifferentiable symmetric duality. *Bull. Aust. Math. Soc.* **1996**, *53*, 177–188. [\[CrossRef\]](#)
26. Gulati, T.R.; Gupta, S.K. Higher-order symmetric duality with cone constraints. *Appl. Math. Lett.* **2009**, *22*, 776–781. [\[CrossRef\]](#)
27. Gulati, T.R.; Gupta, S.K.; Ahmad, I. Second-order symmetric duality with cone constraints. *J. Comput. Appl. Math.* **2008**, *220*, 347–354. [\[CrossRef\]](#)
28. Gulati, T.R.; Ahmad, I.; Husain, I. Second-order symmetric duality with generalized convexity. *Opsearch* **2001**, *38*, 210–222. [\[CrossRef\]](#)
29. Chandra, S.; Goyal, A.; Husain, I. On symmetric duality in mathematical programming with F-convexity. *Optimization* **1998**, *43*, 1–18. [\[CrossRef\]](#)
30. Chandra, S.; Kumar, V. A note on pseudo-invexity and symmetric duality. *Eur. J. Oper. Res.* **1998**, *105*, 626–629. [\[CrossRef\]](#)



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).