

Article

The Properties of a Decile-Based Statistic to Measure Symmetry and Asymmetry

Mohammad Reza Mahmoudi ^{1,2} , Roya Nasirzadeh ², Dumitru Baleanu ^{3,4}  and Kim-Hung Pho ^{5,*} 

¹ Institute of Research and Development, Duy Tan University, Da Nang 550000, Vietnam; mohammadrezamahmoudi@duytan.edu.vn or mahmoudi.m.r@fasau.ac.ir

² Department of Statistics, Faculty of Science, Fasa University, Fasa, Fars 7461686131, Iran; nasirzadeh.roya@fasau.ac.ir

³ Department of Mathematics, Faculty of Art and Sciences, Cankaya University, Ankara 06530, Turkey; dumitru@cankaya.edu.tr

⁴ Institute of Space Sciences, 077125 Bucharest, Romania

⁵ Fractional Calculus, Optimization and Algebra Research Group, Faculty of Mathematics and Statistics, Ton Duc Thang University, Ho Chi Minh City 72915, Vietnam

* Correspondence: phokimhung@tdtu.edu.vn

Received: 20 January 2020; Accepted: 4 February 2020; Published: 18 February 2020



Abstract: This paper studies a simple skewness measure to detect symmetry and asymmetry in samples. The statistic can be obviously applied with only three short central tendencies; i.e., the first and ninth deciles, and the median. The strength of the statistic to find symmetry and asymmetry is studied by employing numerous Monte Carlo simulations and is compared with some alternative measures by applying some simulation studies. The results show that the performance of this statistic is generally good in the simulation.

Keywords: symmetry; asymmetry; measure of skewness; decile; Monte Carlo algorithm

1. Introduction

In scientific studies, the researchers can summarize a given dataset using descriptive statistics. The descriptive statistics contain three known tendencies: central tendencies, dispersion tendencies and shape tendencies [1]. The central and dispersion tendencies, such as mean, median, standard deviation and variance deal with the convenience of the dataset [1–5]. The shape tendencies, such as skewness and kurtosis, are related to the distribution of dataset [6–8]. These measures which may be utilized in divergent disciplines consist of the tests of normality and of the lustiness for normal theoretical procedures. Skewness is often utilized to reference to symmetry. Nevertheless, symmetry is not often perspicuously defined, and it is thought that everybody knows it. There are some definitions about symmetry relying on the disciplines that it is utilized in. In literature, any statement related to the symmetry of a structure has to be done with reference to some rules of symmetry—a score, a line or an axis [9]. In the statistical inference, the meaningful score or axis is taken as the center of a distribution. There are several measures employed to quantify the degree of skewness of a distribution. Assume that μ , m , M , σ , μ_3 , Q_1 and Q_3 , are the mean; median; mode; standard deviation; third centered moment; and the first and the third quartiles, respectively. The statistics introduced for measuring the skewness are Pearson's coefficient of skewness:

$$SK_P = \frac{\mu - M}{\sigma} \quad (1)$$

Pearson's second coefficient of skewness:

$$SK_{P2} = \frac{3(\mu - m)}{\sigma} \quad (2)$$

Yule's coefficient of skewness:

$$SK_Y = \frac{(\mu - m)}{\sigma} \quad (3)$$

the standardized third central moment:

$$\gamma_1 = \frac{\mu_3}{\sigma^3} \quad (4)$$

Bowley's coefficient of skewness:

$$SK_B = \frac{Q_3 + Q_1 - 2m}{Q_3 - Q_1} \quad (5)$$

and three Galip's coefficients of skewness:

$$SK_{G1} = \frac{X_{Max} + X_{min} - 2M}{X_{Max} - X_{min}} \quad (6)$$

$$SK_{G2} = \frac{X_{Max} + X_{min} - 2m}{X_{Max} - X_{min}} \quad (7)$$

$$SK_{G3} = \frac{X_{Max} + X_{min} - 2\mu}{X_{Max} - X_{min}} \quad (8)$$

[9–17].

Although there are numerous different measures, and practical elongations of the above coefficients were proposed afterward, the original measures are still employed to this day, especially γ_1 (or its variants). It is largely utilized in statistical calculation software.

When we face a dataset containing outliers, we need a measure that can carefully consider these outliers. Therefore, probably, the measures that are based on the extreme values (max and min) such as three Galip's coefficients of skewness; are based on the first and the last quartiles (Q_1 and Q_3) such as Bowley's coefficient of skewness; or are based on the first and the last deciles (D_1 and D_9), should be more effective than other methods. The previous studies indicated that the three Galip's coefficients of skewness had the most power to detect symmetry and asymmetry. But the Bowley's coefficient of skewness acted not so well. There is no deep study about the definition of skewness based on deciles and the comparison between them and other alternatives.

In this work, at first, we consider the definition of skewness based on deciles and then study its asymptotic properties, similar to the approach that was applied in [18–23]. Finally, the power of the considered statistic to detect symmetry and asymmetry is compared with the powers of other measures of skewness.

2. Decile-Based Skewness

Let $X_1, \dots, X_n$ be a sample from a distribution F on the real line, and we suppose that F is continuous so that all observations are distinct with probability one. We may then arrange the observations in increasing order without ties, $X_{(1)} < \dots < X_{(n)}$. These variables are called the order statistics, where $X_{(k)}$ is the k^{th} order statistic. For $0 < p < 1$, the p^{th} quantile of F is defined as $x_p = F^{-1}(p)$ and the corresponding sample quantile is defined as $X_{(k)}$ where $k = \lceil np \rceil$, the ceiling of (the smallest integer greater than or equal to np). Let D_1 and D_9 be the first and nine sample deciles (0.1 and 0.9 quantiles), respectively. We consider our statistic for measuring the skewness by

$$SK = \frac{(D_9 - m) - (m - D_1)}{D_9 - D_1} \quad (9)$$

In the following, the asymptotic distribution of the proposed statistic is explored.

Lemma 1. Let $U_1, \dots, U_n$ be independent, identically distributed (iid in short) random variables from $U(0, 1)$ and $U_{(1)} < \dots < U_{(n)}$, which are order statistics of $U_1, \dots, U_n$. If $n \rightarrow \infty$, then

$$\sqrt{n} \begin{pmatrix} U_{[np_1]} - p_1 \\ U_{[np_2]} - p_2 \\ U_{[np_3]} - p_3 \end{pmatrix} \xrightarrow{D} N(\mathbf{0}, \Sigma) \quad (10)$$

where $0 < p_1 < p_2 < p_3 < 1$, and

$$\Sigma = \begin{bmatrix} p_1(1-p_1) & p_1(1-p_2) & p_1(1-p_3) \\ p_1(1-p_2) & p_2(1-p_2) & p_2(1-p_3) \\ p_1(1-p_3) & p_2(1-p_3) & p_3(1-p_3) \end{bmatrix} \quad (11)$$

Proof. Assume that $Y_1, Y_2, \dots$ are iid exponential variables with mean 1 and $S_j = \sum_{i=1}^j Y_i$. Additionally, assume that $\sqrt{n}(\frac{k_1}{n} - p_1) \rightarrow 0$, $\sqrt{n}(\frac{k_2}{n} - p_2) \rightarrow 0$ and $\sqrt{n}(\frac{k_3}{n} - p_3) \rightarrow 0$ as k_1, k_2, k_3 , and $n \rightarrow \infty$. Then by the extension of the results given in [24],

$$\sqrt{n+1} \begin{bmatrix} \frac{1}{n+1} S_{k_1} - p_1 \\ \frac{1}{n+1} (S_{k_2} - S_{k_1}) - (p_2 - p_1) \\ \frac{1}{n+1} (S_{k_3} - S_{k_2}) - (p_3 - p_2) \\ \frac{1}{n+1} (S_{n+1} - S_{k_3}) - (1 - p_3) \end{bmatrix} \xrightarrow{D} N(\mathbf{0}, \Sigma_1),$$

such that

$$\Sigma_1 = \begin{bmatrix} p_1 & 0 & 0 & 0 \\ 0 & p_2 - p_1 & 0 & 0 \\ 0 & 0 & p_3 - p_2 & 0 \\ 0 & 0 & 0 & 1 - p_3 \end{bmatrix}$$

Take $g(x_1, x_2, x_3, x_4) = \frac{1}{x_1 + x_2 + x_3 + x_4} [x_1, x_1 + x_2, x_1 + x_2 + x_3]'$; then, by Cramer's theorem [24],

$$\sqrt{n} \begin{pmatrix} \frac{S_{k_1}}{S_{n+1}} - p_1 \\ \frac{S_{k_2}}{S_{n+1}} - p_2 \\ \frac{S_{k_3}}{S_{n+1}} - p_3 \end{pmatrix} \xrightarrow{D} N(\mathbf{0}, \Sigma)$$

Finally, the proof is completed with the reality that the distribution of $\left(\frac{S_{k_1}}{S_{n+1}}, \frac{S_{k_2}}{S_{n+1}}, \frac{S_{k_3}}{S_{n+1}}\right)'$ given S_{n+1} is the same as the distribution of $(U_{(k_1)}, U_{(k_2)}, U_{(k_3)})'$. $\square$

Corollary 1. Let $X_1, \dots, X_n$ be iid random variables with density and distribution functions f and F , respectively. Additionally, assume that $f(x)$ is continuous and positive in a neighborhood of the quantiles x_{p_1}, x_{p_2} and x_{p_3} with $p_1 < p_2 < p_3$; then,

$$\sqrt{n} \begin{pmatrix} X_{[np_1]} - x_{p_1} \\ X_{[np_2]} - x_{p_2} \\ X_{[np_3]} - x_{p_3} \end{pmatrix} \xrightarrow{D} N(\mathbf{0}, \Sigma^*) \quad (12)$$

where

$$\Sigma^* = \begin{bmatrix} \frac{p_1(1-p_1)}{f^2(x_{p_1})} & \frac{p_1(1-p_2)}{f(x_{p_1})f(x_{p_2})} & \frac{p_1(1-p_3)}{f(x_{p_1})f(x_{p_3})} \\ \frac{p_1(1-p_2)}{f(x_{p_1})f(x_{p_2})} & \frac{p_2(1-p_2)}{f^2(x_{p_2})} & \frac{p_2(1-p_3)}{f(x_{p_2})f(x_{p_3})} \\ \frac{p_1(1-p_3)}{f(x_{p_1})f(x_{p_3})} & \frac{p_2(1-p_3)}{f(x_{p_2})f(x_{p_3})} & \frac{p_3(1-p_3)}{f^2(x_{p_3})} \end{bmatrix} \quad (13)$$

Proof. By applying the transformation $g(y_1, y_2, y_3) = (F^{-1}(y_1), F^{-1}(y_2), F^{-1}(y_3))'$ to the variables $(U_{[np_1]} - p_1, U_{[np_2]} - p_2, U_{[np_3]} - p_3)$ in Lemma 1, the proof will be completed. Be careful that the derivation of g is

$$\dot{g}(y_1, y_2, y_3) = \begin{bmatrix} \frac{1}{f(F^{-1}(y_1))} & 0 & 0 \\ 0 & \frac{1}{f(F^{-1}(y_2))} & 0 \\ 0 & 0 & \frac{1}{f(F^{-1}(y_3))} \end{bmatrix}.$$

□

The asymptotic distribution of SK is provided in the following theorem. This is our major contribution. It is also necessary to infer the skewness of population.

Theorem 1. Let $X_1, \dots, X_n$ be iid random variables with density function f . Additionally, assume that $f(x)$ is continuous and positive in a neighborhood of the quantiles $x_{0.1}, x_{0.5}$ and $x_{0.9}$. Then, the asymptotic distribution of the proposed statistic can be illustrated by

$$T_n = \sqrt{n} \left(SK - \frac{x_{0.9} + x_{0.1} - 2x_{0.5}}{x_{0.9} - x_{0.1}} \right) \xrightarrow{D} N(0, \sigma^2)$$

where

$$\sigma^2 = \frac{1}{(x_{0.9} - x_{0.1})^4} \left[\frac{0.36(x_{0.9} - x_{0.5})^2}{f^2(x_{0.1})} + \frac{(x_{0.9} - x_{0.1})^2}{f^2(x_{0.5})} + \frac{0.36(x_{0.5} - x_{0.1})^2}{f^2(x_{0.9})} - \frac{0.4(x_{0.9} - x_{0.1})(x_{0.9} - x_{0.5})}{f(x_{0.1})f(x_{0.5})} + \frac{0.08(x_{0.5} - x_{0.1})(x_{0.9} - x_{0.5})}{f(x_{0.1})f(x_{0.9})} - \frac{0.4(x_{0.9} - x_{0.1})(x_{0.5} - x_{0.1})}{f(x_{0.5})f(x_{0.9})} \right] \quad (14)$$

Proof. The proof is simply achieved using Cramer's theorem [24] and taking $g(x_1, x_2, x_3) = \frac{x_1 - 2x_2 + x_3}{x_3 - x_1}$. □

Corollary 2. Let $X_1, \dots, X_n$ be iid random variables from $U(0, 1)$; then, the asymptotic distribution of the proposed statistic is given by

$$\sqrt{n}(SK - 0) \xrightarrow{D} N(0, 1.25) \quad (15)$$

These results can be employed to build an asymptotical confidence interval and to check the hypothesis.

2.1. Asymptotic Confidence Interval

Now, T_n can be utilized as a pivotal quantity to build a confidence interval asymptotic to a population's skewness,

$$\left(SK - \frac{\hat{\sigma}}{\sqrt{n}} Z_{\alpha/2}, SK + \frac{\hat{\sigma}}{\sqrt{n}} Z_{\alpha/2} \right) \quad (16)$$

where

$$\hat{\sigma}^2 = \frac{1}{(D_9 - D_1)^4} \left[\frac{0.36(D_9 - m)^2}{f^2(D_1)} + \frac{(D_9 - D_1)^2}{f^2(m)} + \frac{0.36(m - D_1)^2}{f^2(D_9)} - \frac{0.4(D_9 - D_1)(D_9 - m)}{f(D_1)f(m)} + \frac{0.08(m - D_1)(D_9 - m)}{f(D_1)f(D_9)} - \frac{0.4(D_9 - D_1)(m - D_1)}{f(m)f(D_9)} \right] \quad (17)$$

2.2. Hypothesis Testing

Hypothesis testing related to *skewness* is a crucial issue in practical application. For instance, the assumption *Skewness* = 0 is tantamount to the symmetry. Generally, to test $H_0 : \text{Skewness} = \gamma_0$, the test statistic can be

$$T_0 = \sqrt{n} \left(\frac{SK - \gamma_0}{\hat{\sigma}} \right) \quad (18)$$

Similar to the methodology provided in Theorem 1, it can prove that with the null hypothesis, T_0 has, asymptotically, standard normal distribution.

3. Asymptotic Properties of the Proposed Statistic

In this part, many data sets are drawn to analyze the performance of the proposed approach, for distinct symmetric distributions and divergent sample sizes. Firstly, we checked that the given CI and test statistic are truly the asymptotic CI and test statistic. For every parameter, the experiential coverage probability (percentage of runs for which the given CI contains zero (true skewness)) was calculated by relying on 10,000 repetitions using *statistical R 3.6.2* and *SPSS 25* software. In addition, for each repetition, the value of the given test statistic is presented and normal Q–Q plots of the given test statistic are provided. The Shapiro-Wilk's normality test is used to confirm the normality of the given test statistic. The experiential coverage probabilities for divergent parameters are illustrated as in Table 1.

Table 1. The experiential coverage probability of the proposed confidence interval.

Distribution	<i>n</i>					
	50	75	100	200	500	1000
Normal (1,5)	0.9732	0.9734	0.9743	0.9744	0.975	0.9761
t(10)	0.9916	0.9928	0.9934	0.9939	0.9942	0.9947
U(0,1)	0.9485	0.949	0.9491	0.9502	0.9521	0.9569

The results show that the experiential coverage probability of proposed approach is more than nominal level (0.95), especially when the sample sizes grow. In the other hand, we can admit the given CI as the asymptotic CI for the skewness of population. Figure 1 and Table 2 show the Q–Q plots for the standard normal distribution and the results of Shapiro-Wilk's normality test in the test statistic, respectively.

Table 2. Shapiro-Wilk's normality test *p*-value for the given test statistic.

Distribution	<i>n</i>					
	50	75	100	200	500	1000
Normal (1,5)	0.7131	0.7174	0.7899	0.8436	0.9077	0.9213
t(10)	0.433	0.6515	0.781	0.8317	0.9603	0.9945
U(0,1)	0.3144	0.5566	0.6034	0.6219	0.8249	0.9488




Figure 1. The Q–Q plots versus standard normal distribution. Normal distribution: $n = 50$ (a), $n = 1000$ (b). t distribution: $n = 50$ (c), $n = 1000$ (d). Uniform distribution: $n = 50$ (e), $n = 1000$ (f).

It can be then seen that the asymptotic properties are relatively satisfied in all situations (p-value is greater than 5%). Thereafter, it can be seen that our approach is a good choice to build a CI and execute hypothesis testing for the skewness of a population.

4. Comparison with Alternative Measures

To check the performances of the considered statistic, its power to detect asymmetry is compared with the conventional measures of skewness by employing a Monte Carlo simulation. As in Section 3, numerous data sets were drawn to check the performances of the measures, for different asymmetric distributions and different sample sizes using R software. For this purpose, we generated 10,000 samples of size $n = 10, 20, 50$, from a chi-square distribution with m degrees of freedom, $(\chi^2(m))$. We considered three cases: extremely skewed ($m = 1$), moderately skewed ($m = 5$) and slightly skewed ($m = 40$). The powers (at 5% significant level) of different measures to detect asymmetry are summarized in Table 3.

Table 3. The powers of different measures to detect skewness.

Distribution	Measure	<i>n</i>		
		10	20	50
Extremely Skewed	SK	0.798	0.989	1.000
	γ_1	0.687	0.942	1.000
	SK_{G3}	0.817	0.991	1.000
	SK_{G2}	0.834	0.992	1.000
	SK_{G1}	0.461	0.831	0.997
	SK_P	0.200	0.151	0.130
	SK_Y	0.616	0.869	0.999
	SK_{P2}	0.616	0.869	0.999
	SK_B	0.260	0.403	0.711
Moderately Skewed	SK	0.318	0.597	0.945
	γ_1	0.297	0.530	0.889
	SK_{G3}	0.321	0.564	0.911
	SK_{G2}	0.318	0.623	0.941
	SK_{G1}	0.145	0.397	0.814
	SK_P	0.132	0.108	0.100
	SK_Y	0.207	0.344	0.651
	SK_{P2}	0.207	0.344	0.651
	SK_B	0.123	0.156	0.224
Slightly Skewed	SK	0.144	0.163	0.289
	γ_1	0.129	0.165	0.288
	SK_{G3}	0.135	0.175	0.284
	SK_{G2}	0.143	0.153	0.282
	SK_{G1}	0.116	0.136	0.252
	SK_P	0.106	0.103	0.119
	SK_Y	0.120	0.116	0.180
	SK_{P2}	0.120	0.116	0.180
	SK_B	0.117	0.116	0.135

As preliminary results, based on the maximum power, it can be observed that the performances of SK , γ_1 , SK_{G1} , SK_{G2} and SK_{G3} are approximately similar and are more powerful than other methods for all simulated datasets, and are therefore are very promising. The performances of SK_P , SK_{P2} and SK_Y are approximately similar and have the next best ranks, while SK_B has the worst performance in all situations. In general, the measures that are based on the extreme values (maximum and minimum), such as three Galip's coefficients of skewness, and those based on the first and the last deciles (D_1 and D_9), are more effective than other methods, because of their better performances and easy calculations.

5. Discussion

In this work, at first, we considered the definition of skewness based on deciles, and then studied its asymptotic properties. The results showed that the experiential coverage probability of this measure was more than nominal level (0.95), especially when the sample size was increased. The Q-Q plots versus the standard normal distribution and the results of Shapiro-Wilk's normality test verified the

theoretical asymptotic properties. Finally, the power of the considered statistic to detect symmetry and asymmetry was compared with the powers of other measures of skewness. The power study indicated that the performances of decile-based measure and three Galip's coefficients of skewness were approximately similar, and were more powerful than other methods for all simulated datasets, and are therefore promising for application in practice.

6. Conclusions

We presented a simple measure to find skewness in patterns. The new measure relies on a new definition of skewness that contains many outstanding advantages. The proposed coefficient of skewness could be obviously calculated with only three short statistics; i.e., the first and nine deciles and the median. The strength of the proposed statistic to find symmetry and asymmetry was studied by employing numerous Monte Carlo simulations. The results show that the performance of new statistic is generally very good in the simulation. There are many definitions to describe symmetry and asymmetry. To investigate the skewness in datasets including outliers, we should use the measures that consider the effects of outliers. Therefore, probably, the measures that are based on the extreme values (maximum and minimum), such as three Galip's coefficients of skewness; those based on the first and the last quartiles (Q_1 and Q_3), such as Bowley's coefficient of skewness; and those based on the first and the last deciles (D_1 and D_9), are candidates for application. Other studies showed that Galip's coefficients of skewness are more powerful for detecting symmetry and asymmetry. There is no deep study about the definition of skewness based on deciles and a comparison between them and other alternatives. In this work, at first, we considered the definition of skewness based on deciles and then studied its asymptotic properties. Finally, the power of the considered statistic to detect symmetry and asymmetry was compared with the powers of other measures of skewness. For future works, we suggest readers to use a definition of skewness based on combinations of more deciles, not only the first and the ninth deciles. We think this combination will improve the detection of symmetry and asymmetry.

Author Contributions: Conceptualization, M.R.M., R.N., D.B. and K.-H.P.; data curation, M.R.M.; formal analysis, M.R.M., R.N. and K.-H.P.; investigation, M.R.M., R.N. and D.B.; methodology, M.R.M. and K.-H.P.; project administration, D.B.; supervision, M.R.M.; validation, M.R.M.; visualization, M.R.M.; writing—original draft, M.R.M. and R.N.; writing—review and editing, D.B. and K.-H.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Sprinthal, R.C.; Fisk, S.T. *Basic Statistical Analysis*; Prentice Hall: Englewood Cliffs, NJ, USA, 1990.
2. Manikandan, S. Measures of central tendency: Median and mode. *J. Pharmacol. Pharm.* **2011**, *2*, 214. [[CrossRef](#)]
3. Weisberg, H.; Weisberg, H.F. *Central Tendency and Variability*; Sage: Thousand Oaks, CA, USA, 1992.
4. Deshpande, S.; Gogtay, N.J.; Thatte, U.M. Measures of central tendency and dispersion. *J. Assoc. Physicians India* **2016**, *64*, 64–66.
5. Manikandan, S. Measures of dispersion. *J. Pharmacol. Pharm.* **2011**, *2*, 315. [[CrossRef](#)]
6. Kim, T.H.; White, H. On more robust estimation of skewness and kurtosis. *Financ. Res. Lett.* **2004**, *1*, 56–73. [[CrossRef](#)]
7. Oja, H. On location, scale, skewness and kurtosis of univariate distributions. *Scand J. Stat.* **1981**, *1*, 154–168.
8. Wilkins, J.E. A note on skewness and kurtosis. *Ann. Math Stat.* **1944**, *15*, 333–335. [[CrossRef](#)]
9. Murphy, E.A. Skewness and asymmetry of distributions. *Metamedicine* **1982**, *3*, 87–99. [[CrossRef](#)]
10. Arnold, B.C.; Groeneveld, R.A. Skewness and kurtosis orderings: An introduction. *Lect. Notes Monogr. Ser.* **1992**, *22*, 17–24.
11. Arnold, B.C.; Groeneveld, R.A. Measuring skewness with respect to the mode. *Am. Stat.* **1995**, *49*, 34–38.

12. Doane, D.P.; Seward, L.E. Measuring Skewness: A Forgotten Statistic? *J. Stat. Educ.* **2011**, *19*, 1–18. [[CrossRef](#)]
13. García, V.J.; Martel, M.; Vázquez-Polo, F.J. Complementary information for skewness measures. *Stat. Neerl* **2015**, *69*, 442–459. [[CrossRef](#)]
14. Groeneveld, R.A.; Meeden, G. Measuring skewness and kurtosis. *Statistician* **1984**, *33*, 391–399. [[CrossRef](#)]
15. Mahmoudi, M.R.; Nasirzadeh, R.; Mohammadi, M. On the Ratio of Two Independent Skewnesses. *Commun. Stat. Theory Methods* **2019**, *48*, 1721–1727. [[CrossRef](#)]
16. Tabor, J. Investigating the investigative task: Testing for skewness—An investigation of different test statistics and their power to detect skewness. *J. Stat. Educ.* **2010**, *18*, 1–13. [[CrossRef](#)]
17. Tajuddin, I.H. A simple measure of skewness. *Stat. Neerl* **1996**, *50*, 362–366. [[CrossRef](#)]
18. Haghbin, H.; Mahmoudi, M.R.; Shishebor, Z. Large Sample Inference on the Ratio of Two Independent Binomial Proportions. *J. Math. Ext.* **2011**, *5*, 87–95.
19. Mahmoudi, M.R.; Mahmoodi, M. Inference on the Ratio of Variances of Two Independent Populations. *J. Math. Ext.* **2014**, *7*, 83–91.
20. Mahmoudi, M.R.; Mahmoodi, M. Inference on the Ratio of Correlations of Two Independent Populations. *J. Math. Ext.* **2014**, *7*, 71–82.
21. Mahmoudi, M.R.; Maleki, M.; Pak, A. Testing the Difference between Two Independent Time Series Models. *Iran. J. Sci. Technol. Trans. A Sci.* **2017**, *41*, 665–669. [[CrossRef](#)]
22. Mahmoudi, M.R.; Mahmoudi, M.; Nahavandi, E. Testing the Difference between Two Independent Regression Models. *Commun. Stat. Theory Methods* **2016**, *45*, 6284–6289. [[CrossRef](#)]
23. Mahmoudi, M.R.; Behboodian, J.; Maleki, M. Large Sample Inference about the Ratio of Means in Two Independent Populations. *J. Stat. Theory Appl.* **2017**, *16*, 366–374. [[CrossRef](#)]
24. Ferguson, T.S. *A Course in Large Sample Theory*; Chapman & Hall: London, UK, 1996.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).