

## Article

# On the Suppression of the Dark Matter-Nucleon Scattering Cross Section in the $SE_6SSM$

Roman Nevzorov 

I.E.Tamm Theory Department, P.N. Lebedev Physical Institute of the Russian Academy of Sciences,  
53 Leninskiy Prospekt, 119991 Moscow, Russia; nevzorovrb@lebedev.ru

**Abstract:** In the  $E_6$  inspired  $U(1)_N$  extension of the minimal supersymmetric (SUSY) standard model (MSSM), a single discrete  $\tilde{Z}_2^H$  symmetry permits suppressing rapid proton decay and non-diagonal flavor transitions. If matter parity and  $\tilde{Z}_2^H$  symmetry are preserved in this SUSY model ( $SE_6SSM$ ), it may involve two dark matter candidates. In this article, we study a new modification of the  $SE_6SSM$  in which the cold dark matter is composed of gravitino and the lightest neutral exotic fermion. We argue that, in this case, the dark matter nucleon scattering cross-section can be considerably smaller than the present experimental limit.

**Keywords:** unified field theories and models; models beyond the standard model; supersymmetry; cold dark matter



**Citation:** Nevzorov, R. On the Suppression of the Dark Matter-Nucleon Scattering Cross Section in the  $SE_6SSM$ . *Symmetry* **2022**, *14*, 2090. <https://doi.org/10.3390/sym14102090>

Academic Editors: Giuseppe Latino and Jaime Haro Cases

Received: 29 August 2022

Accepted: 7 October 2022

Published: 8 October 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The possible unification of all interactions remains one of the most appealing motivations for studying supersymmetric (SUSY) extensions of the Standard Model (SM). At very high energies, the minimal supersymmetric standard model (MSSM) and its extensions can be embedded into Grand Unified Theories (GUTs). In  $N = 2$  SUSY GUTs based on the  $E_8$  gauge group, all SM bosons and fermions can belong to a single 248 representation of  $E_8$ . This representation involves three  $27 \oplus \overline{27}$ , which are fundamental and anti-fundamental representations of  $E_6$ , adjoint 78 representation of  $E_6$  and 8 components that do not participate in the  $E_6$  gauge interactions. In SUSY GUTs, three fundamental representations of  $E_6$  may include Higgs doublet and three families of the SM fermions. The SM gauge bosons are contained in the adjoint representation of  $E_6$ . It is expected that near the scale  $M_0 \gtrsim M_X \sim 2 \cdot 10^{16}$  GeV the extended SUSY is broken down to  $N = 1$  supersymmetry while the breakdown of  $E_6$  (or  $E_8$ ) gauge group can lead to

$$E_6 \rightarrow SO(10) \times U(1)_\psi \rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \rightarrow SU(3)_C \times SU(2)_W \times U(1)_Y \times U(1)_\psi \times U(1)_\chi, \quad (1)$$

where  $SU(3)_C \times SU(2)_W \times U(1)_Y$  is a SM gauge group.

The exceptional supersymmetric standard model ( $E_6SSM$ ) [1,2] (for recent review, see Ref. [3]) implies that around the GUT scale  $M_X$  rank-6 model with two extra symmetries  $U(1)_\psi$  and  $U(1)_\chi$  is reduced further to an effective rank-5 model based on the SM gauge group together with an additional  $U(1)_N$  factor, i.e.,

$$U(1)_\psi \times U(1)_\chi \rightarrow U(1)_N \times Z_2^M, \quad (2)$$

where  $Z_2^M = (-1)^{3(B-L)}$  is the so-called matter parity, which is a discrete subgroup of  $U(1)_\psi$  and  $U(1)_\chi$ . In Equation (2) the  $U(1)_N$  gauge symmetry is defined as

$$U(1)_N = \frac{1}{4}U(1)_\chi + \frac{\sqrt{15}}{4}U(1)_\psi. \quad (3)$$

In these  $E_6$  inspired  $U(1)_N$  extensions of the MSSM the right-handed neutrinos do not participate in the gauge interactions and may be superheavy [1,2]. Therefore, a see-saw mechanism can be used to shed light on the origin of the mass hierarchy in the lepton sector, providing a comprehensive understanding of the neutrino oscillations data. The successful leptogenesis is a distinctive feature of the  $U(1)_N$  extensions of the MSSM because the heavy Majorana right-handed neutrinos may decay into final states with lepton number  $L = \pm 1$ , creating a lepton asymmetry in the early Universe [4,5].

In the  $E_6$  inspired  $U(1)_N$  extensions of the MSSM gauge anomalies become cancelled if the particle spectrum at low-energies contains complete representations of  $E_6$ . To ensure that the  $E_6$ SSM is anomaly-free one is forced to extend the minimal matter content by extra matter beyond the MSSM which, together with ordinary SM fermions, form three complete 27-plets of  $E_6$  ( $27_i$  with  $i = 1, 2, 3$ ). Each 27-dimensional representations of  $E_6$  involves one generation of ordinary matter, a SM singlet field  $S_i$ , that carries non-zero  $U(1)_N$  charge, up- and down-type Higgs doublets  $H_i^u$  and  $H_i^d$ , as well as charged  $\pm 1/3$  exotic quarks  $D_i$  and  $\bar{D}_i$ . The presence of extra exotic matter may give rise to non-diagonal flavor transitions and rapid proton decay. In the  $E_6$ SSM a set of discrete symmetries can be used to suppress the corresponding operators [1,2].

In this article, we examine the dark matter-nucleon scattering cross section within the modification of the  $E_6$ SSM ( $SE_6$ SSM) [6,7] in which a single discrete  $\tilde{Z}_2^H$  symmetry forbids tree-level flavor-changing transitions, as well as the most dangerous baryon and lepton number violating operators. The conservation of  $\tilde{Z}_2^H$  symmetry and matter parity/ $R$ -parity implies the existence of at least two stable states that may account for all or some of the observed cold dark matter density. At the same time null results from direct detection experiments [8–10] placed stringent limits on the dark matter nucleon scattering cross-section. Here, we explore a new variant of the  $SE_6$ SSM in which two stable states are gravitino and the lightest neutral exotic fermion. In general, the spin-independent part of the dark matter-nucleon scattering cross section can be much larger in this case than the corresponding experimental limit. Nevertheless, the results of our analysis indicate that there is a part of the  $SE_6$ SSM parameter space, in which this cross section is sufficiently strongly suppressed.

The paper is organised as follows. In the next section, we specify a new variant of the  $SE_6$ SSM which is considered in this article. In Section 3, we investigate the dependence of the dark matter-nucleon scattering cross section on the parameters of the  $SE_6$ SSM. Section 4 concludes the paper.

## 2. The $SE_6$ SSM

Since 2006, several modifications of the  $E_6$ SSM have been explored [1,2,6,7,11–19]. The implications of the  $U(1)_N$  extensions of the MSSM were considered for  $Z$ – $Z'$  mixing [20], neutralino sector [20–22], electroweak (EW) symmetry breaking (EWSB) [21,23,24], the renormalization group (RG) flow of couplings [21,25], the renormalization of vacuum expectation values (VEVs) [26,27], non-standard neutrino models [28], and dark matter [19,29,30]. Within the  $E_6$ SSM, the upper bound on the lightest Higgs mass near the quasi-fixed point was examined in [31]. The corresponding quasi-fixed point is an intersection of the invariant and quasi-fixed lines [32,33]. The particle spectrum in the constrained  $E_6$ SSM ( $CE_6$ SSM) and its modifications was analyzed in [34–37]. The degree of fine tuning and threshold corrections were explored in Refs. [38,39] and [40], respectively. Extra exotic matter in the  $E_6$ SSM may result in distinctive LHC signatures [1,2,12,15,41–44] and can give rise to non-standard Higgs decays [7,45,46].

The  $SE_6$ SSM implies that below the scale  $M_X$  three complete 27-plets are accompanied by SM singlet superfield  $\phi$ , which does not participate in the  $E_6$  interactions, and a set of pairs of supermultiplets  $M_l$  and  $\bar{M}_l$ , which belong to additional  $27_l'$  and  $\bar{27}_l'$  representations, respectively. Because  $M_l$  and  $\bar{M}_l$  carry opposite  $SU(3)_C \times SU(2)_W \times U(1)_Y \times U(1)_N$  quantum numbers gauge anomalies still cancel. In the simplest case, the set of  $M_l$  and  $\bar{M}_l$  involves three pairs of  $SU(2)_W$  doublets, i.e.,  $L_4$  and  $\bar{L}_4$ ,  $H_u$  and  $\bar{H}_u$ ,  $H_d$  and  $\bar{H}_d$ , as well as

a pair of superfields,  $S$  and  $\bar{S}$ . The field content of the  $SE_6$ SSM may originate from the  $E_6$  orbifold GUT model in six dimensions in which the appropriate splitting of the bulk  $27'$  supermultiplets can be achieved [6].

The supermultiplets  $\phi$ ,  $S$ ,  $\bar{S}$ ,  $H_u$ ,  $H_d$ ,  $L_4$ , and  $\bar{L}_4$  are required to be even under the  $\tilde{Z}_2^H$  custodial symmetry whereas all other supermultiplets are odd [7]. In the  $SE_6$ SSM superpotential, the  $\tilde{Z}_2^H$  symmetry allows the interactions that comes from  $27'_l \times 27'_m \times 27'_n$  and  $27'_l \times 27_i \times 27_k$  but it forbids all terms which originate from  $27_i \times 27_j \times 27_k$ , where indexes  $l, m, n$  are associated with the supermultiplets  $M_l$  while family indexes  $i, j, k = 1, 2, 3$  run over three generations. Such structure of the superpotential ensures that it does not involve any operators that give rise to rapid proton decay. Because the set of supermultiplets  $M_l$  includes only two Higgs doublets  $H_d$  and  $H_u$ , the up-type quarks couple to just  $H_u$ , whereas the down-type quarks and charged leptons couple to  $H_d$  only. Therefore, the flavor-changing processes are suppressed in the  $SE_6$ SSM at tree-level.

In the simplest scenario  $\bar{H}_u$  and  $\bar{H}_d$  become combined with the superposition of the corresponding components from the  $27_i$ , forming vector-like states with masses of order  $M_X$ . The components of the supermultiplets  $L_4$  and  $\bar{L}_4$ , as well as  $S$  and  $\bar{S}$  are expected to gain the TeV scale masses. The presence of  $L_4$  and  $\bar{L}_4$  at low energies permits the lightest exotic quarks to decay within a reasonable time, as well as facilitates the gauge coupling unification [25] and the generation of the baryon asymmetry of the Universe [5].

Using the method discussed in [47], it was revealed that in the  $E_6$ SSM and its simplest modifications the lightest  $R$ -parity odd states (lightest SUSY particles) have to be lighter than 60–65 GeV [46]. These states are predominantly linear superpositions of the fermion components of the superfields  $S_i$ . Although the couplings of these lightest exotic fermions to the SM particles are very small the lightest SUSY particle (LSP) could account for some of the cold dark matter relic density if the LSP had a mass close to half the  $Z$  boson mass  $M_Z/2$  [46]. However, in this case the SM-like Higgs boson decays mostly into the lightest exotic fermions whereas its other branching ratios are suppressed. LHC experiments have already excluded such a scenario. When the lightest exotic fermions are substantially lighter than  $M_Z$ , the LSP annihilation cross section tends to be too small resulting in too large cold dark matter density. In the simplest phenomenologically viable scenario, the sparticle spectrum of the  $SE_6$ SSM includes the lightest SUSY particles with masses which are much smaller than 1 eV. These lightest exotic fermions form hot dark matter in our Universe. Nevertheless, they give a minor contribution to the total density of dark matter. The existence of neutral fermions with tiny masses may lead to very interesting implications for the neutrino physics [48].

The variant of the  $SE_6$ SSM explored here implies that in addition to all states mentioned above the low-energy matter content of the model involves at least three  $E_6$  singlet superfields  $\phi_i$  which are odd under the  $\tilde{Z}_2^H$  symmetry. This allows to avoid the appearance of the lightest exotic fermions with tiny masses in the particle spectrum. It is expected that the components of the supermultiplets

$$(Q_i, u_i^c, d_i^c, L_i, e_i^c) + (D_i, \bar{D}_i) + S_i + \phi_i + (H_\alpha^u, H_\alpha^d) + L_4 + \bar{L}_4 + S + \bar{S} + H_u + H_d + \phi, \quad (4)$$

where  $\alpha = 1, 2$  and  $i = 1, 2, 3$ , have masses either of the order of 10 TeV or considerably lower. The right-handed neutrino superfields  $N_i^c$  are assumed to be much heavier than 10 TeV. The  $U(1)_Y$  and  $U(1)_N$  charges of all matter supermultiplets in the  $SE_6$ SSM are summarised in Table 1.

**Table 1.** The  $U(1)_Y$  and  $U(1)_N$  charges of matter supermultiplets in the  $SE_6$ SSM. The superfields  $N_i^c$ ,  $\phi_i$ , and  $\phi$  have zero  $U(1)_Y$  and  $U(1)_N$  charges.

	$Q_i$	$u_i^c$	$d_i^c$	$L_i, L_4$	$e_i^c$	$S_i, S$	$H_\alpha^u, H_u$	$H_\alpha^d, H_d$	$D_i$	$\bar{D}_i$	$\bar{L}_4$	$\bar{S}$
$\sqrt{\frac{5}{3}}Q_i^Y$	$\frac{1}{6}$	$-\frac{2}{3}$	$\frac{1}{3}$	$-\frac{1}{2}$	1	0	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{2}$	0
$\sqrt{40}Q_i^N$	1	1	2	2	1	5	−2	−3	−2	−3	−2	−5

Integrating out the right-handed neutrino superfields  $N_i^c$  and neglecting all suppressed non-renormalisable interactions, the low-energy superpotential of the new variant of the SE<sub>6</sub>SSM can be written as

$$W_{\text{SE}_6\text{SSM}} = \lambda S(H_u H_d) - \sigma \phi \bar{S} \bar{S} + \frac{\kappa}{3} \phi^3 + \frac{\mu}{2} \phi^2 + \Lambda \phi + \mu_L L_4 \bar{L}_4 + \tilde{\sigma} \phi L_4 \bar{L}_4 + W_{IH} \\ + \kappa_{ij} S(D_i \bar{D}_j) + g_{ij}^D (Q_i L_4) \bar{D}_j + h_{ia}^E e_i^c (H_a^d L_4) + g_{ij} \phi_i \bar{L}_4 L_j + W_{\text{MSSM}}(\mu = 0). \quad (5)$$

It is worth noting that the extra  $U(1)_N$  gauge symmetry forbids the term  $\mu_0 H_d H_u$  in the superpotential (5), but allows all other terms in  $W_{\text{MSSM}}(\mu = 0)$ . In Equation (5) the part of the superpotential  $W_{IH}$  describes the interactions of  $\phi_i$ ,  $S_i$ ,  $H_\alpha^u$  and  $H_\alpha^d$  with the  $\tilde{Z}_2^H$  even supermultiplets  $\phi$ ,  $S$ ,  $\bar{S}$ ,  $H_u$ , and  $H_d$ .

$$W_{IH} = \tilde{M}_{ij} \phi_i \phi_j + \tilde{\kappa}_{ij} \phi \phi_i \phi_j + \tilde{\lambda}_{ij} \bar{S} \phi_i S_j + \lambda_{\alpha\beta} S(H_\alpha^d H_\beta^u) \\ + \tilde{f}_{ia} S_i(H_\alpha^d H_u) + f_{ia} S_i(H_d H_\alpha^u). \quad (6)$$

The sector responsible for the breakdown of the gauge symmetry in the SE<sub>6</sub>SSM is formed by the scalar components of  $\phi$ ,  $S$ ,  $\bar{S}$ ,  $H_u$ , and  $H_d$ . In the limit  $\sigma \rightarrow 0$  the  $U(1)_N$   $D$ -term contribution to the effective scalar potential can force the minimum of such potential to be along the  $D$ -flat direction [49]. Indeed, if the coupling  $\sigma$  vanishes the part of the scalar potential that depends on  $S$  and  $\bar{S}$  takes the form

$$V_S(S, \bar{S}) = m_S^2 |S|^2 + m_{\bar{S}}^2 |\bar{S}|^2 + \frac{Q_S^2 g_1'^2}{2} (|S|^2 - |\bar{S}|^2)^2, \quad (7)$$

where  $Q_S$  is the  $U(1)_N$  charge of  $S$  and  $\bar{S}$ ,  $g_1'$  is the  $U(1)_N$  gauge coupling whereas  $m_S^2$  and  $m_{\bar{S}}^2$  are the soft SUSY breaking mass parameters. The last term in Equation (7) is associated with the  $U(1)_N$   $D$ -term contribution. For  $\langle S \rangle = \langle \bar{S} \rangle$  the quartic term vanishes. If, in this case  $(m_S^2 + m_{\bar{S}}^2) < 0$ , then the scalar potential (7) has a run-away direction  $\langle S \rangle = \langle \bar{S} \rangle \rightarrow \infty$ . The  $F$ -term contribution to the scalar potential (7) induced by the small coupling  $\sigma$  stabilizes the run-away direction so that the SM singlet superfields acquire VEVs which are much larger than the sparticle mass scale  $M_S$ , i.e.,

$$\langle \phi \rangle \sim \langle S \rangle \simeq \langle \bar{S} \rangle \sim \frac{M_S}{\sigma}, \quad (8)$$

resulting in an extremely heavy  $Z'$  boson. All extra exotic states can be also rather heavy in this limit.

The conservation of  $Z_2^M$  and  $\tilde{Z}_2^H$  symmetries implies that  $R$ -parity and  $Z_2^E$  symmetry are also conserved where  $\tilde{Z}_2^H = Z_2^M \times Z_2^E$  [6]. The transformation properties of different supermultiplets under the  $\tilde{Z}_2^H$ ,  $Z_2^M$  and  $Z_2^E$  symmetries are summarized in Table 2. Here, we assume that gravitino is the lightest  $R$ -parity odd state. Therefore, it has to be stable and potentially contributes to the density of dark matter. There is a large class of models in which gravitino can be much lighter than the superpartners of other particles [50–53]. Recently, the cosmological implications of the gravitino with mass  $m_{3/2} \sim \text{KeV}$  were discussed in Ref. [54]. Since gravitino is the lightest SUSY particle with  $Z_2^E = +1$  the lightest exotic state with  $Z_2^E = -1$  must be absolutely stable as well [6].

**Table 2.** Transformation properties of different supermultiplets under the discrete symmetries  $\tilde{Z}_2^H$ ,  $Z_2^M$  and  $Z_2^E$ . The signs + and – correspond to the states which are even and odd under different  $Z_2$  symmetries.

	$Q_i, u_i^c, d_i^c, L_i, e_i^c, N_i^c$	$\bar{D}_i, D_i, H_\alpha^d, H_\alpha^u, S_i, \phi_i$	$H_d, H_u, S, \bar{S}, \phi$	$L_4, \bar{L}_4$
$\tilde{Z}_2^H$	–	–	+	+
$Z_2^M$	–	+	+	–
$Z_2^E$	+	–	+	–

The scalar components of the supermultiplets  $\phi_i$ ,  $S_i$ ,  $H_\alpha^u$ , and  $H_\alpha^d$  do not develop VEVs. Their fermion components compose the exotic chargino and neutralino states. In the limit, when all components of  $\phi_i$  are considerably heavier than the bosons and fermions from the supermultiplets  $S_i$ ,  $H_\alpha^u$ , and  $H_\alpha^d$ , the superfields  $\phi_i$  can be integrated out so that the part of the SE<sub>6</sub>SSM superpotential  $W_{IH}$  reduces to:

$$W_{IH} \rightarrow \tilde{W}_{IH} \simeq -\tilde{\mu}_{ij} S_i S_j + \lambda_{\alpha\beta} S (H_\alpha^d H_\beta^u) + \tilde{f}_{i\alpha} S_i (H_\alpha^d H_u) + f_{i\alpha} S_i (H_d H_\alpha^u) + \dots \quad (9)$$

Hereafter, we use the field basis in which  $\tilde{\mu}_{ij} = \tilde{\mu}_i \delta_{ij}$  and  $\lambda_{\alpha\beta} = \lambda_{\alpha\alpha} \delta_{\alpha\beta}$ .

In this article, we focus on the scenarios in which the lightest exotic state with  $Z_2^E = -1$  is predominantly formed by the fermion components of the supermultiplets  $H_1^u$  and  $H_1^d$  while all sparticles except gravitino and all other exotic states have masses which are much larger than 1 TeV. If  $H_1^u$  and  $H_1^d$  mostly interact with  $H_u$ ,  $H_d$  and  $S_1$ , whereas all other couplings of  $H_1^u$  and  $H_1^d$  are negligibly small, the part of the mass matrix, that determines the masses of the lightest exotic neutralino states, can be written in the following form:

$$M^{ab} = - \begin{pmatrix} 0 & \mu_{11} & \frac{\tilde{f}_{11}}{\sqrt{2}} v_2 \\ \mu_{11} & 0 & \frac{f_{11}}{\sqrt{2}} v_1 \\ \frac{\tilde{f}_{11}}{\sqrt{2}} v_2 & \frac{f_{11}}{\sqrt{2}} v_1 & \tilde{\mu}_1 \end{pmatrix}, \quad (10)$$

where  $\mu_{11} \simeq \lambda_{11} \langle S \rangle$ ,  $v_1$  and  $v_2$  are the VEVs of the Higgs doublets  $H_d$  and  $H_u$ , i.e.,  $\langle H_d \rangle = v_1/\sqrt{2}$  and  $\langle H_u \rangle = v_2/\sqrt{2}$ . Instead of  $v_1$  and  $v_2$  it is more convenient to use  $\tan \beta = v_2/v_1$  and  $v = \sqrt{v_1^2 + v_2^2} \approx 246$  GeV. Hereafter, we neglect the contribution of loop corrections to the mass matrix (10). In this case, the mass of the charged fermion components of  $H_1^u$  and  $H_1^d$  is determined by  $\mu_{11}$ , i.e.,  $m_{\chi_1^\pm} = \mu_{11}$ .

When  $|\tilde{\mu}_1|$  is much larger than  $|\mu_{11}|$  and  $v$ , the perturbation theory method can be used to diagonalise the mass matrix (10) (see, for example, [55–58]). This method yields

$$\begin{aligned} m_{\chi_1} &\simeq m_{\chi_1^\pm} - \Delta_1, & m_{\chi_2} &\simeq m_{\chi_1^\pm} + \Delta_2, & m_{\chi_2} &\simeq \tilde{\mu}_1 + \Delta_1 + \Delta_2, \\ \Delta_1 &\simeq \frac{(\tilde{f}_{11} v \sin \beta + f_{11} v \cos \beta)^2}{4(\tilde{\mu}_1 - m_{\chi_1^\pm})}, & \Delta_2 &\simeq \frac{(\tilde{f}_{11} v \sin \beta - f_{11} v \cos \beta)^2}{4(\tilde{\mu}_1 + m_{\chi_1^\pm})}. \end{aligned} \quad (11)$$

From Equation (11), it follows that the masses of the lightest exotic chargino and neutralino states in the leading approximation are set by  $\mu_{11}$ . If the mass difference  $m_{\chi_2} - m_{\chi_1}$  is as small as  $O(100$  KeV), the inelastic scattering processes  $\chi_1 N \rightarrow \chi_2 N$ , where  $N$  is a nucleon, may occur. In this article, we restrict our consideration to the part of the SE<sub>6</sub>SSM parameter space where  $m_{\chi_2} - m_{\chi_1} > 200$  MeV. Because of this the inelastic scattering processes do not take place. Moreover, the lifetime of  $\chi_2$  is much smaller than 1 s that allows to preserve the success of the Big Bang Nucleosynthesis (BBN).

In the scenarios under consideration, the contribution of the lightest neutral exotic fermion to the cold dark matter relic density can be estimated using the approximate formula:

$$\Omega_{\tilde{H}} h^2 \simeq 0.1 \left( \frac{\mu_{11}}{1 \text{ TeV}} \right)^2. \quad (12)$$

which was derived in the case of the Higgsino dark matter within the MSSM (see, for example, [59,60]). Because the Planck observations lead to [61]:

$$(\Omega h^2)_{\text{exp}} = 0.1188 \pm 0.0010, \quad (13)$$

in the phenomenologically viable scenarios  $\mu_{11}$  is expected to be lower than 1.1 TeV. If  $\mu_{11} \lesssim 1.1$  TeV then gravitino may account for some or major part of the observed cold dark matter density.

In general, to find a viable cosmological scenario with stable gravitino one has to ensure that the decay products of the lightest unstable  $R$ -parity odd (or exotic) particle  $Y$  do not alter the abundances of light elements induced by BBN. The decays of state  $Y$  change the abundances of light elements the more the longer its lifetime  $\tau_Y$  is. This problem can be evaded if sparticle  $Y$  decays before BBN, i.e.,  $\tau_Y \lesssim 1$  s. The lifetime of state  $Y$  decaying into its SM partner (or lightest exotic fermion) and gravitino can be estimated as [62]:

$$\tau_Y \sim 48\pi \frac{m_{3/2}^2 M_P^2}{m_Y^5}, \quad (14)$$

where  $m_Y$  is its mass and  $M_P = (8\pi G_N)^{-1/2} \simeq 2.4 \cdot 10^{18}$  GeV is the reduced Planck mass. In order to obtain  $\tau_Y \lesssim 1$  sec for  $m_Y \simeq 1$  TeV we should restrict our consideration to  $m_{3/2} \lesssim 1$  GeV.

If gravitinos mostly originate from scattering processes of particles in the thermal bath then their abundance is approximately proportional to the reheating temperature  $T_R$  after inflation. In the leading approximation one finds [63,64]:

$$\Omega_{3/2} h^2 \sim 0.27 \left( \frac{T_R}{10^8 \text{ GeV}} \right) \left( \frac{1 \text{ GeV}}{m_{3/2}} \right) \left( \frac{M_{\tilde{g}}}{1 \text{ TeV}} \right)^2, \quad (15)$$

where  $M_{\tilde{g}}$  is a gluino mass. Taking into account that  $\Omega_{3/2} h^2 \leq 0.12$ , for  $M_{\tilde{g}} \gtrsim 2 - 3$  TeV and  $m_{3/2} \simeq 1$  GeV one obtains an upper bound on the reheating temperature, i.e.,  $T_R \lesssim 10^{6-7}$  GeV [65]. However even for so low reheating temperatures the appropriate amount of the baryon asymmetry can be induced within the  $SE_6$ SSM via the decays of the lightest right-handed neutrino/sneutrino into exotic states [5].

### 3. Dark Matter Nucleon Scattering Cross-section

Since the couplings of gravitino to the SM particles are negligibly small, in the scenarios under consideration the interactions of the cold dark matter with the baryons are determined by the couplings of the lightest neutral exotic fermion  $\chi_1$ . The low energy  $\chi_1$ -quark effective Lagrangian is:

$$\mathcal{L}_{\chi_1 q} = \sum_q \left( a_q \bar{\chi}_1 \chi_1 \bar{q} q + d_q \bar{\chi}_1 \gamma^\mu \gamma_5 \chi_1 \bar{q} \gamma_\mu \gamma_5 q \right). \quad (16)$$

The first term in the brackets gives rise to a spin-independent interaction, while the second one is associated with a spin-dependent interaction.

For spin-independent interactions experiments quote the cross-section to scatter off a nucleon, which is given by [66,67]:

$$\sigma_{SI} = \frac{4m_r^2}{\pi} \frac{(Zf^p + (A-Z)f^n)^2}{A^2}, \quad m_r = \frac{m_{\chi_1} m_N}{m_{\chi_1} + m_N}, \quad (17)$$

where  $A$  and  $Z$  are the nucleon number and charge of the target nucleus, whereas  $f^p$  and  $f^n$  are related to the quantities entering  $\mathcal{L}_{\chi_1 q}$ . To simplify our analysis, we assume that  $f^p \approx f^n \approx f^N$  while:

$$\begin{aligned} \frac{f^N}{m_N} &= \sum_{q=u,d,s} \frac{a_q}{m_q} f_{Tq}^N + \frac{2}{27} \sum_{Q=c,b,t} \frac{a_Q}{m_Q} f_{TQ}^N, \\ m_N f_{Tq}^N &= \langle N | m_q \bar{q} q | N \rangle, \quad f_{TQ}^N = 1 - \sum_{q=u,d,s} f_{Tq}^N. \end{aligned} \quad (18)$$

From Equations (17) and (18), one can see that  $\sigma_{SI}$  depends rather strongly on the hadronic matrix elements, i.e., the coefficients  $f_{Tq}^N$ , which are related to the  $\pi$ -nucleon  $\sigma$



term and the spin content of the nucleon. Here, we set  $f_{Tu}^N \simeq 0.0153$ ,  $f_{Td}^N \simeq 0.0191$ , and  $f_{Ts}^N \simeq 0.0447$ . These values of the hadronic matrix elements are the default values used in micrOMEGAs, as determined in Ref. [68] from lattice results (see also Ref. [69–72]).

Because in the  $SE_6SSM$  the exotic neutralino states do not couple to quarks and squarks, the coefficients  $a_q$  receives only contributions from the  $t$ -channel exchange of CP-even Higgs bosons. In the scenarios under consideration, the mass of the lightest Higgs particle  $m_{h_1}$  is much smaller than the masses of all other Higgs bosons. Therefore, in our analysis we ignore all contributions induced by the heavy Higgs exchange. Moreover, in this case, the lightest Higgs scalar manifests itself in the interactions with the SM fermions and gauge bosons as a SM-like Higgs so that:

$$\frac{a_q}{m_q} \simeq \frac{a_Q}{m_Q} \simeq \frac{g_{h\chi\chi}}{vm_{h_1}^2}. \quad (19)$$

When  $\Delta_1 \ll \mu_{11} \ll \tilde{\mu}_1$  the coupling  $g_{h\chi\chi}$  of the SM-like Higgs to the lightest exotic neutralino is given by:

$$|g_{h\chi\chi}| \simeq \frac{\Delta_1}{v}. \quad (20)$$

The dominant contribution to the coefficients  $d_q$  in Equation (16) comes from  $t$ -channel Z boson exchange. In the field basis  $(\tilde{H}_1^{d0}, \tilde{H}_1^{u0}, \tilde{S}_1)$ , the two lightest exotic neutralinos are made up of the following superposition of interaction states:

$$\chi_\alpha = N_\alpha^1 \tilde{H}_1^{d0} + N_\alpha^2 \tilde{H}_1^{u0} + N_\alpha^3 \tilde{S}_1, \quad (21)$$

where  $\alpha = 1, 2$ . In Equation (21),  $N_i^a$  is the exotic neutralino mixing matrix defined by:

$$N_i^a M^{ab} N_j^b = m_i \delta_{ij}, \quad \text{no sum on } i. \quad (22)$$

where  $M^{ab}$  is  $3 \times 3$  exotic neutralino mass matrix given by Equation (10). Using the above compositions of the lightest exotic neutralino states (21), it is straightforward to derive the couplings of these states to the Z-boson. The part of the Lagrangian that describes the interactions of Z with  $\chi_1$  and  $\chi_2$  can be written as:

$$\mathcal{L}_{Z\chi\chi} = \sum_{\alpha,\beta} \frac{M_Z}{2v} Z_\mu \left( \chi_\alpha^T \gamma_\mu \gamma_5 \chi_\beta \right) R_{Z\alpha\beta}, \quad R_{Z\alpha\beta} = N_\alpha^1 N_\beta^1 - N_\alpha^2 N_\beta^2. \quad (23)$$

Then, the coefficients  $d_q$  and the corresponding cross-section can be presented in the following form:

$$\sigma^{p,n} = \frac{12m_\tau^2}{\pi} \left( \sum_{q=u,d,s} d_q \Delta_q^{p,n} \right)^2, \quad d_q = \frac{T_{3q}}{2v^2} R_{Z11}, \quad (24)$$

where  $T_{3q}$  is the third component of isospin. For fractions of the nucleon spin carried by a given quark  $q$  we use (see also [60])

$$\Delta_u^p = \Delta_d^n = 0.842, \quad \Delta_d^p = \Delta_u^n = -0.427, \quad \Delta_s^p = \Delta_s^n = -0.085. \quad (25)$$

Using the compositions of  $\chi_1$  and  $\chi_2$  one can obtain the analytical expressions for the couplings of these states to the lightest Higgs boson, i.e.,

$$g_{h\alpha\beta} = -\frac{1}{\sqrt{2}} \left( f_{11} N_\alpha^3 N_\beta^2 \cos \beta + \tilde{f}_{11} N_\alpha^3 N_\beta^1 \sin \beta \right). \quad (26)$$

The perturbation theory method allows to compute the entries of the exotic neutralino mixing matrix  $N_i^a$ . Substituting  $N_i^a$  into Equation (26) one can reproduce the analytical

Formula (20) for  $g_{h11} = g_{h\chi\chi}$  and find the approximate expression for  $R_{Z11}$ . If  $\tilde{\mu}_1 \gg \mu_{11} > 0$  and  $\mu_{11}$  is larger than  $\tilde{f}_{11}v \sin \beta$  and  $f_{11}v \cos \beta$ , we get:

$$R_{Z11} \simeq \frac{v^2(f_{11}^2 \cos^2 \beta - \tilde{f}_{11}^2 \sin^2 \beta)}{4\mu_{11}(\tilde{\mu}_1 - \mu_{11})}. \quad (27)$$

Here, we restrict our considerations to low values of  $\tan \beta \simeq 2$ . In this case, the particle spectrum of the SE<sub>6</sub>SSM may include the SM-like Higgs state with mass around 125 GeV if  $\lambda \gtrsim g'_1$ . Indeed, in this SUSY model the upper bound on the lightest Higgs boson mass  $m_{h_1}$  is given by:

$$m_{h_1}^2 \lesssim \frac{\lambda^2}{2} v^2 \sin^2 2\beta + M_Z^2 \cos^2 2\beta + g_1'^2 v^2 (Q_{H_d} \cos^2 \beta + Q_{H_u} \sin^2 \beta)^2 + \tilde{\Delta}, \quad (28)$$

where  $Q_{H_d}$  and  $Q_{H_u}$  are the  $U(1)_N$  charges of  $H_d$  and  $H_u$  whereas  $\tilde{\Delta}$  is the contribution of loop corrections. When  $\lambda \gtrsim \sqrt{2}(M_Z/v) \simeq 0.52 \gtrsim g'_1$  the sum of the first two terms can be larger than  $M_Z^2$  for moderate values of  $\tan \beta$ . As a consequence, for  $\tan \beta \simeq 2$  the 125 GeV Higgs state can be obtained. In this part of the SE<sub>6</sub>SSM parameter space, all Higgs states except the lightest Higgs boson tend to have masses beyond the multi-TeV range and cannot be observed at the LHC experiments [1,2,41,42].

Within the SE<sub>6</sub>SSM the masses of new exotic states including the  $Z'$  boson are set by the VEVs of  $S$  and  $\bar{S}$ . LHC constraints require the extra  $U(1)_N$  gauge boson to be heavier than 4.5 TeV [73,74]. When  $\langle S \rangle \simeq \langle \bar{S} \rangle$  for the mass of the extra vector boson  $M_{Z'}$  one finds:

$$M_{Z'} \approx 2g'_1 Q_S \langle S \rangle. \quad (29)$$

In order to obtain  $M_{Z'} \gtrsim 4.5$  TeV we choose  $\langle S \rangle \simeq \langle \bar{S} \rangle \gtrsim 6$  TeV. The experimental constraints on  $Z - Z'$  mixing are also satisfied in this case. To avoid the experimental lower limit on the mass of the lightest exotic chargino and to ensure that the lightest exotic neutralino state gives rise to the phenomenologically acceptable dark matter density the interval of variation of  $\mu_{11}$  is chosen so that  $\mu_{11} \gtrsim 200$  GeV and  $\mu_{11} \lesssim 1$  TeV. This corresponds to relatively small values of  $\lambda_{11} \lesssim 0.17$ . To simplify our analysis we set  $\tilde{\mu}_1 \simeq 2$  TeV. In addition, we also require the validity of perturbation theory up to the scale  $M_X$  that, in particular, constrains the allowed range of  $f_{11}$  and  $\tilde{f}_{11}$ .

The results of our analysis are summarised in Figures 1 and 2. From the approximate expressions (20) and (27), it follows that the couplings  $g_{h\chi\chi}$  and  $R_{Z11}$  vanish if  $f_{11} \approx -\tilde{f}_{11} \tan \beta$ . In other words, the interactions of the lightest exotic neutralino with the baryons can be extremely weak in some part of the SE<sub>6</sub>SSM parameter space. Our numerical analysis indicates that the perturbation theory method describes the dependence of  $g_{h\chi\chi}$  and  $R_{Z11}$  on the SE<sub>6</sub>SSM parameter quite well when  $\tilde{\mu}_1$  is sufficiently large (see Figures 1 and 2). In particular,  $g_{h\chi\chi}$  and  $\sigma_{SI}$  grow whereas  $R_{Z11}$  and  $\sigma^{p,n}$  diminish with increasing  $\mu_{11}$  from 200 GeV to 1 TeV. Since in the scenario under consideration  $\tilde{f}_{11}$  is chosen to be negative while  $f_{11}$  is positive,  $g_{h\chi\chi}$  and  $R_{Z11}$ , as well as the spin-dependent and spin-independent cross sections decreases when  $f_{11}$  grows.


The calculated values of  $\sigma_{SI}$  and  $\sigma^{p,n}$  should be compared with the corresponding experimental limits. For  $m_{\chi_1} \approx \mu_{11} < 1.1$  TeV the value of  $\Omega_{\tilde{H}} h^2$  can be substantially smaller than  $(\Omega h^2)_{\text{exp}}$  that leads to some reduction in the number density of the lightest neutral exotic fermions. Thereby, the direct detection limits become weakened, i.e.,

$$\sigma_{SI} < \frac{(\Omega h^2)_{\text{exp}}}{\Omega_{\tilde{H}} h^2} (\sigma_{SI})_{\text{exp}}, \quad \sigma^{p,n} < \frac{(\Omega h^2)_{\text{exp}}}{\Omega_{\tilde{H}} h^2} (\sigma^{p,n})_{\text{exp}}, \quad (30)$$


where  $\sigma_{SI}$ ,  $\sigma^{p,n}$  and  $\Omega_{\tilde{H}} h^2$  are computed values of these quantities for each set of the parameters of the SE<sub>6</sub>SSM, while  $(\sigma_{SI})_{\text{exp}}$  and  $(\sigma^{p,n})_{\text{exp}}$  are the experimental limits on the spin-independent and spin-dependent cross-sections at the given mass  $m_{\chi_1}$ . Any set of the



SE<sub>6</sub>SSM parameters that does not satisfy the conditions (30) is basically ruled out. Hereafter, we just compare  $\sigma_{SI}$  and  $\sigma^{p,n}$  with  $(\sigma_{SI})_{\text{exp}}$  and  $(\sigma^{p,n})_{\text{exp}}$ .



**Figure 1.** (Left) The coupling  $g_{h\chi\chi}$  and (Right) the cross-section  $\sigma_{SI}$  as a function of  $f_{11}$  for  $\tilde{f}_{11} = -0.5$ ,  $\tan\beta = 2$ ,  $\tilde{\mu}_1 = 2$  TeV,  $\mu_{11} = 200$  GeV (solid lines), and  $\mu_{11} = 1$  TeV (dashed lines). The dotted lines correspond to the approximate expression for  $g_{h\chi\chi}$  (20).



**Figure 2.** (Left) The coupling  $R_{Z11}$  and (Right) the cross-section  $\sigma^{p,n}$  as a function of  $f_{11}$  for  $\tilde{f}_{11} = -0.5$ ,  $\tan\beta = 2$ ,  $\tilde{\mu}_1 = 2$  TeV,  $\mu_{11} = 200$  GeV (solid lines), and  $\mu_{11} = 1$  TeV (dashed lines). The upper solid and upper dashed lines represent  $\sigma^p$  while lower solid and lower dashed lines are associated with  $\sigma^n$ . The dotted lines correspond to the approximate expression for  $R_{Z11}$  (27).

The spin-independent cross-sections shown in Figure 1 remain always smaller than 60 yb for  $m_{\chi_1} \approx \mu_{11} \approx 200$  GeV and 300 yb for  $m_{\chi_1} \approx \mu_{11} \approx 1$  TeV ( $1 \text{ yb} = 10^{-48} \text{ cm}^2$ ) which are the values of  $(\sigma_{SI})_{\text{exp}}$  obtained by the LUX–ZEPLIN (LZ) experiment [10]. The most stringent experimental upper bound on the WIMP–proton spin-dependent cross-section was set by the PICO-60 experiment [75]. For  $m_{\chi_1} \approx \mu_{11}$  equals to 200 GeV (1 TeV) the value of  $(\sigma^p)_{\text{exp}} \approx 10^5 \text{ zb}$  ( $4 \cdot 10^5 \text{ zb}$ ), where  $1 \text{ zb} = 10^{-45} \text{ cm}^2$ . The WIMP–neutron spin-dependent cross-section is more tightly constrained. If  $m_{\chi_1} \approx \mu_{11} \approx 200$  GeV (1 TeV) the LZ experiment set the experimental limit  $(\sigma^n)_{\text{exp}} \approx 0.9 \cdot 10^4 \text{ zb}$  ( $5 \cdot 10^4 \text{ zb}$ ) [10]. All experimental bounds on the spin-dependent cross sections mentioned here are substantially larger than the values of  $\sigma^{p,n}$  presented in Figure 2.

In the part of the SE<sub>6</sub>SSM parameter space, where  $|g_{h\chi\chi}|$  and  $|R_{Z11}|$  become smaller than  $10^{-3}$ , one cannot ignore quantum corrections to the effective Lagrangian (16) which are induced by the one-loop diagrams involving the electroweak gauge bosons [76,77]. The inclusion of such corrections results in  $\sigma_{SI} \sim 0.1 \text{ yb}$  even when  $|g_{h\chi\chi}| \ll 10^{-3}$  at the tree level [78].

The values of  $\sigma^{p,n}$  and  $\sigma_{SI}$  presented in Figures 1 and 2 are substantially smaller than the maximal possible values of the spin-dependent and spin-independent  $\chi_1$ -nucleon scattering cross-sections in the SE<sub>6</sub>SSM. In the part of the parameter space examined in

Figures 1 and 2, the suppression of  $\sigma^{p,n}$  and  $\sigma_{SI}$  is caused by the large value of  $\tilde{\mu}_1$ , which is associated with the sparticle mass scale, as well as by the partial cancellations of different contributions to  $g_{h\chi\chi}$  and  $R_{Z11}$ . The maximal possible values of  $\sigma^{p,n}$  and  $\sigma_{SI}$  are attained when  $\mu_{11} \simeq \tilde{\mu}_1$  and  $\tilde{f}_{11} \sim f_{11} \sim 1$ . In this case, the spin-independent  $\chi_1$ -nucleon scattering cross section can reach 20–30 zb which is two orders of magnitude larger than the present experimental limit on  $\sigma_{SI}$  [10]. However, even when  $\mu_{11} \simeq \tilde{\mu}_1$ , the desirable suppression of  $\sigma_{SI}$  can be achieved if  $\tilde{f}_{11} \sim f_{11} \ll 0.1$ .

#### 4. Conclusions

To ensure anomaly cancellation the  $U(1)_N$  extensions of the MSSM ( $E_6$ SSM) are expected to include at least three fundamental representations of  $E_6$  below the GUT scale  $M_X$ . These fundamental representations of  $E_6$ , in particular, contain three families of Higgs-like doublets  $H_i^d$  and  $H_i^u$ . One pair of such Higgs-like doublet supermultiplets ( $H_u \equiv H_3^u$  and  $H_d \equiv H_3^d$ ) gains VEVs resulting in the breakdown of the EW symmetry while two other families ( $H_\alpha^d$  and  $H_\alpha^u$ , where  $\alpha = 1, 2$ ) are normally assumed to be inert.

In addition to three 27-plets the low-energy matter content of the  $SE_6$ SSM also involves an  $E_6$  singlet superfield  $\phi$ , a pair of superfields  $S$  and  $\bar{S}$  with opposite  $U(1)_N$  charges, as well as a pair of  $SU(2)_W$  doublets  $L_4$  and  $\bar{L}_4$  with opposite  $SU(2)_W \times U(1)_Y \times U(1)_N$  quantum numbers. The supermultiplets  $S, \bar{S}, L_4$  and  $\bar{L}_4$  can come from additional  $27'_1$  and  $\bar{27}'_1$ . Within the  $SE_6$ SSM a single  $\tilde{Z}_2^H$  symmetry prevents rapid proton decay and tree-level non-diagonal flavor transitions. In this paper, we have considered new variant of this SUSY model in which the low-energy particle spectrum is supplemented by three  $E_6$  singlet superfields  $\phi_i$  which are odd under the  $\tilde{Z}_2^H$  symmetry. This permits to avoid the presence of neutral exotic fermions with tiny masses ( $\lesssim 1$  eV) in the spectrum of the  $SE_6$ SSM. The scalar components of the  $\tilde{Z}_2^H$  even superfields  $\phi, S$  and  $\bar{S}$  may acquire very large VEVs, i.e.,  $\langle \phi \rangle \sim \langle S \rangle \sim \langle \bar{S} \rangle \gg 10$  TeV, breaking the  $U(1)_N$  gauge symmetry and generating masses of all extra exotic fermions which can be much heavier than all SM particles.

The conservation of  $R$ -parity and the  $\tilde{Z}_2^H$  symmetry within the  $SE_6$ SSM ensures that at least two neutral states can be stable, giving rise to dark matter density. Since no firm indication of the presence of dark matter has been observed at the direct detection experiments we assume that one of the stable states is gravitino with mass  $m_{3/2} \lesssim 1$  GeV. Another stable state can be the lightest exotic fermion which is mostly composed of the neutral fermion components of the supermultiplets  $H_1^d$  and  $H_1^u$ . In this case, the lightest and second lightest exotic neutralino states ( $\chi_1$  and  $\chi_2$ ), as well as the lightest exotic chargino  $\chi_1^\pm$  are almost degenerate. If these exotic fermions have masses below 1.1 TeV they can lead to the phenomenologically acceptable density of the dark matter.

The interactions of dark matter with the baryons are defined by the couplings of  $\chi_1$  because the interaction of gravitino with the SM fermions and bosons is extremely weak. In this article, we have explored the scenario in which all scalars and exotic fermions except the lightest Higgs boson, gravitino,  $\chi_1$ ,  $\chi_2$ , and  $\chi_1^\pm$  have masses of the order of a few TeV or even higher. Therefore, the spin-dependent and spin-independent  $\chi_1$ -nucleon cross-sections ( $\sigma^{p,n}$  and  $\sigma_{SI}$ ) are dominated by the  $t$ -channel exchanges of the  $Z$  boson and the lightest Higgs scalar, respectively. Our analysis revealed that for a fixed mass of the lightest exotic fermion  $m_{\chi_1}$  the values of  $\sigma^{p,n}$  and  $\sigma_{SI}$  vanish in the leading approximation if  $f_{11} = -\tilde{f}_{11} \tan \beta$ . As a result for a given  $m_{\chi_1}$  the spin-independent  $\chi_1$ -nucleon cross-section varies from its maximal value, which is about 20–30 zb, to zero. When  $f_{11} \approx -\tilde{f}_{11} \tan \beta$  so that  $|g_{h\chi\chi}| \ll 10^{-3}$  and  $|R_{Z11}| \ll 10^{-3}$ , one cannot ignore the contributions to  $\sigma^{p,n}$  and  $\sigma_{SI}$  induced by the  $t$ -channel exchanges of the heavy Higgs states. Moreover, in this case, one has to take into account the quantum corrections to the effective Lagrangian (16), which is beyond the scope of the present paper. Nevertheless, the results presented in Figures 1 and 2 demonstrate that there is a part of the  $SE_6$ SSM parameter space in which  $\sigma^{p,n}$  and  $\sigma_{SI}$  can be considerably smaller than the corresponding experimental limits. In particular, the spin-independent dark matter nucleon scattering cross section may be less than 10 yb. In the near future, the experiments XENONnT [79], LZ [80], DarkSide-20k

[81], and DARWIN [82] may set even more stringent bounds on the spin-dependent and spin-independent WIMP-nucleon scattering cross-sections constraining further the  $SE_6SSM$  parameter space.

As mentioned before, the scenario under consideration implies that two lightest exotic neutralino states and the lightest exotic chargino must be lighter than 1.1 TeV. Several collider experiments have searched for such particles. However, if the mass splitting between these states is small the decay products of  $\chi_2$  and  $\chi_1^\pm$  are very soft and may escape detection. This happens, for example, with the lightest ordinary neutralino and chargino states within natural SUSY, where the splitting between the heavier neutralino or the chargino and the LSP is at least a few GeV [83–85]. In this case, the results of the search for the lightest exotic fermions depend on  $\Delta = m_{\chi_1^\pm} - m_{\chi_1}$ .

At the LHC the pair production of such states can occur through off-shell  $W$  and  $Z$  bosons. ATLAS excluded the exotic chargino with masses below 193 GeV and 140 GeV for  $\Delta \simeq 4.7$  GeV and  $\Delta \simeq 2$  GeV, respectively [86], while CMS ruled out such chargino with masses below 112 GeV for  $\Delta = 1$  GeV [87]. On the other hand, if  $\Delta \lesssim 150$  MeV the exotic charginos may be long-lived. When their lifetime is longer than the time needed to pass through the detector, these states appear as charged stable massive particles. LHC experiments excluded such charginos with masses below 1090 GeV [88].

The last experimental limit is not applicable in the case of the  $SE_6SSM$  scenario discussed here because  $\Delta$  tends to be larger than 300 MeV [78,89]. The part of the  $SE_6SSM$  parameter space explored in Figures 1 and 2 is associated with relatively small mass splitting between  $\chi_1$ ,  $\chi_2$ , and  $\chi_1^\pm$ . For  $m_{\chi_1^\pm} \simeq \mu_{11} = 200$  GeV the value of  $\Delta \approx \Delta_1$  estimated using Equation (11) changes from 1.67 GeV to 0.42 GeV when  $f_{11}$  increases from 0 to 0.5. As a consequence the lightest exotic chargino decays mainly into hadrons and the lightest exotic neutralino so that it seems to be rather problematic to discover the set of these states at hadron colliders. The discovery prospects for such fermions at future International Linear Collider look more promising (for a review see [90]).

Thus, in this article we argued that within the new variant of the  $SE_6SSM$ , in which the cold dark matter density is formed by two stable states, there are some regions of the parameter space that are safe from all current constraints. The SUSY model under consideration predicts the existence of exotic fermions and exotic scalars which are colored and carry baryon and lepton numbers simultaneously [6]. In collider experiments, such exotic quarks/squarks can only be created in pairs. The stringent LHC lower limits on the masses of scalar leptoquarks [91–93] are not directly applicable in the case of the  $SE_6SSM$ . Indeed, in the  $E_6$  inspired SUSY models ordinary scalar leptoquark can decay either to an up-type quark  $u_i$  and charged lepton  $\ell_j$  or to down-type quark  $d_i$  and left-handed neutrino  $\nu_j$ . On the other hand, in the  $SE_6SSM$  the lightest exotic colored particle  $D_1$  (scalar or fermion) is odd under  $\tilde{Z}_2^E$  symmetry. Therefore, its decay always gives rise to the missing energy and transverse momentum in the final state associated with the lightest exotic fermion, i.e.,

$$D_1 \rightarrow u_i(d_i) + \ell_j(\nu_j) + E_T^{\text{miss}} + X.$$

Here,  $X$  may include extra charged leptons and/or jets that can come from the decays of intermediate states. Although the masses of the exotic colored particles are expected to be in the multi TeV range  $D_1$  can be lighter than 2 TeV. If  $D_1$  is exotic colored fermion the LHC pair production cross section for  $D_1$  decreases from 10 fb to 1 fb when its mass changes from 1.3 TeV to 1.7 TeV [94]. The LHC pair production cross-section for the lightest exotic squarks tends to be an order of magnitude smaller. The pair production of  $D_1$  may lead to some enhancement of the cross sections of  $pp \rightarrow jj\ell^+\ell^- + E_T^{\text{miss}} + X$  and  $pp \rightarrow jj + E_T^{\text{miss}} + X$  if such states are relatively light. The discovery of these exotic colored states will provide a smoking gun signal of the model allowing to distinguish it from other extensions of the SM.

**Funding:** This research received no external funding.

**Data Availability Statement:** Data sharing is not applicable to this article.

**Acknowledgments:** R.N. thanks X. Tata for very valuable comments and remarks. R.N. acknowledges fruitful discussions with S. Khalil, S. Moretti, H. Waltari, and P. V. Zinin.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

- King, S.F.; Moretti, S.; Nevzorov, R. Theory and phenomenology of an exceptional supersymmetric standard model. *Phys. Rev. D* **2006**, *73*, 035009.
- King, S.F.; Moretti, S.; Nevzorov, R. Exceptional supersymmetric standard model. *Phys. Lett. B* **2006**, *634*, 278.
- King, S.F.; Moretti, S.; Nevzorov, R. A Review of the Exceptional Supersymmetric Standard Model. *Symmetry* **2020**, *12*, 557.
- Hambye, T.; Ma, E.; Raidal, M.; Sarkar, U. Allowable low-energy  $E(6)$  subgroups from leptogenesis. *Phys. Lett. B* **2001**, *512*, 373.
- Nevzorov, R. Leptogenesis as an origin of hot dark matter and baryon asymmetry in the  $E_6$  inspired SUSY models. *Phys. Lett. B* **2018**, *779*, 223.
- Nevzorov, R.  $E_6$  inspired supersymmetric models with exact custodial symmetry. *Phys. Rev. D* **2013**, *87*, 015029.
- Athron, P.; Mühlleitner, M.; Nevzorov, R.; Williams, A.G. Non-Standard Higgs Decays in  $U(1)$  Extensions of the MSSM. *J. High Energy Phys.* **2015**, *1501*, 153.
- Aprile, E.; et al. [XENON Collaboration]. Dark Matter Search Results from a One Ton-Year Exposure of XENON1T. *Phys. Rev. Lett.* **2018**, *121*, 111302.
- Meng, Y. et al. [PandaX-4T Collaboration]. Dark Matter Search Results from the PandaX-4T Commissioning Run. *Phys. Rev. Lett.* **2021**, *127*, 261802.
- Aalbers, J.; Akerib, D.S.; Akerlof, C.W.; Musalhi, A.K.A.; Alder, F.; Alqahtani, A.; Alsum, S.K.; Amarasinghe, C.S.; Ames, A.; Anderson, T.J.; et al. First Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment. *arXiv* **2022**, arXiv:2207.03764.
- Howl, R.; King, S.F. Planck Scale Unification in a Supersymmetric Standard Model. *Phys. Lett. B* **2007**, *652*, 331.
- Howl, R.; King, S.F. Minimal  $E_6$  Supersymmetric Standard Model. *J. High Energy Phys.* **2008**, *0801*, 030.
- Howl, R.; King, S.F. Exceptional Supersymmetric Standard Models with non-Abelian Discrete Family Symmetry. *J. High Energy Phys.* **2008**, *0805*, 008.
- Howl, R.; King, S.F. Solving the Flavour Problem in Supersymmetric Standard Models with Three Higgs Families. *Phys. Lett. B* **2010**, *687*, 355.
- Athron, P.; Hall, J.P.; Howl, R.; King, S.F.; Miller, D.J.; Moretti, S.; Nevzorov, R. Aspects of the Exceptional Supersymmetric Standard Model. *Nucl. Phys. Proc. Suppl.* **2010**, *200–202*, 120.
- Hall, J.P.; King, S.F. Bino Dark Matter and Big Bang Nucleosynthesis in the Constrained  $E_6$ SSM with Massless Inert Singlinos. *J. High Energy Phys.* **2011**, *1106*, 006.
- Callaghan, J.C.; King, S.F.  $E_6$  Models from F-theory. *J. High Energy Phys.* **2013**, *1304*, 034.
- Callaghan, J.C.; King, S.F.; Leontaris, G.K. Gauge coupling unification in  $E_6$  F-theory GUTs with matter and bulk exotics from flux breaking. *J. High Energy Phys.* **2013**, *1312*, 037.
- Khalil, S.; Moretti, S.; Rojas-Ciofalo, D.; Waltari, H. Multicomponent dark matter in a simplified  $E_6$ SSM. *Phys. Rev. D* **2020**, *102*, 075039.
- Suematsu, D. Neutralino decay in the mu problem solvable extra  $U(1)$  models. *Phys. Rev. D* **1998**, *57*, 1738.
- Keith, E.; Ma, E. Generic consequences of a supersymmetric  $U(1)$  gauge factor at the TeV scale. *Phys. Rev. D* **1997**, *56*, 7155.
- Keith, E.; Ma, E. Efficacious Extra  $U(1)$  Factor for the Supersymmetric Standard Model. *Phys. Rev. D* **1996**, *54*, 3587.
- Suematsu, D.; Yamagishi, Y. Radiative symmetry breaking in a supersymmetric model with an extra  $U(1)$ . *Int. J. Mod. Phys. A* **1995**, *10*, 4521.
- Daikoku, Y.; Suematsu, D. Mass bound of the lightest neutral Higgs scalar in the extra  $U(1)$  models. *Phys. Rev. D* **2000**, *62*, 095006.
- King, S.F.; Moretti, S.; Nevzorov, R. Gauge coupling unification in the exceptional supersymmetric standard model. *Phys. Lett. B* **2007**, *650*, 57.
- Sperling, M.; Stöckinger, D.; Voigt, A. Renormalization of vacuum expectation values in spontaneously broken gauge theories. *J. High Energy Phys.* **2013**, *1307*, 132.
- Sperling, M.; Stöckinger, D.; Voigt, A. Renormalization of vacuum expectation values in spontaneously broken gauge theories: Two-loop results. *J. High Energy Phys.* **2014**, *1401*, 068.
- Ma, E. Neutrino masses in an extended gauge model with  $E(6)$  particle content. *Phys. Lett. B* **1996**, *380*, 286.
- Hall, J.P.; King, S.F. Neutralino Dark Matter with Inert Higgsinos and Singlinos. *J. High Energy Phys.* **2009**, *0908*, 088.
- Athron, P.; Thomas, A.W.; Underwood, S.J.; White, M.J. Dark matter candidates in the constrained Exceptional Supersymmetric Standard Model. *Phys. Rev. D* **2017**, *95*, 035023.
- Nevzorov, R. Quasifixed point scenarios and the Higgs mass in the  $E_6$  inspired supersymmetric models. *Phys. Rev. D* **2014**, *89*, 055010.
- Nevzorov, R.; Trusov, M.A. Infrared quasifixed solutions in the NMSSM. *Phys. Atom. Nucl.* **2001**, *64*, 1299.
- Nevzorov, R.; Trusov, M.A. Quasifixed point scenario in the modified NMSSM. *Phys. Atom. Nucl.* **2002**, *65*, 335.
- Athron, P.; King, S.F.; Miller, D.J.; Moretti, S.; Nevzorov, R. The Constrained  $E_6$ SSM. *arXiv* **2008**, arXiv:0810.0617.

35. Athron, P.; King, S.F.; Miller, D.J.; Moretti, S.; Nevzorov, R. Constrained Exceptional Supersymmetric Standard Model with a Higgs Near 125 GeV. *Phys. Rev. D* **2012**, *86*, 095003.
36. Athron, P.; Harries, D.; Nevzorov, R.; Williams, A.G.  $E_6$  Inspired SUSY benchmarks, dark matter relic density and a 125 GeV Higgs. *Phys. Lett. B* **2016**, *760*, 19.
37. Athron, P.; Harries, D.; Nevzorov, R.; Williams, A.G. Dark matter in a constrained  $E_6$  inspired SUSY model. *J. High Energy Phys.* **2016**, *1612*, 128.
38. Athron, P.; Binjonaid, M.; King, S.F. Fine Tuning in the Constrained Exceptional Supersymmetric Standard Model. *Phys. Rev. D* **2013**, *87*, 115023.
39. Athron, P.; Harries, D.; Williams, A.G.  $Z'$  mass limits and the naturalness of supersymmetry. *Phys. Rev. D* **2015**, *91*, 115024.
40. Athron, P.; Stöckinger, D.; Voigt, A. Threshold Corrections in the Exceptional Supersymmetric Standard Model. *Phys. Rev. D* **2012**, *86*, 095012.
41. King, S.F.; Moretti, S.; Nevzorov, R. Spectrum of Higgs particles in the ESSM. *arXiv* **2006**, arXiv:hep-ph/0601269.
42. King, S.F.; Moretti, S.; Nevzorov, R.  $E_6$ SSM. *AIP Conf. Proc.* **2007**, *881*, 138.
43. Belyaev, A.; Hall, J.P.; King, S.F.; Svantesson, P. Novel gluino cascade decays in  $E_6$  inspired models. *Phys. Rev. D* **2012**, *86*, 031702.
44. Belyaev, A.; Hall, J.P.; King, S.F.; Svantesson, P. Discovering  $E_6$  supersymmetric models in gluino cascade decays at the LHC. *Phys. Rev. D* **2013**, *87*, 035019.
45. Nevzorov, R.; Pakvasa, S. Exotic Higgs decays in the  $E_6$  inspired SUSY models. *Phys. Lett. B* **2014**, *728*, 210.
46. Hall, J.P.; King, S.F.; Nevzorov, R.; Pakvasa, S.; Sher, M. Novel Higgs Decays and Dark Matter in the  $E_6$ SSM. *Phys. Rev. D* **2011**, *83*, 075013.
47. Hesselbach, S.; Miller, D.J.; Moortgat-Pick, G.; Nevzorov, R.; Trusov, M. Theoretical upper bound on the mass of the LSP in the MNSSM. *Phys. Lett. B* **2008**, *662*, 199.
48. Frere, J.M.; Nevzorov, R.B.; Vysotsky, M.I. Stimulated neutrino conversion and bounds on neutrino magnetic moments. *Phys. Lett. B* **1997**, *394*, 127.
49. Kolda, C.F.; Martin, S.P. Low-energy supersymmetry with D term contributions to scalar masses. *Phys. Rev. D* **1996**, *53*, 3871.
50. Giudice, G.F.; Rattazzi, R. Theories with gauge mediated supersymmetry breaking. *Phys. Rept.* **1999**, *322*, 419.
51. Dubovsky, S.L.; Gorbunov D.S.; Troitsky, S.V. Gauge mechanism of mediation of supersymmetry breaking. *Phys. Usp.* **1999**, *42*, 623.
52. Gherghetta, T.; Pomarol, A. Bulk fields and supersymmetry in a slice of AdS. *Nucl. Phys. B* **2000**, *586*, 141.
53. Gherghetta, T.; Pomarol, A. A Warped supersymmetric standard model. *Nucl. Phys. B* **2001**, *602*, 3.
54. Gu, Y.; Khlopov, M.; Wu, L.; Yang, J.M.; Zhu, B. Light gravitino dark matter: LHC searches and the Hubble tension. *Phys. Rev. D* **2020**, *102*, 115005.
55. Kovalenko, P.A.; Nevzorov, R.B.; Ter-Martirosian, K.A. Masses of Higgs bosons in supersymmetric theories. *Phys. Atom. Nucl.* **1998**, *61*, 812.
56. Nevzorov, R.B.; Trusov, M.A. Particle spectrum in the modified NMSSM in the strong Yukawa coupling limit. *J. Exp. Theor. Phys.* **2000**, *91*, 1079.
57. Nevzorov, R.B.; Ter-Martirosyan, K.A.; Trusov, M.A. Higgs bosons in the simplest SUSY models. *Phys. Atom. Nucl.* **2002**, *65*, 285.
58. Nevzorov, R.; Miller, D.J. Approximate solutions for the Higgs masses and couplings in the NMSSM. *arXiv* **2004**, arXiv:hep-ph/0411275.
59. Arkani-Hamed, N.; Delgado A.; Giudice, G.F. The Well-tempered neutralino. *Nucl. Phys. B* **2006**, *741*, 108.
60. Chalons, G.; Dolan, M.J.; McCabe, C. Neutralino dark matter and the Fermi gamma-ray lines. *J. Cosmol. Astropart. Phys.* **2013**, *02*, 016.
61. The Planck collaboration. Planck 2015 results. XIII. Cosmological parameters. *Astron. Astrophys.* **2016**, *594*, A13.
62. Feng, J.L.; Su, S.; Takayama, F. Supergravity with a gravitino LSP. *Phys. Rev. D* **2004**, *70*, 075019.
63. Bolz, M.; Brandenburg A.; Buchmuller, W. Thermal production of gravitinos. *Nucl. Phys. B* **2001**, *606*, 518.
64. Eberl, H.; Gialamas, I.D.; Spanos, V.C. Gravitino thermal production revisited. *Phys. Rev. D* **2021**, *103*, 075025.
65. Hook, A.; McGehee R.; Murayama, H. Cosmologically Viable Low-energy Supersymmetry Breaking. *Phys. Rev. D* **2018**, *98*, 115036.
66. Ellis, J.R.; Olive, K.A.; Savage, C. Hadronic Uncertainties in the Elastic Scattering of Supersymmetric Dark Matter. *Phys. Rev. D* **2008**, *77*, 065026.
67. Kalinowski, J.; King, S.F.; Roberts, J.P. Neutralino Dark Matter in the USSM. *J. High Energy Phys.* **2009**, *2009*, 066.
68. Bélanger, G.; Boudjema, F.; Pukhov, A.; Semenov, A. micrOMEGAS<sub>3</sub>: A program for calculating dark matter observables. *Comput. Phys. Commun.* **2014**, *185*, 960.
69. Alarcon, J.M.; Martin Camalich J.; Oller, J.A. The chiral representation of the  $\pi N$  scattering amplitude and the pion-nucleon sigma term. *Phys. Rev. D* **2012**, *85*, 051503.
70. Thomas, A.W.; Shanahan, P.E.; Young, R.D. Strangeness in the nucleon: what have we learned? *Nuovo Cim. C* **2012**, *035N04*, 3.
71. Cheng, H.Y.; Chiang, C.W. Revisiting Scalar and Pseudoscalar Couplings with Nucleons. *J. High Energy Phys.* **2012**, *1207*, 009.
72. Alarcon, J.M.; Geng, L.S.; Martin Camalich J.; Oller, J.A. The strangeness content of the nucleon from effective field theory and phenomenology. *Phys. Lett. B* **2014**, *730*, 342.

- 
73. Sirunyan, A.M. et al. [CMS Collaboration]. Search for resonant and nonresonant new phenomena in high-mass dilepton final states at  $\sqrt{s} = 13$  TeV. *J. High Energy Phys.* **2021**, *7*, 208.
  74. Aad, G. et al. [ATLAS Collaboration]. Search for high-mass dilepton resonances using  $139 \text{ fb}^{-1}$  of  $pp$  collision data collected at  $\sqrt{s} = 13$  TeV with the ATLAS detector. *Phys. Lett. B* **2019**, *796*, 68.
  75. Amole, C. et al. [PICO Collaboration]. Dark Matter Search Results from the Complete Exposure of the PICO-60  $\text{C}_3\text{F}_8$  Bubble Chamber. *Phys. Rev. D* **2019**, *100*, 022001.
  76. Hisano, J.; Ishiwata, K.; Nagata, N.; Takesako, T. Direct Detection of Electroweak-Interacting Dark Matter. *J. High Energy Phys.* **2011**, *7*, 005.
  77. Hisano, J.; Ishiwata, K.; Nagata, N. Direct Search of Dark Matter in High-Scale Supersymmetry. *Phys. Rev. D* **2013**, *87*, 035020.
  78. Nagata, N.; Shirai, S. Higgsino Dark Matter in High-Scale Supersymmetry. *J. High Energy Phys.* **2015**, *1*, 029.
  79. Aprile, E. et al. [XENON Collaboration]. Projected WIMP sensitivity of the XENONnT dark matter experiment. *J. Cosmol. Astropart. Phys.* **2020**, *11*, 031.
  80. Akerib, D.S. et al. [LUX-ZEPLIN Collaboration]. Projected WIMP sensitivity of the LUX-ZEPLIN dark matter experiment. *Phys. Rev. D* **2020**, *101*, 052002.
  81. Aalseth, C.E. et al. [DarkSide-20k Collaboration]. DarkSide-20k: A 20 tonne two-phase LAr TPC for direct dark matter detection at LNGS. *Eur. Phys. J. Plus* **2018**, *133*, 131.
  82. Aalbers, J. et al. [DARWIN Collaboration]. DARWIN: towards the ultimate dark matter detector. *J. Cosmol. Astropart. Phys.* **2016**, *11*, 017.
  83. Baer, H.; Barger, V.; Huang, P.; Mustafayev, A.; Tata, X. Radiative natural SUSY with a 125 GeV Higgs boson. *Phys. Rev. Lett.* **2012**, *109*, 161802.
  84. Baer, H.; Barger, V.; Huang, P.; Tata, X. Natural Supersymmetry: LHC, dark matter and ILC searches. *J. High Energy Phys.* **2012**, *5*, 109.
  85. Baer, H.; Barger, V.; Huang, P.; Mickelson, D.; Mustafayev, A.; Tata, X. Radiative natural supersymmetry: Reconciling electroweak fine-tuning and the Higgs boson mass. *Phys. Rev. D* **2013**, *87*, 115028.
  86. Aad, G. et al. [ATLAS Collaboration]. Searches for electroweak production of supersymmetric particles with compressed mass spectra in  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector. *Phys. Rev. D* **2020**, *101*, 052005.
  87. Sirunyan, A.M. et al. [CMS Collaboration]. Search for supersymmetry with a compressed mass spectrum in the vector boson fusion topology with 1-lepton and 0-lepton final states in proton-proton collisions at  $\sqrt{s} = 13$  TeV. *J. High Energy Phys.* **2019**, *8*, 150.
  88. Aaboud, M. et al. [ATLAS Collaboration]. Search for heavy charged long-lived particles in the ATLAS detector in  $36.1 \text{ fb}^{-1}$  of proton-proton collision data at  $\sqrt{s} = 13$  TeV. *Phys. Rev. D* **2019**, *99*, 092007.
  89. Cirelli, M.; Fornengo, N.; Strumia, A. Minimal dark matter. *Nucl. Phys. B* **2006**, *753*, 178.
  90. Baer, H.; Barklow, T.; Fujii, K.; Gao, Y.; Hoang, A.; Kanemura, S.; List, J.; Logan, H.E.; Nomerotski, A.; Perelstein, M.; et al. The International Linear Collider Technical Design Report—Volume 2: Physics. *arXiv* **2013**, arXiv:1306.6352.
  91. Aad, G. et al. [ATLAS Collaboration]. Search for pairs of scalar leptoquarks decaying into quarks and electrons or muons in  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector. *J. High Energy Phys.* **2020**, *10*, 112.
  92. Aad, G. et al. [ATLAS Collaboration]. Search for pair production of third-generation scalar leptoquarks decaying into a top quark and a  $\tau$ -lepton in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector. *J. High Energy Phys.* **2021**, *06*, 179.
  93. Sirunyan, A.M. et al. [CMS Collaboration]. Searches for physics beyond the standard model with the  $M_{T2}$  variable in hadronic final states with and without disappearing tracks in proton-proton collisions at  $\sqrt{s} = 13$  TeV. *Eur. Phys. J. C* **2020**, *80*, 3.
  94. Kang, J.; Langacker, P.; Nelson, B.D. Theory and Phenomenology of Exotic Isosinglet Quarks and Squarks. *Phys. Rev. D* **2008**, *77*, 035003.