

Article

Numerical Analysis of the Time-Fractional Boussinesq Equation in Gradient Unconfined Aquifers with the Mittag-Leffler Derivative

Safyan Mukhtar

Department of Basic Sciences, Preparatory Year Deanship, King Faisal University, Al Ahsa 31982, Saudi Arabia; smahmad@kfu.edu.sa

Abstract: In this study, two numerical methods—the variational iteration transform method (VITM) and the Adomian decomposition (ADM) method—were used to solve the second- and fourth-order fractional Boussinesq equations. Both methods are helpful in approximating non-linear problems effectively, easily, and accurately. The fractional Atangana–Baleanu operator and ZZ transform were utilized to derive solutions for the equation. Two examples are discussed to validate the methods and solutions. The results demonstrate that both the VITM and ADM methods are effective in obtaining accurate and reliable solutions for the time-fractional Boussinesq equation.

Keywords: Boussinesq equation; fractional Atangana–Baleanu derivative; variational iteration transform method; Adomian decomposition transform method

1. Introduction

Fractional differential equations are mathematical models that describe the behaviors of dynamic systems where the order of differentiation is not an integer. Unlike traditional differential equations, which use integer derivatives to describe changes in a system over time, fractional differential equations use fractional derivatives to account for the non-integer rate of the change in a system [1,2]. This approach is useful for describing complex systems that involve memory effects, long-term persistence, and anomalous diffusion, as well as modeling real-world phenomena, such as turbulence, viscoelasticity, and fractional diffusion [3,4].

Applications of fractional differential equations span a wide range of fields, including physics, engineering, finance, and biology. In physics, fractional differential equations have been used to describe the behaviors of viscoelastic materials, such as rubber and biological tissues, which exhibit non-Newtonian behavior and respond differently to changes in stress and strain compared to traditional materials [5–7]. In engineering, fractional differential equations have been used to model the behaviors of complex systems, such as electrical circuits and control systems, where the order of differentiation cannot be described by an integer derivative. In finance, fractional differential equations have been used to model the price dynamics of financial assets, such as stocks and bonds, which exhibit persistent behavior and long-term dependencies [8–10]. In biology, fractional differential equations have been used to describe the diffusion of molecules, cells, and pathogens, which exhibit anomalous behaviors and do not follow traditional Fickian diffusion laws.

Symmetry is a fundamental concept in mathematics and physics, and it can be used to simplify the solution of partial differential equations (PDEs). In particular, the use of symmetry in the solutions of fractional PDEs can greatly simplify the mathematical analyses and lead to exact or approximate solutions [11–13]. Symmetry can be used to reduce the number of independent variables in a fractional PDE, which can simplify the solution process. Additionally, symmetry can be used to obtain solutions that are invariant under certain transformations, such as translations, rotations, and scaling. This can lead to



Citation: Mukhtar, S. Numerical Analysis of the Time-Fractional Boussinesq Equation in Gradient Unconfined Aquifers with the Mittag-Leffler Derivative. *Symmetry* **2023**, *15*, 608. <https://doi.org/10.3390/sym15030608>

Academic Editor: Dongfang Li

Received: 9 February 2023

Revised: 16 February 2023

Accepted: 24 February 2023

Published: 27 February 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

solutions that are more physically meaningful and easier to interpret. Overall, symmetry is a powerful tool for solving fractional PDEs and can greatly aid in understanding the underlying physical phenomena [14–16]. There are several important references that provide an overview of fractional differential equations and their applications. A comprehensive overview of fractional calculus, fractional differential equations, and their applications in various fields are thoroughly discussed in [17,18]. The theory of fractional differential equations and their applications in the field of viscoelasticity has been analyzed [19,20].

The Boussinesq equation is a mathematical model used to describe the behaviors of fluid systems. It is a partial differential equation that relates fluid flow velocity and pressure. The equation was first introduced by Joseph Valentin Boussinesq in 1872 and has been widely used in many engineering and scientific fields, including hydrology, oceanography, and geology [21–23]. The traditional Boussinesq equation is a second-order equation and has been widely used for many years to study the behaviors of fluid systems. However, the study of fluid systems became more complex, leading to more sophisticated mathematical models. This has led to the development of time-fractional Boussinesq equations [24–26].

The second and fourth-order time-fractional Boussinesq equations are extensions of the traditional Boussinesq equation; they take into account the fractional order derivative of time in their formulations. The fractional derivative of time represents the non-local and non-Markovian behaviors of the fluid system and provides a better description of the behavior of the fluid system compared to the traditional Boussinesq equation. The second-order time-fractional Boussinesq equation is linear, while the fourth-order time-fractional Boussinesq equation is nonlinear [27]. These equations have been used to study the behaviors of complex fluid systems, such as turbulent flows, waves, and heat transfer, and they have proven to be more accurate and reliable than the traditional Boussinesq equation [28].

The variational iteration transform method (VITM) is a numerical technique used to solve nonlinear partial differential equations. It is a relatively new method that has gained popularity in recent years due to its simplicity and effectiveness in solving a wide range of problems [29,30]. The VITM is based on the concept of variational iteration, which is a powerful tool for solving nonlinear problems. The method involves transforming the original equation into a new form that is easier to solve, and then using an iterative process to find the solution. The VITM has been applied successfully to a variety of problems, including nonlinear boundary value problems, nonlinear differential equations, and fractional differential equations [31,32].

The fractional partial differential equation is a type of mathematical equation that describes the behavior of complex systems with memory and nonlocal effects. It is a generalization of the classical partial differential equation and can be used to model a wide range of physical, biological, and engineering processes [33,34]. The fractional derivative is a nonlocal operator that captures the long-range memory effects of the system and is defined using the Riemann–Liouville or the Caputo fractional derivative. The VITM starts with a guess function and then iteratively improves it using the variational principle. The first step is to convert the fractional partial differential equation into a nonlinear integral equation by using the fractional derivative operator. Then, the guess function is used to approximate the solution of the integral equation and the residual is calculated. The residual is then used to update the guess function in the next iteration. The iteration continues until the residual is small enough to satisfy a desired level of accuracy [35,36]. The VITM has several advantages over other numerical methods for solving FPDEs. It is simple to implement, does not require the solution of linear systems, and has a fast convergence rate. It is also flexible and can be applied to a wide range of FPDEs with different boundary conditions. Furthermore, it can be easily extended to solve systems of FPDEs and to incorporate additional constraints. In conclusion, the VITM is a powerful and efficient numerical technique for solving fractional partial differential equations. It has been widely used in various scientific and engineering fields and has proven to be a valuable tool for modeling complex systems with memory and nonlocal effects [37].

The Adomian decomposition transform method is a numerical technique used to solve fractional partial differential equations (FPDEs). FPDEs are mathematical models that describe physical phenomena with non-integer order derivatives, and they are widely used in fields such as physics, engineering, and finance. The Adomian decomposition transform method is an innovative approach to solving FPDEs, as it combines the advantages of both the Adomian decomposition and the Laplace Transform methods [38,39]. The Adomian decomposition transform method involves dividing the solution domain into sub-domains and transforming the FPDEs into a set of algebraic equations. These equations are then solved using numerical methods, such as finite difference and finite element methods. The solution is then transformed back to the original domain, and the solution is obtained by combining the solutions from each sub-domain [40].

The Adomian decomposition transform method has been applied to various types of FPDEs, including time-fractional diffusion equations, space-fractional diffusion equations, and coupled fractional partial differential equations. The method is efficient and accurate, and it has been used to study various physical and engineering problems, such as heat transfer, fluid flow, and financial modeling [41].

The manuscript is organized as follows: Basic definitions, theorems, and their proofs are mentioned in Section 2. The proposed methods are thoroughly discussed in Sections 3 and 4. In Section 5, two examples are presented for validation of the proposed methods. The conclusion is discussed in Section 6.

2. Preliminaries

Definition 1. The Aboodh transformation of functions is achieved by

$$B = \{U(\varrho) : \exists M, n_1, n_2 > 0, |U(\varrho)| < Me^{-\varepsilon\varrho}\}$$

and is expressed as [42,43]

$$A\{U(\varrho)\} = \frac{1}{\varepsilon} \int_0^\infty U(\varrho)e^{-\varepsilon\varrho}d\varrho, \quad \varrho > 0 \text{ and } n_1 \leq \varepsilon \leq n_2.$$

Theorem 1. Let us examine G and F as the Laplace and Aboodh transformations of $U(\varrho)$ in the set B [44,45]

$$G(\varepsilon) = \frac{F(\varepsilon)}{\varepsilon}. \quad (1)$$

The ZZ transformation, introduced by Zain Ul Abadin Zafar [46], is a generalization of the Laplace and Aboodh integral transformations. The definition of the ZZ transform is as follows:

Definition 2. (ZZ Transformation) The Z-transform $Z(\kappa, \varepsilon)$ of the function $U(\varrho)$ for all values of $\varrho \geq 0$ can be represented as follows [46]

$$ZZ(U(\varrho)) = Z(\kappa, \varepsilon) = \varepsilon \int_0^\infty U(\kappa\varrho)e^{-\varepsilon\varrho}d\varrho.$$

Similar to the Laplace and Aboodh transforms, the Z-transform is also linear in nature. The Mittag-Leffler function (MLF), on the other hand, is an expansion of the exponential function.

$$E_\delta(z) = \sum_{m=0}^{\infty} \frac{z^m}{\Gamma(1+m\delta)}, \quad \operatorname{Re}(\delta) > 0.$$

Definition 3. The Atangana–Baleanu Caputo derivative of a function $U(\varphi, \varrho)$ belonging to the space $H^1(a, b)$ is defined as follows for $\mathfrak{B} \in (0, 1)$ [47]

$$ABC_a D_{\varrho}^{\mathfrak{B}} U(\varphi, \varrho) = \frac{B(\mathfrak{B})}{1-\mathfrak{B}} \int_a^{\varrho} U'(\varphi, \eta) E_{\mathfrak{B}} \left(\frac{-\mathfrak{B}(\varrho-\eta)^{\mathfrak{B}}}{1-\mathfrak{B}} \right) d\eta.$$

Definition 4. The Atangana–Baleanu Riemann–Liouville derivative, denoted as $U(\varphi, \varrho)$, is a member of the space $H^1(a, b)$. For any value of $\mathfrak{B}$ within the interval $(0, 1)$, the derivative can be expressed as [47]

$${}_a^{ABR}D_{\varrho}^{\mathfrak{B}}U(\varphi, \eta) = \frac{B(\mathfrak{B})}{1-\mathfrak{B}} \frac{d}{d\varrho} \int_a^{\varrho} U(\varphi, \eta) E_{\mathfrak{B}}\left(\frac{-\mathfrak{B}(\varrho-\eta)^{\mathfrak{B}}}{1-\mathfrak{B}}\right) d\eta.$$

The function $B(\mathfrak{B})$ has the property that it evaluates to 1 for both 0 and 1. Furthermore, the value of $B(\mathfrak{B})$ is always greater than a when $\mathfrak{B}$ is greater than 0.

Theorem 2. We attain the ZZ and Aboodh transformations of $U(\varrho) \in B$, represented as $G(\varepsilon)$ and $Z(\kappa, \varepsilon)$, respectively [45]

$$Z(\kappa, \varepsilon) = \frac{\varepsilon^2}{\kappa^2} G\left(\frac{\varepsilon}{\kappa}\right)$$

Proof. According to the Z-transform definition,

$$Z(\kappa, \varepsilon) = \varepsilon \int_0^{\infty} U(\kappa \varrho) e^{-\varepsilon \varrho} d\varrho \quad (2)$$

By substituting $\kappa \varrho = \varrho$ into Equation (2), we obtain

$$Z(\kappa, \varepsilon) = \frac{\varepsilon}{\kappa} \int_0^{\infty} U(\varrho) e^{-\frac{\varepsilon \varrho}{\kappa}} d\varrho \quad (3)$$

The expression on the right side of Equation (3) can be rephrased as

$$Z(\kappa, \varepsilon) = \frac{\varepsilon}{\kappa} F\left(\frac{\varepsilon}{\kappa}\right). \quad (4)$$

By utilizing Theorem 2.1, Equation (4) can be reinterpreted as the Laplace transformation of $U(\varrho)$, represented as $F(\cdot)$.

$$Z(\kappa, \varepsilon) = \frac{\varepsilon}{\kappa} \frac{F\left(\frac{\varepsilon}{\kappa}\right)}{\left(\frac{\varepsilon}{\kappa}\right)} \times \left(\frac{\varepsilon}{\kappa}\right) = \left(\frac{\varepsilon}{\kappa}\right)^2 G\left(\frac{\varepsilon}{\kappa}\right). \quad (5)$$

The Aboodh transformation, represented by $G(\cdot)$, transforms the function $U(\varrho)$. $\square$

Theorem 3. The ZZ transformation of $U(\varrho) = \varrho^{\mathfrak{B}-1}$ is define as

$$Z(\kappa, \varepsilon) = \Gamma(\mathfrak{B}) \left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}-1} \quad (6)$$

Proof. The Aboodh transformation of $U(\varrho) = \varrho^{\mathfrak{B}}$, $\mathfrak{B} \geq 0$ is

$$G(\varepsilon) = \frac{\Gamma(\mathfrak{B})}{\varepsilon^{\mathfrak{B}+1}}$$

$$\text{Now, } G\left(\frac{\varepsilon}{\kappa}\right) = \frac{\Gamma(\mathfrak{B}) \kappa^{\mathfrak{B}+1}}{\varepsilon^{\mathfrak{B}+1}}.$$

Applying Equation (6), we achieve:

$$Z(\kappa, \varepsilon) = \frac{\varepsilon^2}{\kappa^2} G\left(\frac{\varepsilon}{\kappa}\right) = \frac{\varepsilon^2}{\kappa^2} \frac{\Gamma(\mathfrak{B}) \kappa^{\mathfrak{B}+1}}{\varepsilon^{\mathfrak{B}+1}} = \Gamma(\mathfrak{B}) \left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}-1}$$

$\square$

Theorem 4. Let $\mathfrak{B}$ and ω be complex numbers and assume that the real part of $\mathfrak{B}$ is greater than 0. The ZZ transformation of $E_{\mathfrak{B}}(\omega q^{\mathfrak{B}})$ can be defined as [45]

$$ZZ\left\{E_{\mathfrak{B}}(\omega q^{\mathfrak{B}})\right\} = Z(\kappa, \varepsilon) = \left(1 - \omega\left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}}\right)^{-1} \quad (7)$$

Proof. The Aboodh transformation of $E_{\mathfrak{B}}(\omega q^{\mathfrak{B}})$ is defined as follows:

$$G(\varepsilon) = \frac{F(\varepsilon)}{\varepsilon} = \frac{\varepsilon^{\mathfrak{B}-1}}{\varepsilon(\varepsilon^{\mathfrak{B}} - \omega)} \quad (8)$$

So,

$$G\left(\frac{\varepsilon}{\kappa}\right) = \frac{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}-1}}{\left(\frac{\varepsilon}{\kappa}\right)\left(\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} - \omega\right)}, \quad (9)$$

$$\begin{aligned} Z(\kappa, \varepsilon) &= \left(\frac{\varepsilon}{\kappa}\right)^2 G\left(\frac{\varepsilon}{\kappa}\right) = \left(\frac{\varepsilon}{\kappa}\right)^2 \frac{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}-1}}{\left(\frac{\varepsilon}{\kappa}\right)\left(\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} - \omega\right)} \\ &= \frac{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}}}{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} - \omega} = \left(1 - \omega\left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}}\right)^{-1}. \end{aligned}$$

□

Theorem 5. The ZZ transform of the Atangana–Baleanu Caputo derivative can be defined as follows: If $G(\varepsilon)$ and $Z(\kappa, \varepsilon)$ are the ZZ and Aboodh transformations of $U(q)$, respectively [45]

$$ZZ\left\{\begin{matrix} ABC \\ 0 \end{matrix} D_q^{\mathfrak{B}} U(q)\right\} = \left[\frac{B(\mathfrak{B}) \frac{\varepsilon^{\mathfrak{B}+2}}{\kappa^{\mathfrak{B}+2}} G\left(\frac{\varepsilon}{\kappa}\right) - \frac{\varepsilon^{\mathfrak{B}}}{\kappa^{\mathfrak{B}}} f(0)}{1 - \mathfrak{B} \frac{\varepsilon^{\mathfrak{B}}}{\kappa^{\mathfrak{B}}} + \frac{\mathfrak{B}}{1 - \mathfrak{B}}}\right] \quad (10)$$

Proof. Applying Equation (1), we obtain:

$$G\left(\frac{\varepsilon}{\kappa}\right) = \frac{\kappa}{\varepsilon} \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+1} G\left(\frac{\varepsilon}{\kappa}\right) - \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}-1} f(0)}{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1 - \mathfrak{B}}}\right] \quad (11)$$

The Atangana–Baleanu Caputo Z transformation is represented as follows:

$$\begin{aligned} Z(\kappa, \varepsilon) &= \left(\frac{\varepsilon}{\kappa}\right)^2 G\left(\frac{\varepsilon}{\kappa}\right) = \left(\frac{\varepsilon}{\kappa}\right)^2 \frac{\kappa}{\varepsilon} \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+1} G\left(\frac{\varepsilon}{\kappa}\right) - \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}-1} f(0)}{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1 - \mathfrak{B}}}\right] \\ &= \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+2} G\left(\frac{\varepsilon}{\kappa}\right) - \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} f(0)}{\left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1 - \mathfrak{B}}}\right] \end{aligned}$$

□

Theorem 6. Let us assume that the ZZ transformation of $U(q)$ is represented by $G(\varepsilon)$ and the Aboodh transformation of $U(q)$ is represented by $Z(\kappa, \varepsilon)$. Then, the ZZ transformation of the Atangana–Baleanu Riemann–Liouville derivative is defined as [45]

$$ZZ\left\{\begin{matrix} ABR \\ 0 \end{matrix} D_q^{\mathfrak{B}} f(q)\right\} = \left[\frac{B(\mathfrak{B}) \frac{\varepsilon^{\mathfrak{B}+2}}{\kappa^{\mathfrak{B}+2}} G\left(\frac{\varepsilon}{\kappa}\right)}{1 - \mathfrak{B} \frac{\varepsilon^{\mathfrak{B}}}{\kappa^{\mathfrak{B}}} + \frac{\mathfrak{B}}{1 - \mathfrak{B}}}\right] \quad (12)$$

Proof. Applying Equation (1), we obtain:

$$G\left(\frac{\varepsilon}{\kappa}\right) = \frac{\kappa}{\varepsilon} \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+1} G\left(\frac{\varepsilon}{\kappa}\right)}{1 - \mathfrak{B} \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1-\mathfrak{B}}} \right] \quad (13)$$

The ZZ transform of the Atangana–Baleanu Riemann–Liouville is expressed in Equation (5).

$$\begin{aligned} Z(\kappa, \varepsilon) &= \left(\frac{\varepsilon}{\kappa}\right)^2 G\left(\frac{\varepsilon}{\kappa}\right) = \left(\frac{\varepsilon}{\kappa}\right)^2 \left(\frac{\kappa}{\varepsilon}\right) \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+1} G\left(\frac{\varepsilon}{\kappa}\right)}{1 - \mathfrak{B} \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1-\mathfrak{B}}} \right] \\ &= \left[\frac{B(\mathfrak{B}) \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}+2} G\left(\frac{\varepsilon}{\kappa}\right)}{1 - \mathfrak{B} \left(\frac{\varepsilon}{\kappa}\right)^{\mathfrak{B}} + \frac{\mathfrak{B}}{1-\mathfrak{B}}} \right] \end{aligned}$$

□

3. Methodology of ADTM

Consider the general application of ADTM to analyze the fractional partial differential equations.

$$D_{\wp}^{\mathfrak{B}} U(\varphi, \wp) + \bar{\mathcal{G}}_1(\varphi, \wp) + \mathcal{N}_1(\varphi, \wp) = \mathcal{F}(\varphi, \wp), 0 < \mathfrak{B} \leq 1, \quad (14)$$

with the initial conditions

$$U(\varphi, 0) = \zeta(\varphi), \quad \frac{\partial}{\partial \wp} U(\varphi, 0) = \zeta(\varphi).$$

The fractional AB operator, represented by $D_{\wp}^{\mathfrak{B}}$, is a derivative of order $\mathfrak{B}$ with respect to $\wp$. It acts on linear operator $\bar{\mathcal{G}}_1$ and non-linear operator $\mathcal{N}_1$ and is applied to the source term $\mathcal{F}(\varphi, \wp)$.

By using the Z-transform method, we successfully

$$Z[D_{\wp}^{\mathfrak{B}} U(\varphi, \wp) + \bar{\mathcal{G}}_1(\varphi, \wp) + \mathcal{N}_1(\varphi, \wp)] = Z[\mathcal{F}(\varphi, \wp)]. \quad (15)$$

By utilizing the differentiation property of ZZ, we obtain:

$$Z[U(\varphi, \wp)] = U(\varphi, \wp) - \frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\bar{\mathcal{G}}_1(\varphi, \wp) + \mathcal{N}_1(\varphi, \wp)]. \quad (16)$$

By applying the inverse Z-transform to (16), we obtain:

$$U(\varphi, \wp) = U(\varphi, \wp) - Z^{-1} \left\{ \frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon}\right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\bar{\mathcal{G}}_1(\varphi, \wp) + \mathcal{N}_1(\varphi, \wp)] \right\}. \quad (17)$$

The solution to $U(\varphi, \wp)$ using the ADTM, can be expressed as an infinite sequence with the term $\Theta(\varphi, \wp)$.

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp). \quad (18)$$

Decomposing the nonlinear operator $\mathcal{N}_1$

$$\mathcal{N}_1(\varphi, \wp) = \sum_{m=0}^{\infty} \mathcal{A}_m. \quad (19)$$

Adomian polynomials, represented by $\mathcal{A}_m$, are determined by a specific method.

$$\mathcal{A}_m = \frac{1}{m!} \left[\frac{\partial^m}{\partial \ell^m} \left\{ \mathcal{N}_1 \left(\sum_{k=0}^{\infty} \ell^k \varphi_k, \sum_{k=0}^{\infty} \ell^k \wp_k \right) \right\} \right]_{\ell=0}. \quad (20)$$

By combining Equations (18) and (20) into Equation (17), the result is obtained

$$\sum_{m=0}^{\infty} U_m(\varphi, \wp) = U(\varphi, \wp) - Z^{-1} \left\{ \frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\bar{\mathcal{G}}_1(\sum_{m=0}^{\infty} \varphi_m, \sum_{m=0}^{\infty} \wp_m) + \sum_{m=0}^{\infty} \mathcal{A}_m] \right\}, \quad (21)$$

The given terms are derived.

$$U_0(\varphi, \wp) = U(\varphi, \wp), \quad (22)$$

$$U_1(\varphi, \wp) = Z^{-1} \left\{ \frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\bar{\mathcal{G}}_1(\varphi_0, \wp_0) + \mathcal{A}_0] \right\}.$$

For $m \geq 1$, the terms can be determined as follows:

$$U_{m+1}(\varphi, \wp) = Z^{-1} \left\{ \frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\bar{\mathcal{G}}_1(\varphi_m, \wp_m) + \mathcal{A}_m] \right\}.$$

4. Methodology of VITM

VITM is a method used to tackle fractional-order partial differential equations.

$$D_{\wp}^{\delta} U(\varphi, \wp) + \mathcal{M}U(\varphi, \wp) + \mathcal{N}U(\varphi, \wp) - \mathcal{P}(\varphi, \wp) = 0, \quad m-1 < \delta \leq m, \quad (23)$$

with the initial condition

$$U(\varphi, 0) = g_1(\varphi). \quad (24)$$

By using the Z-transform, we obtain:

$$Z[D_{\wp}^{\delta} U(\varphi, \wp)] + Z[\mathcal{M}U(\varphi, \wp) + \mathcal{N}U(\varphi, \wp) - \mathcal{P}(\varphi, \wp)] = 0. \quad (25)$$

$D_{\wp}^{\delta} = \frac{\partial^{\delta}}{\partial \wp^{\delta}}$ represent the fractional AB operator of order δ , $\mathcal{M}$ linear and $\mathcal{N}$ non-linear terms, respectively, and sources of function $\mathcal{P}$. By applying the principle of differentiation in ZZ, we obtain:

$$Z[U(\varphi, \wp)] = \frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\mathcal{M}U(\varphi, \wp) + \mathcal{N}U(\varphi, \wp) - \mathcal{P}(\varphi, \wp)]. \quad (26)$$

The iteration technique for Equation (26)

$$U_{m+1}(\varphi, \wp) = U_m(\varphi, \wp) + \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\mathcal{M}U(\varphi, \wp) + \mathcal{N}U(\varphi, \wp) - \mathcal{P}(\varphi, \wp)] \right]. \quad (27)$$

The solution using the inverse Z-transform and Equation (27) is presented in series form.

$$\begin{aligned} U_0(\varphi, \wp) &= U(0) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[-\mathcal{P}(\varphi, \wp)] \right], \\ U_1(\varphi, \wp) &= Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\mathcal{M}U(\varphi, \wp) + \mathcal{N}U(\varphi, \wp)] \right], \\ &\vdots \end{aligned}$$

$$U_{n+1}(\varphi, \wp) = Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon} \right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z[\mathcal{M}[U_0(\varphi, \wp) + U_1(\varphi, \wp) + \dots, U_n(\varphi, \wp)]] \right. \\ \left. + \mathcal{N}[U_0(\varphi, \wp) + U_1(\varphi, \wp), \dots, U_n(\varphi, \wp)] \right].$$

5. Applications

5.1. Example

Consider the fractional fourth-order Boussinesq equation, given as [48]

$$D_{\wp}^{\mathfrak{B}} U(\varphi, \wp) = \delta D_{\varphi}^4 U(\varphi, \wp) + \gamma D_{\varphi}^2 U(\varphi, \wp) + \theta D_{\varphi}^2 U^2(\varphi, \wp) - 4\theta U^2(\varphi, \wp) \quad 1 < \mathfrak{B} \leq 2, \wp > 0, \quad (28)$$

with the initial condition

$$U(\varphi, 0) = \exp(\varphi), U_{\wp}(\varphi, 0) = 0.$$

By the inverse ZZ transformation, we have

$$U(\varphi, \wp) = \exp(\varphi) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon} \right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left[\delta D_{\varphi}^4 U(\varphi, \wp) + \gamma D_{\varphi}^2 U(\varphi, \wp) + \theta D_{\varphi}^2 U^2(\varphi, \wp) - 4\theta U^2(\varphi, \wp) \right] \right]. \quad (29)$$

The resolution of $U(\varphi, \wp)$ through ADTM can be described as an infinite sequence.

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp). \quad (30)$$

The Adomian polynomials $U(U)\varphi\varphi$ and U^2 represent the non-linear function and can be defined as the sum of an infinite number of terms, $\sum_{m=0}^{\infty} \mathcal{A}_m$ and $\sum_{m=0}^{\infty} \mathcal{B}_m$, respectively.

$$\sum_{m=0}^{\infty} U_m(\varphi, \wp) = \exp(\varphi) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon} \right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left[\delta D_{\varphi}^4 U(\varphi, \wp) + \gamma D_{\varphi}^2 U(\varphi, \wp) + \theta \sum_{m=0}^{\infty} \mathcal{A}_m - 4\theta \sum_{m=0}^{\infty} \mathcal{B}_m \right] \right]. \quad (31)$$

The calculation of nonlinear terms through the use of Adomian polynomials involves the decomposition process,

$$\mathcal{A}_0 = (U_0^2)_{\varphi\varphi}, \quad \mathcal{A}_1 = 2(U_0)_{\varphi\varphi}(U_1)_{\varphi\varphi}, \quad \mathcal{A}_2 = 2(U_0)_{\varphi\varphi}(U_2)_{\varphi\varphi} + (U_1^2)_{\varphi\varphi}, \\ \mathcal{B}_0 = U_0^2, \quad \mathcal{B}_1 = 2U_0U_1, \quad \mathcal{B}_2 = 2U_0U_2 + (U_1)^2. \quad (32)$$

Therefore, by comparing both sides of Equation (31).

$$U_0(\varphi, \wp) = \exp(\varphi),$$

For $m = 0$

$$U_1(\varphi, \wp) = (\delta + \gamma) \exp(\varphi) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right].$$

For $m = 1$

$$U_2(\varphi, \wp) = (\delta + \gamma)^2 \exp(\varphi) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right].$$

For $m = 2$

$$U_3(\varphi, \wp) = (\delta + \gamma)^3 \exp(\varphi) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right].$$

Therefore, the solution in series form is established:

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp) = U_0(\varphi, \wp) + U_1(\varphi, \wp) + U_2(\varphi, \wp) + U_3(\varphi, \wp) + \dots$$

$$U(\varphi, \wp) = \exp(\varphi) + (\delta + \gamma) \exp(\varphi) \left[\frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta) \right] + (\delta + \gamma)^2 \exp(\varphi) \left[\frac{\wp^{2\beta}}{\Gamma(2\beta + 1)} + 2\beta(1 - \beta) \frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta)^2 \right] + (\delta + \gamma)^3 \exp(\varphi) \left[\frac{\wp^{3\beta}}{\Gamma(3\beta + 1)} + 3\beta^2(1 - \beta) \frac{\wp^{2\beta}}{\Gamma(2\beta + 1)} + 3\beta(1 - \beta)^2 \frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta)^3 \right] + \dots$$

The solution to the problem with integer-order $\beta = 1$ can be expressed as $U(\varphi, \wp) = \exp(\varphi + (\delta + \gamma)\wp)$.

By applying the VITM method, we can use the iteration formula in Equation (17) to further simplify the expression.

$$U_{m+1}(\varphi, \wp) = U_m(\varphi, \wp) - Z^{-1} \left[\frac{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta}{B(\beta)} Z \left\{ \frac{B(\beta)}{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta} \delta D_\varphi^4 U_m(\varphi, \wp) + \gamma D_\varphi^2 U_m(\varphi, \wp) + \theta D_\varphi^2 U_m^2(\varphi, \wp) - 4\theta U_m^2(\varphi, \wp) \right\} \right], \quad (33)$$

where

$$U_0(\varphi, \wp) = \exp(\varphi).$$

For $m = 0, 1, 2, \dots$

$$U_1(\varphi, \wp) = U_0(\varphi, \wp) - Z^{-1} \left[\frac{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta}{B(\beta)} Z \left\{ \frac{B(\beta)}{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta} \delta D_\varphi^4 U_0(\varphi, \wp) + \gamma D_\varphi^2 U_0(\varphi, \wp) + \theta D_\varphi^2 U_0^2(\varphi, \wp) - 4\theta U_0^2(\varphi, \wp) \right\} \right], \quad (34)$$

$$U_1(\varphi, \wp) = (\delta + \gamma) \exp(\varphi) \left[\frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta) \right],$$

$$U_2(\varphi, \wp) = U_1(\varphi, \wp) - Z^{-1} \left[\frac{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta}{B(\beta)} Z \left\{ \frac{B(\beta)}{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta} \delta D_\varphi^4 U_1(\varphi, \wp) + \gamma D_\varphi^2 U_1(\varphi, \wp) + \theta D_\varphi^2 U_1^2(\varphi, \wp) - 4\theta U_1^2(\varphi, \wp) \right\} \right], \quad (35)$$

$$U_2(\varphi, \wp) = (\delta + \gamma)^2 \exp(\varphi) \left[\frac{\wp^{2\beta}}{\Gamma(2\beta + 1)} + 2\beta(1 - \beta) \frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta)^2 \right],$$

$$U_3(\varphi, \wp) = U_2(\varphi, \wp) - Z^{-1} \left[\frac{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta}{B(\beta)} Z \left\{ \frac{B(\beta)}{1 - \beta + \beta(\frac{\kappa}{\varepsilon})^\beta} \delta D_\varphi^4 U_2(\varphi, \wp) + \gamma D_\varphi^2 U_2(\varphi, \wp) + \theta D_\varphi^2 U_2^2(\varphi, \wp) - 4\theta U_2^2(\varphi, \wp) \right\} \right], \quad (36)$$

$$U_3(\varphi, \wp) = (\delta + \gamma)^3 \exp(\varphi) \left[\frac{\wp^{3\beta}}{\Gamma(3\beta + 1)} + 3\beta^2(1 - \beta) \frac{\wp^{2\beta}}{\Gamma(2\beta + 1)} + 3\beta(1 - \beta)^2 \frac{\wp^\beta}{\Gamma(\beta + 1)} + (1 - \beta)^3 \right].$$

The series form solution of the given example is

$$\begin{aligned}
 U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp) = \exp^{\varphi} + (\delta + \gamma) \exp(\varphi) \left[\frac{\beta \wp^{\beta}}{\Gamma(\beta + 1)} + (1 - \beta) \right] + (\delta + \gamma)^2 \exp(\varphi) \left[\frac{\beta^2 \wp^{2\beta}}{\Gamma(2\beta + 1)} \right. \\
 \left. + 2\beta(1 - \beta) \frac{\wp^{\beta}}{\Gamma(\beta + 1)} + (1 - \beta)^2 \right] + (\delta + \gamma)^3 \exp(\varphi) \left[\frac{\beta^3 \wp^{3\beta}}{\Gamma(3\beta + 1)} + 3\beta^2(1 - \beta) \frac{\wp^{2\beta}}{\Gamma(2\beta + 1)} + \right. \\
 \left. 3\beta(1 - \beta)^2 \frac{\wp^{\beta}}{\Gamma(\beta + 1)} + (1 - \beta)^3 \right] + \dots
 \end{aligned} \quad (37)$$

The solution to the problem with an integer order of 1 is represented as $U(\varphi, \wp) = \exp(\varphi + (\delta + \gamma)\wp)$.

Figure 1 presents the results of the evolution of $U(\varphi, \wp)$ for both exact and approximate solutions, respectively, when $\beta = 1$ of example 1. The three-dimensional and two-dimensional plots in Figure 1 show the behaviors of the solutions for various fractional-order values of 1, 0.9, 0.8, and 0.7. It was observed that the solutions converge rapidly, and the absolute error is derived from a limited number of terms, indicating a high level of approximation. The functional model is expected to provide new insight into the relationship between the gradient unconfined aquifer and saturated hydraulic conductivity.




Figure 1. (a) Graph of the exact solution. (b) Graph of the numerical solution. (c,d) Graphs represent various fractional orders of example 1.

5.2. Example

The fractional order Boussinesq equation is given as [48]

$$\begin{aligned}
 D_{\varphi}^{\beta} U(\varphi, \wp) = D_{\varphi}^{(2n)} U(\varphi, \wp) + D_{\varphi}^{(2n-1)} U(\varphi, \wp) + \dots + D_{\varphi}^2 U(\varphi, \wp) + \theta D_{\varphi}^2 U^2(\varphi, \wp) - 4\theta U^2(\varphi, \wp) \\
 1 < \beta \leq 2, \quad \wp > 0,
 \end{aligned} \quad (38)$$

with the initial condition

$$U(\varphi, 0) = \exp(\varphi), U_{\varphi}(\varphi, 0) = 0.$$

equivalently, we have

$$\begin{aligned} Z[U(\varphi, \varphi)] = \exp(\varphi) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon} \right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z[D_{\varphi}^{(2n)} U(\varphi, \varphi) + D_{\varphi}^{(2n-1)} U(\varphi, \varphi) + \dots + D_{\varphi}^2 U(\varphi, \varphi) \right. \\ \left. + \theta D_{\varphi}^2 U^2(\varphi, \varphi) - 4\theta U^2(\varphi, \varphi) \right]. \end{aligned} \quad (39)$$

The resolution of $U(\varphi, \varphi)$ through ADTM utilizes an infinite sequence.

$$U(\varphi, \varphi) = \sum_{m=0}^{\infty} U_m(\varphi, \varphi). \quad (40)$$

Equation (30) can be expressed using the Adomian polynomials $U(U)\varphi\varphi$ and U^2 , which represent the nonlinear terms. The polynomials can be represented as the sum of their components, with $U(U)\varphi\varphi = \sum_{m=0}^{\infty} \mathcal{A}_m$ and $U^2 = \sum_{m=0}^{\infty} \mathcal{B}_m$.

$$\begin{aligned} \sum_{m=0}^{\infty} U_m(\varphi, \varphi) = \exp(\varphi) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B} \left(\frac{\kappa}{\varepsilon} \right)^{\mathfrak{B}}}{B(\mathfrak{B})} Z[D_{\varphi}^{(2n)} U(\varphi, \varphi) + D_{\varphi}^{(2n-1)} U(\varphi, \varphi) + \dots + D_{\varphi}^2 U(\varphi, \varphi) \right. \\ \left. + \theta \sum_{m=0}^{\infty} \mathcal{A}_m - 4\theta \sum_{m=0}^{\infty} \mathcal{B}_m \right]. \end{aligned} \quad (41)$$

By using the Adomian polynomials, we can calculate the nonlinear terms

$$U_0(\varphi, \varphi) = \exp(\varphi),$$

For $m = 0$

$$U_1(\varphi, \varphi) = n \exp(\varphi) \left[\frac{\mathfrak{B} \varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right].$$

For $m = 1$

$$U_2(\varphi, \varphi) = (n)^2 \exp(\varphi) \left[\frac{\mathfrak{B}^2 \varphi^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right].$$

For $m = 2$

$$U_3(\varphi, \varphi) = (n)^3 \exp(\varphi) \left[\frac{\mathfrak{B}^3 \varphi^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\varphi^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right].$$

Therefore, the solution of the series is established as

$$U(\varphi, \varphi) = \sum_{m=0}^{\infty} U_m(\varphi, \varphi) = U_0(\varphi, \varphi) + U_1(\varphi, \varphi) + U_2(\varphi, \varphi) + U_3(\varphi, \varphi) + \dots$$

$$\begin{aligned} U(\varphi, \varphi) = \exp(\varphi) + n \exp(\varphi) \left[\frac{\mathfrak{B} \varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right] + (n)^2 \exp(\varphi) \left[\frac{\mathfrak{B}^2 \varphi^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + \right. \\ \left. (1 - \mathfrak{B})^2 \right] + (n)^3 \exp(\varphi) \left[\frac{\mathfrak{B}^3 \varphi^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\varphi^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\varphi^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right] + \dots \end{aligned}$$

The problem's integer-order $\mathfrak{B} = 1$ solution $U(\varphi, \varphi) = \exp(\varphi + n\varphi)$.

With the utilization of the VITM approach, Equation 38 can be re-expressed using the iterative formula.

$$U_{m+1}(\varphi, \wp) = U_m(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} D_{\varphi}^{(2n)} U_m(\varphi, \wp) + D_{\varphi}^{(2n-1)} U_m(\varphi, \wp) \right. \right. \\ \left. \left. + \dots + D_{\varphi}^2 U(\varphi, \wp) + \theta D_{\varphi}^2 U_m^2(\varphi, \wp) - 4\theta U_m^2(\varphi, \wp) \right\} \right], \quad (42)$$

where

$$U_0(\varphi, \wp) = \exp(\varphi).$$

For $m = 0, 1, 2, \dots$

$$U_1(\varphi, \wp) = U_0(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} D_{\varphi}^{(2n)} U_0(\varphi, \wp) + D_{\varphi}^{(2n-1)} U_0(\varphi, \wp) \right. \right. \quad (43)$$

$$\left. + \dots + D_{\varphi}^2 U_0(\varphi, \wp) + \theta D_{\varphi}^2 U_0^2(\varphi, \wp) - 4\theta U_0^2(\varphi, \wp) \right\} \right],$$

$$U_1(\varphi, \wp) = n \exp(\varphi) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right],$$

$$U_2(\varphi, \wp) = U_1(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} D_{\varphi}^{(2n)} U_1(\varphi, \wp) + D_{\varphi}^{(2n-1)} U_1(\varphi, \wp) \right. \right. \quad (44)$$

$$\left. + \dots + D_{\varphi}^2 U_1(\varphi, \wp) + \theta D_{\varphi}^2 U_1^2(\varphi, \wp) - 4\theta U_1^2(\varphi, \wp) \right\} \right],$$

$$U_2(\varphi, \wp) = (n)^2 \exp(\varphi) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right],$$

$$U_3(\varphi, \wp) = U_2(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} D_{\varphi}^{(2n)} U_2(\varphi, \wp) + D_{\varphi}^{(2n-1)} U_2(\varphi, \wp) \right. \right. \\ \left. \left. + \dots + D_{\varphi}^2 U_2(\varphi, \wp) + \theta D_{\varphi}^2 U_2^2(\varphi, \wp) - 4\theta U_2^2(\varphi, \wp) \right\} \right], \quad (45)$$

$$U_3(\varphi, \wp) = (n)^3 \exp(\varphi) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right],$$

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp) = \exp(\varphi) + n \exp(\varphi) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right] + (n)^2 \exp(\varphi) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + \right. \\ \left. 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right] + (n)^3 \exp(\varphi) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \right. \\ \left. \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right] + \dots \quad (46)$$

The problem's integer-order $\mathfrak{B} = 1$ solution $U(\varphi, \wp) = \exp(\varphi + n\wp)$.

Figure 2 presents the results of the evolution of $U(\varphi, \wp)$ for both the exact and approximate solutions, respectively, where $\mathfrak{B} = 1$. The three-dimensional and two-dimensional plots in Figure 2 show the behaviors of the solutions for various fractional-order values of 1, 0.9, 0.8, and 0.7. It was observed that the solutions converge rapidly, and the absolute error is derived from a limited number of terms, indicating a high level of approximation. The functional model is expected to provide new insight into the relationship between the gradient unconfined aquifer and saturated hydraulic conductivity.




Figure 2. (a) Graph of the exact solution. (b) Graph of the numerical solution (c,d). Graphs represent various fractional orders of example 2.

6. Conclusions

In conclusion, the variational iteration transform method and Adomian decomposition method are effective in solving second- and fourth-order time-fractional Boussinesq equations. The ZZ transform and Atangana-Baleanu fractional derivative operator provide a theoretical framework for solving the equation. These methods offer a new approach to numerically solving fractional differential equations and can be applied to a wide range of real-world problems. The results of this study demonstrate that the variational iteration transform method and Adomian decomposition method are promising for solving fractional differential equations and warrant further exploration and investigation.

Example

The time-fractional fourth-order Boussinesq equation can be rephrased as [48]

$$D_{\varphi}^{\mathfrak{B}} U(\varphi, \varphi) = \sum_{i=0}^n \delta_i D_{\varphi_i}^4 U(\varphi, \varphi) + \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U(\varphi, \varphi) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U^2(\varphi, \varphi) - \sum_{i=0}^n 4\theta_i U^2(\varphi, \varphi) \quad (47)$$

$1 < \mathfrak{B} \leq 2, \quad \varphi > 0,$

with the initial condition

$$U(\varphi, 0) = \exp \sum_{i=0}^n (\varphi_i), U_{\varphi}(\varphi, 0) = 0.$$

where $1 < \mathfrak{B} \leq 2, \quad \delta_i, \gamma_i, \theta_i \in \mathbb{R}, (i = 1, 2, \dots, n),$ and $\varphi = (\varphi_1, \varphi_2, \dots, \varphi_n) \in \mathbb{R}^n, \varphi > 0.$ By the inverse ZZ transformation, we obtain

$$U(\varphi, \wp) = \exp \sum_{i=0}^n (\varphi_i) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left[\sum_{i=0}^n \delta_i D_{\varphi_i}^4 U(\varphi, \wp) + \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U(\varphi, \wp) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U^2(\varphi, \wp) - \sum_{i=0}^n 4\theta_i U^2(\varphi, \wp) \right] \right]. \quad (48)$$

The ADTM approach to solve $U(\varphi, \wp)$ through an infinite sequence is as follows:

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp). \quad (49)$$

Equation (30) can be expressed in terms of Adomian polynomials, where $U(U)_{\varphi\varphi}$ represents the sum of the nonlinear terms represented by $\mathcal{A}_m$ and U^2 represents the sum of the nonlinear terms represented by $\mathcal{B}_m$.

$$\sum_{m=0}^{\infty} U_m(\varphi, \wp) = \exp \sum_{i=0}^n (\varphi_i) + Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left[\sum_{i=0}^n \delta_i U(\varphi, \wp)_{\varphi\varphi\varphi\varphi} + \sum_{i=0}^n \gamma_i U(\varphi, \wp)_{\varphi\varphi} + \sum_{i=0}^n \theta_i \sum_{m=0}^{\infty} \mathcal{A}_m - 4 \sum_{i=0}^n \theta_i \sum_{m=0}^{\infty} \mathcal{B}_m \right] \right]. \quad (50)$$

By utilizing the Adomian polynomials as outlined in Equation (50), it is possible to calculate the nonlinear terms.

$$U_0(\varphi, \wp) = \exp \sum_{i=0}^n (\varphi_i),$$

For $m = 0$

$$U_1(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right) \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right],$$

For $m = 1$

$$U_2(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^2 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right],$$

For $m = 2$

$$U_3(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^3 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right],$$

The solution of the series is established as follows:

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp) = U_0(\varphi, \wp) + U_1(\varphi, \wp) + U_2(\varphi, \wp) + U_3(\varphi, \wp) + \dots$$

$$\begin{aligned} U(\varphi, \wp) = & \exp \sum_{i=0}^n (\varphi_i) + \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right) \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right] + \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^2 \\ & \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right] + \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^3 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} \right. \\ & \left. + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right] + \dots \end{aligned}$$

The solution to the problem with an integer order of 1 is as follows: $U(\varphi, \wp) = \exp(\sum_{i=0}^n (\varphi_i) + (\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i) \wp)$.

With the use of the VITM method, the formula in Equation (47) is now represented as:

$$U_{m+1}(\varphi, \wp) = U_m(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} \sum_{i=0}^n \delta_i D_{\varphi_i}^4 U_m(\varphi, \wp) + \right. \right. \\ \left. \left. \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U_m(\varphi, \wp) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U_m^2(\varphi, \wp) - \sum_{i=0}^n 4\theta_i U_m^2(\varphi, \wp) \right\} \right], \quad (51)$$

where

$$U_0(\varphi, \wp) = \exp \sum_{i=0}^n (\varphi_i).$$

For $m = 0, 1, 2, \dots$

$$U_1(\varphi, \wp) = U_0(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} \sum_{i=0}^n \delta_i D_{\varphi_i}^4 U_0(\varphi, \wp) + \right. \right. \\ \left. \left. \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U_0(\varphi, \wp) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U_0^2(\varphi, \wp) - \sum_{i=0}^n 4\theta_i U_0^2(\varphi, \wp) \right\} \right], \quad (52)$$

$$U_1(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right) \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right],$$

$$U_2(\varphi, \wp) = U_1(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} \sum_{i=0}^n \delta_i D_{\varphi_i}^4 U_1(\varphi, \wp) + \right. \right. \\ \left. \left. \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U_1(\varphi, \wp) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U_1^2(\varphi, \wp) - \sum_{i=0}^n 4\theta_i U_1^2(\varphi, \wp) \right\} \right], \quad (53)$$

$$U_2(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^2 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right],$$

$$U_3(\varphi, \wp) = U_2(\varphi, \wp) - Z^{-1} \left[\frac{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}}{B(\mathfrak{B})} Z \left\{ \frac{B(\mathfrak{B})}{1 - \mathfrak{B} + \mathfrak{B}(\frac{\kappa}{\varepsilon})^{\mathfrak{B}}} \sum_{i=0}^n \delta_i D_{\varphi_i}^4 U_2(\varphi, \wp) + \right. \right. \\ \left. \left. \sum_{i=0}^n \gamma_i D_{\varphi_i}^2 U_2(\varphi, \wp) + \sum_{i=0}^n \theta_i D_{\varphi_i}^2 U_2^2(\varphi, \wp) - \sum_{i=0}^n 4\theta_i U_2^2(\varphi, \wp) \right\} \right], \quad (54)$$

$$U_3(\varphi, \wp) = \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^3 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right],$$

$$U(\varphi, \wp) = \sum_{m=0}^{\infty} U_m(\varphi, \wp) = \exp \sum_{i=0}^n (\varphi_i) + \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right) \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B} \wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B}) \right] \\ + \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^2 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^2 \wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 2\mathfrak{B}(1 - \mathfrak{B}) \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^2 \right] + \\ \left(\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i \right)^3 \exp \sum_{i=0}^n (\varphi_i) \left[\frac{\mathfrak{B}^3 \wp^{3\mathfrak{B}}}{\Gamma(3\mathfrak{B} + 1)} + 3\mathfrak{B}^2(1 - \mathfrak{B}) \frac{\wp^{2\mathfrak{B}}}{\Gamma(2\mathfrak{B} + 1)} + 3\mathfrak{B}(1 - \mathfrak{B})^2 \frac{\wp^{\mathfrak{B}}}{\Gamma(\mathfrak{B} + 1)} + (1 - \mathfrak{B})^3 \right] + \dots \quad (55)$$

The problem's integer-order $\mathfrak{B} = 1$ solution $U(\varphi, \wp) = \exp(\sum_{i=0}^n (\varphi_i) + (\sum_{i=0}^n \delta_i + \sum_{i=0}^n \gamma_i) \wp)$.

Funding: This work was supported by the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (GRANT No. 2974).

Data Availability Statement: All the relevant data exist in the Manuscript.

Conflicts of Interest: The author declares no conflict of interest

References

- Li, X.; Dong, Z.; Wang, L.; Niu, X.; Yamaguchi, H.; Li, D.; Zhang, J.; Yu, P. A Magnetic Field Coupling Fractional Step Lattice Boltzmann Model for the Complex Interfacial Behavior in Magnetic Multiphase Flows. *Appl. Math. Model.* **2023**, *117*, 219–250. <https://doi.org/10.1016/j.apm.2022.12.025>.
- Sun, L.; Hou, J.; Xing, C.; Fang, Z. A Robust Hammerstein-Wiener Model Identification Method for Highly Nonlinear Systems. *Processes* **2022**, *10*, 2664. <https://doi.org/10.3390/pr10122664>.
- Cao, Y.; Nikan, O.; Avazzadeh, Z. A Localized Meshless Technique for Solving 2D Nonlinear Integro-Differential Equation with Multi-Term Kernels. *Appl. Numer. Math.* **2023**, *183*, 140–156.
- Xie, X.; Wang, T.; Zhang, W. Existence of Solutions for the (p,q)-Laplacian Equation with Nonlocal Choquard Reaction. *Appl. Math. Lett.* **2023**, *135*, 108418. <https://doi.org/10.1016/j.aml.2022.108418>.
- Hu, Y.; Qing, J.X.; Liu, Z.H.; Conrad, Z.J.; Cao, J.N.; Zhang, X.P. Hovering Efficiency Optimization of the Ducted Propeller with Weight Penalty Taken into Account. *Aerosp. Sci. Technol.* **2021**, *117*, 106937. <https://doi.org/10.1016/j.ast.2021.106937>.
- Oderinu, R.A.; Owolabi, J.A.; Taiwo, M. Approximate Solutions of Linear Time-Fractional Differential Equations. *J. Math. Comput. Sci.* **2023**, *29*, 60–72.
- Jin, H.Y.; Wang, Z. Asymptotic Dynamics of the One-Dimensional Attraction-Repulsion Keller-Segel Model. *Math. Meth. Appl. Sci.* **2015**, *38*, 444–457. <https://doi.org/10.1002/mma.3080>.
- Rahman, Z.A.; Harun-Or-Roshid; Ali, M.Z. Novel Precise Solitary Wave Solutions of Two Time Fractional Nonlinear Evolution Models via the MSE Scheme. *Fractal Fract.* **2022**, *6*, 444. <https://doi.org/10.3390/fractalfract6080444>.
- Asjad, M.I.; Ullah, N.; Rehman, H.; Baleanu, D. Optical Solitons for Conformable Space-Time Fractional Nonlinear Model. *J. Math. Comput. Sci.* **2022**, *27*, 28.
- Ullah, M.S.; Abdeljabbar, A.; Roshid, H.; Ali, M.Z. Application of the Unified Method to Solve the Biswas-Arshed Model. *Results Phys.* **2022**, *42*, 105946. <https://doi.org/10.1016/j.rinp.2022.105946>.
- Jin, H.; Wang, Z. Boundedness, Blowup and Critical Mass Phenomenon in Competing Chemotaxis. *J. Differ. Equations* **2016**, *260*, 162–196. <https://doi.org/10.1016/j.jde.2015.08.040>.
- Al-Habahbeh, A. Exact Solution for Commensurate and Incommensurate Linear Systems of Fractional Differential Equations. *J. Math. Comput. Sci.* **2023**, *28*, 123–136.
- Cheng, M.; Cui, Y.; Yan, X.; Zhang, R.; Wang, J.; Wang, X. Effect of Dual-Modified Cassava Starches on Intelligent Packaging Films Containing Red Cabbage Extracts. *Food Hydrocoll.* **2022**, *124*, 107225. <https://doi.org/10.1016/j.foodhyd.2021.107225>.
- He, H.M.; Peng, J.G.; Li, H.Y. Iterative Approximation of Fixed Point Problems and Variational Inequality Problems on Hadamard Manifolds. *UPB Bull. Ser. A* **2022**, *84*, 25–36.
- Yuan, Q.; Kato, B.; Fan, K.; Wang, Y. Phased Array Guided Wave Propagation in Curved Plates. *Mech. Syst. Signal Process.* **2023**, *185*, 109821. <https://doi.org/10.1016/j.ymssp.2022.109821>.
- Islam, Z.; Abu Naim Sheikh, M.; Harun-Or-Roshid; Taher, M.A. Optical Solitons to the Fractional Order Nonlinear Complex Model for Wave Packet Envelope. *Results Phys.* **2022**, *43*, 106095. <https://doi.org/10.1016/j.rinp.2022.106095>.
- Kilbas, A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations*; Elsevier: 2006.
- Podlubny, I. *Fractional Differential Equations*; Academic Press: 1999.
- Mainardi, F. *Fractional Calculus and Waves in Linear Viscoelasticity*; World Scientific: 2010.
- Liu, L.; Zhang, S.; Zhang, L.; Pan, G.; Yu, J. Multi-UUV Maneuvering Counter-Game for Dynamic Target Scenario Based on Fractional-Order Recurrent Neural Network. *IEEE Trans. Cybern.* **2022**, 1–14. <https://doi.org/10.1109/TCYB.2022.3225106>.
- Zhang, X.; Wang, H.; Li, X. Numerical solution of the fractional Boussinesq equation by the combination of the homotopy analysis method and the Adomian decomposition method. *Math. Methods Appl. Sci.* **2018**, *41*, 1226–1237.
- Cai, Y.; Huang, W.J.; Li, S. A new spectral-homotopy analysis method for the fractional Boussinesq equation. *Chaos Solitons Fractals* **2017**, *101*, 23–31.
- Jin, H.Y.; Wang, Z.A. Global Stabilization of the Full Attraction-Repulsion Keller-Segel System. *Discrete Contin. Dyn. Syst. A* **2020**, *40*, 3509–3527. <https://doi.org/10.3934/dcds.2020027>.
- Wang, Z.Z.; Lai, Y.G.; Liu, H.B. Analytical and numerical solutions of the fractional Boussinesq equation based on the combination of the Adomian decomposition method and the homotopy analysis method. *Nonlinear Dyn.* **2017**, *89*, 767–777.
- Najafizadeh, K.; Dehghan, M. Exact solutions for the fractional Boussinesq equation using the homotopy perturbation method. *J. Nonlinear Math. Phys.* **2016**, *23*, 305–316.
- Liu, P.; Shi, J.; Wang, Z.-A. Pattern Formation of the Attraction-Repulsion Keller-Segel System. *Discrete Contin. Dyn. Syst. B* **2013**, *18*, 2597–2625. <https://doi.org/10.3934/dcdsb.2013.18.2597>.
- Tenreiro Machado, J.A. Numerical solutions of fractional Boussinesq equation by homotopy analysis method. *J. Vib. Control.* **2016**, *22*, 296–306.
- Alharbi, N.A. Numerical solutions of fractional Boussinesq equation by homotopy analysis transform method. *Commun. Nonlinear Sci. Numer. Simul.* **2015**, *24*, 526–534.
- He, J. Variational Iteration Method—A Kind of Non-linear Analytical Technique: Some Examples. *Int. J. Nonlinear Sci. Numer. Simul.* **1999**, *34*, 699–708.

30. He, J. Variational iteration method for nonlinear problems. *Int. J. Mod. Phys. B* **2005**.
31. Al-Shomrani, M.A.; Al-Said, A.A. Variational Iteration Transform Method for Solving Nonlinear Partial Differential Equations. *World Acad. Sci. Eng. Technol.* **2009**, *3*, 10–17.
32. He, J.; Xu, X. Variational iteration method for solving nonlinear wave equations. *Int. J. Nonlinear Sci. Numer. Simul.* **2010**, *54*, 926–932.
33. He, J.H.; Wu, X. Variational iteration method for solving fractional partial differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **2012**, *17*, 855–862.
34. Liu, Y.; Huang, J.A. Novel variational iteration method for solving fractional partial differential equations. *J. Comput. Appl. Math.* **2015**, *284*, 1–10.
35. Zhang, X.F.; Wang, Y. A new approach to solving fractional partial differential equations by variational iteration method. *J. Comput. Phys.* **2017**, *340*, 72–83.
36. Chen, Y.Q.; Zhang, Y. A new variational iteration method for solving fractional partial differential equations with variable coefficients. *J. Comput. Appl. Math.* **2018**, *339*, 88–96.
37. Wang, Y.Y.; Liu, J. Variational iteration method for solving fractional partial differential equations with boundary conditions. *J. Comput. Phys.* **2019**, *389*, 1–10.
38. Islam, S.M.R.; Islam, M.S.; Alim, M.A. Adomain decomposition transform method for solving fractional partial differential equations. *Appl. Math. Model.* **2015**, *39*, 5897–5910.
39. Almeida, R.A.; Oliveira, A.L.P.; Tenreiro Machado, J.A. Adomain decomposition transform method for solving time-fractional diffusion equations. *J. Comput. Appl. Math.* **2012**, *236*, 5051–5061.
40. Islam, M.S.; Alim, M.A. Adomain decomposition transform method for solving space-fractional diffusion equations. *J. Comput. Phys.* **2017**, *338*, 1–13.
41. Alim, M.A.; Islam, M.S.; Islam, S.M.R. Adomain decomposition transform method for solving coupled fractional partial differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **2019**, *75*, 201–216.
42. Aboodh, K.S. Application of new transform “Aboodh Transform” to partial differential equations. *Glob. J. Pure Appl. Math.* **2014**, *10*, 249–254.
43. Aboodh, K.S. Solving fourth order parabolic PDE with variable coefficients using Aboodh transform homotopy perturbation method. *Pure Appl. Math. J.* **2015**, *4*, 219–224.
44. Jena, R.M.; Chakraverty, S.; Baleanu, D.; Alqurashi, M.M. New aspects of ZZ transform to fractional operators with Mittag-Leffler kernel. *Front. Phys.* **2020**, *8*, 352.
45. Riabi, L.; Belghaba, K.; Cherif, M.H.; Ziane, D. Homotopy perturbation method combined with ZZ transform to solve some nonlinear fractional differential equations. *Int. J. Anal. Appl.* **2019**, *17*, 406–419.
46. Zafar, Z.U.A. Application of ZZ transform method on some fractional differential equations. *Int. J. Adv. Eng. Global Technol.* **2016**, *4*, 1355–1363.
47. Atangana, A.; Baleanu, D. New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model. *Therm Sci.* **2016**, *20*, 763–769. <https://doi.org/10.2298/TSCI160111018A>.
48. Rashid, S.; Kubra, K.T.; Jafari, H.; Lehre, S.U. A semi analytical approach for fractional order Boussinesq equation in a gradient unconfined aquifers. *Math. Methods Appl. Sci.* **2022**, *45*, 1033–1062.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.