

Article

The Quantization of Gravity: The Quantization of the Full Einstein Equations

Claus Gerhardt 

Institut für Angewandte Mathematik, Ruprecht-Karls-Universität, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany; gerhardt@math.uni-heidelberg.de

Abstract: We quantized the full Einstein equations in a globally hyperbolic spacetime $N = N^{n+1}$, $n \geq 3$, and found solutions of the resulting hyperbolic equation in a fiber bundle E which can be expressed as a product of spatial eigenfunctions (eigendistributions) and temporal eigenfunctions. The spatial eigenfunctions form a basis in an appropriate Hilbert space while the temporal eigenfunctions are solutions to a second-order ordinary differential equation in \mathbb{R}_+ . In case $n \geq 17$ and provided the cosmological constant Λ is negative, the temporal eigenfunctions are eigenfunctions of a self-adjoint operator \hat{H}_0 such that the eigenvalues are countable and the eigenfunctions form an orthonormal basis of a Hilbert space.

Keywords: quantization of gravity; quantum gravity; black hole; negative cosmological constant; partition function; entropy; temporal eigenfunctions; spatial eigenfunctions

Contents

1. Introduction	1
2. Quantizing the Full Einstein Equations	8
3. Spatial Eigenfunctions	11
4. Temporal Eigenfunctions: The Case $3 \leq n \leq 16$	15
5. Temporal Eigenfunctions: The Case $n \geq 17$	17
5.1. Treating Λ as an Eigenvalue	17
5.2. Treating Λ as a Fixed Cosmological Constant	24
6. Trace Class Estimates for $e^{-\beta \hat{H}_0}$	29
7. Conclusions	32
8. References	33

1. Introduction

General relativity is a Lagrangian theory and the canonical quantization of a Lagrangian theory is performed with the help of the Legendre transformation, which would transform the Lagrangian theory to an equivalent Hamiltonian theory, provided that the Lagrangian is *regular*, i.e., the second derivatives of the Lagrangian with respect to the time derivatives of the variables, which form a bilinear form, should be invertible. The Einstein–Hilbert Lagrangian is not regular. However, in a groundbreaking paper Arnowit, Deser and Misner (ADM) [1] proved that, with the help of a global time function x^0 , the Einstein–Hilbert functional could be expressed in a form which made it possible to define a Hamiltonian H and two constraints, the Hamilton constraint and the diffeomorphism constraint.



Citation: Gerhardt, C. The Quantization of Gravity: The Quantization of the Full Einstein Equations. *Symmetry* **2023**, *15*, 1599. <https://doi.org/10.3390/sym15081599>

Academic Editor: Ignatios Antoniadis

Received: 26 July 2023

Revised: 14 August 2023

Accepted: 16 August 2023

Published: 17 August 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Employing the Hamiltonian one could define the Hamilton equations and, combined with the two constraints, the resulting constrained Hamiltonian system was equivalent to the Einstein equations. Bryce DeWitt used this constrained Hamiltonian system to perform a first canonical quantization of the Einstein equations in [2]. The Hamiltonian H would be transformed to an operator \hat{H} which would act on functions u depending on Riemannian metrics g_{ij} and the Hamilton constraint, which could be expressed as an equation

$$H = 0, \quad (1)$$

would be transformed to the equation

$$\hat{H}u = 0. \quad (2)$$

The last equation is now known as the Wheeler–DeWitt equation. It could at first only be solved in highly symmetric cases like in the quantization of Friedman universes, cf. [3–7] and also the monographs [8,9] and the bibliography therein.

In [10] we quantized a general globally hyperbolic spacetime $N = N^{n+1}$, $n \geq 3$, where n is the space dimension, by using the aforementioned papers [1,2]. In that paper, we first eliminated the diffeomorphism constraint by proving that the Einstein equations, which are the Euler–Lagrange equations of the Einstein–Hilbert functional, are equivalent to the Euler–Lagrange equations which are obtained by only considering Lorentzian metrics which split, i.e., they are of the form

$$d\bar{s}^2 = -w^2(dx^0)^2 + g_{ij}(x^0, x)dx^i dx^j, \quad (3)$$

where the function $w > 0$ and the Riemannian metrics g_{ij} are arbitrary, cf. [10] (Theorem 3.2, p. 8). Let $G_{\alpha\beta}$, $0 \leq \alpha, \beta \leq n$, be the Einstein tensor and Λ a cosmological constant. If only metrics of the form (3) are considered, then the resulting Einstein equations can be split in a tangential part

$$G_{ij} + \Lambda g_{ij} = 0 \quad (4)$$

and a normal part

$$G_{\alpha\beta}v^\alpha v^\beta - \Lambda = 0, \quad (5)$$

where $v = (v^\alpha)$ is a normal vector field to the Cauchy hypersurfaces

$$\{x^0 = t\}, \quad t \in x^0(N). \quad (6)$$

The mixed Einstein equations are trivially satisfied since

$$G_{0j} = g_{0j} = 0. \quad (7)$$

The tangential Einstein equations are equivalent to the Hamilton equations, which are defined by the Hamiltonian H , and the normal equation is equivalent to the Hamilton constraint which can be expressed by the Equation (1).

We also introduced a firm mathematical setting by quantizing a globally hyperbolic spacetime N and working after the quantization in a fiber bundle E with base space S_0 , where S_0 was a Cauchy hypersurface of the quantized spacetime N . The fibers consisted of the Riemannian metrics defined in S_0 . The quantized Hamiltonian \hat{H} was a hyperbolic differential operator of second order in E acting only in the fibers. We solved the Wheeler–DeWitt equation (2) in E , where $u = u(t, x, g_{ij})$, for given initial values, cf. [10] (Theorem 5.4, p. 18). It is worth noting that the Wheeler–DeWitt equation represents a quantization of the Hamilton condition, or equivalently, of the normal Einstein equation. The tangential Einstein equations have been ignored.

In our paper [11] and in the monograph [12], we finally quantized the full Einstein equations by incorporating the Hamilton condition in the Hamilton equations and we quantized this evolution equation. There are two possibilities of how the Hamilton con-

dition can be incorporated into the Hamilton equations, and both modified Hamilton equations combined with the original Hamilton equations are equivalent to the full Einstein equations, cf. [12] (Theorem 1.3.3, p. 13, & equ. 1.6.22, p. 41). After quantization of the modified Hamilton equations, however, the resulting hyperbolic equations are different: one equation, let us call it the first equation to give it a name, is a hyperbolic equation wherein the elliptic parts—two Laplacians with respect to certain metrics—act both in the fibers as well as in the base space of a fiber bundle. The second equation is only a hyperbolic equation in the base space, since the Laplacian acting in the fiber was eliminated by the modification.

The first equation has the form

$$-\Delta u - (n-1)\varphi\tilde{\Delta}u - \frac{n-2}{2}\varphi(R-2\Lambda)u = 0, \quad (8)$$

cf. [11] (equ. (4.51)) or [12] (equ. (1.4.88)), where the embellished Laplacian $\tilde{\Delta}u$ is the Laplacian in the base space \mathcal{S}_0 with respect to the metric g_{ij} if the function

$$u \in C_c^\infty(E, \mathbb{C}) \quad (9)$$

is evaluated at

$$(x, g_{ij}(x)) \in E, \quad (10)$$

or equivalently, after choosing appropriate coordinates in the fibers,

$$\begin{aligned} & \frac{n}{16(n-1)}t^{-m}\frac{\partial}{\partial t}\left(t^m\frac{\partial u}{\partial t}\right) - t^{-2}\Delta_M u \\ & + t^{2-\frac{4}{n}}\left\{-(n-1)\Delta_\sigma u - \frac{n-2}{2}R_\sigma u\right\} + \frac{n-2}{2}t^2\Lambda u = 0, \end{aligned} \quad (11)$$

where

$$m = \frac{(n-1)(n+2)}{2} \quad \wedge \quad n = \dim \mathcal{S}_0. \quad (12)$$

The index σ indicates that the corresponding geometric quantities are defined with respect to the metric $\sigma_{ij} \in M$, where M is the Cauchy hypersurface,

$$M = \{t = 1\}. \quad (13)$$

The term R_σ denotes the scalar curvature of the metric σ_{ij} and Λ is a cosmological constant. By choosing a suitable atlas in the base space \mathcal{S}_0 , cf. Lemma 2 on page 12, each fiber $M(x)$ consists of the positive definite matrices $\sigma_{ij}(x)$ satisfying

$$\det \sigma_{ij}(x) = 1, \quad (14)$$

and hence, it is isometric to the symmetric space

$$SL(n, \mathbb{R})/SO(n) \equiv G/K. \quad (15)$$

cf. [2] (equ.(5.17), p. 1123) and [13] (p. 3).

In [11,12], we could solve the hyperbolic Equation (11) only abstractly. But due to the results in our paper [14], we are now able to apply separation of variables to express the solutions u of (11) as a product of spatial and temporal eigenfunctions, or better, eigendistributions. There are three types of spatial eigenfunctions: first, the eigenfunctions of $-\Delta_M$, for which we choose the elements of the Fourier kernel e_{λ, b_0} such that

$$-\Delta_M e_{\lambda, b_0} = (|\lambda|^2 + |\rho|^2)e_{\lambda, b_0}, \quad (16)$$

see Section 3 on page 11 for details, and then the eigenfunctions of the operator

$$-(n-1)\Delta_\sigma - \frac{n-2}{2}R_\sigma. \quad (17)$$

While the operator in (16) acts in the fibers, and hence, the variables are the metrics $\sigma_{ij} \in M$, the operator in (17) is an elliptic differential operator of second order in \mathcal{S}_0 for a fixed σ_{ij} . Thus, we have to specify a Riemannian metric σ_{ij} in \mathcal{S}_0 which is considered to be important either for physical or mathematical reasons. When a globally hyperbolic spacetime is quantized, then \mathcal{S}_0 is a Cauchy hypersurface, usually a coordinate slice, and it will be equipped with a Riemannian metric χ_{ij} . It can be arranged that an arbitrary Riemannian metric χ_{ij} will be an element of M . Thus, our choice will be provided by the initial Cauchy hypersurface. In [15], we incorporated the standard model into our model; hence, we chose $\mathcal{S}_0 = \mathbb{R}^3$ and $\chi_{ij} = \delta_{ij}$.

When we quantize black holes, Schwarzschild-AdS or Kerr-AdS black holes, the interior region of a black hole can be considered to be a globally hyperbolic spacetime and the slices $\{r = \text{const}\}$ are Cauchy hypersurfaces with induced Riemannian metrics $\chi_{ij}(r)$ (note that here r is a label, not a variable). If the event horizon is characterized by $r = r_0$, we proved that the Riemannian metrics $\chi_{ij}(r)$ converge to a Riemannian metric $\chi_{ij}(r_0)$ in an appropriate coordinate system. Thus, we chose \mathcal{S}_0 to be the event horizon and $\chi_{ij} = \chi_{ij}(r_0)$. Moreover, \mathcal{S}_0 could be written as a product

$$\mathcal{S}_0 = \mathbb{R} \times M_0, \quad (18)$$

where M_0 was a compact Riemannian manifold and χ a product metric

$$\chi = \delta \otimes \bar{\sigma}, \quad (19)$$

where δ is the standard "metric" in \mathbb{R} and $\bar{\sigma}$ a Riemannian metric on M_0 .

Following the lead from the black holes, we shall also assume in case of the quantization of a general globally hyperbolic spacetime $N = N^{n+1}$, $n \geq 3$, that \mathcal{S}_0 is a product

$$\mathcal{S}_0 = \mathbb{R}^{n_1} \times M_0, \quad (20)$$

at least topologically, and that M_0 is a compact manifold of dimension

$$\dim M_0 = n - n_1. \quad (21)$$

If N should be a mathematical model of our universe, then we would choose $n_1 = 3$ and M_0 should be a compact manifold, hidden from our observation, of fairly large dimension. Indeed, we shall see that $n \geq 17$ would be preferable if at the same time the cosmological constant Λ would be negative. Moreover, assuming that N should be equipped with an Einstein metric, we would choose M_0 to be a Calabi–Yau manifold if $\Lambda = 0$, while in the case of $\Lambda < 0$, M_0 should be a Kähler–Einstein space, and if $\Lambda > 0$ then M_0 is supposed to be a round sphere with a given radius. The metric σ which we would use in the definition of the operator (17) would then be

$$\sigma = \chi = \delta \otimes \bar{\sigma}, \quad (22)$$

where δ would be the Euclidean metric in \mathbb{R}^{n_1} and $\bar{\sigma}$ the Riemannian metric in M_0 . The differential operator in (17) would then have the form

$$-(n-1)\Delta_\delta - (n-1)\Delta_{\bar{\sigma}} - \frac{n-2}{2}R_{\bar{\sigma}}, \quad (23)$$

which would have eigenfunctions of the form

$$\zeta\varphi \quad (24)$$

where ζ is an eigenfunction of the Euclidean Laplacian and φ an eigenfunction of the remaining part of the operator. Hence, we would have three types of spatial eigenfunctions which are well-known—both mathematically and physically—and their product will play the part of the spatial eigenfunctions of the hyperbolic Equation (11). The solution u of that equation will then be of the form

$$u = wv\zeta\varphi \quad (25)$$

where

$$v = e_{\lambda, b_0} \circ [g_0] \quad (26)$$

is an eigenfunction of $-\Delta_M$ satisfying

$$-\Delta_M v = (|\lambda|^2 + |\rho|^2)v \quad (27)$$

and

$$v(\chi(x)) = 1 \quad \forall x \in \mathcal{S}_0, \quad (28)$$

for details, we refer to the arguments following Remark 4 on page 13. The function w depends only on t and it will solve a second-order ordinary differential equation (ODE). The functions u will be evaluated at (t, x, χ) . More precisely, we proved the following:

Theorem 1. Assume that \mathcal{S}_0 is a direct product as in (20) endowed with the metric χ in (22). Then, a solution $u = u(x, t, \sigma_{ij})$ of the hyperbolic Equation (11) can be expressed as a product of spatial eigenfunctions $v = v(\sigma_{ij})$, $\zeta = \zeta(y)$, $\varphi_k = \varphi_k(x)$, $k \in \mathbb{N}$, and temporal eigenfunctions $w = w(t)$; u is evaluated at $\sigma_{ij} = \chi_{ij}$, where

$$u = wv\zeta\varphi_k. \quad (29)$$

The temporal eigenfunction w is a solution of the ODE

$$\begin{aligned} & \frac{n}{16(n-1)} t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) + t^{-2} (|\lambda|^2 + |\rho|^2) w \\ & + t^{2-\frac{4}{n}} \{ (n-1) |\tilde{\zeta}|^2 + \bar{\mu}_k \} w + \frac{n-2}{2} t^2 \Lambda w = 0 \end{aligned} \quad (30)$$

in $0 < t < \infty$.

In Section 5 on page 17, we look at the case $n \geq 17$ and $\Lambda < 0$ and prove that the Equation (30) can be considered to be an implicit eigenvalue problem where Λ plays the part of the eigenvalue provided

$$\frac{16(n-1)}{n} |\lambda|^2 < 238. \quad (31)$$

To understand the corresponding theorem, we need a few remarks and definitions. First, we multiply Equation (30) by

$$\frac{16(n-1)}{n}; \quad (32)$$

then, we use the abbreviations

$$\mu_0 = \frac{16(n-1)}{n} (|\lambda|^2 + |\rho|^2), \quad (33)$$

$$m_1 = \frac{16(n-1)}{n} \{ (n-1) |\tilde{\zeta}|^2 + \bar{\mu}_k \} \quad (34)$$

and

$$m_2 = \frac{8(n-1)(n-2)}{n} \quad (35)$$

and define for $w \in C_c^\infty(\mathbb{R}_+)$

$$\hat{B}w = -t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{-2} \mu_0 w. \tag{36}$$

Remark 1. Note that $\mu_0 > 0$, which would in general deprive of success any attempt to solve a meaningful eigenvalue problem for this operator. But if (31) is satisfied and $n \geq 17$, then it is possible to prove the following theorem in Section 5 on page 17.

Theorem 2. There are countably many solutions (Λ_i, w_i) of the implicit eigenvalue problem

$$\hat{B}w_i - m_2 \Lambda_i t^2 w_i = m_1 t^{2-\frac{4}{n}} w_i \tag{37}$$

with eigenfunctions $w_i \in \hat{\mathcal{H}}_2$ such that

$$\Lambda_i < \Lambda_{i+1} < 0 \quad \forall i \in \mathbb{N}, \tag{38}$$

$$\lim_i \Lambda_i = 0, \tag{39}$$

and their multiplicities are one. The transformed eigenfunctions

$$\tilde{w}_i(t) = w_i(\lambda_i^{\frac{n}{4(n-1)}} t), \tag{40}$$

where

$$\lambda_i = (-\Lambda_i)^{-\frac{n-1}{n}}, \tag{41}$$

form a basis of $\hat{\mathcal{H}}_2$ and also of $L^2(\mathbb{R}_+, m)$.

Equation (37) is identical to Equation (30) if Λ is replaced by Λ_i . The vector spaces $\hat{\mathcal{H}}_2$ and $L^2(\mathbb{R}_+, m)$ are Hilbert spaces which are defined later.

However, if we consider $\Lambda < 0$ to be a fixed cosmological constant and not a parameter which can also play the role of an implicit eigenvalue, we have to use a different approach.

First, let us express Equation (30) in the equivalent form

$$\hat{\varphi}_0^{-1} \left\{ -\frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{m-2} \mu_0 w - t^{m+2} m_2 \Lambda w \right\} - \frac{16(n-1)}{n} \{ (n-1) |\zeta|^2 + \bar{\mu}_k \} w = 0, \tag{42}$$

where

$$\hat{\varphi}_0(t) = t^{m+2-\frac{4}{n}} \tag{43}$$

and where we used the definitions (33) and (35). The term

$$(n-1) |\zeta|^2 + \bar{\mu}_k \tag{44}$$

is an eigenvalue of the operator in (23). $|\zeta|^2$ with $\zeta \in \mathbb{R}^{n_1}$ is a continuous eigenvalue while the sequence $\bar{\mu}_k, k \in \mathbb{N}$, satisfies the relations

$$\bar{\mu}_0 < \bar{\mu}_1 \leq \bar{\mu}_2 \leq \dots \tag{45}$$

and

$$\lim_{k \rightarrow \infty} \bar{\mu}_k = \infty. \tag{46}$$

The corresponding eigenfunctions φ_k are smooth and the eigenspaces are finite dimensional.

On the other hand, the operator

$$\hat{H}_0 w \equiv \hat{\varphi}_0^{-1} \left\{ -\frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{m-2} \mu_0 w - t^{m+2} m_2 \Lambda w \right\} \quad (47)$$

is self-adjoint in the Hilbert space $\hat{\mathcal{H}} = L^2(\mathbb{R}_+, d\hat{\mu})$, cf. (242) on page 24, with a complete system of eigenfunctions $w_i, i \in \mathbb{N}$, and corresponding eigenvalues

$$0 < \lambda_0 < \lambda_1 < \lambda_2 < \dots \quad (48)$$

The eigenspaces are all one-dimensional and the ground state w_0 does not change sign, cf. Remark 5 on page 21. Thus, in order to solve Equation (42), we have to find for each pair (w_i, λ_i) eigenvalues $\bar{\mu}_k$ and $\bar{\zeta} \in \mathbb{R}^{n_1}$ such that

$$\frac{16(n-1)}{n} \{ (n-1) |\bar{\zeta}|^2 + \bar{\mu}_k \} = \lambda_i. \quad (49)$$

This is indeed possible provided either $\bar{\mu}_0 \leq 0$ or

$$|\Lambda|^{\frac{n-1}{n}} \geq \bar{\lambda}_0^{-1} \frac{16(n-1)}{n} \bar{\mu}_0, \quad (50)$$

cf. Corollary 1 on page 25. Using the eigenvalues on the left-hand side of (49) and the corresponding eigenfunctions of the operator (23), we then define a self-adjoint operator H_1 in a Hilbert space \mathcal{H} having the same eigenvalues λ_i as \hat{H}_0 but with higher finite multiplicities. Relabelling these eigenvalues to include the multiplicities and denoting them by $\tilde{\lambda}_i$, they satisfy

$$0 < \tilde{\lambda}_0 \leq \tilde{\lambda}_1 \leq \dots \quad (51)$$

and

$$\lim_{i \rightarrow \infty} \tilde{\lambda}_i = \infty. \quad (52)$$

In Section 6 on page 29, we shall prove that the operator $e^{-\beta \hat{H}_0}, \beta > 0$, is of trace class from which we conclude that $e^{-\beta H_1}$ is also of trace class. We are then in a similar situation as in [12] (Chapter 6.5), where we proved the following:

Lemma 1. For any $\beta > 0$, the operator

$$e^{-\beta H_1} \quad (53)$$

is of trace class in \mathcal{H} , i.e.,

$$\text{tr}(e^{-\beta H_1}) = \sum_{i=0}^{\infty} e^{-\beta \tilde{\lambda}_i} < \infty. \quad (54)$$

Let

$$\mathcal{F} \equiv \mathcal{F}_+(\mathcal{H}) \quad (55)$$

be the symmetric Fock space generated by \mathcal{H} and let

$$H = d\Gamma(H_1) \quad (56)$$

be the canonical extension of H_1 to \mathcal{F} . Then,

$$e^{-\beta H} \quad (57)$$

is also of trace class in \mathcal{F}

$$\text{tr}(e^{-\beta H}) = \prod_{i=0}^{\infty} (1 - e^{-\beta \tilde{\lambda}_i})^{-1} < \infty. \quad (58)$$

Remark 2. In [12] (Chapter 6.5), we also used these results to define the partition function Z by

$$Z = \text{tr}(e^{-\beta H}) = \prod_{i=0}^{\infty} (1 - e^{-\beta \tilde{\lambda}_i})^{-1} \quad (59)$$

and the density operator ρ in \mathcal{F} by

$$\rho = Z^{-1} e^{-\beta H} \quad (60)$$

such that

$$\text{tr} \rho = 1. \quad (61)$$

The von Neumann entropy S is then defined by

$$\begin{aligned} S &= -\text{tr}(\rho \log \rho) \\ &= \log Z + \beta Z^{-1} \text{tr}(H e^{-\beta H}) \\ &= \log Z - \beta \frac{\partial \log Z}{\partial \beta} \\ &\equiv \log Z + \beta E, \end{aligned} \quad (62)$$

where E is the average energy

$$E = \text{tr}(H \rho). \quad (63)$$

E can be expressed in the form

$$E = \sum_{i=0}^{\infty} \frac{\tilde{\lambda}_i}{e^{\beta \tilde{\lambda}_i} - 1}. \quad (64)$$

Here, we also set the Boltzmann constant

$$k_B = 1. \quad (65)$$

The parameter β is supposed to be the inverse of the absolute temperature T

$$\beta = T^{-1}. \quad (66)$$

For a more detailed analysis and especially for the dependence on Λ , we refer to [12] (Chapter 6.5).

Remark 3. Let us also mention that we use Planck units in this paper, i.e.,

$$c = G = k_B = \hbar = 1. \quad (67)$$

Moreover, the signature of a Lorentzian metric has the form $(-, +, \dots, +)$.

2. Quantizing the Full Einstein Equations

Let $N = N^{n+1}$, $n \geq 3$, be a globally hyperbolic Lorentzian manifold with metric $\bar{g}_{\alpha\beta}$, $0 \leq \alpha, \beta \leq n$. The Einstein equations are Euler–Lagrange equations of the Einstein–Hilbert functional

$$\int_N (\bar{R} - \Lambda), \quad (68)$$

where \bar{R} is the scalar curvature, Λ a cosmological constant and where we omitted the integration density in the integral. In order to apply a Hamiltonian description of general relativity, one usually defines a time function x^0 and considers the foliation of N given by the slices

$$M(t) = \{x^0 = t\}. \quad (69)$$

We may, without loss of generality, assume that the spacetime metric splits

$$ds^2 = -w^2(dx^0)^2 + g_{ij}(x^0, x)dx^i dx^j, \quad (70)$$

cf. [10] (Theorem 3.2). Then, the Einstein equations also split into a tangential part

$$G_{ij} + \Lambda g_{ij} = 0 \quad (71)$$

and a normal part

$$G_{\alpha\beta}v^\alpha v^\beta - \Lambda = 0, \quad (72)$$

where the naming refers to the given foliation. For the tangential Einstein equations, one can define equivalent Hamilton equations due to the groundbreaking paper by Arnowitt, Deser and Misner [1]. The normal Einstein equations can be expressed by the so-called Hamilton condition

$$\mathcal{H} = 0, \quad (73)$$

where \mathcal{H} is the Hamiltonian used in defining the Hamilton equations. In the canonical quantization of gravity, the Hamiltonian is transformed to a partial differential operator of hyperbolic type $\hat{\mathcal{H}}$ and the possible quantum solutions of gravity are supposed to satisfy the so-called Wheeler–DeWitt equation

$$\hat{\mathcal{H}}u = 0 \quad (74)$$

in an appropriate setting, i.e., only the Hamilton condition (73) has been quantized, or equivalently, the normal Einstein equation, while the tangential Einstein equations have been ignored.

In [10], we solved the Equation (74) in a fiber bundle E with base space \mathcal{S}_0 ,

$$\mathcal{S}_0 = \{x^0 = 0\} \equiv M(0), \quad (75)$$

and fibers $F(x)$, $x \in \mathcal{S}_0$,

$$F(x) \subset T_x^{0,2}(\mathcal{S}_0), \quad (76)$$

the elements of which are the positive definite symmetric tensors of order two, the Riemannian metrics in \mathcal{S}_0 . The hyperbolic operator $\hat{\mathcal{H}}$ is then expressed in the form

$$\hat{\mathcal{H}} = -\Delta - (R - 2\Lambda)\varphi, \quad (77)$$

where Δ is the Laplacian of the DeWitt metric given in the fibers, R the scalar curvature of the metrics $g_{ij}(x) \in F(x)$ and φ is defined by

$$\varphi^2 = \frac{\det g_{ij}}{\det \rho_{ij}}, \quad (78)$$

where ρ_{ij} is a fixed metric in \mathcal{S}_0 such that, instead of densities, we are considering functions. The Wheeler–DeWitt equation could be solved in E but only as an abstract hyperbolic equation. The solutions could not be split in corresponding spatial and temporal eigenfunctions.

The underlying mathematical reason for the difficulty was the presence of the term R in the quantized equation, which prevents the application of separation of variables, since the metrics g_{ij} are the spatial variables. In a recent paper [14], we overcame this difficulty by quantizing the Hamilton equations instead of the Hamilton condition.

As a result, we obtained the equation

$$-\Delta u = 0 \quad (79)$$

in E , where the Laplacian is the Laplacian in (77). The lower order terms of $\hat{\mathcal{H}}$

$$(R - 2\Lambda)\varphi \quad (80)$$

were eliminated during the quantization process. However, Equation (79) is only valid provided $n \neq 4$, since the resulting equation actually looks like

$$-\left(\frac{n}{2} - 2\right)\Delta u = 0. \quad (81)$$

This restriction seems to be acceptable, since n is the dimension of the base space \mathcal{S}_0 which, by general consent, is assumed to be $n = 3$. The fibers add additional dimensions to the quantized problem, namely

$$\dim F = \frac{n(n+1)}{2} \equiv m + 1. \quad (82)$$

The fiber metric, the DeWitt metric, which is responsible for the Laplacian in (79), can be expressed in the form

$$ds^2 = -\frac{16(n-1)}{n}dt^2 + \varphi G_{AB}d\zeta^A d\zeta^B, \quad (83)$$

where the coordinate system is

$$(\zeta^a) = (\zeta^0, \zeta^A) \equiv (t, \zeta^A). \quad (84)$$

The (ζ^A) , $1 \leq A \leq m$, are coordinates for the hypersurface

$$M \equiv M(x) = \{(g_{ij}) : t^4 = \det g_{ij}(x) = 1, \forall x \in \mathcal{S}_0\}. \quad (85)$$

We also assumed that $\mathcal{S}_0 = \mathbb{R}^n$ and that the metric ρ_{ij} in (78) is the Euclidean metric δ_{ij} . It is well-known that M is a symmetric space

$$M = SL(n, \mathbb{R})/SO(n) \equiv G/K. \quad (86)$$

It is also easily verified that the induced metric of M in E is isometric to the Riemannian metric of the coset space G/K .

Now, we were in a position to use separation of variables, namely we wrote a solution of (79) in the form

$$u = w(t)v(\zeta^A), \quad (87)$$

where v is a spatial eigenfunction of the induced Laplacian of M

$$-\Delta_M v \equiv -\Delta v = (|\lambda|^2 + |\rho|^2)v \quad (88)$$

and w is a temporal eigenfunction satisfying the ODE

$$\ddot{w} + mt^{-1}\dot{w} + \mu_0 t^{-2}w = 0 \quad (89)$$

with

$$\mu_0 = \frac{16(n-1)}{n}(|\lambda|^2 + |\rho|^2). \quad (90)$$

The eigenfunctions of the Laplacian in G/K are well-known and we chose the kernel of the Fourier transform in G/K in order to define the eigenfunctions. This choice also allowed us to use Fourier quantization similar to the Euclidean case such that the eigenfunctions are transformed into Dirac measures and the Laplacian into a multiplication operator in Fourier space.

In the present paper, we want to quantize the full Einstein equations by using a previous result, cf. [11] (Theorem 3.2) or [12] (Theorem 1.3.4), where we proved that the full Einstein equations are equivalent to the Hamilton equations and a scalar evolution equation, which we obtained by incorporating the Hamilton condition into the right-hand

side of the second Hamilton equations and we quantized this evolution equation in fiber bundle E with base space \mathcal{S}_0 and fibers

$$F(x) \in T_x^{0,2}(\mathcal{S}_0), \quad \forall x \in \mathcal{S}_0, \quad (91)$$

cf. (76).

The quantization of the scalar evolution equation then yielded the following hyperbolic equation in E :

$$-\Delta u - (n-1)\varphi\tilde{\Delta}u - \frac{n-2}{2}\varphi(R-2\Lambda)u = 0, \quad (92)$$

cf. [11] (equ. (4.51)) or [12] (equ. (1.4.88)) where the embellished Laplacian $\tilde{\Delta}u$ is the Laplacian in the base space \mathcal{S}_0 with respect to the metric g_{ij} if the function

$$u \in C_c^\infty(E, \mathbb{C}) \quad (93)$$

is evaluated at

$$(x, g_{ij}(x)) \in E. \quad (94)$$

Let us recall that the time function t in (84) is defined by

$$t^2 = \varphi \quad (95)$$

and that t is independent of x , cf. [11] (Lemma 4.1, p. 726), and, furthermore, that the fiber elements $g_{ij}(x)$ can be expressed as

$$g_{ij}(x) = t^{\frac{4}{n}}\sigma_{ij}(x), \quad (96)$$

where the metrics $\sigma_{ij}(x)$ are elements of the fibers of the subbundle

$$E_1 = \{t = 1\} \subset E \quad (97)$$

with fibers

$$M(x) \subset F(x) \quad \forall x \in \mathcal{S}_0 \quad (98)$$

consisting of metrics $\sigma_{ij}(x)$ satisfying

$$\det \sigma_{ij}(x) = \det \rho_{ij}(x) \quad \forall x \in \mathcal{S}_0. \quad (99)$$

Now, combining (96), the definition of the fiber metric (83) and the relation between the scalar curvatures of conformal metrics the hyperbolic Equation (92) can be expressed in the form

$$\begin{aligned} & \frac{n}{16(n-1)}t^{-m}\frac{\partial}{\partial t}\left(t^m\frac{\partial u}{\partial t}\right) - t^{-2}\Delta_M u \\ & + t^{2-\frac{4}{n}}\left\{-(n-1)\Delta_\sigma u - \frac{n-2}{2}R_\sigma u\right\} + \frac{n-2}{2}t^2\Lambda u = 0, \end{aligned} \quad (100)$$

where the index σ indicates that the corresponding geometric quantities are defined with respect to the metric σ_{ij} .

In the following sections, we shall solve Equation (100) by employing separation of variables to obtain corresponding spatial and temporal eigenfunctions or eigendistributions.

3. Spatial Eigenfunctions

Let us first look for spatial eigenfunctions of the operators

$$-\Delta_M \quad (101)$$

and

$$-(n-1)\Delta_\sigma - \frac{n-2}{2}R_\sigma. \quad (102)$$

In the case of the Laplacian in (101), we would want to use the fact that each Cauchy hypersurface $M(x)$ is isometric to the symmetric space

$$SL(n, \mathbb{R})/SO(n) \equiv G/K \quad (103)$$

provided

$$\det \rho_{ij}(x) = 1. \quad (104)$$

In our former papers [14,15], we had chosen $\mathcal{S}_0 = \mathbb{R}^n$ and

$$\rho_{ij} = \delta_{ij}, \quad (105)$$

i.e., the condition (104) had been automatically satisfied by choosing Euclidean coordinates. However, for the quantization of black holes, this choice will not be possible since \mathcal{S}_0 will then be the event horizon equipped with a non-flat metric.

To overcome this difficulty, we need the following lemma:

Lemma 2. *Let \mathcal{S}_0 be a Riemannian manifold of dimension $n \geq 2$ and of class $C^{k,\alpha}$ for $0 \leq k \in \mathbb{N}$ and $0 < \alpha < 1$, where $C^{k,\alpha}$ are the usual Hölder spaces, and let ρ_{ij} be a metric of class $C^{k,\alpha}$ in \mathcal{S}_0 ; then, there exists an atlas $\{(x_\beta, U_\beta)\}$ of $C^{k+1,\alpha}$ charts such that the metric ρ_{ij} expressed in an arbitrary chart (x_β, U_β) satisfies*

$$\det \rho_{ij}(x) = 1 \quad \forall x \in x_\beta(U_\beta) \subset \mathbb{R}^n. \quad (106)$$

Proof. We first prove (106) locally. Let ρ_{ij} be a local expression of ρ in coordinates $x = (x^i)$ and let $\tilde{x} = \tilde{x}(x)$ be a coordinate transformation and $\tilde{\rho}_{kl}$ be the corresponding expression for the metric ρ ; then,

$$\tilde{\rho}_{kl} = \rho_{ij} \frac{\partial x^i}{\partial \tilde{x}^k} \frac{\partial x^j}{\partial \tilde{x}^l} \quad (107)$$

and

$$\det \tilde{\rho}_{kl} = \det \rho_{ij} \left| \frac{\partial x}{\partial \tilde{x}} \right|^2, \quad (108)$$

where

$$\left| \frac{\partial x}{\partial \tilde{x}} \right| = \det \frac{\partial x^i}{\partial \tilde{x}^k}, \quad (109)$$

the Jacobi determinant.

Let the coordinates $x = (x^i)$ be defined in an open set $\Omega \subset \mathbb{R}^n$ with boundary $\partial\Omega \in C^{k+1,\alpha}$; then, due to a result of Dacorogna and Moser, there exists a diffeomorphism $y = y(x)$, $y \in C^{k+1,\alpha}(\bar{\Omega}, \mathbb{R}^n)$ such that

$$\begin{aligned} \left| \frac{\partial y}{\partial x} \right| &= \lambda \sqrt{\det \rho_{ij}} && \text{in } \Omega, \\ y(x) &= x && \text{in } \partial\Omega, \end{aligned} \quad (110)$$

where

$$\lambda = \frac{\int_{\Omega} dx}{\int_{\Omega} \sqrt{\det \rho_{ij}} dx}, \quad (111)$$

cf. [16] (Theorem 1' and Remark, p. 4).

Hence, the diffeomorphism

$$\tilde{x} = \lambda^{\frac{1}{n}} y \quad (112)$$

satisfies

$$\left| \frac{\partial \tilde{x}}{\partial x} \right| = \sqrt{\det \rho_{ij}}, \quad (113)$$

or equivalently,

$$\det \tilde{\rho}_{kl} = \det \rho_{ij} \left| \frac{\partial x}{\partial \tilde{x}} \right|^2 = 1, \quad (114)$$

where $\tilde{\rho}_{kl}$ are the coordinate expressions of ρ in the coordinates \tilde{x} .

From the local result, we easily infer the existence of an atlas consisting of local charts with that property. \square

Thus, we are able to identify the fiber $M(x)$ with the symmetric space G/K in (103) and we may choose the elements of the Fourier kernel e_{λ, b_0} as eigenfunctions of $-\Delta_M$ such that

$$-\Delta_M e_{\lambda, b_0} = (|\lambda|^2 + |\rho|^2) e_{\lambda, b_0}, \quad (115)$$

see [17] (Chapter III) and [14] (Section 5) for details, where

$$|\rho|^2 = \frac{1}{12}(n-1)^2 n, \quad (116)$$

cf. [14] (equ. (5.40)). Here, λ is an abbreviation for $\lambda\alpha$, where $\alpha \in (\mathbb{R}^{n-1})^*$ is a character representing an elementary graviton and $\lambda \in \mathbb{R}_+$. There are

$$\alpha = \begin{cases} \alpha_i, & 1 \leq i \leq n-1 \\ \alpha_{ij}, & 1 \leq i < j \leq n \end{cases} \quad (117)$$

special characters. These characters are normalized to have $\|\alpha\| = 1$. They correspond to the degrees of freedom in choosing the entries of a metric g_{ij} satisfying

$$\det g_{ij} = 1. \quad (118)$$

Remark 4. Due to the scalar curvature term R_σ in Equation (102), it is evident that spatial eigenfunctions for this operator cannot be defined on the full subbundle E_1 , cf. (97) on page 11, but only for a fixed metric $\sigma_{ij} \in M$, if $R_\sigma = \text{const}$ maybe for that class of metrics. However, in general, we cannot assume that the scalar curvature is constant, since we shall have to pick a metric χ_{ij} that is a natural metric determined by the underlying spacetime which has been quantized. In the case of a black hole, χ_{ij} will be a metric on the event horizon. Now, let us recall that χ_{ij} should belong to fibers of the subbundle E_1 ; hence, we have to choose ρ_{ij} , which is still arbitrary but fixed, to be equal to χ_{ij}

$$\rho_{ij} = \chi_{ij}. \quad (119)$$

Thus, we evaluate the spatial eigenfunctions at

$$(x, \chi_{ij}(x)) \quad \forall x \in \mathcal{S}_0, \quad (120)$$

especially also e_{λ, b_0} , i.e.,

$$e_{\lambda, b_0}(\chi_{ij}(x)) \quad (121)$$

may not depend on x explicitly. Now, it is well known that

$$e_{\lambda, b_0}(\delta_{ij}(x)) = 1 \quad \forall x \in \mathcal{S}_0 \quad (122)$$

and the Laplacian Δ_M is invariant under the action of G on M . The action of $g \in M$ on $\sigma \in M$ is defined by

$$[g]\sigma = g\sigma g^*, \quad (123)$$

where g^* is the transposed matrix. Since every $\sigma \in M$ is also an element of G , we conclude by choosing

$$g = g_0 \equiv \sqrt{\chi^{-1}}, \quad (124)$$

that

$$[g_0]\chi = \text{id} = (\delta_{ij}), \quad (125)$$

and, furthermore, that the function

$$v = e_{\lambda, b_0} \circ [g_0] \quad (126)$$

is an eigenfunction of $-\Delta_M$ satisfying

$$-\Delta_M v = (|\lambda|^2 + |\rho|^2)v \quad (127)$$

and

$$v(\chi(x)) = 1 \quad \forall x \in \mathcal{S}_0. \quad (128)$$

Let us summarize these results in

Theorem 3. Let e_{λ, b_0} be an eigenfunction of $-\Delta_M$ as in (115) and let g_0 be defined as in (124); then,

$$v = e_{\lambda, b_0} \circ [g_0] \quad (129)$$

is an eigenfunction of $-\Delta_M$ satisfying (127) as well as (128).

Next, let us consider the operator in (102) with $\sigma = \chi$. We furthermore assume that \mathcal{S}_0 is a direct product,

$$\mathcal{S}_0 = \mathbb{R}^{n_1} \times M_0, \quad (130)$$

where M_0 is a smooth, compact and connected manifold of dimension $n - n_1$,

$$\dim M_0 = n - n_1 \equiv n_0. \quad (131)$$

The metric χ_{ij} is then supposed to be a metric product,

$$\chi = \delta \otimes \bar{\sigma}, \quad (132)$$

where δ is the Euclidean metric in \mathbb{R}^{n_1} and $\bar{\sigma}$ a Riemannian metric in M_0 . In case of a black hole, n_1 will be equal to 1.

Since the scalar curvature of the product metric χ is equal to the scalar curvature of $\bar{\sigma}$,

$$R_\chi = R_{\bar{\sigma}}, \quad (133)$$

the operator in (102) can be expressed in the form

$$-(n-1)\Delta_\delta - (n-1)\Delta_{\bar{\sigma}} - \frac{n-2}{2}R_{\bar{\sigma}}. \quad (134)$$

Hence, the corresponding eigenfunctions can be written as a product

$$\zeta \varphi, \quad (135)$$

where ζ is defined in \mathbb{R}^{n_1} ,

$$\zeta(y) = e^{i\langle \xi, y \rangle} \quad \zeta, y \in \mathbb{R}^{n_1}, \quad (136)$$

such that

$$-\Delta_\delta \zeta = |\xi|^2 \zeta, \quad (137)$$

while $\varphi \in C^\infty(M_0)$ is an eigenfunction of the operator

$$A = -(n-1)\Delta_{\bar{\sigma}} - \frac{n-2}{2}R_{\bar{\sigma}}. \quad (138)$$

Since M_0 is compact, it is well-known that A is self-adjoint with countably many eigenvalues $\bar{\mu}_k, k \in \mathbb{N}$, which are ordered

$$\bar{\mu}_0 < \bar{\mu}_1 \leq \bar{\mu}_2 \leq \dots \tag{139}$$

satisfying

$$\lim_{k \rightarrow \infty} \bar{\mu}_k = \infty. \tag{140}$$

The corresponding eigenfunctions φ_k are smooth and the eigenspaces are finite dimensional. The eigenspace belonging to $\bar{\mu}_0$ is one-dimensional and φ_0 never vanishes, i.e., if we consider φ_0 to be real-valued, it will either be strictly positive or negative.

Let us summarize the results we proved so far in the following theorem:

Theorem 4. Assume that S_0 is a direct product as in (130) endowed with the metric χ in (132). Then, a solution $u = u(x, t, \sigma_{ij})$ of the hyperbolic Equation (100) on page 11 can be expressed as a product of spatial eigenfunctions $v = v(\sigma_{ij}), \zeta = \zeta(y), \varphi_k = \varphi_k(x), k \in \mathbb{N}$, and temporal eigenfunctions $w = w(t); u$ is evaluated at $\sigma_{ij} = \chi_{ij}$, where

$$u = wv\zeta\varphi_k. \tag{141}$$

The temporal eigenfunction w is a solution of the ODE

$$\begin{aligned} \frac{n}{16(n-1)} t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) + t^{-2} (|\lambda|^2 + \rho^2) w \\ t^{2-\frac{4}{n}} \{ (n-1) |\zeta|^2 + \bar{\mu}_k \} w + \frac{n-2}{2} t^2 \Lambda w = 0 \end{aligned} \tag{142}$$

in $0 < t < \infty$.

In the next sections, we shall solve the ODE and shall also show that, for large $n, n \geq 17$ and negative Λ w can be chosen to be an eigenfunction of a self-adjoint operator where the cosmological constant plays the role of an implicit eigenvalue.

4. Temporal Eigenfunctions: The Case $3 \leq n \leq 16$

Let us first divide Equation (142) by $\frac{n}{16(n-1)}$ to obtain what we consider to be a normal form

$$\begin{aligned} t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) + t^{-2} \frac{16(n-1)}{n} (|\lambda|^2 + \rho^2) w \\ t^{2-\frac{4}{n}} \frac{16(n-1)}{n} \{ (n-1) |\zeta|^2 + \bar{\mu}_k \} w + \frac{16(n-1)}{n} \frac{n-2}{2} t^2 \Lambda w = 0 \end{aligned} \tag{143}$$

Using the abbreviations

$$\mu_0 = \frac{16(n-1)}{n} (|\lambda|^2 + |\rho|^2), \tag{144}$$

$$m_1 = \frac{16(n-1)}{n} \{ (n-1) |\zeta|^2 + \bar{\mu}_k \} \tag{145}$$

and

$$m_2 = \frac{8(n-1)(n-2)}{n} \tag{146}$$

we can rewrite Equation (143) in the form

$$t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) + t^{-2} \mu_0 w + t^{2-\frac{4}{n}} m_1 w + t^2 m_2 \Lambda w = 0. \tag{147}$$

We shall use two different approaches in solving this ODE depending on the sign of

$$\mu_0 - \frac{(m-1)^2}{4}. \quad (148)$$

Let us recall that

$$m = \frac{(n-1)(n+2)}{2}. \quad (149)$$

and

$$\rho^2 = \frac{(n-1)^2 n}{12}. \quad (150)$$

One can easily verify that

$$\frac{16(n-1)}{n} \rho^2 - \frac{(m-1)^2}{4} = \begin{cases} > 1, & 3 \leq n \leq 16, \\ < -238, & 17 \leq n. \end{cases} \quad (151)$$

Hence, in the case $3 \leq n \leq 16$, the term in (148) will be strictly larger than 1 for all values of $|\lambda|$ and, in the case $n \geq 17$, strictly negative for small values of $|\lambda|$, or more precisely, for all

$$\frac{16(n-1)}{n} |\lambda|^2 < 238. \quad (152)$$

Let us first consider the case $3 \leq n \leq 16$ and let us rewrite Equation (147) in the form

$$\ddot{w} + mt^{-1}w + t^{-2}\{\mu_0 + m_2 t^{4-\frac{4}{n}} + m_3 \Lambda t^4\}w = 0 \quad \forall t > 0. \quad (153)$$

Then, we look at the more general equation

$$\ddot{w} + mt^{-1}w + t^{-2}(\mu_0 + q_0(t))w = 0 \quad \forall t > 0, \quad (154)$$

for which we proved in [18] (Theorem 1.1) the following theorem:

Theorem 5. *Let us assume that the constants m, μ_0 and the real function $q_0 \in C^1(\mathbb{R}_+)$ have the properties*

$$m > 1, \quad (155)$$

$$1 < \mu_0 - \frac{(m-1)^2}{4} \equiv 1 + \gamma, \quad \gamma > 0, \quad (156)$$

and

$$\lim_{t \rightarrow 0} q_0(t) = 0. \quad (157)$$

Then, any non-trivial solution w of (154) satisfies

$$\lim_{t \rightarrow 0} (|w|^2 + t^2 |\dot{w}|^2) = \infty \quad (158)$$

as well as

$$\limsup_{t \rightarrow 0} |w|^2 = \infty. \quad (159)$$

We also described the oscillation behavior of w near $t = 0$, which can be considered to be a big bang of the solutions, as well as to be asymptotically equal to the oscillations of the solutions of the ODE

$$\ddot{w} + mt^{-1}w + \mu_0 t^{-2}w = 0 \quad \forall t > 0, \quad (160)$$

cf. [18] (Theorem 3.2). The solutions of the above equation are

$$w(t) = t^{-\frac{(m-1)}{2}} e^{i\mu \log t}, \quad \mu > 0, \tag{161}$$

where

$$\mu^2 = \mu_0 - \frac{(m-1)^2}{4}, \tag{162}$$

see [14] (equ. (273)).

5. Temporal Eigenfunctions: The Case $n \geq 17$

5.1. Treating Λ as an Eigenvalue

Now, let us consider the case $n \geq 17$ assuming in addition that (152) on page 16 is satisfied such that

$$\bar{\mu} \equiv \mu_0 - \frac{(m-1)^2}{4} < 0 \tag{163}$$

and also that

$$\Lambda < 0. \tag{164}$$

The last two assumptions shall allow us to consider (147) on page 15 as an implicit eigenvalue equation where Λ plays the role of the eigenvalue. We shall prove that the corresponding operator is self-adjoint with a pure point spectrum provided the constant m_1 in (147), which is defined by (145), is strictly positive. This can easily be arranged by choosing a $|\zeta|$ large enough. Notice also that at most finitely, many eigenvalues $\bar{\mu}_k$ are negative.

Equation (147) can be written in the equivalent form

$$-t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{-2} \mu_0 w - t^2 m_2 \Lambda w = t^{2-\frac{4}{n}} m_1 w \quad \forall t > 0. \tag{165}$$

We have a similar equation, or, since the constants, m_1, m_2 , are not specified and their actual positive values are irrelevant, an identical equation already solved by spectral analysis in [3] (Sections 4 and 6). Therefore, we shall only outline the proof by giving the necessary definitions and stating the results but referring the actual proofs to the old paper.

Closely related to Equation (165) is the following equation:

$$-t^{-1} \frac{\partial}{\partial t} \left(t \frac{\partial u}{\partial t} \right) - t^{-2} \bar{\mu} u - t^2 m_2 \Lambda u = t^{2-\frac{4}{n}} m_1 u \quad \forall t > 0, \tag{166}$$

where $\bar{\mu}$ is defined in (163). If $w \in C^2(\mathbb{R}_+^*)$ is a solution of (165), then

$$u = t^{\frac{m-1}{2}} w \tag{167}$$

is a solution of (166) and vice versa, as can be easily verified. The operator

$$Bu = -t^{-1} \frac{\partial}{\partial t} \left(t \frac{\partial u}{\partial t} \right) - t^{-2} \bar{\mu} u \tag{168}$$

is known as a Bessel operator.

Definition 1. Let $I = (0, \infty)$ and let $q \in \mathbb{R}$. Then, we define

$$L^2(I, q) = \{ u \in L^2_{\text{loc}}(I, \mathbb{R}) : \int_I r^q |u|^2 < \infty \}. \tag{169}$$

$L^2(I, q)$ is a Hilbert space with scalar product

$$\langle u_1, u_2 \rangle_q = \int_I r^q u_1 u_2, \tag{170}$$

but let us emphasize that we shall apply this definition only for $q \neq 2$. The scalar product $\langle \cdot, \cdot \rangle_2$ will be defined differently.

We consider real-valued functions for simplicity but we could just as well allow for complex-valued functions with the standard scalar product, or more precisely, sesquilinear form.

Definition 2. For functions $w, u \in C_c^\infty(I)$ define the operators

$$\hat{A}_1 w = -t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{-2} \mu_0 w - t^2 m_2 \Lambda w \tag{171}$$

and

$$A_1 u = -t^{-1} \frac{\partial}{\partial t} \left(t \frac{\partial u}{\partial t} \right) - t^{-2} \bar{\mu} u - t^2 m_2 \Lambda u, \tag{172}$$

as well as the scalar product

$$\langle u_1, u_2 \rangle_2 = \langle Bu_1 + t^2 m_2 u_1, u_2 \rangle_1 \quad \forall u_1, u_2 \in C_c^\infty(I). \tag{173}$$

The right-hand side of (173) is an integral. Integrating by parts, we deduce

$$\langle u_1, u_2 \rangle_2 = \int_I (t \dot{u}_1 \dot{u}_2 - \bar{\mu} t^{-1} u_1 u_2 + t^3 m_2 u_1 u_2), \tag{174}$$

i.e., the scalar product is indeed positive definite due to the assumption (163). Let us define the norm

$$\|u\|_2^2 = \langle u, u \rangle_2 \quad \forall u \in C_c^\infty(I) \tag{175}$$

and the Hilbert space $\mathcal{H}_2 = \mathcal{H}_2(I)$ as the closure of $C_c^\infty(I)$ with respect to the norm $\|\cdot\|_2$.

Proposition 1. The functions $u \in \mathcal{H}_2$ have the properties

$$u \in C^0([0, \infty)), \tag{176}$$

$$|u(t)| \leq c \|u\|_2 \quad \forall t \in I, \tag{177}$$

where $c = c(\bar{\mu}, m_2, |\Lambda|)$,

$$\lim_{t \rightarrow 0} u(t) = 0 \tag{178}$$

and

$$|u(t)| \leq c \|u\|_2 t^{-1} \quad \forall t \in I, \tag{179}$$

where c is a different constant depending on $\bar{\mu}, m_2$ and $|\Lambda|$.

Proof. Let us first assume $u \in C_c^\infty(I)$ and let $\delta > 0$; then,

$$u^2(\delta) = 2 \int_0^\delta \dot{u} u \leq \int_0^\delta t |\dot{u}|^2 + \int_0^\delta t^{-1} |u|^2. \tag{180}$$

This estimate is also valid for any $u \in \mathcal{H}_2$ by approximation, which in turn implies the relations (177), (178) and also (176) since u is certainly continuous in I .

It remains to prove (179). Let $u \in \mathcal{H}_2$ and define $v = v(\tau)$ by

$$v(\tau) = u(\tau^{-1}), \tag{181}$$

where $\tau = t^{-1}$ for all $t > 0$. Applying simple calculus arguments, we then obtain

$$\int_0^\infty \{ \tau |v'|^2 - \bar{\mu} \tau^{-1} |v|^2 + m_2 \tau^{-5} |v|^2 \} d\tau = \|u\|_2^2 \tag{182}$$

as well as

$$\int_0^\infty \{\tau|v'|^2 - \bar{\mu}\tau^{-1}|v|^2\}d\tau = \int_0^\infty \{t|u'|^2 - \bar{\mu}t^{-1}|u|^2\}dt. \tag{183}$$

Moreover, first assuming, as before, that u and hence v are test functions, we argue as in (180) that, for any $\delta > 0$,

$$\begin{aligned} v^2(\delta) &= 2 \int_0^\delta v'v \leq 2 \left(\int_0^\delta \tau|v'|^2 \right)^{\frac{1}{2}} \left(\int_0^\delta \tau^{-1}|v|^2 \right)^{\frac{1}{2}} \\ &\leq 2 \left(\int_0^\delta \tau|v'|^2 \right)^{\frac{1}{2}} \left(\int_0^\delta \tau^{-5}|v|^2 \right)^{\frac{1}{2}} \delta^2 \\ &\leq c\|u\|_2^2 \delta^2, \end{aligned} \tag{184}$$

where we used (182) for the last inequality and where $c = c(\bar{\mu}, m_2)$. Setting $\delta = t^{-1}$ for arbitrary $t > 0$, we have proved the estimate (179) for test functions and hence for arbitrary $u \in \mathcal{H}_2$. □

We are now ready to solve the Equation (166) as an implicit eigenvalue equation. First, we need

Lemma 3. *Let K be the quadratic form*

$$K(u) = m_1 \int_I t^{3-\frac{4}{n}} u^2; \tag{185}$$

then, K is compact in \mathcal{H}_2 , i.e.,

$$u_i \xrightarrow{\mathcal{H}_2} u \implies K(u_i) \rightarrow K(u), \tag{186}$$

and positive definite, i.e.,

$$K(u) > 0 \quad \forall u \neq 0. \tag{187}$$

For a proof, we refer to [3] (Lemma 6.8). Then, we look at the eigenvalue problem for $u \in \mathcal{H}_2$

$$Bu + m_2 t^2 u = \lambda m_1 t^{2-\frac{4}{n}} u, \tag{188}$$

or equivalently,

$$\tilde{B}(u, v) \equiv \langle Bu + m_2 t^2 u, v \rangle_1 = \lambda K(u, v) \quad \forall v \in \mathcal{H}_2, \tag{189}$$

where $K(u, v)$ is the bilinear form associated with K .

Theorem 6. *The eigenvalue problem (189) has countably many solutions (λ_i, \tilde{u}_i) , $\tilde{u}_i \in \mathcal{H}_2$, with the properties*

$$\lambda_i < \lambda_{i+1} \quad \forall i \in \mathbb{N}, \tag{190}$$

$$\lim_i \lambda_i = \infty, \tag{191}$$

$$K(\tilde{u}_i, \tilde{u}_j) = \delta_{ij}. \tag{192}$$

The pairs (λ_i, \tilde{u}_i) are recursively defined by the variational problems

$$\lambda_0 = \tilde{B}(\tilde{u}_0) = \inf \left\{ \frac{\tilde{B}(u)}{K(u)} : 0 \neq u \in \mathcal{H}_2 \right\} \tag{193}$$

and for $i > 0$

$$\lambda_i = \tilde{B}(\tilde{u}_i) = \inf \left\{ \frac{\tilde{B}(u)}{K(u)} : 0 \neq u \in \mathcal{H}_2, K(u, u_j) = 0, 0 \leq j \leq i-1 \right\}. \quad (194)$$

The (\tilde{u}_i) form a Hilbert space basis in \mathcal{H}_2 and in $L^2(1, 3 - \frac{4}{n})$, the eigenvalues are strictly positive and the eigenspaces are one-dimensional.

Proof. This theorem is well-known and goes back to the book of Courant–Hilbert [19], though in a general separable Hilbert space the eigenvalues are not all positive and the eigenspaces are only finite dimensional. For a proof in the general case, we refer to [20] (Theorem 1.6.3, p. 37).

The positivity of the eigenvalues in the above theorem is evident and the fact that the eigenspaces are one-dimensional is proved by contradiction. Thus, suppose there exist an eigenvalue $\lambda = \lambda_i$ and two corresponding linearly independent eigenfunctions $u_1, u_2 \in \mathcal{H}_2$. Then, for any $t_0 > 0$, there would exist an eigenfunction $u \in \mathcal{H}_2$ with eigenvalue λ satisfying $v(t_0) = 0$ and Equation (188). Multiplying this equation by u and integrating the result in the interval $(0, t_0)$ with respect to the measure $t dt$ we obtain

$$\int_0^{t_0} -\bar{\mu}t^{-1}u^2 \leq t_0^{4-\frac{4}{n}} \int_0^{t_0} \lambda m_1 t^{-1}u^2, \quad (195)$$

where we used

$$1 \leq \frac{t_0}{t}, \quad \forall t \in (0, t_0), \quad (196)$$

yielding a contradiction if t_0 is sufficiently small. \square

The functions

$$u_i(t) = \tilde{u}_i(\lambda_i^{-\frac{n}{4(n-1)}} t) \quad (197)$$

then satisfy the equation

$$Bu_i + m_2 \lambda_i^{-\frac{n}{n-1}} t^2 u_i = m_1 t^{2-\frac{4}{n}} u_i \quad (198)$$

and they are mutually orthogonal with respect to the bilinear form

$$\int_1 t^3 uv, \quad (199)$$

as one can easily verify. Furthermore, the following lemma is valid:

Lemma 4. Let $(\lambda, u) \in \mathbb{R}_+^* \times \mathcal{H}_2$, be a solution of

$$Bu + m_2 \lambda^{-\frac{n}{n-1}} t^2 u = m_1 t^{2-\frac{4}{n}} u; \quad (200)$$

then, there exists i such that

$$\lambda = \lambda_i \quad \wedge \quad u \in \langle u_i \rangle. \quad (201)$$

Proof. Define

$$\tilde{u}(t) = u(\lambda^{\frac{n}{4(n-1)}} t); \quad (202)$$

then, the pair (λ, \tilde{u}) is a solution of the Equation (188), hence the result. \square

Thus, we have proved

Theorem 7. There are countably many solutions (Λ_i, u_i) of the implicit eigenvalue problem

$$Bu_i - m_2 \Lambda_i t^2 u_i = m_1 t^{2-\frac{4}{n}} u_i \quad (203)$$

with eigenfunctions $u_i \in \mathcal{H}_2$ such that

$$\Lambda_i < \Lambda_{i+1} < 0 \quad \forall i \in \mathbb{N}, \quad (204)$$

$$\lim_i \Lambda_i = 0, \quad (205)$$

and their multiplicities are one. The transformed eigenfunctions

$$\tilde{u}_i(t) = u_i(\lambda_i^{\frac{n}{4(n-1)}} t), \quad (206)$$

where

$$\lambda_i = (-\Lambda_i)^{-\frac{n-1}{n}}, \quad (207)$$

form a basis of \mathcal{H}_2 and also of $L^2(I, 1)$.

Remark 5. The eigenfunctions \tilde{u}_0 resp. u_0 corresponding to the smallest eigenvalues λ_0 resp. Λ_0 do not change sign in I , since

$$\tilde{B}(|u|) \leq \tilde{B}(u) \quad \forall u \in \mathcal{H}_2, \quad (208)$$

in view of (168), and hence we deduce that $|\tilde{u}_0|$ is also an eigenfunction with eigenvalue λ_0 , i.e., we may assume $\tilde{u}_0 \geq 0$. But if \tilde{u}_0 would vanish in a $t_0 > 0$, then its derivative \tilde{u}'_0 would also vanish in t_0 , yielding \tilde{u}_0 , which would completely vanish and represent a contradiction.

In Definition 2, we defined the operators A_1 and \hat{A}_1 . The operator A_1 can be expressed with the help of the Bessel operator B as

$$A_1 u = B u - t^2 m_2 \Lambda u. \quad (209)$$

Let us express \hat{A}_1 similarly as

$$\hat{A}_1 w = \hat{B} w - t^2 m_2 \Lambda w, \quad (210)$$

where

$$\hat{B} w = -t^{-m} \frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{-2} \mu_0 w. \quad (211)$$

We claim that B and \hat{B} are unitarily equivalent.

Lemma 5. Let φ be the linear map from $L^2(I, m)$ to $L^2(I, 1)$ defined by

$$\varphi(w) = u = t^{\frac{m-1}{2}} w. \quad (212)$$

Then, φ is unitary and, if B resp. \hat{B} are defined in $C_c^\infty(I)$, the relation

$$\hat{B} = \varphi^{-1} \circ B \circ \varphi \quad (213)$$

is valid.

Since we assume for simplicity the Hilbert spaces to be real Hilbert spaces, it would be better to call the map φ *orthogonal*, but the result would be the same if we would consider complex-valued functions and the corresponding scalar products.

For the simple proof of the lemma, we refer to [3] (Lemma 4.1). Moreover, for any measurable function $f = f(t)$, we have

$$\langle f \varphi(w), \varphi(v) \rangle_1 = \langle f w, v \rangle_m \quad \forall v, w \in C_c^\infty(I). \quad (214)$$

Hence, we infer

$$\begin{aligned} \langle \varphi(w), \varphi(v) \rangle_2 &= \langle B\varphi(w) + t^2 m_2 \varphi(w), \varphi(v) \rangle_1 \\ &= \langle \hat{B}w + t^2 m_2 w, v \rangle_m \quad \forall v, w \in C_c^\infty(I), \end{aligned} \tag{215}$$

$$\langle \hat{B}w, v \rangle_m = \langle B\varphi(w), \varphi(v) \rangle_1 \tag{216}$$

and we deduce, by setting $u = \varphi(w) = t^{-\frac{m-1}{2}} w$, that

$$\langle \hat{B}w, w \rangle_m = \langle Bu, u \rangle_1 = \int_I (t|\dot{u}|^2 - \bar{\mu}t^{-1}|u|^2) > 0, \tag{217}$$

or equivalently,

$$\int_I t^m |\dot{w}|^2 = \int_I (t|\dot{u}|^2 - \bar{\mu}t^{-1}|u|^2) + \mu_0 \int_I t^{m-2}|w|^2 \quad \forall w \in C_c^\infty(I). \tag{218}$$

Let us recall that $\bar{\mu} < 0$ and $\mu_0 > 0$.

Remark 6. Defining the Hilbert space $\hat{\mathcal{H}}_2$ by

$$\hat{\mathcal{H}}_2 = \{ w = t^{-\frac{m-1}{2}} u : u \in \mathcal{H}_2 \} \tag{219}$$

with norm

$$\|w\|_2 = \|u\|_2 \tag{220}$$

and the quadratic form \hat{K} by

$$\hat{K}(w) = \langle m_1 t^{2-\frac{4}{n}} w, w \rangle_m = \langle m_1 t^{2-\frac{4}{n}} u, u \rangle_1 = K(u) \quad \forall w \in \hat{\mathcal{H}}_2 \tag{221}$$

it is fairly easy to verify that all results in Theorem 6 remain valid if $B, \tilde{B}, K, \mathcal{H}_2$ are replaced by $\hat{B}, \hat{\tilde{B}}, \hat{K}, \hat{\mathcal{H}}_2$. The eigenvalues λ_i are identical and the eigenfunctions are related by

$$\tilde{w}_i = t^{-\frac{m-1}{2}} \tilde{u}_i, \tag{222}$$

i.e.,

$$\hat{B}\tilde{w}_i + t^2 m_2 t^2 \tilde{w}_i = \lambda_i m_1 t^{2-\frac{4}{n}} \tilde{w}_i. \tag{223}$$

Similarly, the transformed eigenfunctions u_i in Theorem 7 correspond to

$$w_i = t^{-\frac{m-1}{2}} u_i \tag{224}$$

satisfying

$$\hat{B}w_i - m_2 \Lambda_i t^2 u_i = m_1 t^{2-\frac{4}{n}} w_i, \tag{225}$$

which is the original ODE (147) on page 15 with $\Lambda = \Lambda_i$.

For completeness, let us restate Theorem 7 in the new setting

Theorem 8. There are countably many solutions (Λ_i, w_i) of the implicit eigenvalue problem

$$\hat{B}w_i - m_2 \Lambda_i t^2 w_i = m_1 t^{2-\frac{4}{n}} w_i \tag{226}$$

with eigenfunctions $w_i \in \hat{\mathcal{H}}_2$ such that

$$\Lambda_i < \Lambda_{i+1} < 0 \quad \forall i \in \mathbb{N}, \tag{227}$$

$$\lim_i \Lambda_i = 0, \tag{228}$$

and their multiplicities are one. The transformed eigenfunctions

$$\tilde{w}_i(t) = w_i(\lambda_i^{\frac{n}{4(n-1)}} t), \tag{229}$$

where

$$\lambda_i = (-\Lambda_i)^{-\frac{n-1}{n}}, \tag{230}$$

form a basis of $\hat{\mathcal{H}}_2$ and also of $L^2(I, m)$.

Finally, let us show how the eigenvalue Equations (188) resp. (223) can be considered to be eigenvalue equations of an essentially self-adjoint operator in an appropriate Hilbert space. We shall first demonstrate it for the Equation (188).

Let $\varphi_0(t)$ be defined by

$$\varphi_0(t) = m_1 t^{3-\frac{4}{n}} \quad \forall t \in I \tag{231}$$

and define the Hilbert space \mathcal{H} as $L^2(I, d\mu)$ with respect to the measure

$$d\mu = \varphi_0 dt. \tag{232}$$

Moreover, denote the scalar product in \mathcal{H} by $\langle \cdot, \cdot \rangle$ and the corresponding norm by $\|\cdot\|$. Note that, in view of (185),

$$\langle u, v \rangle = K(u, v). \tag{233}$$

The operator

$$Au = \varphi_0^{-1} \left\{ - \left(\frac{\partial}{\partial t} \left(t \frac{\partial u}{\partial t} \right) - t^{-1} \bar{\mu} u + t^3 m_2 u \right) \right\} \quad \forall u \in C_c^\infty(I) \tag{234}$$

is densely defined and symmetric in \mathcal{H} such that

$$\langle Au, v \rangle = \langle u, v \rangle_2 \quad \forall u, v \in C_c^\infty(I) \tag{235}$$

The above relation is also valid for all $u, v \in \mathcal{H}_2$ by partial integration. Hence, the domain $D(A)$ of A is contained in \mathcal{H}_2 . In view of Equation (188), we infer

$$A\tilde{u}_i = \lambda_i \tilde{u}_i, \quad \forall i \in \mathbb{N}, \tag{236}$$

i.e., \tilde{u}_i is an eigenfunction of A in the classical sense. Since A is symmetric A is closable. Let \bar{A} be the closure of A . If \bar{A} is surjective

$$R(\bar{A}) = \mathcal{H}, \tag{237}$$

then \bar{A} is self-adjoint and A is essentially self-adjoint. These are well-known facts. Let us prove (237) for convenience.

Lemma 6. \bar{A} is surjective.

Proof. First, we observe that $R(A)$ is dense in \mathcal{H} due to (236). Indeed, the eigenfunctions $(\tilde{u}_i), i \in \mathbb{N}$, are complete and the eigenvalues are strictly positive, cf. Theorem 6.

Next, let $v \in \mathcal{H}$ be arbitrary and let $u_i \in D(A)$ be a sequence such that

$$Au_i \rightarrow v; \tag{238}$$

then,

$$\begin{aligned}\lambda_0 \|u_i - u_j\|^2 &= \lambda_0 \langle u_i - u_j, u_i - u_j \rangle \leq \langle A(u_i - u_j), u_i - u_j \rangle \\ &\leq \|A(u_i - u_j)\| \|u_i - u_j\|,\end{aligned}\quad (239)$$

where $0 < \lambda_0$ is the smallest eigenvalue, cf. (193). Hence,

$$\lambda_0 \|u_i - u_j\| \leq \|A(u_i - u_j)\|, \quad (240)$$

i.e., (u_i) is a Cauchy sequence which implies $v \in R(\bar{A})$, completing the proof of the lemma. \square

In case of Equation (223), we define $\hat{\varphi}_0(t)$ by

$$\hat{\varphi}_0(t) = m_1 t^{m+2-\frac{4}{n}} \quad \forall t \in I \quad (241)$$

and define the Hilbert space $\hat{\mathcal{H}}$ as $L^2(I, d\hat{\mu})$ with respect to the measure

$$d\hat{\mu} = \hat{\varphi}_0 dt. \quad (242)$$

Moreover, denote the scalar product in $\hat{\mathcal{H}}$ by $\langle\langle \cdot, \cdot \rangle\rangle$ and the corresponding norm by $\|\cdot\|$. Note that, in view of (221),

$$\langle\langle w, v \rangle\rangle = \hat{K}(w, v) \quad (243)$$

The operator

$$\hat{A}w = \hat{\varphi}_0^{-1} \left\{ - \left(\frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{m-2} \mu_0 w + t^{m+2} m_2 w \right) \right\} \quad \forall w \in C_c^\infty(I) \quad (244)$$

is densely defined and symmetric in $\hat{\mathcal{H}}$ such that

$$\langle\langle \hat{A}w_1, w_2 \rangle\rangle = \langle Au_1, u_2 \rangle = \langle u_1, u_2 \rangle_2 \quad \forall w_1, w_2 \in C_c^\infty(I), \quad (245)$$

where

$$u_i = \varphi(w_i) \equiv t^{\frac{m-1}{2}} w_i, \quad i = 1, 2, \quad (246)$$

cf. the definition of φ in Lemma 5 and also the Equation (215). If Equation (245) would be valid for all $w_1, w_2 \in D(\hat{A})$, then \hat{A} and A would be unitarily equivalent, since φ is evidently a unitary (orthogonal) map between $\hat{\mathcal{H}}$ and \mathcal{H} .

This is indeed the case as one can easily infer from Remark 6; hence,

$$\hat{A}\tilde{w}_i = \lambda_i \tilde{w}_i, \quad (247)$$

where

$$\tilde{w}_i = t^{-\frac{m-1}{2}} \tilde{u}_i \quad (248)$$

and \tilde{u}_i an eigenfunction A with eigenvalue λ_i . The domain of \hat{A} satisfies

$$D(\hat{A}) = \varphi^{-1}(D(A)). \quad (249)$$

5.2. Treating Λ as a Fixed Cosmological Constant

If we want to define a partition function and entropy for our quantum system, we have to consider Λ to be a fixed cosmological constant and not a parameter which can also

play the role of an implicit eigenvalue. Our approach to solve the ODE (147) on page 15, then, is similar but different. First, let us express Equation (147) in the equivalent form

$$\hat{\varphi}_0^{-1} \left\{ -\frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{m-2} \mu_0 w - t^{m+2} m_2 \Lambda w \right\} - \frac{16(n-1)}{n} \{ (n-1)|\zeta|^2 + \bar{\mu}_k \} w = 0, \tag{250}$$

where

$$\hat{\varphi}_0(t) = t^{m+2-\frac{4}{n}} \tag{251}$$

and where we used the definition (145) on page 15 of m_1 . The term

$$(n-1)|\zeta|^2 + \bar{\mu}_k \tag{252}$$

is an eigenvalue of the operator in (134) on page 14. $|\zeta|^2$ with $\zeta \in \mathbb{R}^{m_1}$ is a continuous eigenvalue while the sequence $\bar{\mu}_k, k \in \mathbb{N}$ satisfies the relations (139) and (140). The operator

$$\hat{H}_0 w \equiv \hat{\varphi}_0^{-1} \left\{ -\frac{\partial}{\partial t} \left(t^m \frac{\partial w}{\partial t} \right) - t^{m-2} \mu_0 w - t^{m+2} m_2 \Lambda w \right\} \tag{253}$$

is identical to the operator \hat{A} defined in (244) if $\Lambda = -1$. The properties we proved for \hat{A} are also valid for \hat{H}_0 by simply replacing $-m_2 \Lambda$ by a positive constant m'_2 . Thus, we know that \hat{H}_0 is essentially self-adjoint in the Hilbert space $\hat{\mathcal{H}} = L^2(I, d\hat{\mu})$, cf. (242) with a complete system of eigenfunctions $w_i, i \in \mathbb{N}$, and corresponding eigenvalues

$$0 < \lambda_0 < \lambda_1 < \lambda_2 < \dots \tag{254}$$

The eigenspaces are all one-dimensional and the ground state w_0 does not change sign, cf. Remark 5 on page 21.

Note that we denote the eigenfunctions by w_i and not by \tilde{w}_i since they will not be transformed to obtain the final solutions of the ODE. Instead, they will be the solutions of the ODE satisfying

$$\hat{H}_0 w_i = \lambda_i w_i \quad \forall i \in \mathbb{N}. \tag{255}$$

But w_i is a solution of the ODE (250) if and only if there exist j and ζ such that

$$\lambda_i = \frac{16(n-1)}{n} \{ (n-1)|\zeta|^2 + \bar{\mu}_j \} \tag{256}$$

Evidently, the previous equation can only be satisfied for all λ_i if

$$\lambda_0 \geq \frac{16(n-1)}{n} \bar{\mu}_0. \tag{257}$$

In [12] (Lemma 6.4.9, p. 172), we proved the following lemma:

Lemma 7. *Let λ_i be the temporal eigenvalues depending on $\Lambda < 0$ and let $\bar{\lambda}_i$ be the corresponding eigenvalues for*

$$\Lambda = -1; \tag{258}$$

then,

$$\lambda_i = \bar{\lambda}_i |\Lambda|^{\frac{n-1}{n}}. \tag{259}$$

Thus, we deduce

Corollary 1. Suppose that $\bar{\mu}_0 > 0$ and define $\Lambda_0 < 0$ by

$$|\Lambda_0|^{\frac{n-1}{n}} = \bar{\lambda}_0^{-1} \frac{16(n-1)}{n} \bar{\mu}_0; \quad (260)$$

then, the inequality (257) is satisfied provided

$$|\Lambda| \geq |\Lambda_0|. \quad (261)$$

The inequality (257) is always satisfied if $\bar{\mu}_0 \leq 0$.

The eigenvalues on the right-hand side of Equation (256), i.e., the sum inside the braces, are the eigenvalues of the operator defined in (134) on page 14 which can be written as the sum

$$-(n-1)\Delta_\delta + A, \quad (262)$$

where A is a uniformly elliptic operator on a compact Riemannian manifold, cf. Equation (138) on page 14. Hence, we can interpret the right-hand side of (256) as eigenvalues of the operator

$$H_1 = -\frac{16(n-1)^2}{n}\Delta_\delta + \frac{16(n-1)}{n}A. \quad (263)$$

To facilitate a comparison with former results in [12] (Sections 6.4 and 6.5), let us define

$$\tilde{A} = \frac{16(n-1)}{n}A \quad (264)$$

and

$$\tilde{\mu}_j = \frac{16(n-1)}{n}\bar{\mu}_j; \quad (265)$$

then, \tilde{A} has the same eigenfunctions as A with eigenvalues $\tilde{\mu}_j$ instead of $\bar{\mu}_j$ and the condition (256) can be rephrased in the form

$$\lambda_i = \frac{16(n-1)^2}{n}|\zeta|^2 + \tilde{\mu}_j \quad (266)$$

and the inequality (257) can now be expressed as

$$\lambda_0 \geq \tilde{\mu}_0. \quad (267)$$

In [12] (equ. (6.4.67), p.166), we considered an operator H_1 which was similarly defined as the operator in (263); the only difference was that the Laplacian Δ_δ was defined in \mathbb{R} , i.e., the dimension n_1 was equal to one. In this case, it is fairly simple to determine the tempered eigendistributions ζ_{ijk} in $\mathcal{S}'(\mathbb{R})$ satisfying

$$-\zeta_{ijk}'' = \omega_{ij}^2 \zeta_{ijk}, \quad k = 1, 2, \quad (268)$$

where

$$\zeta_{ij1}(\tau) = \frac{1}{\sqrt{2\pi}} e^{i\omega_{ij}\tau} \quad (269)$$

and

$$\zeta_{ij2}(\tau) = \frac{1}{\sqrt{2\pi}} e^{-i\omega_{ij}\tau}, \quad (270)$$

where

$$\omega_{ij} \geq 0 \quad (271)$$

is defined by the relation

$$\lambda_i = \tilde{\mu}_j + \frac{16(n-1)^2}{n}\omega_{ij}^2. \quad (272)$$

In the higher dimensional case, $n_1 > 1$, we have a whole continuum of vectors $\zeta \in \mathbb{R}^{n_1}$ satisfying (266), and hence, a whole continuum of eigendistributions which we cannot handle—neither physically nor mathematically. Therefore, let us pick a finite numbers of unit vectors $\zeta_k \in \mathbb{R}^{n_1}, 1 \leq k \leq k_1$ which are fixed. Then, the eigendistributions are defined by

$$\zeta_{ijk}(y) = (2\pi)^{-\frac{n_1}{2}} e^{i\omega_{ij}\langle \zeta_k, y \rangle}, \quad 1 \leq k \leq k_1, \tag{273}$$

where

$$\lambda_i = \tilde{\mu}_j + \frac{16(n-1)^2}{n} \omega_{ij}^2 \tag{274}$$

if $\tilde{\mu}_j < \lambda_i$. We consider the eigendistributions ζ_{ijk} to be mutually orthogonal since their Fourier transforms

$$\hat{\zeta}_{ijk} = \delta_{\omega_{ij}\zeta}, \tag{275}$$

which are Dirac measures, have disjoint supports.

Now, we are able to define the eigenfunctions of the operator H_1 in (263).

Theorem 9. Let $\varphi_j \in L^2(M)$ be the mutually orthogonal unit eigenvectors of \tilde{A} with corresponding eigenvalues $\tilde{\mu}_j$ and assume either that $\tilde{\mu}_0 \leq 0$ or that Λ satisfies the condition (261) in Corollary 1. Then, for any eigenvalue λ_i , we define

$$N_i = \{j \in \mathbb{N} : \tilde{\mu}_j \leq \lambda_i\} \tag{276}$$

and $\omega_{ijk} \geq 0$ such that

$$\frac{16(n-1)^2}{n} \omega_{ijk}^2 + \tilde{\mu}_j = \lambda_i, \quad 1 \leq k \leq k_1, \tag{277}$$

provided $\tilde{\mu}_j < \lambda_i$. If $\tilde{\mu}_j = \lambda_i$, then we choose $\omega_{ijk} = 0$ and the multiplicity will be only the multiplicity of $\tilde{\mu}_j$.

Note that

$$0 \in N_i \quad \forall i \in \mathbb{N}, \tag{278}$$

since

$$\tilde{\mu}_0 \leq \tilde{\lambda}_0, \tag{279}$$

For $j \in N_i$, define the eigenfunctions v_{ijk} of H_1 by

$$v_{ijk} = \zeta_{ijk} \varphi_j, \tag{280}$$

where this distinction only occurs if

$$\tilde{\mu}_j < \lambda_i, \tag{281}$$

such that

$$H_1 v_{ijk} = \lambda_i v_{ijk}. \tag{282}$$

Remark 7. H_1 has the same eigenvalues λ_i as \hat{H}_0 but with finite multiplicities $m(\lambda_i)$ in general different from one which can be estimated from above by

$$m(\lambda_i) \leq k_1 \text{ card } N_i \equiv k_1 n(\lambda_i). \tag{283}$$

Recall that we labelled the eigenvalues $\tilde{\mu}_j$ by including their multiplicities, cf. (139) on page 15. Hence, if

$$\tilde{\mu}_j < \lambda_i \quad \forall j \in N_i \tag{284}$$

then

$$m(\lambda_i) = k_1 n(\lambda_i). \tag{285}$$

Let us now define a separable Hilbert space \mathcal{H} such that H_1 is essentially self-adjoint in \mathcal{H} and its eigenvectors with eigenvalues λ_i form an ONB, an orthonormal basis.

First we declare the countable eigenvectors in (282) to be mutually orthogonal unit vectors and we consider them to be the Hamel basis of the complex vector space \mathcal{H}' . Since the basis vectors are mutually orthogonal unit vectors, they also define a unique hermitian scalar product in \mathcal{H}' . Let \mathcal{H} be the completion of \mathcal{H}' with respect to that scalar product. Since the eigenvalues λ_i are positive and bounded from below by λ_0 , we could prove in [12] (Lemma 6.5.1, p. 174) the following lemma:

Lemma 8. *The linear operator H_1 with domain \mathcal{H}' is essentially self-adjoint in \mathcal{H} . Let \bar{H}_1 be its closure; then, the only eigenvectors of \bar{H}_1 are those of H_1 .*

Remark 8. *In the following, we shall write H_1 instead of \bar{H}_1 and we also let $\tilde{\lambda}_i$ be a relabelling of the eigenvalues λ_i of H_1 to include the multiplicities.*

In [12] (Lemma 6.5.3, p. 175), we also proved

Lemma 9. *For any $\beta > 0$, the operator*

$$e^{-\beta H_1} \quad (286)$$

is of trace class in \mathcal{H} , i.e.,

$$\text{tr}(e^{-\beta H_1}) = \sum_{i=0}^{\infty} e^{-\beta \tilde{\lambda}_i} < \infty. \quad (287)$$

Let

$$\mathcal{F} \equiv \mathcal{F}_+(\mathcal{H}) \quad (288)$$

be the symmetric Fock space generated by \mathcal{H} and let

$$H = d\Gamma(H_1) \quad (289)$$

be the canonical extension of H_1 to \mathcal{F} . Then,

$$e^{-\beta H} \quad (290)$$

is also of trace class in \mathcal{F}

$$\text{tr}(e^{-\beta H}) = \prod_{i=0}^{\infty} (1 - e^{-\beta \tilde{\lambda}_i})^{-1} < \infty, \quad (291)$$

where $\tilde{\lambda}_i$ is a relabelling of the eigenvalues λ_i to include the multiplicities.

The proof relies on the fact that a temporal Hamiltonian H_0 , which is similarly defined as the operator \hat{H}_0 in (253), has these properties.

For the present operator \hat{H}_0 , it is also valid that $e^{-\beta \hat{H}_0}$ is of trace class and the proof of this property is very similar to the proof we gave in [12] (Theorem 6.2.8, p. 148); however, the structure of the operator in (253) is slightly different so that we cannot simply refer to the previous result. We shall give a proof in the next section instead.

Remark 9. *In [12] (Chapter 6.5), we used these results to define the partition function Z by*

$$Z = \text{tr}(e^{-\beta H}) = \prod_{i=0}^{\infty} (1 - e^{-\beta \tilde{\lambda}_i})^{-1} \quad (292)$$

and the density operator ρ in \mathcal{F} by

$$\rho = Z^{-1} e^{-\beta H} \quad (293)$$

such that

$$\text{tr } \rho = 1. \tag{294}$$

The von Neumann entropy S is then defined by

$$\begin{aligned} S &= -\text{tr}(\rho \log \rho) \\ &= \log Z + \beta Z^{-1} \text{tr}(He^{-\beta H}) \\ &= \log Z - \beta \frac{\partial \log Z}{\partial \beta} \\ &\equiv \log Z + \beta E, \end{aligned} \tag{295}$$

where E is the average energy

$$E = \text{tr}(H\rho). \tag{296}$$

E can be expressed in the form

$$E = \sum_{i=0}^{\infty} \frac{\tilde{\lambda}_i}{e^{\beta \tilde{\lambda}_i} - 1}. \tag{297}$$

Here, we also set the Boltzmann constant

$$k_B = 1. \tag{298}$$

The parameter β is supposed to be the inverse of the absolute temperature T

$$\beta = T^{-1}. \tag{299}$$

For a more detailed analysis and especially for the dependence on Λ , we refer to [12] (Chapter 6.5).

6. Trace Class Estimates for $e^{-\beta H_0}$

Let us first consider the operator

$$H_0 u = \varphi_0^{-1} \left\{ -\left(\frac{\partial}{\partial t}\right)\left(t\frac{\partial u}{\partial t}\right) - t^{-1}\bar{\mu}u + t^3 m_2 |\Lambda|u \right\} \quad \forall u \in C_c^\infty(I) \tag{300}$$

which is unitarily equivalent to the operator in (253) on page 25. H_0 is essentially self-adjoint in

$$\mathcal{H} = L^2(\mathbb{R}_+, d\mu), \tag{301}$$

where

$$d\mu = \varphi_0 dt \tag{302}$$

with

$$\varphi_0(t) = t^{3-\frac{4}{n}}. \tag{303}$$

We shall use the same symbol for its closure, i.e., we shall assume that H_0 is self-adjoint in \mathcal{H} with eigenvectors $u_i \in \mathcal{H}_2$, cf. the remarks following (236) on page 23, and with eigenvalues λ_i satisfying the statements in Theorem 6 on page 19, where now we denote the eigenvectors by u_i , since they will not be transformed.

Remark 10. The norm

$$\langle H_0 u, u \rangle^{\frac{1}{2}} \tag{304}$$

is equivalent to the norm $\|u\|_2$ in \mathcal{H}_2 , cf. (174) and (175) on page 18.

Let us also assume that all Hilbert spaces are complex vector spaces with a positive definite sesquilinear form (hermitian scalar product).

We shall now prove that

$$e^{-\beta H_0}, \quad \beta > 0, \quad (305)$$

is of trace class in \mathcal{H} . The proof is essentially the proof given in [12] (Chapter 6.2) with the necessary modifications due to the different structure of the operator.

First, we need two lemmata:

Lemma 10. *The embedding*

$$j : \mathcal{H}_2 \hookrightarrow \mathcal{H}_0 = L^2(\mathbb{R}_+, d\tilde{\mu}), \quad (306)$$

where

$$d\tilde{\mu} = (1+t)^{-2} dt, \quad (307)$$

is Hilbert–Schmidt, i.e., for any ONB (e_i) in \mathcal{H}_2 , the sum

$$\sum_{i=0}^{\infty} \|j(e_i)\|_0^2 < \infty \quad (308)$$

is finite, where $\|\cdot\|_0$ is the norm in \mathcal{H}_0 . The square root of the left-hand side of (308) is known as the Hilbert–Schmidt norm $|j|$ of j and it is independent of the ONB.

Proof. Let $w \in \mathcal{H}_2$; then, assuming that w is real-valued,

$$\begin{aligned} |w(t)|^2 &= 2 \int_0^t \dot{w} w \leq \int_0^\infty t |\dot{w}|^2 + \int_0^\infty t^{-1} |w|^2 \\ &\leq c \|w\|_2^2 \end{aligned} \quad (309)$$

for all $t > 0$, where $\|\cdot\|_2$ is the norm in \mathcal{H}_2 . To derive the last inequality in (309), we used (174) and (163) on page 17. The estimate

$$|w(t)| \leq c \|w\|_2 \quad \forall t > 0 \quad (310)$$

is of course also valid for complex-valued functions from which we infer that, for any $t > 0$, the linear form

$$w \rightarrow w(t), \quad w \in \mathcal{H}_2, \quad (311)$$

is continuous; hence, it can be expressed as

$$w(t) = \langle \varphi_t, w \rangle, \quad (312)$$

where

$$\varphi_t \in \mathcal{H}_2 \quad (313)$$

and

$$\|\varphi_t\|_2 \leq c. \quad (314)$$

Now, let

$$e_i \in \mathcal{H}_2 \quad (315)$$

be an ONB; then,

$$\sum_{i=0}^{\infty} |e_i(t)|^2 = \sum_{i=0}^{\infty} |\langle \varphi_t, e_i \rangle|^2 = \|\varphi_t\|_2^2 \leq c^2. \quad (316)$$

Integrating this inequality over \mathbb{R}_+ with respect to $d\tilde{\mu}$, we infer

$$\sum_{i=0}^{\infty} \int_0^\infty |e_i(t)|^2 d\tilde{\mu} \leq c^2 \quad (317)$$

completing the proof of the lemma. \square

Lemma 11. *Let u_i be the eigenfunctions of H_0 ; then, there exist positive constants c and γ such that*

$$\|u_i\|_2 \leq c|1 + \lambda_i|^\gamma \|u_i\|_0 \quad \forall i \in \mathbb{N}, \tag{318}$$

where $\|\cdot\|_0$ is the norm in \mathcal{H}_0 .

Proof. We have

$$\langle H_0 u_i, u_i \rangle = \lambda_i \langle u_i, u_i \rangle \tag{319}$$

and hence, in view of Remark 10,

$$\begin{aligned} \|u_i\|_2^2 &\leq c_1 \lambda_i \int_0^\infty \varphi_0(t) |u_i|^2 \\ &\leq c_1 \lambda_i \left\{ \int_0^1 \varphi_0(t) |u_i|^2 + c_2 \int_1^\infty t^{3-\frac{4}{n}} |u_i|^2 \right\}. \end{aligned} \tag{320}$$

To estimate the second integral in the braces, let us define $p = 3$ and observe that

$$3 - \frac{4}{n} \leq p - \frac{p}{n}, \tag{321}$$

and hence,

$$t^{3-\frac{4}{n}} \leq t^{p-\frac{p}{n}} \quad \forall t \geq 1. \tag{322}$$

Then, choosing small positive constants δ and ϵ , we apply Young's inequality, with

$$q = \frac{p}{p - p\delta} = \frac{1}{1 - \delta} \tag{323}$$

and

$$q' = \delta^{-1} \tag{324}$$

to estimate the integral from above by

$$\begin{aligned} &\frac{1}{q} \epsilon^q \int_1^\infty \left\{ t^{p-\frac{p}{n}} (1+t)^{\frac{p}{n}-p\delta} \right\}^q |u_i|^2 \\ &+ \frac{1}{q'} \epsilon^{-q'} \int_1^\infty (1+t)^{-(\frac{p}{n}-p\delta)q'} |u_i|^2. \end{aligned} \tag{325}$$

Choosing, now, δ so small such that

$$\left(\frac{p}{n} - p\delta\right)\delta^{-1} > 2 \tag{326}$$

the preceding integrals can be estimated from above by

$$\frac{1}{q} \epsilon^q \int_1^\infty (1+t)^p |u_i|^2 + \frac{1}{q'} \epsilon^{-q'} \int_0^\infty (1+t)^{-2} |u_i|^2 \tag{327}$$

which in turn can be estimated by

$$\frac{1}{q} \epsilon^q c \|u_i\|_2^2 + \frac{1}{q'} \epsilon^{-q'} \|u_i\|_0^2, \tag{328}$$

in view of Remark 10.

The first integral in the braces on the right-hand side of (320) can be estimated by

$$\begin{aligned} \int_0^1 \varphi_0(t)|u_i|^2 &\leq \frac{1}{2}c\epsilon^2 \int_0^1 |u_i|^2 \\ &\quad + \frac{1}{2}\epsilon^{-2} \int_0^\infty (1+t)^{-2}|u_i|^2 \\ &\leq \tilde{c}\epsilon^2 \|u_i\|_2^2 + \frac{1}{2}\epsilon^{-2} \|u_i\|_0^2. \end{aligned} \tag{329}$$

Choosing now ϵ, γ and c appropriately, the result follows. \square

We are now ready to prove:

Theorem 10. *Let $\beta > 0$; then, the operator*

$$e^{-\beta H_0} \tag{330}$$

is of trace class in \mathcal{H} , i.e.,

$$\text{tr}(e^{-\beta H_0}) = \sum_{i=0}^\infty e^{-\beta \lambda_i} = c(\beta) < \infty. \tag{331}$$

Proof. In view of Lemma 10, the embedding

$$j : \mathcal{H}_2 \hookrightarrow \mathcal{H}_0 \tag{332}$$

is Hilbert–Schmidt. Let

$$u_i \in \mathcal{H} \tag{333}$$

be an ONB of eigenfunctions; then,

$$\begin{aligned} e^{-\beta \lambda_i} &= e^{-\beta \lambda_i} \|u_i\|^2 \leq e^{-\beta \lambda_i} c \lambda_i^{-1} \|u_i\|_2^2 \\ &\leq e^{-\beta \lambda_i} \lambda_i^{-1} c |\lambda_i + 1|^{2\gamma} \|u_i\|_0^2, \end{aligned} \tag{334}$$

in view of (318), but

$$\|u_i\|_0^2 = \|u_i\|_2^2 \|\tilde{u}_i\|_0^2 \leq c \lambda_i \|\tilde{u}_i\|_0^2, \tag{335}$$

where

$$\tilde{u}_i = u_i \|u_i\|_2^{-1} \tag{336}$$

is an ONB in \mathcal{H}_2 , yielding

$$\sum_{i=0}^\infty e^{-\beta \lambda_i} \leq c_\beta \sum_{i=0}^\infty \|\tilde{u}_i\|_0^2 < \infty, \tag{337}$$

since j is Hilbert–Schmidt. Here, we also used that $\lambda_0 > 0$. \square

Since the operator in (253) on page 25 has the same eigenvalues as the operator in (300), we have also proved the following:

Theorem 11. *The operator \hat{H}_0 in (253), which is self-adjoint in the Hilbert space $\hat{\mathcal{H}}$, has the property that*

$$e^{-\beta \hat{H}_0}, \quad \beta > 0, \tag{338}$$

is of trace class in $\hat{\mathcal{H}}$.

7. Conclusions

We quantized the full Einstein equations and found solutions to the resulting hyperbolic equation in a fiber bundle E which can be expressed as a product of spatial

eigenfunctions (eigendistributions) and temporal eigenfunctions. The spatial eigenfunctions form a basis in an appropriate Hilbert space, while the temporal eigenfunctions are solutions to a second-order ODE in \mathbb{R}_+ .

The base space \mathcal{S}_0 with dimension $n \geq 3$ is a Cauchy hypersurface of the quantized spacetime N . The solutions u of the hyperbolic equation in E are evaluated at $(t, x, \chi(x))$, where χ is the metric of the Cauchy hypersurface. The main assumptions for proving the existence of spatial eigenfunctions that also form a basis of a Hilbert space is that \mathcal{S}_0 is a metric product as described in (130) and (132) on page 14, where the compact part M_0 of the product may in general be hidden from observations. In case of Schwarzschild and Kerr-AdS black holes being considered in [21,22], these assumptions are satisfied.

For large n , $n \geq 17$ and negative Λ , the temporal eigenfunctions are also the eigenfunctions of a self-adjoint operator, the eigenvalues are countable and either Λ plays the role of an implicit eigenvalue, cf. Theorem 9 on page 27, or $\Lambda < 0$ is considered to be a fixed cosmological constant, in which case the temporal eigenfunctions are eigenfunctions of a self-adjoint operator \hat{H}_0 and a subset of the spatial eigenfunctions are eigenfunctions of a self-adjoint operator H_1 acting in \mathcal{S}_0 such that \hat{H}_0 and H_1 have the same eigenvalues but with different multiplicities. The operators

$$e^{-\beta\hat{H}_0} \quad \wedge \quad e^{-\beta H_1} \quad (339)$$

are of trace class in their respective Hilbert spaces and also in the corresponding symmetric Fock spaces. The latter result makes it possible to define a partition function Z , a density operator ρ , the von Neumann entropy S and the average energy E of the quantum system, cf. Lemma 9 on page 28 and [12] (Chapter 6.5).

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declares no conflicts of interest.

References

1. Arnowitt, R.; Deser, S.; Misner, C.W. The dynamics of general relativity, In *Gravitation: An Introduction to Current Research*; Witten, L., Ed.; John Wiley: New York, NY, USA, 1962; pp. 227–265.
2. DeWitt, B.S. Quantum Theory of Gravity. I. The Canonical Theory. *Phys. Rev.* **1967**, *160*, 1113–1148. [[CrossRef](#)]
3. Gerhardt, C. Quantum cosmological Friedman models with an initial singularity. *Class. Quantum Gravity* **2009**, *26*, 015001. [[CrossRef](#)]
4. Kiefer, C.; Sandhöfer, B. Quantum cosmology. *Z. Für Naturforschung A* **2022**, *77*, 543–559. [[CrossRef](#)]
5. Misner, C.W. Quantum Cosmology. I. *Phys. Rev.* **1969**, *186*, 1319–1327. [[CrossRef](#)]
6. Moniz, P.V. (Ed.) *Quantum Cosmology*; MDPI: Basel, Switzerland, 2022. [[CrossRef](#)]
7. Unruh, W. G. Unimodular theory of canonical quantum gravity. *Phys. Rev. D* **1989**, *40*, 1048–1052. [[CrossRef](#)] [[PubMed](#)]
8. Kiefer, C. *Quantum Gravity*, 2nd ed.; International Series of Monographs on Physics; Oxford University Press: Oxford, UK, 2007.
9. Thiemann, T. Modern canonical quantum general relativity. In *Cambridge Monographs on Mathematical Physics*; With a foreword by Chris Isham; Cambridge University Press: Cambridge, UK, 2007.
10. Gerhardt, C. The quantization of gravity in globally hyperbolic spacetimes. *Adv. Theor. Math. Phys.* **2013**, *17*, 1357–1391.
11. Gerhardt, C. The quantization of gravity. *Adv. Theor. Math. Phys.* **2018**, *22*, 709–757.
12. Gerhardt, C. *The Quantization of Gravity*, 1st ed.; Fundamental Theories of Physics; Springer: Cham, Switzerland, 2018; Volume 194. [[CrossRef](#)]
13. Jorgenson, J.; Lang, S. *Spherical Inversion on $SL_n(\mathbb{R})$* ; Springer: New York, NY, USA, 2001. [[CrossRef](#)]
14. Gerhardt, C. The quantization of gravity: Quantization of the Hamilton equations. *Universe* **2021**, *7*, 91. [[CrossRef](#)]
15. Gerhardt, C. A unified quantization of gravity and other fundamental forces of nature. *Universe* **2022**, *8*, 404. [[CrossRef](#)]
16. Dacorogna, B.; Moser, J. On a partial differential equation involving the jacobian determinant. *Ann. De L'I.H.P. Anal. Non Linéaire* **1990**, *7*, 1–26. [[CrossRef](#)]
17. Helgason, S. Geometric analysis on symmetric spaces. *Math. Surv. Monogr.* **1994**, *39*. [[CrossRef](#)]
18. Gerhardt, C. A Unified Quantization of Gravity and Other Fundamental Forces of Nature Implies a Big Bang on the Quantum Level. 2023. Available online: https://www.researchgate.net/publication/367523922_A_UNIFIED_QUANTIZATION_OF_GRAVITY_AND_OTHER_FUNDAMENTAL_FORCES_OF_NATURE_IMPLIES_A_BIG_BANG_ON_THE_QUANTUM_LEVEL?channel=doi&linkId=63d66bb064fc860638f89b35&showFulltext=true (accessed on 15 August 2023).

19. Courant, R.; Hilbert, D. *Methoden der mathematischen Physik. I*; Dritte Auflage, Heidelberger Taschenbücher, Band 30; Springer-Verlag: Berlin, German, 1968. [[CrossRef](#)]
20. Gerhardt, C. *Partial Differential Equations II*, Lecture Notes, University of Heidelberg, 2013. Available online: <http://www.math.uni-heidelberg.de/studinfo/gerhardt/PDE2.pdf> (accessed on 15 August 2023).
21. Gerhardt, C. The quantization of a black hole. *arXiv* **2016**, arXiv:1608.08209.
22. Gerhardt, C. The quantization of a Kerr-AdS black hole. *Adv. Math. Phys.* **2018**, *2018*, 4328312. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.