



The Neutron Skin of ⁴⁸Ca and ²⁰⁸Pb: A Critical Analysis

Francesca Sammarruca 匝

Physics Department, University of Idaho, Moscow, ID 83844-0903, USA; fsammarr@uidaho.edu

Abstract: We discuss the neutron skins of ⁴⁸Ca and ²⁰⁸Pb. We review and critically examine modern predictions and empirical constraints, with special attention to the different interpretations of the findings from the PREX-II experiment and the recently reported value of the neutron skin in ⁴⁸Ca extracted from the CREX experiment. We argue that, in the spirit of the ab initio philosophy, the path to understanding the behavior of dense neutron-rich matter must not circumvent fundamental nuclear forces. Based only on that argument, a thick neutron skin in ²⁰⁸Pb is highly unlikely.

Keywords: neutron skins; neutron matter; neutron density; equation of state; symmetry energy; chiral effective field theory

1. Introduction

The existence of the neutron skin is a remarkable feature of neutron-rich nuclei. Because neutrons do not bind, neutron excess, typically measured by the isospin asymmetry $\alpha = (N - Z)/A$, is a destabilizing effect in a nucleus. As a consequence, some of the excess neutrons are "pushed out" from the (neutron-enriched) core and form the skin. The same physics plays an important role in neutron stars, which are supported against gravitational collapse by the outward pressure existing in dense systems with high neutron concentration. Studies of nuclear interactions in systems with high or extreme neutron to proton ratio are crucial for understanding the neutron driplines, the location of which is not well known. The new Facility for Rare Isotope Beams (FRIB), operational since May 2022, is expected to increase the number of known rare isotopes from 3000 to about 6000 [1].

The role of microscopic nuclear physics is to predict observables based on fundamental nuclear forces derived from first principles. Comparison with measurements is, then, a true test of the predictive power of the theory. The neutron skin is not a "conventional" observable, in that it is not measured directly. Instead, it is extracted from measurements of observables that are sensitive to the neutron density distribution in nuclei. Naturally, the neutron density cannot be probed with electron scattering in the same way as for proton densities, and so different methods must be employed, such as experiments with hadronic probes or measurements that exploit the neutron's *weak charge*. Those will be reviewed in Section 2.

We recall that the neutron skin is defined as

$$S = < r^2 >_n^{1/2} - < r^2 >_p^{1/2}, \tag{1}$$

that is, the difference between the root mean square radii of the neutron and the proton distributions.

The common goal of experimentalists and theorists is to shed light on fundamental questions. In the case of neutron skins, the physics one wants to pin down concern the aforementioned pressure that determines the spatial extension of the neutron skin, information contained in the equation of state (EoS) of neutron-rich matter. The importance of this quantity cannot be overstated, given that its relevance extends from nuclei to compact astrophysical systems. These connections will be elucidated in Section 3.



Citation: Sammarruca, F. The Neutron Skin of ⁴⁸Ca and ²⁰⁸Pb: A Critical Analysis. *Symmetry* **2024**, *16*, 34. https://doi.org/10.3390/ sym16010034

Academic Editors: Martin V. Ivanov and Mitko K. Gaidarov

Received: 3 November 2023 Revised: 12 December 2023 Accepted: 20 December 2023 Published: 27 December 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). The purpose of this article is to review and critically examine modern predictions and empirical constraints as well as the different points of view currently being debated. A special focus is placed on the recently reported value of the neutron skin in ⁴⁸Ca extracted from the CREX experiment [2], relative to the findings from the PREX-II [3] experiment. We end with thoughts and suggestions on the best way forward to strengthen the link between experiment and *ab initio* theory.

2. Extracting Neutron Skins from Experiments

2.1. Brief Review of Useful Facts

We will go through the main steps leading to the energy per nucleon in neutron-rich matter and related expressions, because they will be useful for the present discussion. We introduce the energy per nucleon, $e(\rho, \alpha)$, in an infinite system of nucleons at density ρ and isospin asymmetry $\alpha = \frac{\rho_n - \rho_p}{\rho}$ —namely, the EoS of neutron-rich matter—and expand this quantity with respect to the isospin asymmetry parameter, α :

$$e(\rho,\alpha) = e(\rho,0) + \frac{1}{2} \left(\frac{\partial^2 e(\rho,\alpha)}{\partial \alpha^2}\right)_{(\alpha=0)} \alpha^2 + \mathcal{O}(\alpha^4) .$$
⁽²⁾

Neglecting terms of order $\mathcal{O}(\alpha^4)$, Equation (2) takes the well-known form:

$$e(\rho, \alpha) \approx e_0(\rho) + e_{sym}(\rho) \alpha^2 .$$
(3)

Note that $e_{sym}(\rho) = \frac{1}{2} \left(\frac{\partial^2 e(\rho, \alpha)}{\partial \alpha^2} \right)_{\alpha=0}$ and $e_0(\rho) = e(\rho, 0)$, the EoS of isospin-symmetric nuclear matter. Within the quadratic approximation applied in Equation (3), the symmetry energy becomes the difference between the energy per neutron in neutron matter (NM) and the energy per nucleon in symmetric nuclear matter (SNM):

$$e_{sym}(\rho) = e_n(\rho) - e_0(\rho) , \qquad (4)$$

where $e_n(\rho) = e(\rho, 1)$, the energy per neutron in pure NM.

We recall that $e_0(\rho)$ exhibits a minimum at a density approximately equal to the average central density of nuclei, ρ_0 , a reflection of the saturating nature of the nuclear force. Next, we expand the symmetry energy about the saturation point:

$$e_{sym}(\rho) \approx e_{sym}(\rho_0) + L \, \frac{\rho - \rho_0}{3\rho_0} + \frac{K}{2} \frac{(\rho - \rho_0)^2}{(3\rho_0)^2} \,,$$
 (5)

where the expansion parameters are obviously related to the first and higher-order derivatives of $e_{sym}(\rho)$. *L* is a measure of the slope of the symmetry energy at saturation:

$$L = 3\rho_0 \left(\frac{\partial e_{sym}(\rho)}{\partial \rho}\right)_{\rho_0}.$$
 (6)

Furthermore, from Equations (4) and (6), we see that L is a measure of the slope of the NM EoS at saturation density, since the SNM EoS has a vanishing slope at that point.

Using the relation between pressure and energy density, we define the symmetry pressure:

$$P_{sym}(\rho) = \rho^2 \frac{\partial e_{sym}}{\partial \rho} = P_{NM}(\rho) - P_{SNM}(\rho) .$$
⁽⁷⁾

If the derivative is evaluated at or very near ρ_0 , the symmetry pressure is essentially the pressure in NM because the pressure in SNM vanishes at saturation. Then:

$$P_{NM}(\rho_0) = \left(\rho^2 \frac{\partial e_n(\rho)}{\partial \rho}\right)_{\rho_0}.$$
(8)

From Equations (6) and (8), it is clear that the slope parameter L is a measure of the pressure in NM around saturation density:

$$P_{NM}(\rho_0) = \rho_0 \frac{L}{3} . (9)$$

It is then easy to understand how the symmetry energy slope, essentially a pressure gradient acting on excess neutrons, determines the formation and size of the neutron skin. Therefore, constraints on *L* have the ability to provide constraints on the skin, and vice versa.

2.2. Experiments and Phenomenological Analyses

Indirect measurements of the neutron skin in ²⁰⁸Pb and ⁴⁸Ca have been performed using a variety of techniques, such as those listed below. Parity-violating electron scattering will be addressed separately. Some representative measurements are:

- Proton-nucleus elastic scattering [4–7];
- Polarized proton-nucleus elastic scattering [8,9];
- Pionic probes [10,11];
- Coherent pion photoproduction [12,13];
- Antiprotonic atom data [14–16];
- Electric dipole polarizability [17–20];
- Pygmy dipole resonances [21];
- Interaction cross sections with microscopic optical potentials [22];
- (α, α') giant dipole resonance (GDR) [23];
- *α*-particle scattering [24].

Constraints directly on the symmetry energy and its density dependence have also been sought through a variety of techniques, such as: data on nuclear masses across the periodic table [25], giant dipole resonance energies [26], electric dipole polarizability [27], measurements of directed and elliptic flows in heavy ion (HI) collisions [28], isobaric analog states [29], isospin diffusion in HI collisions [30], neutron and proton transverse emission ratio measurements [31], and HI collisions at intermediate energies [32].

In Table 1, we summarize values for ²⁰⁸Pb and ⁴⁸Ca neutron skins deduced from the indicated experiments. There is a considerable spread, as the result of a multitude of methods and theoretical input over decades. Analyses of hadronic scattering experiments, in particular, require modeling of the nuclear potential. Based on previous measurements of the skin in ⁴⁸Ca, we see no strong reasons to deem the CREX result surprising or unexpected, whereas the opposite is true for ²⁰⁸Pb. These observations are well captured in Figure 1, the content of which we have extracted from Figure 8 of ref. [33]. On the Calcium side, from left to right, the data points are from refs. [4,7–11,24]. On the Lead side, in the same order, the first two points correspond to refs. [10,12]. The third point is at 0.18 ± 0.06 fm, as in ref. [33], whereas, from the same references, [14,15], we read the values reported in Table 1. The remaining points are from refs. [4–6,8,10]. A comment is in place with regard to the values from ref. [4], 3rd and 6th data points, at 0.098 ± 0.043 fm and 0.119 ± 0.045 fm, for ⁴⁸Ca and ²⁰⁸Pb, respectively. From ref. [4], we read the smaller values shown in Table 1.

For ⁴⁸Ca (²⁰⁸Pb), the last point on the right is the result of CREX [2] (PREX-II [3]).

We end this section with Figure 2, showing our predictions of the neutron skin for selected nuclei as a function of the isospin asymmetry [34], confirming a nearly linear relation between these two quantities and, thus, motivating the expectation that the skins of ⁴⁸Ca and ²⁰⁸Pb should be close. The boundaries of the gray area are linear fits to the data from Ref. [35].

Table 1. Values of the neutron skins in ⁴⁸Ca and in ²⁰⁸Pb from a variety of experimental methods. It is our understanding that the quoted errors refer to one standard deviation. For the values taken from ref. [12] and ref. [15], the first and second uncertainties are statistical and systematic errors, respectively. In ref. [14], the first uncertainty is the experimental error, while the second originates from theoretical modeling of the experimental charge densities. Finally, in the result from ref. [18], the first two uncertainties are the experimental and the theoretical error, respectively, and the third one stands for an estimated uncertainty in the symmetry energy at saturation.

Type of Measurement	Extracted Neutron Skin in ⁴⁸ Ca	Extracted Neutron Skin in ²⁰⁸ Pb
Proton-nucleus scattering [4]	0.056 - 0.102	0.083 - 0.111
Proton-nucleus scattering [5]		$0.211 \begin{array}{c} +0.054 \\ -0.063 \end{array}$
Proton-nucleus scattering [6]		0.20 ± 0.04
Proton-nucleus scattering [7]	0.10 ± 0.03	
Polarized proton-nucleus scattering [8]	0.23 ± 0.05	0.16 ± 0.05
Polarized proton-nucleus scattering [9]	$0.168 \ ^{+0.025}_{-0.028}$	
Polarized proton-nucleus scattering [5]		$0.211 \substack{+0.054 \\ -0.063}$
Pionic probes [10]	0.13 ± 0.06	0.11 ± 0.06
Pionic probes [11]	0.11 ± 0.04	
Coherent π photoproduction [12]		$0.15\pm 0.03^{+0.01}_{-0.03}$
Coherent π photoproduction [13]		$0.20 \substack{+0.01 \\ -0.03}$
Antiprotonic atoms [14]		$0.20 (\pm 0.04) (\pm 0.05)$
Antiprotonic atoms [15]		$0.16 (\pm 0.02) (\pm 0.04)$
Antiprotonic atoms [16]		0.15 ± 0.02
Electric dipole polarizability [17]		0.13 - 0.19
Electric dipole polarizability [18]		$0.165 (\pm 0.09) (\pm 0.013) (\pm 0.021)$
Electric dipole polarizability		
<i>via</i> polarized scattering at forward angle [19]		$0.156 \ {}^{+0.025}_{-0.021}$
Electric dipole polarizability [20]	0.14 - 0.20	
Pygmy dipole resonances [21]		0.18 ± 0.035
Interaction cross sections [22]	0.105 ± 0.06	
(<i>α</i> , <i>α'</i>) GDR 120 MeV [23]		0.19 ± 0.09
α -particle scattering [24]	0.171 ± 0.05	



Figure 1. (Color online) An overview of experimental constraints, taken from ref. [33] (see text for explanation). On the ⁴⁸Ca (²⁰⁸Pb) side, the last point on right is the result of CREX [2] (PREX-II [3]). The horizontal lines mark the weighted means of the experiments \pm one standard deviation, not including parity-violating electron scattering.



Figure 2. (Color online) Red bars: Neutron skin of ⁵⁸Ni, ²⁷Al, ⁵⁹Co, ⁹⁰Zr, ⁴⁸Ca, and ²⁰⁸Pb, in order of increasing isospin asymmetry. The figure is taken from ref. [34]. The shaded area is bounded by linear fits to the data [35].

2.3. Parity-Violating Electron Scattering

The parity-violating electron scattering asymmetry, A_{PV} , is defined for a spin-zero nucleus as

$$A_{PV} = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L} , \qquad (10)$$

where $\sigma_{R(L)}$ is the elastic cross section for right (left) handed electrons [36]. A_{PV} is proportional to the ratio of weak ($F_W(q)$) to charge ($F_{ch}(q)$) form factors, whith q the fourmomentum transfer. $F_{ch}(q)$ is taken from existing measurements and $F_W(q)$ is extracted from the measured A_{PV} . We recall that the weak and charge form factors are the Fourier transforms of the weak charge density and the charge density, respectively:

$$F_W(q) = \frac{1}{Q_W} \int d^3r j_0(qr) \rho_W(r) , \qquad (11)$$

where Q_W is the weak charge of the nucleus, and

$$F_{ch}(q) = \frac{1}{Z} \int d^3r j_0(qr) \rho_{ch}(r) .$$
 (12)

In Equations (11) and (12), $j_0(qr)$ is the zeroth-order spherical Bessel function. In Equation (11), a form is assumed for $\rho_W(r)$ and the radius parameter of the density function is adjusted to reproduce the experimental A_{PV} . A form must also be assumed for ρ_{ch} in Equation (12), but its parametrization relies upon well-established measurements from charged electron scattering. The CREX result is found insensitive to the assumed form for the weak charge density [2].

3. The Status of ab initio Theory

3.1. Development of Microscopic Nuclear Forces

Our still-incomplete knowledge of nuclear forces is the result of decades of struggle. Currently, the optimal approach to the construction of nuclear forces is based on the understanding that the energy scale determines the appropriate degrees of freedom of the theory—the central concept to the development of chiral effective field theory (EFT) [37,38]. Here, we provide only a brief summary of the strongest features of chiral EFT.

Chiral EFT allows the development of nuclear interactions as an expansion where theoretical uncertainties can be assessed at each order. The organizational scheme that controls the expansion is known as "power counting". The crucial point is that chiral EFT maintains consistency with the underlying fundamental theory of strong interactions, quantum chromodynamics (QCD), through the symmetries and symmetry breaking mechanisms of the low-energy QCD Lagrangian. The first step towards the development of an EFT is the identification of a "soft scale" and a "hard scale", which is suggested by the hadron spectrum, observing the large separation between the mass of the pion and the mass of the vector meson ρ . It is therefore natural to identify the pion mass and the ρ mass (approximately 1 GeV) with the soft and the hard scale, respectively. Moreover, since quarks and gluons are ineffective degrees of freedom in the low-energy regime, pions and nucleons can be taken as the appropriate degrees of freedom of the EFT. Having identified pions and nucleons as the appropriate degrees of freedom of the EFT, one can proceed to construct the Lagrangian of the effective theory:

$$\mathcal{L}_{eff} = \mathcal{L}_{\pi\pi} + \mathcal{L}_{\pi N} + \mathcal{L}_{NN} + \dots, \qquad (13)$$

which is then expanded in terms of a natural parameter, identified with the ratio of the "soft scale" over the "hard scale", $\frac{Q}{\Lambda_{\chi}}$. *Q* is of the order of the pion mass, whereas Λ_{χ} is the energy scale of chiral symmetry breaking, approximately 1 GeV. The contributions to the effective Lagrangian are arranged according to the power-counting scheme, with increasing order resulting in smaller terms. While the expansion itself is, of course, infinite, at each order we are assured that the number of terms is finite and the contributions well defined. The combination of meson–theoretic NN potentials augmented with selected three-nucleon forces (3NF) is an outdated paradigm, from which it is essentially impossible to estimate the theoretical uncertainty of a prediction.

With chiral EFT, there has been enormous progress in the understanding and development of few-nucleon forces. We are on the right path, but there is still much to accomplish. Convergence at next-to-next-to-next-to leading order (N^3LO) needs to be on more robust grounds. Furthermore, there are indications that specific components of the 3NF at next-tonext-to-next-to-leading order (N^4LO) have the potential to solve some outstanding problems in microscopic nuclear structure [39].

3.2. Neutron Skin Predictions

We show in Table 2 recent predictions for the skins in ⁴⁸Ca and in ²⁰⁸Pb. Some comments are in place. In both refs. [40,41], the nature of the nucleon–nucleon (NN) chiral potentials, N²LO_{sat} (nucleon–nucleon potential at next-to-next-to leading order fitted also to the saturation properties of nuclear matter), and $\Delta N^2 LO_{GO}$ (nucleon-nucleon potential at next-to-next-to leading order including the Δ -isobar developed by groups in Gothenburg and Oak Ridge), is such that the results are not truly ab initio. Note, also, that the value for ²⁰⁸Pb from ref. [41] is not a prediction, but was obtained using the linear regression the authors constructed from the skins of lighter nuclei.

Nucleus	Predicted Skin	Source	
⁴⁸ Ca	0.120-0.150	Ref. [40]	
⁴⁸ Ca	0.141-0.187	Ref. [42] 68% CR	
⁴⁸ Ca	0.123-0.199	Ref. [42] 90% CR	
²⁰⁸ Pb	0.139-0.200	Ref. [42] 68% CR	
²⁰⁸ Pb	0.120-0.221	Ref. [42] 90% CR	
⁴⁸ Ca	0.114-0.186	Ref. [41]	
²⁰⁸ Pb	0.184-0.236	Ref. [41]	
⁴⁸ Ca	0.12-0.15	Ref. [34]	
²⁰⁸ Pb	0.13-0.17	Ref. [34]	
⁴⁸ Ca	0.176 ± 0.018	Ref. [43]	
²⁰⁸ Pb	0.168 ± 0.022	Ref. [43]	

Table 2. Status of ab initio predictions for the skins (in fm) in 48 Ca and in 208 Pb. The two results from ref. [42] for both 48 Ca and in 208 Pb show acceptable ranges within 68% and 90% of the credibility region (CR). The last two entries were obtained from density functional theory [43].

Figure 3 shows the neutron skin in ⁴⁸Ca vs. the one in ²⁰⁸Pb. All values, except for the green bars, have been extracted from Figure 5 of ref. [2]. The PREX-II and PREX-I combined experimental result is shown by the blue bar, while the red vertical bar is the CREX result.

The gray circles (pink diamonds) show results from a variety of relativistic (non-relativistic) density functionals, which give values of the *L* parameter ranging from small and negative to large and positive. Coupled cluster (CC) [40,41] and dispersive optical model (DOM) predictions [44] are also displayed. Our predictions are shown by the green bars [34]. There exist, of course, relativistic mean-field (RMF) models that agree with the PREX result, and, correspondingly, generate values for ⁴⁸Ca that are also on the larger side.





We conclude that a value between 0.212 fm and 0.354 fm (0.283 \pm 0.071) for the skin of ²⁰⁸Pb is outside the boundaries set by microscopic theory. Simultaneous consistency with both CREX and PREX seems to be a challenge even for phenomenology. On the other hand, PREX aside, a small skin for ⁴⁸Ca does not appear to be peculiar based on the facts we reviewed above, see also Figure 1. At the same time, it is important to keep in mind that an experimental measurement has a probability of approximately 68% (95%) to be within \pm one standard deviation (\pm two standard deviations). That is, it is possible that the PREX value is the result of a large statistical fluctuation.

4. Further Discussion

Irrespective of the inconsistency between PREX and CREX, there are far-reaching questions one must consider with regard to a thick skin in ²⁰⁸Pb and its ramifications.

4.1. Direct Urca Processes

We define the total energy per baryon in β -equilibrated matter (in absence of muons) as

$$e_T(\rho, Y_p) = e_0(\rho) + e_{sym} (1 - 2Y_p)^2 + e_e + \sum_{i=n,p} Y_i \cdot m_i .$$
(14)

The last term accounts for the baryon rest masses (in units of energy), while e_e is the electron energy. The particle fractions are

$$Y_i = \frac{\rho_i}{\rho} , \qquad (15)$$

and the chemical potentials are given by:

$$\mu_i = \frac{\partial \epsilon_i}{\partial \rho_i} = \frac{\partial e_i}{\partial Y_i} , \qquad (16)$$

where i = n, p, e. Naturally, we impose the constraints of fixed baryon density, Equation (17), and global charge neutrality, Equation (18):

$$\rho_p + \rho_n = \rho \quad \Rightarrow \quad Y_p + Y_n = 1,$$
(17)

$$\rho_p = \rho_e \quad \Rightarrow \quad Y_p = Y_e \,. \tag{18}$$

The energy of ultrarelativistic electrons is easily written as

$$e_e = \frac{\hbar c}{4\pi^2} \,\rho^{\frac{1}{3}} (3\pi^2 Y_p)^{\frac{4}{3}} \,, \tag{19}$$

and thus

$$\frac{\partial e_e}{\partial Y_p} = \frac{\hbar c}{3\pi^2} (3\pi^2)^{\frac{4}{3}} (\rho Y_p)^{\frac{1}{3}} .$$
(20)

Noting that the β -equilibrium condition is

$$\frac{\partial e_p}{\partial Y_p} = \frac{\partial e_n}{\partial Y_n} - \frac{\partial e_e}{\partial Y_e} , \qquad (21)$$

and using Equations (14) and (21), we obtain:

$$4 e_{sym}(\rho) (1 - 2 \cdot Y_p) = \hbar c (3\pi^2)^{\frac{1}{3}} (\rho Y_p)^{\frac{1}{3}} , \qquad (22)$$

where we have neglected the mass difference between the proton and the neutron.

If ρ_{DU} is the density at which Y_p is equal to the value needed for direct Urca processes (DU), about 1/9, the following relation holds:

$$e_{sym}(\rho_{DU}) = \hbar c \frac{9}{28} (\pi^2/3)^{\frac{1}{3}} (\rho_{DU})^{1/3} .$$
⁽²³⁾

This simple relation can be quite insightful. If, for example, ρ_{DU} is close to saturation density, the symmetry energy would be over 50 MeV at that point. If $\rho_{DU} = \frac{2}{3}\rho_0$, the symmetry energy at that density would be over 40 MeV. In microscopic predictions, DU processes are more likely to open at a few to several times normal density (based on projections, since chiral EFT predictions cannot be extended to such high densities). Using PREX II constraints, the Urca threshold is found to be approximately $1.5\rho_0$ or just above 0.2 fm^{-3} [45]. From Equation (23), the value of the symmetry energy at $1.5\rho_0$ is then about 58 MeV, clearly indicating an unusually steep density dependence, taking the PREX II value of 38.1 ± 4.7 MeV at saturation.

In ref. [45], the authors develop different parametrizations of RMF models constrained by CREX results, PREX II results, or a combination of both, and report that the direct Urca threshold density decreases from 0.71 fm⁻³ to 0.21 fm⁻³ as the skin of ²⁰⁸Pb increases from 0.13 fm to 0.28 fm, with expected implications for neutron star cooling processes. To the best of our knowledge, there is no evidence for rapid neutrino cooling *via* direct Urca for low-mass (low central density) neutron stars. In fact, from analyses including luminosities and ages determined from observations of isolated neutron stars [46], one may conclude that the direct Urca process, although possible at the central densities of neutron stars with masses between 1.7 and 2.0 solar masses, is unlikely around $M = 1.7M_{\odot}$, and is likely, but not the principal cooling mechanism, for the $M = 2.0M_{\odot}$ stars.

Inclusion of heavy baryons (such as hyperons and Δ 's) in stellar matter would of course change the above discussion. Applications of modern approaches to hypernuclear stars and their cooling processes are at an early stage. See ref. [47] for a recent review of this active and promising field.

4.2. Impact of the Isovector Component of the Free-Space Nuclear Force

The importance of a realistic isovector component of the nuclear force (that is, carefully calibrated through free-space NN data), on the density dependence of the symmetry energy has been demonstrated [42,48]. Relaxing the constraint of accurate phase shifts for the isospin-1 *S* and *P* waves leads to drastic variations of the pressure in NM [48] and, consequently, the neutron skin.

For the purpose of the present discussion, it is useful to recall that, typically, the isovector sector of RMF functionals includes the coupling of the nucleon to the isovector ρ -meson and a nonlinear $\omega - \rho$ isoscalar–isovector cross-coupling term. In the models of Ref. [49], designed to reconcile the PREX and CREX results, the novel feature is the inclusion of the scalar isovector δ -meson (better known as the a_0 -meson). The isospin splitting resulting from the isovector nature of the new contribution opens an additional fitting degree of freedom [49].

As mesons heavier than the pion do not enter the development of the NN force in chiral EFT, we will discuss this important point from the perspective of quantitative one-boson-exchange potentials (OBEP), as we already did over 10 years ago [50]. Studies of the ρ and δ contributions to the potential part of the symmetry energy in quantum hydrodynamics (QHD) models can be found in refs. [30-35] of ref. [49], where those contributions are shown to be opposite in sign and very large in magnitude around normal density. Thus, in QHD-inspired models, the interplay between ρ and δ is treated as the equivalent, in the isovector channel, of the interplay between σ and ω in the isoscalar channel. In contrast, the role of the δ -meson in meson theory is subtle (although important) and it is seen in the difference of its contributions to isospin-1 or isospin-0 partial waves, particularly 1S_0 and 3S_1 . The dramatic difference between the description of the isovector channel in QHD-based models or realistic meson models originate from several sources, including the absence of the pion in QHD, and the fact that meson contributions in a microscopic approach are iterated, and thus reduced by Pauli blocking.

In ref. [50], we have shown the difference between the potential energy contributions to NM and SNM from the isovector mesons, to estimate the effect of each meson on the potential energy part of the symmetry energy. The impact of the pion on the symmetry energy is by far the largest. On the other hand, mean-field theories are generally pionless, because the bulk of the attraction–repulsion balance needed for a realistic description of nuclear matter can be technically obtained from σ and ω only, an observation that is at the very foundation of the Walecka model [51]. On the other hand, in any theory of nuclear forces, the pion is the most important ingredient. Chiral symmetry is spontaneously broken in low-energy QCD and the pion emerges as the Goldstone boson of this symmetry breaking. Moreover, NN scattering data, and most definitely the deuteron, cannot be described without the pion.

In conclusion, developing additional density functionals with unrealistic isovector contributions for the purpose of reconciling the parity-violating experiments does not provide more clarity. The road to understanding the behavior of dense neutron-rich matter must go through ab initio nuclear forces.

5. Conclusions

The above has been a critical analysis of the available facts on the neutron skins in ²⁰⁸Pb and ⁴⁸Ca, from both the experimental and the theoretical side.

We have pointed out and discussed problematic aspects that arise in conjunction with a stiff EoS and the corresponding thick skin in ²⁰⁸Pb. We emphasized that such thick skin is outside the boundaries of ab initio nuclear theory.

On the other hand, in the context of existing measurements and microscopic predictions, the CREX result for the skin of ⁴⁸Ca does not appear to be an outlier.

We wait with excitement for the MREX experiment at the MESA accelerator, which promises to measure the neutron skin of ²⁰⁸Pb with *ultimate precision* [52].

Laboratory neutron skin measurements have important implications for neutron star properties, such as radius, tidal deformability, and cooling processes, and are complementary to astronomical observations. In the "multi-messanger" era, it is especially important to keep a broad view and, at the same time, stay in close touch with first principles, as we study the remarkable connection between microscopic physics and "telescope physics" enabled by the equation of state. **Funding:** This work was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Award Number DE-FG02-03ER41270.

Data Availability Statement: All results presented in this paper are publicly available through the cited references.

Conflicts of Interest: The author declares no conflict of interest.

References

- 1. Smits, O.R.; Düllmann, C.E.; Indelicato, P.; Nazarewicz, W.; Schwerdtfeger, P. The quest for superheavy elements and the limit of the periodic table. *Nat. Rev. Phys.* 2023, 1–13. https://doi.org/10.1038/s42254-023-00668-y
- Adhikari, D.; Albataineh, H.; Androic, D.; Aniol, K.A.; Armstrong, D.S.; Averett, T.; Gayoso, C.A.; Barcus, S.K.; Bellini, V.; Beminiwattha, R.S.; et al. (CREX Collaboration). Precision Determination of the Neutral Weak Form Factor of ⁴⁸Ca. *Phys. Rev. Lett.* 2022, 129, 042501.
- 3. Reed, B.T.; Fattoyev, F.J.; Horowitz, C.J.; Piekarewicz, J. Implications of PREX-II on the equation of state of neutron-rich matter. *Phys. Rev. Lett.* **2021**, *126*, 172503.
- 4. Clark, B.C.; Kerr, L.J.; Hama, S. Neutron densities from a global analysis of medium-energy proton-nucleus elastic scattering. *Phys. Rev. C* 2003, *67*, 054605.
- Zenhiro, J.; Sakaguchi, H.; Murakami, T.; Yosoi, M.; Yasuda, Y.; Terashima, S.; Iwao, Y.; Takeda, H.; Itoh, M.; Yoshida, H.P.; et al. Neutron density distributions of ^{204,206,208}Pb deduced via proton elastic scattering at Ep = 295 MeV. *Phys. Rev. C* 2010, *82*, 044611.
- 6. Starodubnsky, V.E.; Hintz, N.M. Extraction of neutron densities from elastic proton scattering by ^{206,207,208}Pb at 650 MeV. *Phys. Rev. C* **1994**, *49*, 2118.
- 7. Shlomo, S.; Schaeffer, R. The difference between neutron and proton radii in the Ca isotopes. *Phys. Lett. B* 1979, 83, 5.
- 8. Ray, L. Neutron isotopic density differences deduced from 0.8 GeV polarized proton elastic scattering. Phys. Rev. C 1979, 19, 1855.
- 9. Zenhiro, J.; Sakaguchi, H.; Terashima, S.; Uesaka, T.; Hagen, G.; Itoh, M.; Murakami, T.; Nakatsugawa, Y.; Ohnishi, T.; Sagawa, H.; et al. Direct determination of the neutron skin thicknesses in ^{40,48}Ca from proton elastic scattering at Ep = 295 MeV. *arXiv* 2018, arXiv:1810.11796.
- 10. Friedman, E. Neutron skins of ²⁰⁸Pb and ⁴⁸Ca from pionic probes. *Nucl. Phys. A* 2012, 896, 46.
- 11. Gibbs, W.R.; Dedonder, J.-P. Neutron radii of the calcium isotopes. *Phys. Rev. C* 1992, 46, 1825.
- 12. Tarbert, C.M.; Watts, D.P.; Glazier, D.I.; Aguar, P.; Ahrens, J.; Annand, J.R.M.; Arends, H.J.; Beck, R.; Bekrenev, V.; Boillat, B.; et al. Neutron Skin of ²⁰⁸Pb from Coherent Pion Photoproduction. *Phys. Rev. Lett.* **2014**, *112*, 242502.
- 13. Zana, L.; for the A2 Collaboration at MAMI. Neutron skins from Coherent Pion Photoproduction. Epj Web Conf. 2015, 96, 01036.
- 14. Brown, B.A.; Shen, G.; Hillhouse, G.C.; Meng, J.; Trzcinska, A. Neutron skin deduced from antiprotonic atom data. *Phys. Rev. C* **2007**, *76*, 034305.
- 15. Klos, B.; Trzcinska, A; Jastrzebski, J.; Czosnyka, T.; Kisielinski, M.; Lubinski, P.; Napiorkowski, P.; Pienkowski, L.; Hartmann, F.J.; Ketzer, B.; et al. Neutron density distributions from antiprotonic ²⁰⁸Pb and ²⁰⁹Bi atoms. *Phys. Rev. C* **2007**, *76*, 014311.
- 16. Trzcinska, A.; Jastrzebski, J.; Lubinski, P.; Hartmann, F.J.; Schmidt, R.; von Egidy, T.; Kłos, B. Neutron Density Distributions Deduced from Antiprotonic Atoms. *Phys. Rev. Lett.* **2001**, *87*, 082501.
- Roca-Maza, X.; Viñas, X.; Centelles, M.; Agrawal, B.K.; Colò, G.; Paar, N.; Piekarewicz, J.; Vretenar, D. Neutron skin thickness from the measured electric dipole polarizability in ⁶⁸Ni, ¹²⁰Sn, and ²⁰⁸Pb. *Phys. Rev. C* 2015, 92, 064304.
- Roca-Maza, X.; Brenna, M.; Colò, G.; Centelles, M.; Viñas, X.; Agrawal, B.K.; Paa, N.; Vretenar, D.; Piekarewicz, J. Electric dipole polarizability in ²⁰⁸Pb: Insights from the droplet model. *Phys. Rev. C* 2013, *88*, 024316.
- 19. Tamii, A.; Complete Electric Dipole Response and the Neutron Skin in ²⁰⁸Pb. *Phys. Rev. Lett.* **2011**, 107, 062502.
- Birkham, J.; Miorelli, M.; Bacca, S.; Bassauer, S.; Bertulani, C. A.; Hagen,G.; Matsubara, H.; von Neumann-Cosel, P.; Papenbrock, T.; Pietralla, N. et al. Electric Dipole Polarizability of ⁴⁸Ca and Implications for the Neutron Skin. *Phys. Rev. Lett.* 2017, 118, 252501.
- 21. Klimkiewicz, A.; Paar, N.; Adrich, P.; Fallot, M.; Boretzky, K.; Aumann, T.; Cortina-Gil, D.; Pramanik, U.D.; Elze, T.W.; Emling, H.; et al. Nuclear symmetry energy and neutron skins derived from pygmy dipole resonances. *Phys. Rev. C* 2007, *76*, 051603.
- 22. Matsuzaki, M.; Yahiro, M. Neutron skin of ⁴⁸Ca deduced from interaction cross section. *Mod. Phys. Lett. A* 2022, 37, 2250123.
- 23. Krasznahorkay, A.; Balanda, A.; Bordewijk, J.A.; Brandenburg, S.; Harakeh, M.N.; Kalantar-Nayestanaki, N.; Nyakó, B.M.; Timár, J.; van der Woude, A. Excitation of the isovector GDR by inelastic α scattering as a measure of the neutron skin of nuclei. *Nucl. Phys. A* 1994, 567, 521.
- 24. Gils, H.J.; Rebel, H.; Friedman, E. Isotopic and isotonic differences between a particle optical potentials and nuclear densities of 1*f*_{7/2} nuclei. *Phys. Rev. C* **1984**, *29*, 1295.
- 25. Mondal, C.; Agrawal, B.K.; De, J.N. Constraining the symmetry energy content of nuclear matter from nuclear masses: A covariance analysis. *Phys. Rev. C* 2015, *92*, 024302.
- 26. Trippa, L.; Colò, G.; Vigezzi, E. Giant dipole resonance as a quantitative constraint on the symmetry energy. *Phys. Rev. C* 2008, 77, 061304(R).
- 27. Zhang, Z.; Chen, L.-W. Electric dipole polarizability in ²⁰⁸Pb as a probe of the symmetry energy and neutron matter around $\rho_0/3$. *Phys. Rev. C* **2015**, *92*, 031301.

- Russotto, P.; Gannon, S.; Kupny, S.; Lasko, P.; Acosta, L.; Adamczyk, M.; Al-Ajlan, A.; Al-Garawi, M.; Al-Homaidhi, S.; Amorini, F.; et al. Results of the ASY-EOS experiment at GSI: The symmetry energy at suprasaturation density. *Phys. Rev. C* 2016, 94, 034608.
- 29. Danielewicz, P.; Lee, J. Symmetry energy II: Isobaric analog states. Nucl. Phys. A 2014, 922, 1.
- 30. Tsang, M.B.; Liu, T.X.; Shi, L.; Danielewicz, P.; Gelbke, C.K.; Liu, X.D.; Lynch, W.G.; Tan, W.P.; Verde, G.; Wagner, A.; et al. Isospin Diffusion and the Nuclear Symmetry Energy in Heavy Ion Reactions. *Phys. Rev. Lett.* **2004**, *92*, 062701.
- Famiano, M.A.; Liu, T.; Lynch, W.G.; Mocko, M.; Rogers, A.M.; Tsang, M.B.; Wallace, M.S.; Charity, R.J.; Komarov, S.; Sarantites, D.G.; et al. Neutron and proton transverse emission ratio measurements and the density dependence of the asymmetry term of the nuclear equation of state. *Phys. Rev. Lett.* 2006, 97, 052701.
- 32. Yong, G.-C.; Guo, Y.-F. Probing High-density Symmetry Energy Using Heavy-ion Collisions at Intermediate Energies. *Nucl. Phys. Rev.* **2020**, *37*, 136.
- 33. Lattimer, J.M. Constraints on Nuclear Symmetry Energy Parameters. arXiv 2023, arXiv:2301.03666.
- 34. Sammarruca, F. Neutron skin systematics from microscopic equations of state. Phys. Rev. C 2022, 105, 064303.
- 35. Swiatecki, W.J.; Trzcinska, A.; Jastrzebski, J. Difference of the root-mean-square sizes of neutron and proton distributions in nuclei: Comparison of theory with data. *Phys. Rev. C* 2005, *71*, 047301.
- 36. Donnelly, T.W.; Dubach, J.; Sick, I. Isospin dependences in parity-violating electron scattering. Nucl. Phys. A 1989, 503, 589.
- 37. Weinberg, S. Nuclear forces from chiral lagrangians. *Phys. Lett. B* 1990, 251, 288.
- 38. Weinberg, S. Three-body interactions among nucleons and pions. Phys. Lett. B 1992, 295, 114.
- 39. Machleidt, R. What is ab initio? *arXiv* **2023**, arXiv:2307.06416.
- Hagen, G.; Ekström, A.; Forssén, C.; Jansen, G.R.; Nazarewicz, W.; Papenbrock, T.; Wendt, K.A.; Bacca, S.; Barnea, N.; Carlsson, B.; et al. Neutron and weak-charge distributions of the ⁴⁸Ca nucleus. *Nat. Phys.* **2016**, *12*, 186.
- 41. Novario, S.J.; Lonardoni, D.; Gandolfi, S.; Hagen, G. Trends of Neutron Skins and Radii of Mirror Nuclei from First Principles. *Phys. Rev. Lett.* **2023**, 130, 032501.
- 42. Hu, B.; Jiang, W.; Miyagi, T.; Sun, Z.; Ekström, A.; Forssen, C.; Hagen, G.; Holt, J.D.; Papenbrock, T.; Stroberg, S.R.; Vernon, I. Ab initio predictions link the neutron skin of ²⁰⁸Pb to nuclear forces. *Nat. Phys.* **2022**, *18*, 1196.
- 43. Piekarewicz, J.; Agrawal, B.K.; Colo, G.; Nazarewicz, W.; Paar, N.; Reinhard, P.G.; Roca-Maza, X.; Vretenar, D. Electric dipole polarizability and the neutron skin. *Phys. Rev. C* **2012**, *85*, 041302R.
- Atkinson, M.C.; Mahzoon, M.H.; Keim, M.A.; Bordelon, B.A.; Pruitt, C.D.; Charity, R.J.; Dickhoff, W.H. Dispersive optical model analysis of ²⁰⁸Pb generating a neutron-skin prediction beyond the mean field. *Phys. Rev. C* 2020, 101, 044303.
- 45. Kumar, M.; Kumar, S.; Thakur, V.; Kumar, R.; Agrawal, B.K.; Dhiman, S.K. CREX- and PREX-II-motivated relativistic interactions and their implications for the bulk properties of nuclear matter and neutron stars. *Phys. Rev. C* **2023**, *107*, 055801.
- 46. Beloin, S.; Han, S.; Steiner, A. W.; Odbadrakh, K. Simultaneous fitting of neutron star structure and cooling data. *Phys. Rev. C* **2019**, *100*, 055801.
- 47. Sedrakian, A.; Li, J.J.; Weber, F. Heavy Baryons in Compact Stars. Prog. Part. Nucl. Phys. 2023, 131, 104041.
- Sammarruca, F. The Symmetry Energy: Current Status of *Ab Initio* Predictions vs. Empirical Constraints. *Symmetry* 2023, *15*, 450.
 Reed, B.T.; Fattoyev, F.J.; Horowitz, C.J.; Piekarewicz, J. Density Dependence of the Symmetry Energy in the Post PREX-CREX Era. *arXiv* 2023, arXiv:2305.19376.
- 50. Sammarruca, F. Contribution of isovector mesons to the symmetry energy in a microscopic model. Phys. Rev. C 2011, 84, 044307.
- 51. Serot, B.D.; Walecka, J.D. Recent Progress in Quantum Hadrodynamics. Int. J. Mod. Phys. 1997, E6, 515.
- 52. Becker, D.; Bucoveanu, R.; Grzesik, C.; Imai, K.; Kempf, R.; Molitor, M.; Tyukin, A.; Zimmermann, M.; Armstrong, D.; Aulenbacher, K.; et al. The P2 Experiment—A future high-precision measurement of the electroweak mixing angle at low momentum transfer. *EPJ A* **2018**, *54*, 208.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.