

Article

Grammatikopoulosite, NiVP, a New Phosphide from the Chromitite of the Othrys Ophiolite, Greece

Luca Bindi ¹^(D), Federica Zaccarini ^{2,*}, Elena Ifandi ^{3,4}^(D), Basilios Tsikouras ³^(D), Chris Stanley ⁵^(D), Giorgio Garuti ² and Daniela Mauro ⁶

- ¹ Dipartimento di Scienze della Terra, Università degli Studi di Firenze, I-50121 Florence, Italy; luca.bindi@unifi.it
- ² Department of Applied Geological Sciences and Geophysics, University of Leoben, A-8700 Leoben, Austria; giorgio.garuti1945@gmail.com
- ³ Faculty of Science, Physical and Geological Sciences, Universiti Brunei Darussalam, BE 1410 Gadong, Brunei Darussalam; selena.21@windowslive.com (E.I.); basilios.tsikouras@ubd.edu.bn (B.T.)
- ⁴ Department of Geology, Section of Earth Materials, University of Patras, 265 00 Patras, Greece
- ⁵ Department of Earth Sciences, Natural History Museum, London SW7 5BD, UK; c.stanley@nhm.ac.uk
- ⁶ Dipartimento di Scienze della Terra, Università degli Studi di Pisa, I-56126 Pisa, Italy; daniela.mauro@dst.unipi.it
- * Correspondence: federica.zaccarini@unileoben.ac.at; Tel.: +43-(0)3842-402-6218

Received: 22 December 2019; Accepted: 27 January 2020; Published: 31 January 2020



Abstract: Grammatikopoulosite, NiVP, is a new phosphide discovered in the podiform chromitite and hosted in the mantle sequence of the Othrys ophiolite complex, central Greece. The studied samples were collected from the abandoned chromium mine of Agios Stefanos. Grammatikopoulosite forms small crystals (from 5 µm up to about 80 µm) and occurs as isolated grains. It is associated with nickelphosphide, awaruite, tsikourasite, and an undetermined V-sulphide. It is brittle and has a metallic luster. In plane-polarized light, it is creamy-yellow, weakly bireflectant, with measurable but not discernible pleochroism and slight anisotropy with indeterminate rotation tints. Internal reflections were not observed. Reflectance values of mineral in air (R_1 , R_2 in %) are: 48.8–50.30 at 470 nm, 50.5–53.5 at 546 nm, 51.7–55.2 at 589 nm, and 53.2–57.1 at 650 nm. Five spot analyses of grammatikopoulosite give the average composition: P 19.90, S 0.41, Ni 21.81, V 20.85, Co 16.46, Mo 16.39, Fe 3.83, and Si 0.14, total 99.79 wt %. The empirical formula of grammatikopoulosite—based on $\Sigma(V + Ni + Co + Mo + Fe + Si) = 2$ apfu, and taking into account the structural results—is $(Ni_{0.57}Co_{0.32}Fe_{0.11})_{\Sigma 1.00}(V_{0.63}Mo_{0.26}Co_{0.11})_{\Sigma 1.00}(P_{0.98}S_{0.02})_{\Sigma 1.00}$. The simplified formula is (Ni,Co)(V,Mo)P and the ideal formula is NiVP, which corresponds to Ni 41.74%, V 36.23%, P 22.03%, total 100 wt %. The density, calculated on the basis of the empirical formula and single-crystal data, is 7.085 g/cm³. The mineral is orthorhombic, space group *Pnma*, with a = 5.8893(8), b = 3.5723(4), c = 6.8146(9) Å, V = 143.37(3) Å³, and Z = 4. The mineral and its name have been approved by the Commission of New Minerals, Nomenclature and Classification of the International Mineralogical Association (IMA 2019-090). The mineral honors Tassos Grammatikopoulos, geoscientist at the SGS Canada Inc., for his contribution to the economic mineralogy and mineral deposits of Greece.

Keywords: grammatikopoulosite; phosphide; chromitite; Agios Stefanos mine; Othrys; ophiolite; Greece

1. Introduction

Natural phosphides are very rare phases, representing only 3% of the minerals approved by the International Mineralogical Association (IMA) ([1] and references therein). Most of these natural phosphides have been discovered and described in meteorites. Only recently have several new



phosphides been discovered in terrestrial rocks, most of which are associated with the pyrometamorphic rocks of the Hatrurim Formation, Southern Levant [2]. Phosphides are documented in podiform chromitites hosted in the mantle sequence of ophiolite complexes of Greece and Russia [3–6] but are very rare. The phosphide found in the Russian chromitite was found in situ, in contact with serpentine of the altered interstitial silicates. The phosphides associated with the Greek chromitite were documented in heavy concentrates. Recently, the new mineral tsikourasite (Mo₃Ni₂P_{1 + x} (x < 0.25) was discovered in heavy mineral concentrates from a chromitite from the Othrys ophiolite (Greece)) [1]. Further investigation of the same mineral concentrates led to the discovery of a second new mineral. Quantitative chemical analysis and crystal structure proved that the studied phase is a new phosphide, characterized by the simplified formula (Ni,Co)(V,Mo)P and by the ideal formula NiVP, which corresponds to Ni 41.74%, V 36.23%, P 22.03%, total 100 wt %. The mineral is orthorhombic, space group *Pnma*, with a = 5.8893(8), b = 3.5723(4), c = 6.8146(9) Å, and Z = 4. The mineral and its name have been approved by the Commission of New Minerals, Nomenclature and Classification of the International Mineralogical Association (IMA 2019-090). The mineral honors Tassos Grammatikopoulos (b. 1966), a geoscientist at the SGS Canada Inc., for his contribution to the investigation of economic mineralogy and mineral deposits of Greece. Holotype material is deposited in the Mineralogical Collection of the Museo di Storia Naturale, Università di Pisa, Via Roma 79, Calci (Pisa, Italy), under catalogue number 19,911.

2. Geological Background and Occurrence of Grammatikopoulosite

Grammatikopoulosite was discovered in a heavy mineral concentrate, which was prepared from chromitite specimens hosted in the mantle sequence of the Mesozoic Othrys ophiolite, central Greece (Figure 1A–C).

Othrys ophiolite is a complete but dismembered suite (Mirna Group) and consists of three structural units: the uppermost succession with variably serpentinized peridotites, which is structurally bounded by an ophiolite mélange; the intermediate Kournovon dolerite, including cumulate gabbro and local rhyolite; and the lower Sipetorrema Pillow Lava unit including also basaltic flows, siltstones, and chert. The Mirna Group constituted multiple inverted thrust sheets [7], which were eventually obducted onto the Pelagonian Zone during the Late Jurassic–Early Cretaceous [8–14]. Three types of basalts with different geochemical signatures have been described: (i) alkaline within-plate (WPB), (ii) normal-type mid-ocean ridge (N-MORB), and (iii) low-K tholeiite (L-KT). A biostratigraphic investigation indicated that radiolarites associated with N-MORB were deposited in the Middle and Late Triassic. Radiolarites deposited over the L-KT basalts are Early Carnian–Middle Norian–Late Norian in age [15]. N-MORB erupted during the Middle–Late Triassic period. The L-KT basalts erupted during the Middle–Late Triassic period. The L-KT basalts erupted during the Middle–Late Triassic period. The L-KT basalts or continent transition zone, close to the rifted continental margin. Finally, the alkaline WPB are interpreted to have formed in oceanic seamounts or in the ocean–continent transition zone adjacent to the rifted continental margin [15].

Electron microprobe analyses of the assemblages of these chromitites revealed that the studied spinel-supergroup minerals are magnesiochromites [1,5,6]. However, spinel mineral chemistry is rather heterogeneous with Cr_2O_3 (44.96–51.64 wt %), Al_2O_3 (14.18–20.78 wt %), MgO (13.34–16.84 wt %), and FeO (8.3–13.31 wt %). The calculated Fe₂O₃ ranges from 6.72 to 9.26 wt %. The amounts of MnO (0.33–0.60 wt %), V_2O_3 (0.04–0.30 wt %), ZnO (up to 0.07 wt %), and NiO (0.03–0.24 wt %) exhibit minor variations. The TiO₂ content is low (0.03–0.23 wt %), in agreement with the typical value reported for the mantle-hosted podiform chromites. The Cr/(Cr + Al) ratios of the investigated magnesiochromite are lower than those of chromites formed in the supra-subduction zone (SSZ) and those related to boninitic melt [1,5,6].



Figure 1. Location of the Othrys complex in Greece: (**A**) general geological map of the Othrys ophiolite showing the location of the Agios Stefanos chromium mine; (**B**) and (**C**) detailed geological setting of the Agios Stefanos area (modified after [5,10,16,17]).

3. Analytical Methods

The processing and recovery of the heavy minerals were carried out by treating about 10 kg of massive chromitite at SGS Mineral Services, Canada, following the procedure described by several authors [1,5,6,18]. Approximately 10 kg of chromitite were collected in the abandoned mine of Agios Stefanos. Subsequently, they were crushed and blended to generate a homogenous composite sample. About 500 g of the composite sample were riffled and stage crushed to a P80 (80% passing) of 75 μ m. After, the sample was treated with heavy liquid (the density of the heavy liquids was 3.1 g/cm³) to separate the heavy minerals. This procedure produced a heavy and light mineral concentrate. The heavy fraction was selected and processed with a superpanner. This method is designed for small amounts of sample and is closely controlled, leading to very effective separation. It consists of a tapering triangular deck with a "V" shape cross section. The table reproduces the concentrating action of a gold pan. Initially, the sample is swirled to stratify the minerals. Then, the heaviest minerals settle to the bottom and are deposited on the deck surface. The less dense material moves towards the top, overlying the heavy minerals. The operation of the deck is then changed to a rapid reciprocal motion, with an appropriate "end-knock" at the up-slope end of the board, and a steady flow of wash water is introduced. The "end-knock" forces the heavy minerals to migrate to the up-slope end of the deck. The wash water carries the light minerals to move to the narrower, down-slope end of the deck. The heaviest fractions were split into the heaviest fraction (tip) followed by a less dense fraction (middling). The "tip" and the "middlings" of the superpanner are the densest fractions and included liberated grains of chromite, sulphides, alloys, and phosphides, while the lighter tail consisted of a particle mixture of chromite and silicates. No source of contamination is likely during

are listed in Table 1.

sample collection and subsequent treatment. The heavy minerals were prepared in epoxy blocks, and then polished for mineralogical examination. Quantitative chemical analyses and acquisition of back-scattered electron images of grammatikopoulosite were performed with a JEOL JXA-8200 electron microprobe, installed in the E. F. Stumpfl laboratory, Leoben University, Austria, operating in Wavelength Dispersive Spectrometry (WDS) mode. Major and minor elements were determined at a 20 kV accelerating voltage and a 10 nA beam current, with 20 s as the counting time for the peak and 10 s for the backgrounds. The beam diameter was about 1 μ m in size. For the WDS analyses, the following lines and diffracting crystals were used: P, S, Si = (K α , PETJ), V, Fe, Co, Ni = (K α , LIFH), and Mo = (L α , PETJ). The following standards were selected: synthetic Ni₃P for Ni and P, molybdenite for S and Mo, synthetic Fe₃P for Fe, synthetic metallic vanadium for V, skutterudite for Co, quartz for Si, and chromite for Cr. The ZAF correction method was applied. Automatic corrections were performed for interferences P-Mo, Cr-V, and Mo-S. Representative analyses of grammatikopoulosite

 Table 1. Chemical data (wt % of elements) for grammatikopoulosite.

Sample	Р	S	Ni	\mathbf{V}	Со	Mo	Fe	Si	Total
VP40-1	20.38	0.41	21.98	21.02	16.33	16.72	3.82	0.14	100.79
VP40-2	19.83	0.42	21.70	20.48	16.66	16.36	3.83	0.14	99.41
VP40-3	19.65	0.39	21.72	20.73	16.51	16.35	3.86	0.13	99.33
VP40-4	19.65	0.40	21.95	21.05	16.37	16.31	3.85	0.13	99.71
VP40-5	20.01	0.41	21.69	20.98	16.45	16.20	3.78	0.16	99.67
average	19.90	0.41	21.81	20.85	16.46	16.39	3.83	0.14	99.79

Single-crystal and powder X-ray diffraction data were collected at the University of Florence using a Bruker D8 Venture equipped with a Photon II CCD detector, with graphite-monochromatized MoK α radiation ($\lambda = 0.71073$ Å) using a crystal fragment hand-picked from the polished section under a reflected light microscope (cif file, see Supplementary Materials). The crystal (about 80 µm in size) was carefully and repeatedly washed in acetone. It did not show any other visible phase attached to the surface. Single-crystal X-ray diffraction intensity data were integrated and corrected for standard Lorentz polarization factors with the software package *Apex3* [19,20].

The reflectance measurements on grammatikopoulosite were carried out using a WTiC standard and a J&M TIDAS diode array spectrophotometer at the Natural History Museum of London, UK.

4. Physical and Optical Properties

More than 30 grains of grammatikopoulosite were found in the studied polished sections. Grammatikopoulosite occurs as anhedral to subhedral grains. Most of them were less than 10 μ m in size and only two of them were up to about 80 μ m. Grammatikopoulosite consists of single (Figure 2A) or poly-phase grains associated with other minerals, such as tsikourasite, nickelphosphide, awaruite, and a V-sulphide, which likely represents another new mineral (Figure 2B,C). Compositions of these minerals have been reported in previous papers [5,6]. In plane-polarized light, grammatikopoulosite is creamy-yellow, weakly bireflectant, with measurable but not discernible pleochroism and slight anisotropy with indeterminate rotation tints. Internal reflections were not observed. Reflectance values of the mineral in air (*R* in %) are reported in Table 2 and in Figure 3. Density was not measured because of the small amount of available material and the presence of fine intergrowths with awaruite. The calculated density is equal to 7.085 g·cm⁻³, based on the empirical composition and unit–cell volume refined from single-crystal XRD data.



Figure 2. Digital image in reflected plane-polarized light (A,B) and a back-scattered electron image (C) showing grammatikopoulosite from the chromitite of Agios Stefanos. Abbreviations: Grm = grammatikopoulosite, Tsk = tsikourasite, VS = V-sulphide, Aw = awaruite, Npd = nickelphosphide, Epx = epoxy. Scale bar = 50 microns.



Figure 3. Reflectance data for grammatikopoulosite.

λ nm	R ₁	λ nm	R ₂
400	47.6	400	48.8
420	47.9	420	49.1
440	48.3	440	49.4
460	48.6	470	49.9
470	48.8	470	50.3
480	49.0	480	50.7
500	49.4	500	51.5
520	49.9	520	52.4
540	50.3	540	53.3
546	50.5	546	53.5
560	50.9	560	54.1
580	51.4	580	54.9
589	51.7	589	55.2
600	51.9	600	55.5
620	52.4	620	56.2
640	53.0	640	56.8
650	53.2	650	57.1
680	53.8	680	58.0
700	54.2	700	58.6

Table 2. Reflectance values for grammatikopoulosite. The values required by the Commission on Ore Mineralogy are given in bold.

5. Chemical Composition and X-Ray Crystallography

Chemical composition and X-ray data reveal that the empirical formula of grammatikopoulosite, based on $\Sigma(V + Ni + Co + Mo + Fe + Si) = 2$ apfu and taking into account the structural results (see below), is $(Ni_{0.57}Co_{0.32}Fe_{0.11})_{\Sigma 1.00}(V_{0.63}Mo_{0.26}Co_{0.11})_{\Sigma 1.00}(P_{0.98}S_{0.02})_{\Sigma 1.00}$. The simplified formula is (Ni,Co)(V,Mo)P and the ideal formula is NiVP, which corresponds to the composition of Ni = 41.74%, V = 36.23%, and P 22.03% (total 100 wt %).

A small grammatikopoulosite grain (about 80 µm in size) was handpicked from the polished section under a reflected light microscope and mounted on a 5 µm diameter carbon fiber, which was, in turn, attached to a glass rod in preparation for the single-crystal X-ray diffraction study. The fragment consists of crystalline grammatikopoulosite associated with minor, fine-grained polycrystalline awaruite. A total of 493 unique reflections were collected up to $2\theta = 80.33^{\circ}$. The mineral is orthorhombic, space group *Pnma*, with a = 5.8893(8), b = 3.5723(4), c = 6.8146(9) Å, and Z = 4. Given the similarity in unit–cell values and space groups, the structure was refined starting from the atomic coordinates reported for allabogdanite [21] using the software Shelxl-97 [22]. The site occupancy factor (s.o.f.) at the two cation sites was allowed to vary (Ni vs. Mo) using scattering curves for neutral atoms taken from the International Tables for Crystallography [23], leading to 27.5 and 28.4 e^- for M1 and M2 sites, respectively. The P site was found to be fully occupied (P vs. structural vacancy) by phosphorous (site scattering = $15.0 e^{-}$) and fixed accordingly. Taking into account the site distribution observed in florenskyite [24], allabogdanite [21], and andreyivanovite [25], and in most Co₂Si-structure-type synthetic compounds [26], we assigned Ni, Fe, and Co to fill the *M*1 site (i.e., Ni_{0.57}Co_{0.32}Fe_{0.11}) and all the other elements (i.e., $V_{0.63}Mo_{0.26}Co_{0.11}$) to the M2 site. The mean electron numbers calculated with such a site distribution were identical (27.36 and $28.38 e^{-}$) to those obtained in the refinement. For this reason, the subsequent cycles of refinement were run with the above constrained site populations, yielding an $R_1 = 0.0276$ for 465 reflections with $Fo > 4\sigma(Fo)$ and $R_1 = 0.0291$ for all the 493 independent reflections and 19 parameters. Refined atomic coordinates and isotropic displacement parameters are given in Table 3, whereas selected bond distances are reported in Table 4.

X-ray powder diffraction data for grammatikopoulosite (Table 5) were obtained with a Bruker D8 Venture equipped with a Photon II CCD detector, with graphite-monochromatized CuK α radiation ($\lambda = 1.54138$ Å). The least squares refinement gave the following values: a = 5.8088(2), b = 3.5993(2), c = 6.8221(3) Å, and V = 142.634(8) Å³. The calculated powder diffraction pattern obtained using the site occupancies and atomic coordinates is reported in Table 3.

Atom	Site Occupancy	x/a	y/b	z/c	$U_{\rm iso}$
<i>M</i> 1	Ni _{0.57} Co _{0.32} Fe _{0.11}	0.35709(6)	$\frac{1}{4}$	0.93703(5)	0.00578(10)
M2	V _{0.63} Mo _{0.26} Co _{0.11}	0.47087(6)	$\frac{1}{4}$	0.33109(5)	0.00595(9)
Р	P _{1.00}	0.23639(12)	$\frac{1}{4}$	0.62449(10)	0.00547(13)

Table 3. Atoms, site occupancies, atom coordinates, and isotropic displacement parameters ($Å^2$) for grammatikopoulosite.

Table 4. Selected bond distances (Å) for grammatikopoulosite.

Atoms	Bond Distance
<i>M</i> 1–P	2.2453(8)
M1–P (×2)	2.2639(5)
М1-Р	2.2728(8)
M1–M1 (×2)	2.6000(6)
M1–M2 (×2)	2.7280(5)
M1–M2 (×2)	2.7487(5)
M1-M2	2.7677(5)
M1-M2	2.7696(5)
М2-Р	2.4299(8)
M2–P (×2)	2.5008(6)
M2–P (×2)	2.5811(6)
M2–M1 (×2)	2.7280(5)
M2–M1 (×2)	2.7487(5)
M2–M1	2.7677(5)
M2–M1	2.7696(5)
M2–M2 (×2)	2.9339(6)

Table 5. Observed and calculated X-ray powder diffraction data (d in Å) for grammatikopoulosite. The strongest observed reflections are given in bold.

Indices	1	_	2	
hkl	d _{obs}	I _{obs}	d _{calc}	I _{calc}
101	4.43	10	4.4559	14
002	-	-	3.4073	5
102	2.950	20	2.9493	19
111	2.785	25	2.7872	24
201	2.699	5	2.7031	5
112	2.273	60	2.2743	65
210	2.269	10	2.2722	8
201	2.230	10	2.2279	9
211	2.157	100	2.1555	100
103	2.118	25	2.1194	27
013	1.915	15	1.9168	14
301	1.888	10	1.8864	12
113	1.824	15	1.8227	20
020	1.784	20	1.7861	21
004	1.702	10	1.7036	10
302	1.700	15	1.7010	22
213	1.608	10	1.6065	7
114	1.489	5	1.4878	6
303	1.482	5	1.4853	6
400	1.470	5	1.4723	6
123	1.367	10	1.3658	7
322	1.233	10	1.2318	9
314	1.211	5	1.2106	6
215	1.170	10	1.1688	9
511	1.102	5	1.1038	6
513	1.005	5	1.0035	5

1 = observed diffraction pattern; 2 = calculated diffraction pattern obtained with the atomic coordinates reported in Table 3 (only reflections with $I_{rel} \ge 4$ are listed).

6. Description of the Structure and Relations to Other Species

In the structure of grammatikopoulosite (Figure 4), *M*1 links four P atoms and eight *M*2 (Figure 5—left), whereas *M*2 links five P, six *M*1, and two *M*2 (Figure 5—right). The metal–phosphorous distances are much shorter in the *M*1 coordination sphere than in that of *M*2 (Table 4). Interestingly, if only the *M*–P distances are considered in the coordination polyhedra of the *M* atoms, *M*1P₄ tetrahedra (Figure 6) forming corner-sharing chains along the **b**-axis or *M*2P₅ square pyramids forming zig-zag chains along the **a**-axis (Figure 7) can be observed.



Figure 4. The crystal structure of grammatikopoulosite projected down [10]. Blue, green, and red circles refer to *M*1, *M*2, and P, respectively.



Figure 5. Coordination environment of *M*1 (left) and *M*2 (right) sites in the crystal structure of grammatikopoulosite. Colors and symbols as in Figure 4.



Figure 6. Polyhedral representation of the $M1P_4$ tetrahedra in the crystal structure of grammatikopoulosite. Colors and symbols as in Figure 4.



Figure 7. Polyhedral representation of the M2P₅ distorted pyramids in the crystal structure of grammatikopoulosite. Colors and symbols as in Figure 4.

It was reported that the departure from the hexagonal, high-temperature *P*-62*m* structure (barringerite structure) was linked to some ordering between the metals at the *M* sites. This was not immediately evident in both allabogdanite [21] and andreyivanovite [25] because of the close site scattering of Fe, Ni, and Cr. On the contrary, the ordering is evident if we take into consideration florenskyite [24] and grammatikopoulosite with ideal formulae FeTiP and NiVP, respectively. Ti and V preferentially enter the larger *M*2 site in agreement with their larger metallic radius (1.47 and 1.35 Å, respectively [27]).

Grammatikopoulosite does not correspond to any valid or invalid unnamed mineral [28] and it belongs to the group of natural phosphides with the Co₂Si orthorhombic *Pnma* structure, i.e., florenskyite [24], allabogdanite [21], and andreyivanovite [25].

7. Discussion and Genetical Implications

Grammatikopoulosite, similar to tsikourasite, is natural in origin despite the fact that both phases were found in heavy mineral concentrates and not in situ [1]. This assumption is supported by the following observations: during the sampling, the only tool used to collect the chromitite was a steel hammer; the concentrates were obtained in a laboratory that uses a grinder manufactured by TM Engineering, with media Alloy 1 composed of Cr and Mo, and does not contain Ni, V, Co, Fe, or P; the used material is very hard and is recognized to be best for applications where abrasion and impact is the norm; and the chemical composition of grammatikopoulosite indicates crystallization under reducing conditions, similar to what has been previously proposed for other phosphides found in ophiolitic chromitites [1,3–6]. According to literature data, the crystallization temperature of most of the terrestrial phosphides is generally comprised between 700 °C and 1150 °C [4]. However, a conclusive model to explain the origin of grammatikopoulosite cannot be provided yet, since it may have a terrestrial origin or represent fragments of a meteorite. The following models, which imply the crystallization in a terrestrial environment from low to high temperatures, can be suggested for the origin of grammatikopoulosite: (i) alteration of chromitite during sub-oceanic or on-land serpentinization at temperatures between 400 °C and 150 °C, which is a typical reducing process [29]; (ii) reaction of the chromitite with high-temperature reducing fluids in the mantle; and (iii) interaction of the chromitite and their host serpentinized peridotites with a surface lightning strike. Phosphide minerals occurring in ophiolitic chromitites in Othrys and Gerakini-Ormylia (Greece), as well as in Alapaevsk (Russia), are closely associated with highly reducing phases such as awaruite, typically forming during serpentinization [1,3-6]. Likewise, grammatikopoulosite is also associated with awaruite and other reducing phases, likely suggesting a low-temperature origin during serpentinization. This hypothesis is strongly supported by the discovery of a Ni-phosphide in situ in

the serpentine-rich matrix of the Alapaevsk chromitite of Russia [3]. Recently, Etiope et al. [30,31] have shown that the Agios Stefanos chromitite is the source rock of abiotic CH₄ measured in the west Othrys springs. These authors argued that CH₄ formation took place at temperatures below 150 °C via Sabatier reaction during oceanic and on-land serpentinization. This observation provides further evidence that reducing conditions can be achieved at a low temperature during the alteration of mantle-derived rocks. Alternatively, Xiong et al. [32] suggested that reducing conditions can be achieved in mantle rocks, including chromitites, in the shallow lithosphere due to the interaction of mantle-derived fluids enriched in CH₄ and H₂ with basaltic melt. However, no phosphides are reported in the ultra-reduced, high-temperature mineralogical assemblage described by Xiong et al. [32]. Natural phosphides and other ultra-reduced minerals have been rarely described in few fulgurites [33,34] but they are very common accessory phases in meteorites [21,24,25,35–39] or. The rare and localized occurrence of the Othrys phosphides may provide some support to the two last models. However, the probability to have collected a fragment of a meteorite or a fulgurite in the Othrys ophiolite during the sampling of the studied chromitite seems improbable, and for the discrimination of the meteoritic materials, a more detailed isotopic study may be needed. To conclude, an indisputable genetic model to explain how grammatikopoulosite is crystallized still needs further investigation.

Supplementary Materials: The following are available online at http://www.mdpi.com/2075-163X/10/2/131/s1, CIF: grammatikopoulosite.

Author Contributions: L.B. and F.Z. wrote the manuscript; F.Z. and L.B. performed the chemical analyses and the diffraction experiments, respectively; E.I. and B.T. provided the concentrate sample and information on the sample provenance and petrography of Othrys chromitite; G.G. and D.M. discussed the chemical data; and C.S. obtained the optical data. All authors provided support in the data interpretation and revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The authors are grateful to the University Centrum for Applied Geosciences (UCAG) for the access to the E.F. Stumpfl electron microprobe laboratory. The authors also thank SGS Mineral Services, Canada, for performing the concentrate sample. S. Karipi is thanked for collecting the sample and participating in the field work. L. Bindi thanks MIUR, project "TEOREM deciphering geological processes using Terrestrial and Extraterrestrial ORE Minerals", prot. 2017AK8C32 (PI: Luca Bindi).

Acknowledgments: The authors acknowledge Ritsuro Miyawaki, Chairman of the CNMNC and its members for helpful comments on the submitted new mineral proposal. We are very grateful to the four anonymous referees for constructive comments that improved the manuscript. Many thanks are due to the editorial staff of *Minerals* and to the consulted referees. This paper is dedicated to the memory of our friend and colleague Demetrios.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- 1. Zaccarini, F.; Bindi, L.; Ifandi, E.; Grammatikopoulos, T.; Stanley, C.; Garuti, G.; Mauro, D. Tsikourasite, $Mo_3Ni_2P_{1+x}$ (x < 0.25), a new phosphide from the chromitite of the othrys ophiolite, Greece. *Minerals* **2019**, *9*.
- 2. Britvin, S.N.; Murashko, M.N.; Vapnik, Y.; Polekhovsky, Y.S.; Krivovichev, S.V. Earth's phosphides in levant and insights into the source of Archean prebiotic phosphorus. *Sci. Rep.* **2015**, *5*. [CrossRef]
- 3. Zaccarini, F.; Pushkarev, E.; Garuti, G.; Kazakov, I. Platinum-group minerals and other accessory phases in chromite deposits of the Alapaevsk ophiolite, Central Urals, Russia. *Minerals* **2016**, *6*, 108. [CrossRef]
- 4. Sideridis, A.; Zaccarini, F.; Grammatikopoulos, T.; Tsitsanis, P.; Tsikouras, B.; Pushkarev, E.; Garuti, G.; Hatzipanagiotou, K. First occurrences of Ni-phosphides in chromitites from the ophiolite complexes of Alapaevsk, Russia and GerakiniOrmylia, Greece. *Ofioliti* **2018**, *43*, 75–84.
- Ifandi, E.; Zaccarini, F.; Tsikouras, B.; Grammatikopoulos, T.; Garuti, G.; Karipi, S.; Hatzipanagiotou, K. First occurrences of Ni-V-Co phosphides in chromitite of Agios Stefanos mine, Othrys ophiolite, Greece. *Ofioliti* 2018, 43, 131–145.
- Zaccarini, F.; Ifandi, E.; Tsikouras, B.; Grammatikopoulos, T.; Garuti, G.; Mauro, D.; Bindi, L.; Stanley, C. Occurrences of of new phosphides and sulfide of Ni, Co, V, and Mo from chromitite of the Othrys ophiolite complex (Central Greece). *Per. Ital. Mineral.* 2019, 88. [CrossRef]

- 7. Smith, A.G.; Rassios, A. The evolution of ideas for the origin and emplacement of the western Hellenic ophiolites. *Geol. Soc. Am. Spec. Pap.* **2003**, *373*, 337–350.
- 8. Hynes, A.J.; Nisbet, E.G.; Smith, G.A.; Welland, M.J.P.; Rex, D.C. Spreading and emplacement ages of some ophiolites in the Othris region (eastern central Greece). *Z. Deutsch Geol. Ges.* **1972**, *123*, 455–468.
- 9. Smith, A.G.; Hynes, A.J.; Menzies, M.; Nisbet, E.G.; Price, I.; Welland, M.J.; Ferrière, J. The stratigraphy of the Othris Mountains, eastern central Greece: A deformed Mesozoic continental margin sequence. *Eclogue Geol. Helv.* **1975**, *68*, 463–481.
- Rassios, A.; Smith, A.G. Constraints on the formation and emplacement age of western Greek ophiolites (Vourinos, Pindos, and Othris) inferred from deformation structures in peridotites. In *Ophiolites and Oceanic Crust: New Insights from Field Studies and the Ocean Drilling Program*; Dilek, Y., Moores, E., Eds.; Geological Society of America: Boulder, CO, USA, 2001; pp. 473–484.
- 11. Barth, M.G.; Mason, P.R.D.; Davies, G.R.; Drury, M.R. The Othris Ophiolite, Greece: A snapshot of subduction initiation at a mid-ocean ridge. *Lithos* **2008**, *100*, 234–254. [CrossRef]
- 12. Barth, M.; Gluhak, T. Geochemistry and tectonic setting of mafic rocks from the Othris Ophiolite, Greece. *Contrib. Mineral. Petrol.* **2009**, *157*, 23–40. [CrossRef]
- 13. Dijkstra, A.H.; Barth, M.G.; Drury, M.R.; Mason, P.R.D.; Vissers, R.L.M. Diffuse porous melt flow and melt-rock reaction in the mantle lithosphere at a slow-spreading ridge: A structural petrology and LA-ICP-MS study of the Othris Peridotite Massif (Greece). *Geochem. Geophys. Geosyst.* **2003**, *4*. [CrossRef]
- 14. Magganas, A.; Koutsovitis, P. Composition, melting and evolution of the upper mantle beneath the Jurassic Pindos ocean inferred by ophiolitic ultramafic rocks in East Othris, Greece. *Int. J. Earth Sci.* **2015**, *104*, 1185–1207. [CrossRef]
- 15. Bortolotti, V.; Chiari, M.; Marcucci, M.; Photiades, A.; Principi, G.; Saccani, E. New geochemical and age data on the ophiolites from the Othrys area (Greece): Implication for the Triassic evolution of the Vardar ocean. *Ofioliti* **2008**, *33*, 135–151.
- Economou, M.; Dimou, E.; Economou, G.; Migiros, G.; Vacondios, I.; Grivas, E.; Rassios, A.; Dabitzias, S. Chromite deposits of Greece. In *Chromites, UNESCO's IGCP197 Project Metallogeny of Ophiolites*; Petrascheck, W., Karamata, S., Eds.; Theophrastus Publ. S.A.: Athens, Greece, 1986; pp. 129–159.
- Garuti, G.; Zaccarini, F.; Economou-Eliopoulos, M. Paragenesis and composition of laurite from chromitites of Othrys (Greece): Implications for Os-Ru fractionation in ophiolite upper mantle of the Balkan Peninsula. *Mineral. Depos.* 1999, 34, 312–319. [CrossRef]
- Tsikouras, B.; Ifandi, E.; Karipi, S.; Grammatikopoulos, T.A.; Hatzipanagiotou, K. Investigation of platinum-group minerals (PGM) from Othrys chromitites (Greece) using superpanning concentrates. *Minerals* 2016, *6*, 94. [CrossRef]
- 19. Bruker. *APEX3*; Bruker AXS Inc.: Madison, WI, USA, 2016; Available online: https://www.bruker.com/products/x-ray-diffraction-and-elemental-analysis/single-crystal-x-ray-diffraction/sc-xrd-software/apex3.html (accessed on 31 January 2020).
- 20. Bruker. *SAINT and SADABS*; Bruker AXS Inc.: Madison, WI, USA, 2016. Available online: https://www.bruker.com/products/x-ray-diffraction-and-elemental-analysis/single-crystal-x-ray-diffraction/sc-xrd-software/apex3.html (accessed on 31 January 2020).
- 21. Britvin, S.N.; Rudashevskii, N.S.; Krivovichev, S.V.; Burns, P.C.; Polekhovsky, Y.S. Allabogdanite, (Fe,Ni)₂P, a new mineral from the Onello meteorite: The occurrence and crystal structure. *Am. Mineral.* **2002**, *87*, 1245–1249. [CrossRef]
- 22. Sheldrick, G.M. A short history of SHELX. Acta Crystallogr. 2008, A64, 112–122. [CrossRef]
- 23. Wilson, A.J.C. *International Tables for Crystallography: Mathematical, Physical, and Chemical Tables;* International Union of Crystallography: Chester, UK, 1992; Volume 3.
- 24. Ivanov, A.V.; Zolensky, M.E.; Saito, A.; Ohsumi, K.; Yang, S.V.; Kononkova, N.N.; Mikouchi, T. Florenskyite, FeTiP, a new phosphide from the Kaidun meteorite, Locality: Kaidun chondritic meteorite, South Yemen. *Am. Mineral.* **2000**, *85*, 1082–1086. [CrossRef]
- 25. Zolensky, M.; Gounelle, M.; Mikouchi, T.; Ohsumi, K.; Le, L.; Hagiya, K.; Tachikawa, O. Andreyivanovite: A second new phosphide from the Kaidun meteorite. *Am. Mineral.* **2008**, *93*, 1295–1299. [CrossRef]
- 26. Fruchart, R.; Roger, A.; Sénateur, J.P. Crystallographic and magnetic properties of solid solutions of the phosphides M2P, M = Cr, Mn, Fe, Co, and N. *J. Appl. Phys.* **1969**, *40*, 1250–1257. [CrossRef]
- 27. Wells, A.F. Structural Inorganic Chemistry, 5th ed.; Clarendon Press: Oxford, UK, 1984; p. 1288.

- Smith, D.G.W.; Nickel, E.H. A system for codification for unnamed minerals: Report of the subcommittee for unnamed minerals of the IMA commission on new minerals, nomenclature and classification. *Can. Mineral.* 2007, 45, 983–1055. [CrossRef]
- 29. Malvoisin, B.; Chopin, C.; Brunet, F.; Matthieu, E.; Galvez, M.E. Low-temperature Wollastonite formed by carbonate reduction: A marker of serpentinite redox conditions. *J. Petrol.* **2012**, *53*, 159–176. [CrossRef]
- 30. Etiope, G.; Tsikouras, B.; Kordella, S.; Ifandi, E.; Christodoulou, D.; Papatheodorou, G. Methane flux and origin in the Othrys ophiolite hyperalkaline springs, Greece. *Chem. Geol.* **2013**, 347, 161–174. [CrossRef]
- 31. Etiope, G.; Ifandi, E.; Nazzari, M.; Procesi, M.; Tsikouras, B.; Ventura, G.; Steele, A.; Tardini, R.; Szatmari, P. Widespread abiotic methane in chromitites. *Sci. Rep.* **2018**, *8*, 8728. [CrossRef]
- 32. Xiong, Q.; Griffin, W.L.; Huang, J.X.; Gain, S.E.M.; Toledo, V.; Pearson, N.J.; O'Reilly, S.Y. Super-reduced mineral assemblages in "ophiolitic" chromitites and peridotites: The view from Mount Carmel. *Eur. J. Mineral.* **2017**, *29*, 557–570. [CrossRef]
- 33. Pasek, M.A.; Hammeijer, J.P.; Buick, R.; Gull, M.; Atlas, Z. Evidence for reactive reduced phosphorus species in the early Archean ocean. *Proc. Natural Acad. Sci. USA* **2013**, *110*, 100089–100094. [CrossRef]
- 34. Ballhaus, C.; Wirth, R.; Fonseca, R.O.C.; Blanchard, H.; Pröll, W.; Bragagni, A.; Nagel, T.; Schreiber, A.; Dittrich, S.; Thome, V.; et al. Ultra-high pressure and ultra-reduced minerals in ophiolites may form by lightning strikes. *Geochem. Perspec. Lett.* **2017**, *5*, 42–46. [CrossRef]
- 35. Buseck, P.R. Phosphide from meteorites: Barringerite, a new iron-nickel mineral. *Science* **1969**, *165*, 169–171. [CrossRef]
- Britvin, S.N.; Kolomensky, V.D.; Boldyreva, M.M.; Bogdanova, A.N.; Krester, Y.L.; Boldyreva, O.N.; Rudashevsky, N.S. Nickelphosphide (Ni,Fe)₃P—The nickel analogue of schreibersite. *Zap. Vserossi. Mineral. Obschch.* 1999, 128, 64–72.
- 37. Ma, C.; Beckett, J.R.; Rossman, G.R. Monipite, MoNiP, a new phosphide mineral in a Ca-Al-rich inclusion from the Allende meteorite. *Am. Mineral.* **2014**, *99*, 198–205. [CrossRef]
- Pratesi, G.; Bindi, L.; Moggi-Cecchi, V. Icosahedral coordination of phosphorus in the crystal structure of melliniite, a new phosphide mineral from the Northwest Africa 1054 acapulcoite. *Am. Mineral.* 2006, *91*, 451–454. [CrossRef]
- 39. Skala, R.; Cisarova, I. Crystal structure of meteoritic schreibersites: Determination of absolute structure. *Phys. Chem. Mineral.* **2005**, *31*, 721–732. [CrossRef]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).