

Article

Pseudo Overlap Functions, Fuzzy Implications and Pseudo Grouping Functions with Applications

Xiaohong Zhang ^{1,2,*}, Rong Liang ¹ , Humberto Bustince ³, Benjamin Bedregal ⁴ , Javier Fernandez ³, Mengyuan Li ¹ and Qiqi Ou ¹

¹ Department of Mathematics, Shaanxi University of Science & Technology, Xi'an 710021, China

² Shaanxi Joint Laboratory of Artificial Intelligence, Shaanxi University of Science & Technology, Xi'an 710021, China

³ Departamento de Estadística, Matemáticas e Informática, Universidad Pública de Navarra, Campus Arrosadia s/n, 31006 Pamplona, Spain

⁴ Departamento de Informática e Matemática Aplicada, Universidade Federal do Rio Grande do Norte, Natal 59078-970, Brazil

* Correspondence: zhangxiaohong@sust.edu.cn

Abstract: Overlap and grouping functions are important aggregation operators, especially in information fusion, classification and decision-making problems. However, when we do more in-depth application research (for example, non-commutative fuzzy reasoning, complex multi-attribute decision making and image processing), we find overlap functions as well as grouping functions are required to be commutative (or symmetric), which limit their wide applications. For the above reasons, this paper expands the original notions of overlap functions and grouping functions, and the new concepts of pseudo overlap functions and pseudo grouping functions are proposed on the basis of removing the commutativity of the original functions. Some examples and construction methods of pseudo overlap functions and pseudo grouping functions are presented, and the residuated implication (co-implication) operators derived from them are investigated. Not only that, some applications of pseudo overlap (grouping) functions in multi-attribute (group) decision-making, fuzzy mathematical morphology and image processing are discussed. Experimental results show that, in many application fields, pseudo overlap functions and pseudo grouping functions have greater flexibility and practicability.

Keywords: fuzzy logic; information fusion; pseudo overlap function; pseudo t-norm; fuzzy implication

MSC: 03B52; 03E72; 47S40



Citation: Zhang, X.; Liang, R.; Bustince, H.; Bedregal, B.; Fernandez, J.; Li, M.; Ou, Q. Pseudo Overlap Functions, Fuzzy Implications and Pseudo Grouping Functions with Applications. *Axioms* **2022**, *11*, 593. <https://doi.org/10.3390/axioms11110593>

Academic Editor: Oscar Castillo

Received: 5 October 2022

Accepted: 23 October 2022

Published: 26 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The aggregation function (or aggregation operator) is an important concept in decision theory, information fusion and fuzzy inference systems etc. (see [1,2]). The process of combining several (numerical) values into a single representative one is called aggregation, and the function performing this process is called aggregation function. As a special aggregation operator, the concept of overlap functions (OF for short) is proposed in [3] by scholars such as H. Bustince. Since then, OFs as well as associated grouping functions have become a new hot direction, with rich research achievements (see [4–10]), and have been applied in many fields, such as multi-attribute (group) decision-making, rule-based classification and image processing etc. (see [11–13]).

In recent years, in order to expand the application scope of OFs and grouping functions, many scholars have put forward some more general concepts, such as \tilde{r} -(quasi-)overlap functions [14] etc. However, all of them keep the requirement of commutativity (symmetry). This paper mainly removes the commutativity of OFs as well as grouping functions, thus

studies more general aggregation functions, that is, pseudo overlap functions as well as pseudo grouping functions.

It is necessary to emphasize the motivation for studying pseudo overlap (grouping) functions. Our main starting points include the following aspects:

(1) For aggregation functions, the commutativity is not always necessary, for example, a copula is a kind of aggregation function which is closely related to overlap function, but a copula could be non-commutative (see [15]). So, it is natural to investigate non-commutative overlap (grouping) functions;

(2) From the perspective of fuzzy logic theory, overlap functions are closely related to t-norms and fuzzy reasoning. The notion of t-norm is generalized to the non-commutative case, that is, pseudo t-norm, which plays an important role in non-commutative fuzzy logic (see [16–18]). Therefore, and in the same way, it becomes natural to study pseudo overlap (grouping) functions;

(3) The overlap functions can be applied to multi-attribute (group) decision problems, but when multiple attributes (or multiple decision makers) have different importance, overlap functions may display some limitations due to the commutativity. Then, it is necessary to extend overlap function to non-commutative case. In fact, I. A. Da Silva et al. constructed a kind of weighted average operators based on n-dimensional overlap functions (see Theorem 3.1 in [13]), which are non-commutative overlap functions when a weak condition is attached;

(4) In the theory of mathematical morphology, t-norms and fuzzy implication operators are the basic tools (see [19–23]). Similar to fuzzy morphology based on t-norms, we can construct a new mathematical morphology based on overlap functions. More generally, since the theory of mathematical morphology endowing non-commutative operators has wider applicability (see [24,25]), the establishment of corresponding fuzzy erosion and dilation operators considering pseudo overlap (grouping) functions can expand the existing fuzzy mathematical morphology theory;

(5) Moreover, grouping functions can be applied to image segmentation [11,12], but for image processing, the target object and background are usually unequal. In order to express such inequality (or asymmetry), we naturally select non-commutative operators, that is, pseudo grouping functions. We have reason to believe that we can obtain a better effect when we use pseudo grouping functions to edge extraction and threshold segmentation of images.

Based on the above considerations, we will study the basic properties and some applications of pseudo overlap (grouping) functions in detail. This paper is organized as follows. In Section 2, we review some preliminary concepts (pseudo t-norm, pseudo t-conorm, copula, overlap function, grouping function, fuzzy implication and residuated lattice). In Section 3, we give the notion of (n-dimension) pseudo overlap functions as well as some concrete examples, provide some construction methods, illustrate the relevant conclusions of residuated implications induced by pseudo overlap functions, and analyze the relationship between overlap functions and residuated lattices. In Section 4, we investigate pseudo grouping functions and co-implications derived from them. Section 5 presents some applications of pseudo overlap (grouping) functions in multi-attribute group decision making (MAGDM for short), fuzzy mathematical morphology and image processing. Finally, conclusions are declared in Section 6.

2. Preliminaries

In this part we state a few basic notions as well as results.

Definition 1 ([16,26]). A bivariate operator T on $[0, 1]$ is known as a pseudo t-norm when and only when:

(pt1) T meets associativity;

(pt2) T is increasing concerning both arguments, i.e., $T(x, z) \leq T(y, z)$ as well as $T(z, x) \leq T(z, y)$

when taking random $x, y, z \in [0, 1]$, $x \leq y$;

(pt3) T has unit element 1, that is, $T(c, 1) = c = T(1, c)$, when taking random $c \in [0, 1]$.

Definition 2 ([16]). The bivariate operator S on $[0, 1]$ is known as a pseudo t -conorm when and only when:

(ps1) S meets associativity;

(ps2) S is increasing about both arguments, i.e., $S(x, z) \leq S(y, z)$ as well as $S(z, x) \leq S(z, y)$ when taking random $x, y, z \in [0, 1]$, $x \leq y$;

(ps3) S has unit element 0, i.e., $S(x, 0) = x = S(0, x)$, when taking random $x \in [0, 1]$.

Definition 3 ([27]). The binary operation C on $[0, 1]$ is known as a copula when C meets conditions as below, for arbitrary $k, k', l, l' \in [0, 1]$ satisfying $k \leq k'$ as well as $l \leq l'$:

(C1) $C(k, l) + C(k', l') \geq C(k, l') + C(k', l)$;

(C2) $C(k, 0) = C(0, k) = 0$;

(C3) $C(l, 1) = C(1, l) = l$.

Definition 4 ([28]). An algebra $L = (L, \wedge, \vee, \otimes, \rightarrow, \rightsquigarrow, 0, 1)$ is known as a residuated lattice when meets the requirements as below:

(L1) $(L, \wedge, \vee, 0, 1)$ is a lattice with 0 as the lower bound and 1 as the upper bound;

(L2) $(L, \otimes, 1)$ is a monoid with 1 as neutral element;

(L3) $x \otimes y \leq z$ iff $x \leq y \rightarrow z \Leftrightarrow y \leq x \rightsquigarrow z$, when taking arbitrary $x, y, z \in L$.

Definition 5 ([4]). A bivariate mapping I on $[0, 1]$ is known as a fuzzy implication when taking random $u, v, w \in [0, 1]$, I meets:

(I1) monotonic decreasing concerning the first element: $I(v, w) \leq I(u, w)$ when $u \leq v$;

(I2) monotonic increasing concerning the second element: $I(u, v) \leq I(u, w)$ when $v \leq w$;

(I3) three boundary conditions: $I(0, 0) = 1, I(1, 1) = 1, I(1, 0) = 0$.

Next, we give some properties of fuzzy implication:

Definition 6 ([4]). A fuzzy implication I on $[0, 1]$ satisfies:

(NP) Neutral property, i.e., $I(1, e) = e, \forall 0 \leq e \leq 1$.

(EP) Exchange property, i.e., $I(x, I(y, z)) = I(y, I(x, z)), \forall x, y, z \in [0, 1]$.

(IP) Identity property, i.e., $I(x, x) = 1, \forall x \in [0, 1]$.

(LOP) Left ordering property, i.e., $x \leq y \Rightarrow I(x, y) = 1, \forall x, y \in [0, 1]$.

(ROP) Right ordering property, i.e., $I(s, t) = 1 \Rightarrow s \leq t, \forall s, t \in [0, 1]$.

(OP) Ordering property, i.e., $I(s, t) = 1 \Leftrightarrow s \leq t, \forall s, t \in [0, 1]$.

(CB) Consequent boundary, i.e., $t \leq I(s, t), \forall s, t \in [0, 1]$.

(SIB) Sub-iterative boolean property, i.e., $I(s, I(s, t)) \geq I(s, t), \forall s, t \in [0, 1]$.

(IB) Iterative Boolean property, i.e., $I(s, I(s, t)) = I(s, t), \forall s, t \in [0, 1]$.

(SBC) Strong boundary condition for 0, i.e., $i \neq 0 \Rightarrow I(i, 0) = 0, \forall i \in [0, 1]$.

(LBC) Left boundary condition, i.e., $I(0, a) = 1, \forall a \in [0, 1]$.

(RBC) Right boundary condition, i.e., $I(b, 1) = 1, \forall b \in [0, 1]$.

(EP1) Exchange property for 1, i.e., $I(l, I(m, n)) = 1 \Rightarrow I(m, I(l, n)) = 1, \forall l, m, n \in [0, 1]$.

(PEP) Pseudo exchange property, i.e., $I(p, r) \geq q \Leftrightarrow I(q, r) \geq p, \forall p, q, r \in [0, 1]$.

Definition 7 ([3,4]). A bivariate function O on $[0, 1]$ is said to be an overlap function (OF for short) when O meets statements as below:

(O1) It meets commutativity;

(O2) Its value is 0 iff $xy = 0$;

(O3) Its value is 1 iff $x = y = 1$;

(O4) It meets monotonic increasing property;

(O5) It meets continuity.

Definition 8 ([5]). A binary operation G on $[0, 1]$ is said to be a grouping function when G meets statements as follows:

- (G1) It meets commutativity;
- (G2) Its value is 0 when and only when values of x and y are 0;
- (G3) Its value is 1 when and only when at least one of x and y is 1;
- (G4) It meets monotonic increasing property;
- (G5) It meets continuity.

Theorem 1 ([3]). An operator $O: [0, 1]^2 \rightarrow [0, 1]$ meets (O1) \sim (O5) when and only when there exist bivariate operations h, g on $[0, 1]$ with

$$O(x, y) = \frac{h(x, y)}{h(x, y) + g(x, y)},$$

where

- (1) h and g satisfy symmetry;
- (2) h satisfies monotonic increasing property and g satisfies monotone decreasing property;
- (3) the value of h is 0 when and only when at least one of x and y has a value of 0;
- (4) the value of g is 0 when and only when values of x and y are 1;
- (5) h as well as g satisfy continuity.

Definition 9 ([29–31]). A binary operation $C: [0, 1]^2 \rightarrow [0, 1]$ is said to be a co-implication if it satisfies:

- (C1') decreasing about its first element;
- (C2') increasing about its second variable;
- (C3') $C(0, 0) = 0$;
- (C4') $C(a, b) = 0$ when $a = 1$ as well as $b = 1$;
- (C5') $C(c, d) = 1$ when $c = 0$ as well as $d = 1$.

Definition 10 ([4]). An n -dimension mapping A on $[0, 1]$ is known as an aggregation operation when requirements as below are satisfied:

- (A1) A is monotonically increasing concerning all elements: for every $i \in \{1, 2, \dots, n\}$, $A(a_1, \dots, a_n) \leq A(a_1, \dots, a_{i-1}, y, a_{i+1}, \dots, a_n)$ when $a_i \leq y$;
- (A2) A meets requirements: (i) $A(a_1, \dots, a_n) = 0$ when $a_i = 0$ ($i = 1, \dots, n$) and (ii) $A(a_1, \dots, a_n) = 1$ when $a_i = 1$ ($i = 1, \dots, n$).

3. Pseudo Overlap Functions and Their Residuated Implications

In this section, we consider existing outcomes of residuated implication to introduce the concept of pseudo overlap functions and residuated implication derived from them. Then, we elaborate the relationship between pseudo overlap functions and continuous pseudo t-norms and copulas, and expand the dimension to explain the definition of multidimensional pseudo overlap functions. After that, we also illustrate the general construction method of pseudo overlap functions, and finally provide some examples to explain in detail.

Definition 11. A bivariate mapping $O: [0, 1]^2 \rightarrow [0, 1]$ is called a pseudo overlap function (briefly POF) when it meets statements as follows:

- (O1') The value of O is 0 iff at least one of the two variables has a value of 0;
- (O2') The value of O is 1 iff values of two variables are 1;
- (O3') O meets monotonic increment;
- (O4') O meets continuity.

Obviously, every overlap function is a pseudo overlap function. Now, we will provide some examples of POFs.

Example 1. (1) The operation $O: [0, 1]^2 \rightarrow [0, 1]$ defined, when taking arbitrary $a, b \in [0, 1]$, as

$$O(a, b) = a^p b^q, p, q > 0 \quad (1)$$

is a POF. When $p = q$, O is an OF.

(2) An operation $O: [0, 1]^2 \rightarrow [0, 1]$ defined, when taking random $x, y \in [0, 1]$, as

$$O(x, y) = \min\{x^p, y^q\}, p, q > 0 \quad (2)$$

is a POF. When $p = q$, O is an OF.

(3) A mapping $O: [0, 1]^2 \rightarrow [0, 1]$ defined, when taking random $x, y \in [0, 1]$, as

$$O(x, y) = \begin{cases} \frac{5xy+17}{22}, & \text{if } (x, y) \in A_1 \\ \frac{90xy}{56}, & \text{if } (x, y) \in A_2 \\ \frac{9x}{8}, & \text{if } (x, y) \in A_3 \\ \frac{2y-2}{11} + \frac{105x}{88}, & \text{if } (x, y) \in A_4 \\ \frac{9y}{7}, & \text{if } (x, y) \in A_5 \\ \frac{7x-7}{44} + \frac{205y}{154}, & \text{if } (x, y) \in A_6 \end{cases} \quad (3)$$

is a pseudo overlap function, where $A_1 = \{(x, y) | 0.8 \leq x \leq 1, 0.7 \leq y \leq 1, (x, y) \neq (0.8, 0.7)\}$, $A_2 = \{(x, y) | 0 \leq x \leq 0.8, 0 \leq y \leq 0.7\}$, $A_3 = \{(x, y) | 0 \leq x < 0.8, 0.7 < y \leq 1, y \leq 1 - \frac{3x}{8}\}$, $A_4 = \{(x, y) | 0 \leq x < 0.8, 0.7 < y \leq 1, y > 1 - \frac{3x}{8}\}$, $A_5 = \{(x, y) | 0.8 < x \leq 1, 0 \leq y < 0.7, y \leq \frac{7}{2}(1-x)\}$, $A_6 = \{(x, y) | 0.8 < x \leq 1, 0 \leq y < 0.7, y > \frac{7}{2}(1-x)\}$, specific interval distribution is as follows (see Figure 1).




Figure 1. Diagram of (3) in Example 1.

Lemma 1 ([16]). Given a pseudo t -norm T on $[0, 1]$ meeting continuity, then it meets commutativity, i.e., it is a t -norm.

Theorem 2. Let $O: [0, 1]^2 \rightarrow [0, 1]$ be a POF. Then

- (1) if O is commutative, then it is an overlap function.
- (2) if O is pseudo t -norm, then it is a positive continuous t -norm (where, “positive” means $x \neq 0$ and $y \neq 0 \Rightarrow O(x, y) > 0$).
- (3) if O is pseudo t -norm, then it is a copula if and only if it satisfies Lipschitz property with constant 1 (that is, $\forall x_1, x_2, y_1, y_2 \in [0, 1], |O(x_1, y_1) - O(x_2, y_2)| \leq |x_1 - x_2| + |y_1 - y_2|$).

Proof. (1) It follows from Definitions 7 and 11;

(2) If O is pseudo t -norm, since O is a pseudo overlap function, by Definition 11 (O4’), O is continuous. Using Lemma 1, we know that O is commutative. By (1), O meets (O1) ~ (O5). Thus, O is a t -norm satisfying continuity and has no nontrivial zero factors;

(3) Suppose that O is a pseudo overlap function and pseudo t -norm. By (2) we know that O

is a positive continuous t-norms. Applying Theorem 1 in [15], we can get that O is a copula if and only if it satisfies Lipschitz property with constant 1. \square

According to the proof of Theorem 2 in [15], we get that every positive copula is a pseudo overlap function. Therefore, we obtain the diagram in Figure 2 with the relationship between some concepts.




Figure 2. Intersection of the main classes of aggregation functions considered in this paper.

The following example shows that there are some POFs and pseudo t-norms which are not copulas.

Example 2. The operation $O: [0, 1]^2 \rightarrow [0, 1]$ defined, when taking arbitrary $x, y \in [0, 1]$, as

$$O(x, y) = \begin{cases} 0.5xy & \text{if } (x, y) \in [0, 0.5]^2, \\ x \cdot 2^{-4(1-y)} & \text{if } (x, y) \in [0, 0.5] \times (0.5, 1], \\ y \cdot 2^{-4(1-x)} & \text{if } (x, y) \in (0.5, 1] \times [0, 0.5], \\ 2^{4x+4y-7} & \text{if } (x, y) \in (0.5, 1]^2 \text{ and } x + y < 1.5, \\ x + y - 1 & \text{if } (x, y) \in (0.5, 1]^2 \text{ and } x + y \geq 1.5. \end{cases} \quad (4)$$

Then, O meets $(O1') \sim (O4')$ as well as being a t-norm, but it is not a copula (see [32,33]).

Definition 12. The n -dimension ($n \geq 2$) mapping O on $[0, 1]$ is known as an n -ary pseudo overlap function when properties as below are established:

- $(O1^n)$ $O(x_1, \dots, x_n) = 0$ when and only when there is at least a certain x_i ($i = 1, \dots, n$) with a value of 0;
- $(O2^n)$ The value of O is 1 when and only when the values of its all elements are 1;
- $(O3^n)$ O meets monotonic increment;
- $(O4^n)$ O is continuous.

Example 3. (1) An operation $O: [0, 1]^n \rightarrow [0, 1]$ formulated, when taking arbitrary $x_1, x_2, \dots, x_n \in [0, 1]$, as

$$O(x_1, \dots, x_n) = \prod_{i=1}^{n-1} x_i \cdot x_n^2 \quad (5)$$

is an n -dimension pseudo overlap function.

(2) The operator $O: [0, 1]^n \rightarrow [0, 1]$ given, when taking arbitrary $x_1, x_2, \dots, x_n \in [0, 1]$, as

$$O(x_1, \dots, x_n) = \min\{x_1, x_2, \dots, x_{n-1}, \sqrt{x_n}\} \quad (6)$$

is an n -dimension pseudo overlap function.

Next, we provide some construction methods and theorems of POFs.

Theorem 3. The operation $O: [0, 1]^2 \rightarrow [0, 1]$ is a POF when and only when there exist bivariate operators f, g on $[0, 1]$ with

$$O(x, y) = \frac{f(x, y)}{f(x, y) + g(x, y)},$$

where

- (1) f meets monotonic increment and g meets monotonic decreasing property;
- (2) $f(x, y) = 0$ when and only when $x = 0$ or $y = 0$;
- (3) $g(x, y) = 0$ when and only when $x = y = 1$;
- (4) f as well as g meeting continuity.

Proof. (\Leftarrow) By (2), we get: $xy = 0$ iff $f(x, y) = 0 \Leftrightarrow O(x, y) = 0$, i.e., the mapping O satisfies (O1').

By (3), we get: $xy = 1 \Leftrightarrow g(x, y) = 0 \Leftrightarrow O(x, y) = 1$, i.e., the mapping O satisfies (O2').

By (1), assume that $x_1 \leq x_2$, then for any $y \in [0, 1]$, $f(x_1, y)g(x_2, y) \leq f(x_2, y)g(x_1, y)$. Then we consider $f(x_1, y)f(x_2, y)$ as the common factor, we get $f(x_1, y)(f(x_2, y) + g(x_2, y)) \leq f(x_2, y)(g(x_1, y) + f(x_1, y))$, i.e., $O(x_1, y) \leq O(x_2, y)$. Since an analogous calculation holds for the other variable, O is non-decreasing, that is, the mapping O satisfies (O3').

By (4), we know that O is continuous, i.e., the mapping O satisfies (O4').

(\Rightarrow) Consider that O meets (O1') \sim (O4'), and assume $f(x, y) = O(x, y)$ and $g(x, y) = 1 - O(x, y)$. Then the function $O(x, y) = \frac{f(x, y)}{f(x, y) + g(x, y)}$ is well-defined. Besides, it is obvious that requirements (1) \sim (4) hold. \square

Proposition 1. Let $\varphi_1, \varphi_2, \varphi_3: [0, 1] \rightarrow [0, 1]$ be continuous and monotonically increasing operations satisfying $\varphi_i(x) = 0$ when and only when $x = 0$, $\varphi_i(x) = 1$ when and only when $x = 1$ ($i = 1, 2, 3$), and let O be a 2-dimension pseudo overlap function. Then the operation $O^{\varphi_1, \varphi_2, \varphi_3}$ is defined as:

$$O^{\varphi_1, \varphi_2, \varphi_3}(x, y) = \varphi_1(O(\varphi_2(x), \varphi_3(y)))$$

also meets (O1') \sim (O4').

Proof. Obviously, $O^{\varphi_1, \varphi_2, \varphi_3}$ meets (O3') and (O4'). We just need to reveal that two boundary conditions are true. Suppose first that $O^{\varphi_1, \varphi_2, \varphi_3}(x, y) = 0$. According to the properties that are met by φ_1 , it is obvious that this can be established when and only when $O(\varphi_2(x), \varphi_3(y)) = 0$. However, because O is a POF, when and only when $\varphi_2(x) = 0$ or $\varphi_3(y) = 0$ when and only when $xy = 0$. Analogously, another condition (O2') can be proven. \square

Proposition 2. Given POFs O_1, \dots, O_n , let $M: [0, 1]^n \rightarrow [0, 1]$ be a continuous aggregate operation meeting $M(x_1, \dots, x_n) = 0$ only if $x_i = 0$ for some $i = 1, \dots, n$ and $M(x_1, \dots, x_n) = 1$ only if $x_i = 1$ for a certain $i = 1, \dots, n$. Then operator $O(x, y) = M(O_1(x, y), \dots, O_n(x, y))$ fits (O1') \sim (O4').

Proof. Clearly, the operator meets (O3') and (O4'). (O1') and (O2') are proven as below. We can discover that, when $M(O_1(x, y), \dots, O_n(x, y)) = 0$, for a certain $i \in \{1, \dots, n\}$ it is clear that $O_i(x, y) = 0$. Because O_i is a POF, it means $x = 0$ or $y = 0$. On the contrary, if $x = 0$ or $y = 0$ then $O_i(x, y) = 0$ when taking arbitrary $i \in \{1, \dots, n\}$, as well as thus $O(x, y) = M(0, \dots, 0) = 0$. Analogously, another (O2') can be proven. \square

Corollary 1. Given an n -ary aggregation operation M on $[0, 1]$ satisfying continuity and $\min \leq M \leq \max$. Let O_1, \dots, O_n be POFs, then operation $O(x, y) = M(O_1(x, y), \dots, O_n(x, y))$ meets (O1') \sim (O4').

Proof. From $\min(x_1, \dots, x_n) \leq M(x_1, \dots, x_n)$, we get that $M(x_1, \dots, x_n) = 0$ implies $\min(x_1, \dots, x_n) = 0$, i.e., $x_j = 0$ for a certain $j = 1, \dots, n$. Thus, it satisfies the condition required in Proposition 2. Another condition is verified similarly making use of the non-equality $M \leq \max$. \square

Proposition 3. Given a POF O , as well as T being a positive continuous pseudo t -norm. Further, (1) the mapping $O_T: [0, 1]^2 \rightarrow [0, 1]$ formulated by $O_T(x, y) = O(x, y)T(x, y)$ meets $(O1') \sim (O4')$; (2) for arbitrary pseudo t -norm $T_1: [0, 1]^2 \rightarrow [0, 1]$ with continuity and no nontrivial zero factors, the mapping $O_{TT_1}: [0, 1]^2 \rightarrow [0, 1]$ given by $O_{TT_1}(x, y) = T_1(O(x, y), T(x, y))$ is a pseudo overlap function.

Proof. (1) Evidently, O_T is continuous and monotonous. Then $O_T(c, d) = 0$ iff $O(c, d)T(c, d) = 0$ iff $cd = 0$, $O_T(c, d) = 1$ iff $O(c, d) = 1$ and $T(c, d) = 1$ iff $cd = 1$ are established, so O_T is a POF;
(2) It is clear that O_{TT_1} is continuous and monotonous. Then $O_{TT_1}(c, d) = 0 \Leftrightarrow T_1(O(c, d), T(c, d)) = 0$ iff $O(c, d) = 0 \vee T(c, d) = 0 \Leftrightarrow cd = 0$, $O_{TT_1}(c, d) = 1 \Leftrightarrow T_1(O(c, d), T(c, d)) = 1 \Leftrightarrow O(c, d) = 1 \wedge T(c, d) = 1 \Leftrightarrow cd = 1$ are established, so O_{TT_1} is a pseudo overlap function. \square

Theorem 4. Given POFs $O_1, O_2, \dots, O_n: [0, 1]^2 \rightarrow [0, 1]$, w_1, w_2, \dots, w_n on $[0, 1]$ satisfying $\sum_{i=1}^n w_i = 1$, then operation O formulated by $O(x, y) = \sum_{i=1}^n w_i O_i(x, y)$ meets $(O1') \sim (O4')$.

Proof. Evidently, O is continuous and monotonous. Due to $O(x, y) = 0 \Leftrightarrow \sum_{i=1}^n w_i O_i(x, y) = 0 \Leftrightarrow \forall i = 1, 2, \dots, n, w_i O_i(x, y) = 0$, and $\sum_{i=1}^n w_i = 1$, so $\exists k \in \{1, 2, \dots, n\}$, s.t. $w_k \neq 0$, i.e., $O_k(x, y) = 0 \Leftrightarrow xy = 0$. Additionally, due to $O(x, y) = 1$ iff $\sum_{i=1}^n w_i O_i(x, y) = 1 = \sum_{i=1}^n w_i$, so $\sum_{i=1}^n w_i O_i(x, y) - \sum_{i=1}^n w_i = 0$, i.e., $\sum_{i=1}^n w_i [O_i(x, y) - 1] = 0$. Because $w_i [O_i(x, y) - 1] \leq 0$, $w_i [O_i(x, y) - 1] = 0$. Similarly, due to $\sum_{i=1}^n w_i = 1 \neq 0$, so $\exists k \in \{1, 2, \dots, n\}$, s.t. $w_k \neq 0$, then $O_k(x, y) - 1 = 0 \Leftrightarrow O_k(x, y) = 1 \Leftrightarrow xy = 1$. Therefore, O is a POF. \square

Theorem 5. Given a POF O on $[0, 1]$, as well as $T_1, T_2: [0, 1]^2 \rightarrow [0, 1]$ are pseudo t -norms without divisors of zero and satisfy continuity, then the mapping O given by $O(u, v) = O(T_1(u, v), T_2(u, v))$ is a pseudo overlap function.

Proof. One easily verifies that O is continuous and monotonous. Then $O(u, v) = 0$ iff $O(T_1(u, v), T_2(u, v)) = 0$ iff $T_1(u, v)T_2(u, v) = 0 \Leftrightarrow uv = 0$, and $O(u, v) = 1$ iff $O(T_1(u, v), T_2(u, v)) = 1 \Leftrightarrow T_1(u, v)T_2(u, v) = 1$ iff $T_1(u, v) = T_2(u, v) = 1 \Leftrightarrow uv = 1$. \square

In the existing literature, some scholars have shown that fuzzy conjunctions can induce residual implication, such as t -norms, t -conorms and overlap functions etc. Additionally, some conjunctions can also induce two residuated implications by removing commutativity, such as pseudo t -norms and pseudo t -conorms. Moreover, since the function can still induce fuzzy implication without commutativity, we can define two residuated implications which satisfy the residual property induced by pseudo overlap function, namely residuated implication $R_O^{(1)}, R_O^{(2)}$.

Definition 13. Let O be a POF on $[0, 1]$. Two bivariate mappings $R_O^{(1)}$ and $R_O^{(2)}$ on $[0, 1]$ are called left (right) residuated implications, when taking any $x, y \in [0, 1]$:

$$R_O^{(1)}(x, y) = \sup\{z \in [0, 1] | O(z, x) \leq y\}, \quad (7)$$

$$R_O^{(2)}(x, y) = \sup\{z \in [0, 1] | O(x, z) \leq y\}. \quad (8)$$

Obviously, $R_O^{(1)} = R_O^{(2)}$ if the considered pseudo overlap function is commutative. Binary functions $R_O^{(1)}$ and $R_O^{(2)}$ are called residuals associated with pseudo overlap function O on the first and second variables, respectively.

Theorem 6. Let O be a POF on $[0, 1]$; statements as follows are equivalent:

- (1) O is infinitely \vee -distributive in its first variable;
- (2) $O(x, y) \leq z$ when and only when $x \leq R_O^{(1)}(y, z)$ when taking arbitrary $x, y, z \in [0, 1]$; (RP1)
- (3) $O(R_O^{(1)}(x, y), x) \leq y$ for any $x, y \in [0, 1]$;
- (4) $R_O^{(1)}(x, y) = \max\{z \in [0, 1] | O(z, x) \leq y\}$ when taking any $x, y \in [0, 1]$.

Moreover, conditions as below are equivalent:

- (1') O is infinitely \vee -distributive in its second variable;
- (2') $O(x, y) \leq z$ iff $y \leq R_O^{(2)}(x, z)$ for arbitrary $0 < x, y, z < 1$; (RP2)
- (3') $O(x, R_O^{(2)}(x, y)) \leq y$ when taking any $x, y \in [0, 1]$;
- (4') $R_O^{(2)}(x, y) = \max\{z \in [0, 1] | O(x, z) \leq y\}$ when taking any $x, y \in [0, 1]$.

Proof. By Theorem 4.1 in [34] (or Theorem 3.1 in [35]), we get that the conditions (1) \sim (4) are equivalent. By Theorem 4.2 in [34], we get that the conditions (1') \sim (4') are equivalent. \square

Proposition 4. Given two pseudo overlap functions O_1, O_2 , let $R_{O_1}^{(1)}, R_{O_1}^{(2)}$ be the residuated implications induced by O_1 and $R_{O_2}^{(1)}, R_{O_2}^{(2)}$ be the residuated implications induced by O_2 . Then,

- (1) $O_1 \leq O_2$ if and only if $R_{O_2}^{(1)} \leq R_{O_1}^{(1)}$, if and only if $R_{O_2}^{(2)} \leq R_{O_1}^{(2)}$.
- (2) $O_1 = O_2$ if and only if $R_{O_2}^{(1)} = R_{O_1}^{(1)}$, if and only if $R_{O_2}^{(2)} = R_{O_1}^{(2)}$.

Proof. (1) By the definition of residuated implication, $R_{O_1}^{(1)} = \sup\{z | O_1(z, x) \leq y\}$, $R_{O_1}^{(2)} = \sup\{z | O_1(x, z) \leq y\}$, $R_{O_2}^{(1)} = \sup\{z | O_2(z, x) \leq y\}$, $R_{O_2}^{(2)} = \sup\{z | O_2(x, z) \leq y\}$. We hold that $O_1 \leq O_2$

$$\begin{aligned} &\Leftrightarrow (\forall x, y, z \in [0, 1], O_1(z, x) \leq y \text{ when } O_2(z, x) \leq y) \\ &\Leftrightarrow (\forall x, y \in [0, 1], \{z \in [0, 1] | O_2(z, x) \leq y\} \subseteq \{z \in [0, 1] | O_1(z, x) \leq y\}) \\ &\Leftrightarrow (\forall x, y \in [0, 1], \sup\{z \in [0, 1] | O_2(z, x) \leq y\} \leq \sup\{z \in [0, 1] | O_1(z, x) \leq y\}) \\ &\Leftrightarrow (R_{O_2}^{(1)} \leq R_{O_1}^{(1)}). \end{aligned}$$

Similarly, we have $O_1 \leq O_2 \Leftrightarrow R_{O_2}^{(2)} \leq R_{O_1}^{(2)}$.

(2) It follows from (1). \square

In the following table, we provide some examples of pseudo overlap functions and their residuated implications (see Table 1).

- In Table 1, ① is $\min\{\max\{1 - x, \frac{y - (1 - \alpha)(x - 1)}{\alpha x^2 + 1 - \alpha}\}, \frac{y}{\alpha x^2}\}$,
- ② is $\min\{\max\{1 - x, \frac{\alpha - 1 + \sqrt{(1 - \alpha)^2 + 4\alpha x[(1 - \alpha)(1 - x) + y]}}{2\alpha x}\}, \sqrt{\frac{y}{\alpha x}}\}$,
- ③ is $\frac{x + \alpha x - \alpha x^2}{3(\alpha x - \alpha x^2)} + \sqrt[3]{\sqrt{\gamma} + \delta} + \sqrt[3]{-\sqrt{\gamma} + \delta}$, where $\gamma = (\frac{y}{2(\alpha x - \alpha x^2)} - \frac{(x + \alpha x - \alpha x^2)^3}{27(\alpha x - \alpha x^2)^3})^2 - \frac{(x + \alpha x - \alpha x^2)^6}{729(\alpha x - \alpha x^2)^6}$ and $\delta = \frac{(x + \alpha x - \alpha x^2)^3}{27(\alpha x - \alpha x^2)^3} - \frac{y}{2(\alpha x - \alpha x^2)}$,
- ④ is $\frac{x + \alpha x + \sqrt{\alpha^2 x^4 - 2\alpha^2 x^3 + \alpha^2 x^2 - 2\alpha x^3 + 2\alpha x^2 + 4\alpha xy - 4\alpha y + x^2 - \alpha x^2}}{2(\alpha x - \alpha x^2)}$.

In the following, some properties that pseudo overlap functions and residuated implications $R_O^{(1)}, R_O^{(2)}$ satisfy are presented.

Proposition 5. Let $O: [0, 1]^2 \rightarrow [0, 1]$ be a POF. Then statements as below hold:

- (1) $R_O^{(1)}, R_O^{(2)}$ satisfy (NP) if and only if 1 is the neutral element of O ;
- (2) $R_O^{(2)}$ satisfies (EP) if and only if O is associative, i.e., $O(r, O(s, t)) = O(s, O(r, t))$, for arbitrary $r, s, t \in [0, 1]$;
- (3) $R_O^{(2)}$ satisfies (IP) when and only when O satisfies $O(x, 1) \leq x$, for arbitrary $x \in [0, 1]$;
- (4) $R_O^{(2)}$ satisfies (LOP) when and only when O satisfies $O(x, 1) \leq x$, for arbitrary $x \in [0, 1]$;
- (5) $R_O^{(2)}$ satisfies (ROP) when and only when O satisfies $O(m, 1) \geq m$, for arbitrary $m \in [0, 1]$;

- (6) $R_O^{(2)}$ satisfies (OP) when and only when O satisfies $O(x, 1) = x, \forall x \in [0, 1]$;
(7) $R_O^{(2)}$ satisfies (CB) if $O(p, q) \leq \min\{p, q\}$;
(8) $R_O^{(1)}$ satisfies (SIB) if and only if $R_O^{(1)}$ satisfies (CB), $R_O^{(2)}$ satisfies (SIB) if and only if $R_O^{(2)}$ satisfies (CB);
(9) $R_O^{(1)}, R_O^{(2)}$ satisfies (IB) if $O(x, y) = \min\{x, y\}$;
(10) $R_O^{(1)}, R_O^{(2)}$ satisfies (SBC), (LBC) and (RBC);
(11) $R_O^{(1)}, R_O^{(2)}$ satisfies (CB) if O has unit element 1.

Table 1. Examples of pseudo overlap functions and their residuated implications.

Pseudo Overlap Function		Residuated Implications
(1)	$O(x, y) = \min\{x^p, y^q\}, p, q > 0$	$R_O^{(1)}(x, y) = \begin{cases} 1, & x^q \leq y \\ \sqrt[q]{y}, & x^q > y \end{cases}, R_O^{(2)}(x, y) = \begin{cases} 1, & x^p \leq y \\ \sqrt[p]{y}, & x^p > y \end{cases}$
(2)	$O(x, y) = \frac{2x^p y^q}{1+x^p y^q}, p, q > 0$	$R_O^{(1)}(x, y) = \begin{cases} 1, & y \geq (2-y)x^q \\ \sqrt[p]{\frac{y}{(2-y)x^q}}, & \text{otherwise} \end{cases},$ $R_O^{(2)}(x, y) = \begin{cases} 1, & y \geq (2-y)x^p \\ \sqrt[p]{\frac{y}{(2-y)x^p}}, & \text{otherwise} \end{cases}$
(3)	$O(x, y) = x^p y^q, p, q > 0$	$R_O^{(1)}(x, y) = \begin{cases} 1, & x^q \leq y \\ \sqrt[q]{\frac{y}{x^q}}, & x^q > y \end{cases}, R_O^{(2)}(x, y) = \begin{cases} 1, & x^p \leq y \\ \sqrt[p]{\frac{y}{x^p}}, & x^p > y \end{cases}$
(4)	$O(x, y) = \begin{cases} \frac{(\alpha+\beta)xy}{\alpha x + \beta y}, & \alpha x + \beta y \neq 0 \\ 0, & \alpha x + \beta y = 0 \end{cases}$ $\alpha, \beta > 0$	$R_O^{(1)}(x, y) = \begin{cases} 1, & (\alpha + \beta)x - \alpha y \leq \beta xy \\ \frac{\beta xy}{(\alpha + \beta)x - \alpha y}, & \text{otherwise} \end{cases},$ $R_O^{(2)}(x, y) = \begin{cases} 1, & (\alpha + \beta)x - \beta y \leq \alpha xy \\ \frac{\alpha xy}{(\alpha + \beta)x - \beta y}, & \text{otherwise} \end{cases}$
(5)	$O(x, y) = \max\{\min\{x, \frac{y}{2}\}, x + y - 1\}$	$R_O^{(1)}(x, y) = \begin{cases} 1, & y \geq x \\ y - x + 1, & \frac{x}{2} \leq y < x \\ y, & y < \frac{x}{2} \end{cases},$ $R_O^{(2)}(x, y) = \begin{cases} 1, & y \geq x \\ \min\{2y, y - x + 1\}, & y < x \end{cases}$
(6)	$O(x, y) = \min\{x, y\} \max\{x^2, y\}$	$R_O^{(1)}(x, y) = \begin{cases} 1, & x \leq y \\ x, & x^2 \leq y < x \\ \frac{y}{x}, & y < x^2 \end{cases}, R_O^{(2)}(x, y) = \begin{cases} 1, & y \geq x \\ \frac{y}{x}, & x^2 \leq y < x \\ \min\{\frac{y}{x^2}, \sqrt{y}\}, & y < x^2 \end{cases}$
(7)	$O(x, y) = \begin{cases} \frac{xy}{\gamma x + (1-\gamma)y}, & \gamma x + (1-\gamma)y \neq 0 \\ 0, & \gamma x + (1-\gamma)y = 0 \end{cases}$ $0 < \gamma < 1$	$R_O^{(1)}(x, y) = \begin{cases} 1, & x \leq \gamma y + (1-\gamma)xy \\ \frac{(1-\gamma)xy}{x - \gamma y}, & \text{otherwise} \end{cases},$ $R_O^{(2)}(x, y) = \begin{cases} 1, & x \leq (1-\gamma)y + \gamma xy \\ \frac{\gamma xy}{x - (1-\gamma)y}, & \text{otherwise} \end{cases}$
(8)	$O(x, y) = \alpha xy^2 + (1-\alpha) \max\{0, x + y - 1\},$ $0 < \alpha < 1$	$R_O^{(1)}(x, y) = \begin{cases} 1, & \alpha x^2 + (1-\alpha)x \leq y \\ \textcircled{1}, & \text{otherwise} \end{cases}, R_O^{(2)}(x, y) = \begin{cases} 1, & x \leq y \\ \textcircled{2}, & x > y \end{cases}$
(9)	$O(x, y) = x^2 y + \alpha x^2 y(1-x)(1-y),$ $\alpha \in [-1, 1]$	$R_O^{(1)}(x, y) = \begin{cases} 1, & x \leq y \\ \textcircled{3}, & x > y \end{cases}, R_O^{(2)}(x, y) = \begin{cases} 1, & x^2 \leq y \\ \textcircled{4}, & x^2 > y \end{cases}$

Proof. (1) (\Rightarrow) Suppose for arbitrary $x \in [0, 1]$, $R_O^{(1)}(1, x) = \sup\{z | O(z, 1) \leq x\} = x$, so for a random $x \in [0, 1]$, one has $O(x, 1) \leq x$. If we take some x_0 in $[0, 1]$, one has $O(x_0, 1) < x_0$, then taking $z = O(x_0, 1)$. According to RP1, $z < x_0 \leq R_O^{(1)}(1, z)$, which is contradiction. Similarly, $R_O^{(2)}(1, x) = \sup\{z | O(1, z) \leq x\} = x \Rightarrow O(1, x) \leq x$. According to RP2, $O(1, x) = x$ when taking each $x \in [0, 1]$. So O has unit element 1.

(\Leftarrow) Suppose $O(x, 1) = x$ as well as $O(1, x) = x$, for arbitrary $x \in [0, 1]$. So $R_O^{(1)}(1, x) =$

- $\sup\{z|O(z,1) \leq x\} = \sup\{z|z \leq x\} = x$, $R_O^{(2)}(1,x) = \sup\{z|O(1,z) \leq x\} = \sup\{z|z \leq x\} = x$;
- (2) For all $r,s,t \in [0,1]$, suppose $O(r,O(s,t)) = O(s,O(r,t)) = a$. According to RP2, $O(r,O(s,t)) = a$ and $O(s,O(r,t)) = a \Leftrightarrow O(s,t) = R_O^{(2)}(r,a)$ and $O(r,t) = R_O^{(2)}(s,a) \Leftrightarrow t = R_O^{(2)}(s,R_O^{(2)}(r,a))$ and $t = R_O^{(2)}(r,R_O^{(2)}(s,a))$, i.e., $R_O^{(2)}(r,R_O^{(2)}(s,a)) = R_O^{(2)}(s,R_O^{(2)}(r,a))$;
- (3) When taking random $x \in [0,1]$, $R_O^{(2)}(x,x) = \sup\{z|O(x,z) \leq x\} = 1 \Leftrightarrow O(x,1) \leq x$;
- (4) (\Rightarrow) For an arbitrary $x \in [0,1]$, due to $x \leq x$, $R_O^{(2)}(x,x) = \sup\{z|O(x,z) \leq x\} = 1 \Rightarrow O(x,1) \leq x$.
- (\Leftarrow) If we take random $x,y \in [0,1]$, then because $O(x,z) \leq O(x,1) \leq x \leq y$ when taking each $z \in [0,1]$, $R_O^{(2)}(x,y) = \sup\{z|O(x,z) \leq y\} = 1$;
- (5) (\Rightarrow) For any $m \in [0,1]$, $R_O^{(2)}(m,O(m,1)) = \sup\{z \in [0,1]|O(m,z) \leq O(m,1)\} = 1$, so $m \leq O(m,1)$.
- (\Leftarrow) Assume $O(m,1) \geq m$, for arbitrary $m \in [0,1]$. $R_O^{(2)}(m,n) = \sup\{z \in [0,1]|O(m,z) \leq n\} = 1 \Rightarrow O(m,1) \leq n \Rightarrow m \leq O(m,1) \leq n$, i.e., $m \leq n$;
- (6) Obviously, it can be obtained from the above two certificates;
- (7) For arbitrary $p,q \in [0,1]$, $O(p,q) \leq \min\{p,q\} \Rightarrow O(p,q) \leq q \Rightarrow q \in \{z \in [0,1]|O(p,z) \leq q\} \Rightarrow q \leq \sup\{z \in [0,1]|O(p,z) \leq q\} = R_O^{(2)}(p,q)$;
- (8) When taking random $x,y \in [0,1]$, because O fits monotonic increment, we have $y \leq R_O^{(1)}(x,y) \Leftrightarrow y \leq \sup\{z \in [0,1]|O(z,x) \leq y\} \Leftrightarrow \{z \in [0,1]|O(z,x) \leq y\} \subseteq \{z \in [0,1]|O(z,x) \leq R_O^{(1)}(x,y)\} \Leftrightarrow \sup\{z \in [0,1]|O(z,x) \leq y\} \leq \sup\{z \in [0,1]|O(z,x) \leq R_O^{(1)}(x,y)\} \Leftrightarrow R_O^{(1)}(x,y) \leq R_O^{(1)}(x,R_O^{(1)}(x,y))$. Similarly, $y \leq R_O^{(2)}(x,y) \Leftrightarrow R_O^{(2)}(x,R_O^{(2)}(x,y)) \leq R_O^{(2)}(x,y)$;
- (9) When taking random $x,y \in [0,1]$, $R_O^{(1)}(x,y) = \sup\{z \in [0,1]| \min(z,x) \leq y\} = \begin{cases} 1, & x \leq y \\ y, & x > y \end{cases}$ when $O(x,y) = \min\{x,y\}$. Then, whenever $x \leq y$, $R_O^{(1)}(x,R_O^{(1)}(x,y)) = R_O^{(1)}(x,1) = 1 = R_O^{(1)}(x,y)$. On the other side, $R_O^{(1)}(x,R_O^{(1)}(x,y)) = R_O^{(1)}(x,y)$ when $x > y$. Similarly, $R_O^{(2)}(x,R_O^{(2)}(x,y)) = R_O^{(2)}(x,y)$;
- (10) When taking random $x \in [0,1]$, suppose $x \neq 0$, $R_O^{(1)}(x,0) = \sup\{z \in [0,1]|O(z,x) \leq 0\}$, since $O(z,x) \geq 0$, then $O(z,x) = 0 \Rightarrow z = 0$, i.e., $R_O^{(1)}(x,0) = 0$. When taking a random $x,y \in [0,1]$, obviously, $R_O^{(1)}(0,y) = \sup\{z \in [0,1]|O(z,0) \leq y\} = 1$, $R_O^{(1)}(x,1) = \sup\{z \in [0,1]|O(z,x) \leq 1\} = 1$. Similarly, $R_O^{(2)}(x,0) = 0$, $R_O^{(2)}(0,y) = 1$ and $R_O^{(2)}(x,1) = 1$;
- (11) Suppose O has 1 as a neutral element, i.e., for any $y \in [0,1]$, $O(1,y) = y = O(y,1)$. Since O is increasing, $O(y,x) \leq O(y,1) = y$ and $O(x,y) \leq O(1,y) = y$. According to RP1 and RP2, we have that $y \leq R_O^{(1)}(x,y), y \leq R_O^{(2)}(x,y)$. \square

Remark 1. Since the two functions $R_O^{(1)}$ and $R_O^{(2)}$ are induced by the pseudo overlap function without commutativity, the above proposition can also expand some properties, such as:

- (2') $R_O^{(1)}$ satisfies (EP) when and only when O satisfies $O(O(x,y),z) = O(O(x,z),y)$;
- (3') $R_O^{(1)}$ satisfies (IP) when and only when O satisfies $O(1,x) \leq x, \forall x \in [0,1]$;
- (4') $R_O^{(1)}$ satisfies (LOP) when and only when O satisfies $O(1,x) \leq x, \forall x \in [0,1]$;
- (5') $R_O^{(1)}$ satisfies (ROP) when and only when O satisfies $O(1,x) \geq x, \forall x \in [0,1]$;
- (6') $R_O^{(1)}$ satisfies (OP) when and only when O satisfies $O(1,x) = x, \forall x \in [0,1]$.

The proof is similar to the above proposition.

4. Pseudo Grouping Functions and Their Residuated Co-Implications

In this section, we show notions of pseudo grouping functions as well as discuss residuated co-implications (the notion of co-implication was first proposed by B. De Baets

and J. Fodor in [29]; it is also called deresiduum, see [35]). In addition, we also provide the general construction method of pseudo grouping functions. Finally, we provide some detailed examples.

Definition 14. The binary mapping $G: [0, 1]^2 \rightarrow [0, 1]$ is known as a pseudo grouping function (briefly PGF) when G meets the requirements as below:

- (G1') The value of G is 0 when and only when values of two elements are 0;
- (G2') The value of G is 1 when and only when at least one of x and y has a value of 1;
- (G3') G meets monotonic increment;
- (G4') G meets continuity.

Observe that a PGF can be obtained by duality from a pseudo overlap function. We have the following basic result.

Proposition 6. Let O be a pseudo overlap function and N a continuous negation such that $N(x) = 0$ when and only when $x = 1$ and $N(x) = 1$ when and only when $x = 0$. Then operation $G(x, y) = N(O(N(x), N(y)))$ is a pseudo grouping function. In particular, G is a PGF when and only when $O(x, y) = 1 - G(1 - x, 1 - y)$ meets (O1') \sim (O4').

Proof. Continuity as well as monotonicity are straightforward. Moreover, $G(x, y) = 0$ iff $N(O(N(x), N(y))) = 0$ iff $O(N(x), N(y)) = 1$ iff $N(x)N(y) = 1$ iff $x = y = 0$. Additionally, the other property is analogous. In particular, if we consider the negation $N(x) = 1 - x$, we have the result as follows: G is a PGF when and only when $O(x, y) = 1 - G(1 - x, 1 - y)$ fits (O1') \sim (O4'). \square

Now, we provide some examples of PGFs.

Example 4. (1) The operator $G: [0, 1]^2 \rightarrow [0, 1]$ defined, when taking random $x, y \in [0, 1]$, as

$$G(x, y) = 1 - (1 - x)(1 - y)^2 \quad (9)$$

is a PGF;

(2) An operation $G: [0, 1]^2 \rightarrow [0, 1]$ formulated, when taking arbitrary $x, y \in [0, 1]$, as

$$G(x, y) = 1 - \min\{(1 - x)^p, (1 - y)^q\}, p, q > 0 \quad (10)$$

is a PGF.

Definition 15. An n -dimension ($n \geq 2$) operator G on $[0, 1]$ is known as an n -dimension pseudo grouping function when the following properties are established:

- (1) $G(x_1, \dots, x_n) = 0$ when and only when all values of x_i are 0 ($i = 1, \dots, n$);
- (2) $G(x_1, \dots, x_n) = 1$ when and only when there are some i ($i = 1, \dots, n$) so that $x_i = 1$;
- (3) G meets monotonic increment;
- (4) G meets continuity.

Then we will provide some examples of n -dimensional PGFs.

Example 5. (1) The operator $G: [0, 1]^n \rightarrow [0, 1]$ formulated, when taking random $x_1, x_2, \dots, x_n \in [0, 1]$, as

$$G(x, y) = 1 - \prod_{i=1}^{n-1} (1 - x_i) \cdot (1 - x_n)^2 \quad (11)$$

is an n -dimensional PGF.

(2) An operation $G: [0, 1]^n \rightarrow [0, 1]$ formulated, when taking random $x_1, x_2, \dots, x_n \in [0, 1]$, as

$$G(x, y) = 1 - \min\{1 - x_1, 1 - x_2, \dots, 1 - x_{n-1}, \sqrt{1 - x_n}\} \quad (12)$$

is an n -dimensional PGF.

Proposition 7. Let G_1, G_2 be two operations satisfying $(G1') \sim (G4')$. Then, $\max(G_1, G_2)(x, y) = \max(G_1(x, y), G_2(x, y))$ as well as $\min(G_1, G_2)(x, y) = \min(G_1(x, y), G_2(x, y))$ meet $(G1') \sim (G4')$.

Proposition 8. Let $\varphi_1, \varphi_2, \varphi_3: [0, 1] \rightarrow [0, 1]$ be monotonous mappings satisfying continuity and $\varphi_i(x) = 0 \Leftrightarrow x = 0, \varphi_i(x) = 1 \Leftrightarrow x = 1$ ($i = 1, 2, 3$). Let $G: [0, 1]^2 \rightarrow [0, 1]$ be a PGF. Then mapping $G^{\varphi_1, \varphi_2, \varphi_3}$, defined as

$$G^{\varphi_1, \varphi_2, \varphi_3}(x, y) = \varphi_1(G(\varphi_2(x), \varphi_3(y))),$$

also meets $(G1') \sim (G4')$.

Proposition 9. Let G_1, \dots, G_n be PGFs as well as F on $[0, 1]$ be an aggregate operator satisfying continuity, $F(x_1, \dots, x_n) = 0$ only if $x_i = 0$ for a few $i = 1, \dots, n$ as well as $F(x_1, \dots, x_n) = 1$ only if $x_i = 1$ for a few $i = 1, \dots, n$. Then operation $G(c, d) = F(G_1(c, d), \dots, G_n(c, d))$ meets $(G1') \sim (G4')$.

Proposition 10. Given a continuous aggregate operation $F: [0, 1]^n \rightarrow [0, 1]$, and it meets $\min(i_1, \dots, i_n) \leq F(i_1, \dots, i_n) \leq \max(i_1, \dots, i_n)$. Let G_1, \dots, G_n be PGFs, then operation $G(c, d) = F(G_1(c, d), \dots, G_n(c, d))$ meets $(G1') \sim (G4')$.

Corollary 2. Given two pseudo grouping functions G_1, G_2 , then function G defined as $G(x, y) = \alpha G_1(x, y) + (1 - \alpha)G_2(x, y)$ is also a pseudo grouping function, where $\alpha \in [0, 1]$.

Proposition 11. Let G be a pseudo grouping function, and S a continuous pseudo t -conorm without divisors of zero. Then

- (1) The bivariate mapping G_S on $[0, 1]$ formulated as $G_S(e, f) = G(e, f)S(e, f)$ is a PGF;
- (2) When taking arbitrary positive pseudo t -conorm S' on $[0, 1]$ satisfying continuity, the bivariate mapping $G_{SS'}$ on $[0, 1]$ formulated as $G_{SS'}(e, f) = S'(G(e, f), S(e, f))$ is a PGF.

Theorem 7. Let $G_1, G_2, \dots, G_n: [0, 1]^2 \rightarrow [0, 1]$ be pseudo grouping functions, and $w_1, w_2, \dots, w_n \in [0, 1]$ satisfy $\sum_{i=1}^n w_i = 1$; then the mapping G given by $G(x, y) = \sum_{i=1}^n w_i G_i(x, y)$ is a PGF.

Theorem 8. Given a pseudo grouping function G on $[0, 1]$, and $S_1, S_2: [0, 1]^2 \rightarrow [0, 1]$ are continuous pseudo t -conorms, as well as $(\forall x, y \in [0, 1], i = 1, 2) S_i(x, y) = 1$ implying $x = 1$ or $y = 1$. Then the function G , formulated as $G(x, y) = G(S_1(x, y), S_2(x, y))$, meets $(G1') \sim (G4')$.

The proofs of the Theorem 7 and Theorem 8 are consistent with those of Theorem 4 and Theorem 5, respectively, which are related to the pseudo overlap function. Similarly, we obtain the residuated co-implications induced from PGFs.

Definition 16. Given a PGF G on $[0, 1]$. The following $R_G^{(1)}$ and $R_G^{(2)}: [0, 1]^2 \rightarrow [0, 1]$ are called two residuated implications, when taking random $x, y \in [0, 1]$:

$$R_G^{(1)}(x, y) = \inf\{z \in [0, 1] | G(z, x) \geq y\}, \quad (13)$$

$$R_G^{(2)}(x, y) = \inf\{z \in [0, 1] | G(x, z) \geq y\}. \quad (14)$$

Obviously, $R_G^{(1)} = R_G^{(2)}$ when the pseudo grouping function G is commutative. The functions $R_G^{(1)}$ and $R_G^{(2)}$ are called residuated co-implications associated with the pseudo grouping function G on the first and second variables, respectively.

Theorem 9. Let G be a PGF on $[0, 1]$, statements as follows are equivalent:

- (1) G is infinitely \wedge -distributive in its first variable;
- (2) $G(x, y) \geq z$ when and only when $x \geq R_G^{(1)}(y, z)$ for arbitrary $x, y, z \in [0, 1]$;
- (3) $G(R_G^{(1)}(x, y), x) \geq y$ for any $x, y \in [0, 1]$;
- (4) $R_G^{(1)}(x, y) = \min\{z \in [0, 1] | G(z, x) \geq y\}$ when taking any $x, y \in [0, 1]$.

Similarly, the following statements are equivalent:

- (1') G is infinitely \wedge -distributive in its second variable;
- (2') $G(x, y) \geq z$ iff $y \geq R_G^{(2)}(x, z)$ when taking arbitrary $x, y, z \in [0, 1]$;
- (3') $G(x, R_G^{(2)}(x, y)) \geq y$ for arbitrary $x, y \in [0, 1]$;
- (4') $R_G^{(2)}(x, y) = \min\{z \in [0, 1] | G(x, z) \geq y\}$ for any $x, y \in [0, 1]$.

Proof. By Theorem 4.3 in [34] (or Theorem 3.4 in [35]), we get that the conditions (1) ~ (4) are equivalent. In the same way, we get that the conditions (1') ~ (4') are equivalent. \square

Proposition 12. Let G_1, G_2 be two PGFs, $R_{G_1}^{(1)}, R_{G_1}^{(2)}$ be residuated co-implications induced by G_1 and $R_{G_2}^{(1)}, R_{G_2}^{(2)}$ are residuated co-implications induced by G_2 . Then

- (1) $G_1 \leq G_2$ if and only if $R_{G_2}^{(1)} \leq R_{G_1}^{(1)}$, $G_1 \leq G_2$ if and only if $R_{G_2}^{(2)} \leq R_{G_1}^{(2)}$.
- (2) $G_1 = G_2$ if and only if $R_{G_2}^{(1)} = R_{G_1}^{(1)}$ if and only if $R_{G_2}^{(2)} = R_{G_1}^{(2)}$.

The concrete examples of pseudo grouping functions and residuated co-implications corresponding to them are as follows (see Table 2).

Table 2. Examples of pseudo grouping functions and their residuated co-implications.

Pseudo Grouping Function		Residuated Co-Implications	
(1)	$G(x, y) = 1 - \min\{(1-x)^p, (1-y)^q\},$ $p, q > 0$	$R_G^{(1)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[q]{1-y} \\ 1 - \sqrt[p]{1-y}, & \text{otherwise} \end{cases}$	$R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[p]{1-y} \\ 1 - \sqrt[q]{1-y}, & \text{otherwise} \end{cases}$
(2)	$G(x, y) = 1 - \frac{2(1-x)^p(1-y)^q}{1+(1-x)^p(1-y)^q},$ $p, q > 0$	$R_G^{(1)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[q]{\frac{1-y}{1+y}} \\ 1 - \sqrt[p]{\frac{1-y}{(1-x)^q(1+y)}}, & \text{otherwise} \end{cases}$	$R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[p]{\frac{1-y}{1+y}} \\ 1 - \sqrt[q]{\frac{1-y}{(1-x)^p(1+y)}}, & \text{otherwise} \end{cases}$
(3)	$G(x, y) = 1 - (1-x)^p(1-y)^q, p, q > 0$	$R_G^{(1)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[q]{1-y} \\ 1 - \sqrt[p]{\frac{1-y}{(1-x)^q}}, & \text{otherwise} \end{cases}$	$R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt[p]{1-y} \\ 1 - \sqrt[q]{\frac{1-y}{(1-x)^p}}, & \text{otherwise} \end{cases}$
(4)	$G(x, y) = \begin{cases} 1, & \alpha(1-x) + \beta(1-y) = 0 \\ 1 - \frac{(\alpha+\beta)(1-x)(1-y)}{\alpha(1-x) + \beta(1-y)}, & \text{otherwise} \end{cases}$ $\alpha, \beta > 0$	$R_G^{(1)}(x, y) = \begin{cases} 0, & [(\alpha+\beta) - \beta(1-y)](1-x) \leq \alpha(1-y) \\ 1 - \frac{\beta(1-x)(1-y)}{(\alpha+\beta)(1-x) - \alpha(1-y)}, & \text{otherwise} \end{cases}$	$R_G^{(2)}(x, y) = \begin{cases} 0, & [(\alpha+\beta) - \alpha(1-y)](1-x) \leq \beta(1-y) \\ 1 - \frac{\alpha(1-x)(1-y)}{(\alpha+\beta)(1-x) - \beta(1-y)}, & \text{otherwise} \end{cases}$
(5)	$G(x, y) = 1 - \max\{\min\{1-x, \frac{1-y}{2}\}, 1-x-y\}$	$R_G^{(1)}(x, y) = \begin{cases} 0, & y \leq x \\ \frac{1-x}{2}, & x < y \leq \frac{x+1}{2} \\ y, & y > \frac{x+1}{2} \end{cases}$	$R_G^{(2)}(x, y) = \begin{cases} 0, & y \leq \max\{x, \frac{1}{2}\} \\ 2y-1, & y > \frac{1}{2} \\ y-x, & y > x \end{cases}$

Table 2. Cont.

Pseudo Grouping Function	Residuated Co-Implications
(6) $G(x, y) = 1 - \min\{1 - x, 1 - y\}$ $\max\{(1 - x)^2, 1 - y\}$	$R_G^{(1)}(x, y) = \begin{cases} 0, & x \geq y \\ \max\{\frac{y-x}{1-x}, 1 - \sqrt{1-y}\}, & x < y \end{cases}$ $R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq y \\ 1 - \frac{1-y}{1-x}, & 1 - \sqrt{1-y} \leq x < y \\ 1 - \frac{1-y}{(1-x)^2}, & x < 1 - \sqrt{1-y} \end{cases}$
(7) $G(x, y) = \begin{cases} 1, \alpha(1-x) + (1-\alpha)(1-y) = 0 \\ 1 - \frac{(1-x)(1-y)}{\alpha(1-x) + (1-\alpha)(1-y)}, \text{ otherwise} \\ \alpha \in (0, 1) \end{cases}$	$R_G^{(1)}(x, y) = \begin{cases} 0, x + \alpha(1-y) + (1-\alpha)(1-x)(1-y) \geq 1 \\ 1 - \frac{(1-\alpha)(1-x)(1-y)}{(1-x) - \alpha(1-y)}, \text{ otherwise} \end{cases}$ $R_G^{(2)}(x, y) = \begin{cases} 0, x + \alpha(1-x)(1-y) + (1-\alpha)(1-y) \geq 1 \\ 1 - \frac{\alpha(1-x)(1-y)}{(1-x) - (1-\alpha)(1-y)}, \text{ otherwise} \end{cases}$
(8) $G(x, y) = 1 - \alpha(1-x)(1-y)^2 - (1-\alpha)$ $\max\{0, 1 - x - y\}, 0 < \alpha < 1$	$R_G^{(1)}(x, y) = \begin{cases} 0, & 1 - \alpha(1-x)^2 - (1-\alpha)(1-x) \geq y \\ \textcircled{5}, & \text{ otherwise} \end{cases}$ $R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq y \\ \textcircled{6}, & x < y \end{cases}$
(9) $G(x, y) = 1 - (1-x)^2(1-y) - \alpha xy$ $(1-x)^2(1-y), \alpha \in [-1, 1]$	$R_G^{(1)}(x, y) = \begin{cases} 0, & x \geq y \\ \textcircled{7}, & x < y \end{cases}$ $R_G^{(2)}(x, y) = \begin{cases} 0, & x \geq 1 - \sqrt{1-y} \\ \textcircled{8}, & x < 1 - \sqrt{1-y} \end{cases}$

In Table 2, $\textcircled{5}$ is $\max\{\min\{1 - x, 1 - \frac{1-y+(1-\alpha)x}{\alpha(1-x)^2+1-\alpha}\}, 1 - \frac{1-y}{\alpha(1-x)^2}\}$,
 $\textcircled{6}$ is $\max\{\min\{1 - x, 1 - \frac{\alpha-1+\sqrt{(1-\alpha)^2+4\alpha(1-x)[1-y+(1-\alpha)x]}}{2\alpha(1-x)}\}, 1 - \sqrt{\frac{1-y}{\alpha(1-x)}}\}$,
 $\textcircled{7}$ is $\frac{x+2\alpha x-2\alpha x^2-1}{3(\alpha x-\alpha x^2)} + \sqrt[3]{p+q} + \sqrt[3]{p-q}$, where p is
 $\frac{y-x}{2(\alpha x-\alpha x^2)} - \frac{(x+2\alpha x-2\alpha x^2-1)^3}{27(\alpha x-\alpha x^2)^3} + \frac{(2x+\alpha x-\alpha x^2-2)(x+2\alpha x-2\alpha x^2-1)}{6(\alpha x-\alpha x^2)^2}$ and q is
 $\sqrt{\left(\frac{2x+\alpha x-\alpha x^2-2}{3(\alpha x-\alpha x^2)} - \frac{(x+2\alpha x-2\alpha x^2-1)^2}{9(\alpha x-\alpha x^2)^2}\right)^3 + p^2}$,
 $\textcircled{8}$ is $\frac{\alpha x^3+2x+\alpha x-x^2-2\alpha x^2-1+\sqrt{v}}{2(\alpha x^3+\alpha x-2\alpha x^2)}$, where v is $(2\alpha x^2 - \alpha x^3 - 2x + x^2 - \alpha x + 1)^2 - 4(\alpha x^3 + \alpha x - 2\alpha x^2)(2x - x^2 - y)$.

5. Some Applications of Pseudo Overlap Functions

In this section, we reveal some applications of POFs (PGFs) in MAGDM, fuzzy morphology and image processing.

5.1. Applications of Pseudo Overlap Functions in Multi-Attribute Group Decision Making

5.1.1. Pseudo Overlap Functions Generated by n-Dimensional Overlap Functions

Overlap functions have been used in multiple-attribute decision making, but because they are commutative, they cannot be directly applied to the decision problems in which the importance degree of different attributes is different. At the same time, in group decision-making, if the importance of different decision experts is different, the overlap function cannot be directly used to aggregate their decision information.

Interestingly, we can construct some pseudo overlap functions from an n-dimensional overlap function and a set of weights (importance degree), thus applying pseudo overlap functions to solving multi-attribute (group) decision problems with preferences. According to reference [13], we get the theorem as below.

Theorem 10. Let O be an n -ary overlap function as well as $w = (w_1, \dots, w_n)^T$ be a weight vector satisfying for every $j = 1, \dots, n$, $w_j \neq 0$ and $\sum_{j=1}^n w_j = 1$. If O is strict, i.e., it satisfies

$O(y, x_2, \dots, x_n) > O(x_1, x_2, \dots, x_n)$ if $\prod_{j=1}^n x_j > 0$ and $y > x_1$, then the function O_w defined by $O_w(x_1, \dots, x_n) = \frac{O(w_1 x_1, \dots, w_n x_n)}{O(w_1, \dots, w_n)}$ is a POF.

Proof. It is a corollary from Theorem 3.1 in [13]. \square

Example 6. Let $w_1 = (0.3, 0.3, 0.4)$. The following $O_i (i = 1, 2, 3, 4, 5)$ are pseudo overlap functions obtained based on the different three-dimensional overlap functions with weight w_1 .

(1) Given an overlap function $O(x_1, x_2, x_3) = 0$ (when $x_1 + x_2 + x_3 = 0$) and $\frac{3x_1 x_2 x_3}{x_1 + x_2 + x_3}$ (otherwise), it is strict, so the function $O_1(x_1, x_2, x_3) = 0$ (when $0.3x_1 + 0.3x_2 + 0.4x_3 = 0$) and $\frac{x_1 x_2 x_3}{0.3x_1 + 0.3x_2 + 0.4x_3}$ (otherwise) is a POF by Theorem 10;

(2) Given a strict overlap function $O(x_1, x_2, x_3) = (x_1 x_2 x_3)^{\frac{x_1 + x_2 + x_3}{3}}$, the function $O_2(x_1, x_2, x_3) = x_1 x_2 x_3 (0.3x_1 + 0.3x_2 + 0.4x_3)$ is a POF by Theorem 10;

(3) Given a strict overlap function $O(x_1, x_2, x_3) = 0$ (when $x_1 x_2 x_3 = 0$) and $\frac{3x_1 x_2 x_3}{x_1 x_2 + x_2 x_3 + x_1 x_3}$ (otherwise), the function $O_3(x_1, x_2, x_3) = 0$ (when $x_1 = x_2 = x_3 = 0$) and $\frac{0.33x_1 x_2 x_3}{0.12x_2 x_3 + 0.12x_1 x_3 + 0.09x_1 x_2}$ (otherwise) is a POF by Theorem 10;

(4) Given an overlap function $O(x_1, x_2, x_3) = \prod_{i=1}^3 x_{(i)}^{2i}$, in which $x_{(1)}, x_{(2)}, x_{(3)}$ is the substitution of (x_1, x_2, x_3) satisfying $x_{(1)} \geq x_{(2)} \geq x_{(3)}$, O is strict, so the function,

$$O_4(x_1, x_2, x_3) = \frac{x_{(1')}^2 x_{(2')}^4 x_{(3')}^6}{0.4^2 0.3^4 0.3^6},$$

is a POF by Theorem 10, where $x_{(1')}, x_{(2')}, x_{(3')}$ is a permutation of $(0.3x_1, 0.3x_2, 0.4x_3)$ satisfying $x_{(1')} \geq x_{(2')} \geq x_{(3')}$;

(5) Given an overlap function $O(x_1, x_2, x_3) = \prod_{i=1}^3 x_{(i)}^{\frac{1}{2i}}$, in which $x_{(1)}, x_{(2)}, x_{(3)}$ is a substitution of (x_1, x_2, x_3) satisfying $x_{(1)} \leq x_{(2)} \leq x_{(3)}$, O is strict, so function

$$O_5(x_1, x_2, x_3) = \frac{x_{(1')}^{\frac{1}{2}} x_{(2')}^{\frac{1}{4}} x_{(3')}^{\frac{1}{6}}}{0.3^{\frac{1}{2}} 0.3^{\frac{1}{4}} 0.4^{\frac{1}{6}}}$$

is a POF by Theorem 10, where $x_{(1')}, x_{(2')}, x_{(3')}$ is the permutation of $(0.3x_1, 0.3x_2, 0.4x_3)$ such that $x_{(1')} \leq x_{(2')} \leq x_{(3')}$.

Example 7. Let $w_2 = (0.1, 0.1, 0.2, 0.2, 0.2, 0.2)$. The following $O_i (i = 6, 7, 8, 9, 10)$ are pseudo overlap functions obtained based on the different six-dimensional overlap functions with weight w_2 .

(1) Given a strict overlap function $O(x_1, x_2, x_3, x_4, x_5, x_6) = \frac{6x_1 x_2 x_3 x_4 x_5 x_6}{x_1 + x_2 + x_3 + x_4 + x_5 + x_6}$ (when $\sum_{i=1}^6 x_i \neq 0$)

and 0 (when $\sum_{i=1}^6 x_i = 0$), then the function

$$O_6(x_1, x_2, x_3, x_4, x_5, x_6) = \begin{cases} 0, & \text{if } 0.1x_1 + 0.1x_2 + 0.2x_3 + 0.2x_4 + 0.2x_5 + 0.2x_6 = 0 \\ \sqrt{\frac{x_1 x_2 x_3 x_4 x_5 x_6}{0.1x_1 + 0.1x_2 + 0.2x_3 + 0.2x_4 + 0.2x_5 + 0.2x_6}}, & \text{otherwise} \end{cases}$$

is also a POF by Theorem 10;

(2) Given a strict overlap function $O(x_1, x_2, x_3, x_4, x_5, x_6) = x_1 x_2 x_3 x_4 x_5 x_6^{\frac{x_1 + x_2 + x_3 + x_4 + x_5 + x_6}{6}}$, then the function

$$O_7(x_1, x_2, x_3, x_4, x_5, x_6) = x_1 x_2 x_3 x_4 x_5 x_6 (0.1x_1 + 0.1x_2 + 0.2x_3 + 0.2x_4 + 0.2x_5 + 0.2x_6)$$

is a POF by Theorem 10;

(3) Given a strict overlap function $O(x_1, x_2, x_3, x_4, x_5, x_6) = \frac{6}{\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} + \frac{1}{x_4} + \frac{1}{x_5} + \frac{1}{x_6}}$ ($x_i \neq 0, i =$

$1, \dots, 6$) and 0 (when $\prod_{i=1}^6 x_i = 0$), then function $O_8(x_1, x_2, x_3, x_4, x_5, x_6) = \frac{8}{\frac{2}{x_1} + \frac{2}{x_2} + \frac{1}{x_3} + \frac{1}{x_4} + \frac{1}{x_5} + \frac{1}{x_6}}$

(when $\prod_{i=1}^6 x_i \neq 0$) and 0 (when $\prod_{i=1}^6 x_i = 0$), is a POF by Theorem 10;

(4) Given a strict overlap function $O(x_1, x_2, x_3, x_4, x_5, x_6) = \prod_{i=1}^6 x_{(i)}^{2i}$, $x_{(1)}, \dots, x_{(6)}$ is the permutation of $(x_1, x_2, x_3, x_4, x_5, x_6)$ such that $x_{(1)} \geq x_{(2)} \geq x_{(3)} \geq x_{(4)} \geq x_{(5)} \geq x_{(6)}$, then the function

$$O_9(x_1, x_2, x_3, x_4, x_5, x_6) = \frac{x_{(1')}^2 x_{(2')}^4 x_{(3')}^6 x_{(4')}^8 x_{(5')}^{10} x_{(6')}^{12}}{0.2^2 0.2^4 0.2^6 0.2^8 0.1^{10} 0.1^{12}}$$

is a POF by Theorem 10, in which $x_{(1')}, \dots, x_{(6')}$ is the substitution of $(0.1x_1, 0.1x_2, 0.2x_3, 0.2x_4, 0.2x_5, 0.2x_6)$ such that $x_{(1')} \geq x_{(2')} \geq x_{(3')} \geq x_{(4')} \geq x_{(5')} \geq x_{(6')}$;

(5) Given a strict overlap function $O(x_1, x_2, x_3, x_4, x_5, x_6) = \prod_{i=1}^6 x_{(i)}^{\frac{1}{2i}}$, where $x_{(1)}, \dots, x_{(6)}$ is the permutation of $(x_1, x_2, x_3, x_4, x_5, x_6)$ such that $x_{(1)} \leq x_{(2)} \leq x_{(3)} \leq x_{(4)} \leq x_{(5)} \leq x_{(6)}$, then the function

$$O_{10}(x_1, x_2, x_3, x_4, x_5, x_6) = \frac{x_{(1')}^{\frac{1}{2}} x_{(2')}^{\frac{1}{4}} x_{(3')}^{\frac{1}{6}} x_{(4')}^{\frac{1}{8}} x_{(5')}^{\frac{1}{10}} x_{(6')}^{\frac{1}{12}}}{0.1^{\frac{1}{2}} 0.1^{\frac{1}{4}} 0.2^{\frac{1}{6}} 0.2^{\frac{1}{8}} 0.2^{\frac{1}{10}} 0.2^{\frac{1}{12}}}$$

is a POF by Theorem 10, in which $x_{(1')}, \dots, x_{(6')}$ is the substitution of $(0.1x_1, 0.1x_2, 0.2x_3, 0.2x_4, 0.2x_5, 0.2x_6)$ such that $x_{(1')} \leq x_{(2')} \leq x_{(3')} \leq x_{(4')} \leq x_{(5')} \leq x_{(6')}$.

5.1.2. MAGDM on Account of Pseudo Overlap Functions

A solution for an MAGDM matter is a measure to select a relatively better scheme from a list of them while regarding some attributes of the alternatives as well as the view of a panel of experts. In such problems, a limited set $X = \{x_1, \dots, x_n\}$ including practicable schemes, $U = \{u_1, \dots, u_m\}$ including attributes as well as a set $d = \{d_1, \dots, d_t\}$ containing policy makers are given. The measure must select a better scheme that meets all attributes. Moreover, every policy maker d_k determines matrix $S^{(k)} = (s_{ij}^{(k)})_{n \times m}$, in which each line stands for a scheme and each rank represents an attribute. In classical decision problems, according to the view of the policy makers d_k , if an alternative x_i has the attribute u_j , the value of the position $s_{ij}^{(k)}$ of $S^{(k)}$ is 1, otherwise it is 0. For some fuzzy attributes, the value of the position $s_{ij}^{(k)}$ is the grade of membership, that is, a value in $[0, 1]$ standing for the degree to which the scheme x_i belongs to a fuzzy set related to the attribute u_j . Additionally, all attributes are classified into two classes—benefit and cost—and we use I to represent the index set of the benefit attributes.

Similar to the method in [13], we present a solution for the problem as below:

(1) Standardize the decision matrix $S^{(k)}$ to a normal decision matrix $N^{(k)} = (n_{ij}^{(k)})_{n \times m}$,

$$\text{where } n_{ij}^{(k)} = \begin{cases} s_{ij}^{(k)}, & j \in I \\ 1 - s_{ij}^{(k)}, & j \notin I \end{cases}$$

(2) Given a t-dimensional POF O , the standard decision matrices are aggregated into the overall decision matrix with O as follows: $c_{ij} = O(n_{ij}^{(1)}, \dots, n_{ij}^{(t)})$;

(3) For every scheme x_i , in order to calculate the group totality preference value gtp_i , we aggregate its membership degrees to each attribute by using an m-dimensional pseudo overlap function O' as follows: $gtp_i = O'(c_{i1}, \dots, c_{im})$;

(4) Array all schemes in descending order according to group totality preference values and select the top scheme.

5.1.3. Living Example

The living example in [13] is regarded to demonstrate our approach.

Assume an investor plans to invest a portion of capital in one corporation. Finally, he narrows the range to six corporations: x_1 stands for the chemical corporation, x_2 stands for the food corporation, x_3 stands for the computer corporation, x_4 stands for the car

corporation, x_5 stands for the furniture corporation, x_6 stands for the pharmaceutical corporation. The investor is assisted by three experts (e_1 , e_2 and e_3). The experts establish some attributes utilized to estimate investments.

Three beneficial attributes as below:

(u_1) short period interests; (u_2) mid-term interests; (u_3) long-term interests;

The lossy attributes as below:

(u_4) investment hazard; (u_5) investment trouble; (u_6) other investment disadvantages.

The decision matrices of the three experts are as follows, see Tables 3–5.

Table 3. Appraisal of expert e_1 .

$S^{(1)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.7	0.8	0.6	0.7	0.5	0.9
x_2	0.8	0.6	0.9	0.7	0.6	0.7
x_3	0.5	0.4	0.8	0.3	0.8	0.8
x_4	0.6	0.7	0.6	0.7	0.8	0.6
x_5	0.9	0.8	0.4	0.7	0.7	0.8
x_6	0.8	0.3	0.7	0.7	0.6	0.7

Table 4. Appraisal of expert e_2 .

$S^{(2)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.6	0.8	0.5	0.6	0.4	0.8
x_2	0.7	0.6	0.8	0.6	0.7	0.7
x_3	0.7	0.6	0.8	0.7	0.8	0.8
x_4	0.6	0.7	0.5	0.6	0.8	0.7
x_5	0.7	0.8	0.7	0.7	0.6	0.8
x_6	0.6	0.4	0.8	0.7	0.6	0.7

Table 5. Appraisal of expert e_3 .

$S^{(3)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.7	0.6	0.6	0.6	0.4	0.7
x_2	0.7	0.6	0.7	0.6	0.6	0.7
x_3	0.6	0.5	0.8	0.5	0.8	0.8
x_4	0.6	0.7	0.7	0.5	0.8	0.6
x_5	0.7	0.8	0.6	0.7	0.6	0.8
x_6	0.4	0.5	0.9	0.7	0.6	0.6

These decision matrices are normalized respectively to obtain three normal decision matrices $N^{(1)}$, $N^{(2)}$ and $N^{(3)}$, which are described as below, see Tables 6–8.

Table 6. Normalized decision matrix of expert e_1 .

$N^{(1)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.7	0.8	0.6	0.3	0.5	0.1
x_2	0.8	0.6	0.9	0.3	0.4	0.3
x_3	0.5	0.4	0.8	0.7	0.2	0.2
x_4	0.6	0.7	0.6	0.3	0.2	0.4
x_5	0.9	0.8	0.4	0.3	0.3	0.2
x_6	0.8	0.3	0.7	0.3	0.4	0.3

Table 7. Normalized decision matrix of expert e_2 .

$N^{(2)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.6	0.8	0.5	0.4	0.6	0.2
x_2	0.7	0.6	0.8	0.4	0.3	0.3
x_3	0.7	0.6	0.8	0.3	0.2	0.2
x_4	0.6	0.7	0.5	0.4	0.2	0.3
x_5	0.7	0.8	0.7	0.3	0.4	0.2
x_6	0.6	0.4	0.8	0.3	0.4	0.3

Table 8. Normalized decision matrix of expert e_3 .

$N^{(3)}$	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.7	0.6	0.6	0.4	0.6	0.3
x_2	0.7	0.6	0.7	0.4	0.4	0.3
x_3	0.6	0.5	0.8	0.5	0.2	0.2
x_4	0.6	0.7	0.7	0.5	0.2	0.4
x_5	0.7	0.8	0.6	0.3	0.4	0.2
x_6	0.4	0.5	0.9	0.3	0.4	0.4

Next, because three experts with the weights $w = (0.3, 0.3, 0.4)$, we aggregate the standard decision matrices of them by using three-dimensional pseudo overlap function O_1 in Example 6 to obtain the overall decision matrix, as shown in Table 9 below.

Table 9. Overall decision matrix.

C	u_1	u_2	u_3	u_4	u_5	u_6
x_1	0.439	0.533	0.316	0.130	0.316	0.029
x_2	0.537	0.360	0.638	0.130	0.130	0.090
x_3	0.350	0.240	0.640	0.210	0.040	0.040
x_4	0.360	0.490	0.344	0.146	0.040	0.130
x_5	0.580	0.640	0.295	0.090	0.130	0.040
x_6	0.331	0.146	0.622	0.090	0.160	0.106

Then we obtain the group totality preference vector GTP by considering O_6 in Example 7, in which $w = (0.1, 0.1, 0.2, 0.2, 0.2, 0.2)$. The outcome is shown in Table 10.

Table 10. Group totality preference value.

GTP	x_1	x_2	x_3	x_4	x_5	x_6
gtp_i	0.0184	0.0255	0.0086	0.0146	0.0148	0.0137

On account of Table 10, we get the array of the schemes as below:

$$x_2 \succ x_1 \succ x_5 \succ x_4 \succ x_6 \succ x_3.$$

Of course, we can get other rankings by using other different pseudo overlap functions in Examples 6 and 7. Table 11 contains eleven rankings obtained by the pseudo overlap functions and the result of four methods in [13] for this matter.

Table 11. Summary of the permutations gained with presented means as well as other methods.

Three-Dimensional Functions	Six-Dimensional Functions	Ranking
	Maximum	$x_2 \succ x_1 \succ x_6 \succ x_5 \succ x_4 \succ x_3$
	Minimum	$x_2 \succ x_1 \succ x_6 \succ x_4 = x_5 \succ x_3$
	WHD	$x_5 \succ x_3 \succ x_2 \succ x_4 \succ x_6 \succ x_1$
	AOWAD	$x_5 \succ x_2 \succ x_4 \succ x_3 \succ x_6 \succ x_1$
O_1	O_6	$x_2 \succ x_1 \succ x_5 \succ x_4 \succ x_6 \succ x_3$
O_1	O_7	$x_2 \succ x_1 \succ x_5 \succ x_6 \succ x_4 \succ x_3$
O_1	O_8	$x_2 \succ x_6 \succ x_4 \succ x_5 \succ x_1 \succ x_3$
O_2	O_6	$x_2 \succ x_1 \succ x_4 \succ x_5 \succ x_6 \succ x_3$
O_2	O_7	$x_2 \succ x_1 \succ x_5 \succ x_6 \succ x_4 \succ x_3$
O_2	O_8	$x_2 \succ x_6 \succ x_4 \succ x_5 \succ x_1 \succ x_3$
O_3	O_6	$x_2 \succ x_1 \succ x_5 \succ x_4 \succ x_6 \succ x_3$
O_3	O_7	$x_2 \succ x_1 \succ x_5 \succ x_6 \succ x_4 \succ x_3$
O_3	O_8	$x_2 \succ x_5 \succ x_4 \succ x_1 \succ x_6 \succ x_3$
O_4	O_9	$x_2 \succ x_6 \succ x_1 \succ x_4 \succ x_5 \succ x_3$
O_5	O_{10}	$x_2 \succ x_5 \succ x_4 \succ x_1 \succ x_6 \succ x_3$

The above table shows that fifteen methods return ten different rankings.

In addition, when there are no exact weights, other non-commutative pseudo overlap functions (which cannot be directly generated by the method in Theorem 10) can also

be selected. Non-commutative functions represent the different importance of different attributes or experts, but the importance degrees are not expressed by specific numbers, which are hidden in the functions (in this case, we can think that the weights can vary according to the attribute value). For the example above, we can use the following non-commutative pseudo overlap functions:

$$\begin{aligned}
 O_{11}(x_1, x_2, x_3) &= x_1^{\frac{1}{6}} x_2^{\frac{1}{4}} x_3^{\frac{1}{2}} \\
 O_{12}(x_1, x_2, x_3) &= \begin{cases} 0 & \text{if } x_1 = x_2 = x_3 = 0 \\ \frac{6x_1x_2x_3}{3x_1+2x_2+x_3} & \text{otherwise} \end{cases} \\
 O_{13}(x_1, x_2, x_3) &= \frac{2x_1^{\frac{1}{4}}x_2^{\frac{1}{3}}x_3^{\frac{1}{2}}}{1+x_1^{\frac{1}{4}}x_2^{\frac{1}{3}}x_3^{\frac{1}{2}}} \\
 O_{14}(x_1, x_2, x_3, x_4, x_5, x_6) &= x_1^{\frac{1}{12}}x_2^{\frac{1}{10}}x_3^{\frac{1}{8}}x_4^{\frac{1}{6}}x_5^{\frac{1}{4}}x_6^{\frac{1}{2}} \\
 O_{15}(x_1, x_2, x_3, x_4, x_5, x_6) &= \begin{cases} 0, \text{ if } x_i = 0 (i = 1, 2, \dots, 6) \\ \frac{10x_1x_2x_3x_4x_5x_6}{4x_1+2x_2+x_3+x_4+x_5+x_6}, & \text{otherwise} \end{cases} \\
 O_{16}(x_1, x_2, x_3, x_4, x_5, x_6) &= \frac{2x_1^{\frac{1}{7}}x_2^{\frac{1}{6}}x_3^{\frac{1}{5}}x_4^{\frac{1}{4}}x_5^{\frac{1}{3}}x_6^{\frac{1}{2}}}{1+x_1^{\frac{1}{7}}x_2^{\frac{1}{6}}x_3^{\frac{1}{5}}x_4^{\frac{1}{4}}x_5^{\frac{1}{3}}x_6^{\frac{1}{2}}}
 \end{aligned}$$

The results are shown in Table 12 below.

Analyzing Table 12, it is obvious that the nine schemes revert to four various arrangements. Among them, most (seven) think the best alternative is x_2 and nine think that x_3 is the worst alternative. Moreover, rankings in Tables 11 and 12 can be fused and further analyzed by referring to the method in [36], which is omitted here.

Through the above comparative analysis, we show that the pseudo overlap functions can not only aggregate multiple information, but also indicate the importance of different information (including the importance of attributes and the importance of experts). They are more flexible than the overlap functions, and convenient for decision makers for choosing good alternatives.

Table 12. Summary of the permutations gained with suggested means.

Three-Dimensional Functions	Six-Dimensional Functions	Ranking
O_{11}	O_{14}	$x_6 \succ x_2 \succ x_4 \succ x_1 \succ x_5 \succ x_3$
O_{11}	O_{15}	$x_2 \succ x_1 \succ x_6 \succ x_4 \succ x_5 \succ x_3$
O_{11}	O_{16}	$x_2 \succ x_6 \succ x_1 \succ x_4 \succ x_5 \succ x_3$
O_{12}	O_{14}	$x_2 \succ x_6 \succ x_4 \succ x_1 \succ x_5 \succ x_3$
O_{12}	O_{15}	$x_2 \succ x_1 \succ x_6 \succ x_4 \succ x_5 \succ x_3$
O_{12}	O_{16}	$x_2 \succ x_6 \succ x_1 \succ x_4 \succ x_5 \succ x_3$
O_{13}	O_{14}	$x_6 \succ x_2 \succ x_4 \succ x_1 \succ x_5 \succ x_3$
O_{13}	O_{15}	$x_2 \succ x_1 \succ x_6 \succ x_4 \succ x_5 \succ x_3$
O_{13}	O_{16}	$x_2 \succ x_6 \succ x_1 \succ x_4 \succ x_5 \succ x_3$

5.2. Application of Pseudo Overlap Functions in Fuzzy Mathematical Morphology

The method of analyzing object shape in images using different neighborhood transformations based on set theory is known as mathematical morphology. At present, there is a mathematical morphology for binary images and a fuzzy mathematical morphology for gray images. Some authors improve fuzzy mathematical morphology by quoting t-norms and fuzzy implications when defining the most basic fuzzy dilation and fuzzy erosion operations (see [37]). We will introduce pseudo overlap functions and pseudo

grouping functions into fuzzy mathematical morphology, and use the new operators in fuzzy mathematical morphology to extract the edge of gray images.

5.2.1. Fuzzy Mathematical Morphology Based on Pseudo Overlap Functions

A, B denote fuzzy sets on the referential S , and the membership function values corresponding to x are $A(x)$ as well as $B(x)$, where A represents fuzzy set of the gray image to be operated, and B represents the fuzzy set of a smaller structural element. In fuzzy mathematical morphology, the fuzzy expansion operation $D(A, B)$ as well as the corrosion operation $E(A, B)$ of gray image A by structural element B are as follows:

$$D(A, B)(y) = \sup_{x \in \{x | x \in B, x+y \in A\}} \{ \min[A(y+x), B(x)] \} \forall y \in A$$

$$E(A, B)(y) = \inf_{x \in \{x | x \in B, x+y \in A\}} \{ \max[A(y+x), B(x)] \} \forall y \in A.$$

When a pseudo overlap function is applied to fuzzy mathematical morphology, the pseudo overlap function is used to replace the intersection operation in fuzzy dilation, and its corresponding pseudo grouping function is used to replace the union operation in fuzzy erosion. At this time, the gray image A is subjected to the fuzzy expansion operation $D(A, B)$ and the fuzzy corrosion operation $E(A, B)$ performed by the structural element B , which are defined as follows, respectively:

$$D(A, B)(y) = \sup_{x \in \{x | x \in B, x+y \in A\}} \{ O[A(y+x), B(x)] \} \quad (15)$$

$$E(A, B)(y) = \inf_{x \in \{x | x \in B, x+y \in A\}} \{ G[A(y+x), B(x)] \} \quad (16)$$

5.2.2. Experimental Framework

In the original fuzzy mathematical morphology, after the image is blurred, t-norms and fuzzy implications are often used to expand and erode the image, and then de-blur it. However, we use pseudo overlap functions and pseudo grouping functions as fuzzy expansion and erosion operators, convex combine their results, and then defuzzify it. The specific steps are as follows.

(1) Fuzzy grayscale image

The grayscale image shown in Figure 3 is selected as the image to be processed, which is a single channel image with a resolution of 1200 * 675.



Figure 3. Grayscale image.

Set M represents the set of all points $x = (m, n)$ in this image, and N represents the pixel value of each point in set M , at this time, there is a surjection $f: M \rightarrow N$. Where M is a point set with 1200 * 675 points, $m = 1, 2, \dots, 1200, n = 1, 2, \dots, 675, N \subseteq Z$, and $\forall x \in M, f(x) \subseteq [0, 255]$.

Note that function $g: N \rightarrow A$ is the membership function of pixel value a in B :

$$g(a) = \frac{a - X_{\min}}{X_{\max} - X_{\min}} \quad a \in B,$$

where X_{min} and X_{max} are the maximum and minimum values in set N respectively; then the mapping $gf: M \rightarrow A$ is the function $A(x)$ of the membership of each point x in the gray image M .

(2) Fuzzy dilation and fuzzy erosion operation

After obtaining the fuzzy set A of the gray image, the structure operators of fuzzy dilation and fuzzy erosion are B_D and B_E respectively.

$$B_D = \{0.9/(-1, -1), 0.9/(0, -1), 0.9/(1, -1), 0.9/(-1, 0), 1.0/(0, 0), 0.9/(1, 0), 0.9/(-1, 1), 0.9/(0, 1), 0.9/(1, 1)\};$$

$$B_E = \{0.9/(-1, -1), 0.1/(0, -1), 0.1/(1, -1), 0.1/(-1, 0), 0.2/(0, 0), 0.1/(1, 0), 0.1/(-1, 1), 0.1/(0, 1), 0.1/(1, 1)\}.$$

Fuzzy set A is fuzzily dilated by structural operator B_D through formula (15) to obtain fuzzy set $D(A, B_D)$, and is fuzzily eroded by structural operator B_E through formula (16) to obtain fuzzy set $E(A, B_E)$. The POF O as well as its corresponding pseudo grouping function G selected in the formula are $O(a, b) = a^2b$, $G(a, b) = 1 - (1 - a)^2(1 - b)$.

(3) Defuzzification convex combination result

Firstly, the results of dilation and erosion are obtained through the convex combination formula shown below to obtain the fuzzy set C .

$$C(y) = 0.9D(A, B_D)(y) + 0.1E(A, B_E)(y) \quad \forall y \in A.$$

Then, the fuzzy set C is defuzzified to obtain the number set M' .

$$M'(y) = \lfloor y * 255 \rfloor \quad \forall y \in C, \quad (17)$$

where $\lfloor \cdot \rfloor$ is rounding down.

After the fuzzy expansion and fuzzy erosion results of the original intensity of the image are convex combined, the convex combination results are de-blurred to obtain M' , as the grayscale image shown in Figure 4.



Figure 4. Convex combination result after defuzzification.

On the other hand, when the de-blurring operation shown in formula (17) is directly performed on $D(A, B_D)$ and $E(A, B_E)$, the results of dilation and erosion of the original image are shown in Figure 5 and Figure 6, respectively.



Figure 5. Result of dilation of the original image.



Figure 6. Result of erosion of the original image.

5.2.3. Analysis of Experimental Results

In order to present the actual application effect, we used the edge extraction operation in mathematical morphology to extract the edges of objects in the original gray image M and the gray image M' processed by fuzzy mathematical morphology based on pseudo overlap functions. The principle of edge extraction of a binary image by mathematical morphology is to dilate the foreground of the binary image, and then make a difference between the dilation result and the original image to obtain the edge of the binary image.

Firstly, binarization is performed on the gray image M and M' , respectively, to obtain the binary images M_b , M'_b , as shown in Figures 7 and 8 below.




Figure 7. Binarization result of gray image M .




Figure 8. Binarization result of gray image M' .

We take the binary structure operator B_b as follows:

$$B_b = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} \quad (18)$$

Then we perform edge extraction operations on binary images M_b and M'_b , respectively: $\text{Edge1} = D(M'_b) - M'_b$, $\text{Edge2} = D(M_b) - M_b$.

The results of edge extraction for the original gray image and the gray image processed by fuzzy mathematical morphology with pseudo overlap functions are revealed in Figure 9 and Figure 10 respectively.




Figure 9. Result of edge extraction for the original gray image.




Figure 10. Result of edge extraction for the gray image processed by fuzzy mathematical morphology.

According to Figures 9 and 10, the edge graph obtained by our method (Figure 10) has fewer unorganized points, that is, the gray image processed by fuzzy mathematical morphology with pseudo overlap functions can achieve better image edges through edge extraction in mathematical morphology. This is because the dilation and erosion operations of fuzzy mathematical morphology with pseudo overlap function expand the contrast of foreground and background in the gray image from two aspects: reducing the background gray value and increasing the foreground gray value, which can be seen from Figures 5 and 6. Moreover, after introducing convex combination, the advantages of dilation and erosion are combined to further expand the contrast of foreground and background in the gray image. This allows us to extract the foreground with a more accurate threshold in the binarization operation, thus reducing noise similar to that in Figure 9.

5.3. Application of Pseudo Grouping Functions in Image Processing

In image segmentation, one of the most commonly used methods is threshold segmentation. In threshold processing, different targets of a picture are described by the gray level of every pixel. This technology mainly looks for a threshold t , so that the pixels with an intensity less than or equal to t are classified as the backdrop of the picture, while pixels with an intensity greater than t are classified as targets, and the converse is also true ([38]). Aranzazu Jurio et al. proposed a threshold algorithm based on the convex combination of the fuzzy method and grouping function in reference [12]. Our work is to introduce the pseudo grouping function to calculate the grouping value. Finally, we use the improved threshold algorithm to perform threshold segmentation on a gray image and compare it with Otus's algorithm, one of the most commonly used thresholding methods.

In this section, we show the performance of thresholding segmentation with the convex combination of pseudo grouping functions and the convex combination of grouping functions on 10 T1 weighted MRI images (see Figure 11). The MR cerebrum information as well as hand-operated segmentations originated from the morphometric analysis center of Massachusetts General Hospital. We evaluated the quality of excision by comparing with the ideal manually segmented images obtained on the same web page (see Figure 12). The purpose of this image segmentation was to divide each pixel in the cerebrum into two kinds—gray matter and white matter. In fact, it is a part of brain region volume analysis, which plays a great role in assessing the development of illness, for instance, Alzheimer's disease, epilepsy or schizophrenia ([39,40]).




Figure 11. Original images.




Figure 12. Ideal manual segmentation image.

To improve the effect of thenconvex combination thresholding results of grouping functions on image segmentation, we consider using pseudo grouping functions instead of grouping functions. Because theoretically a grouping function is a special case of a pseudo grouping function, pseudo grouping functions give a wider function selection range and have a greater chance of finding a better function, which may improve the accuracy of thresholding image segmentation. We confirm this below.

The specific content of the experiment is that we randomly selected the grouping functions in advance and segmented the 10 images with their convex combination, which achieved a certain segmentation effect. Then, the convex combination of pseudo grouping functions was also used for segmentation to compare with it. Finally, we illustrate the superiority of the proposed method by the percentage of good classified pixels.

In the thresholding process of Aranzazu jurio et al., a fuzzy set $\mu_{Q_{B_t}}(q)$ related to the background and a fuzzy set $\mu_{Q_{O_t}}(q)$ related to the image target are constructed by using the strict equivalence function for a fixed gray level $t(t = 0, 1, \dots, L - 1)$, for each gray level $q(q = 0, 1, \dots, L - 1)$, $h(q)$ is expressed as quantity of pixels with pixel intensity q .

When calculating the maximum grouping value using the convex combination of grouping functions $G_{G_{comb}}(x, y)$, select an appropriate threshold for each image:

$$t^* = \arg \max_t \sum_{q=0}^{L-1} G_{G_{comb}}(\mu_{Q_{B_t}}(q), \mu_{Q_{O_t}}(q)) \cdot h(q), \quad (19)$$

when the convex combination of pseudo grouping functions $PG_{G_{comb}}(x, y)$ is introduced for the thresholding operation, the threshold t^* related to the maximum grouping value is taken as the best threshold. The formula of the improved algorithm is:

$$t^* = \arg \max_t \sum_{q=0}^{L-1} PG_{G_{comb}}(\mu_{Q_{B_t}}(q), \mu_{Q_{O_t}}(q)) \cdot h(q). \quad (20)$$

Then, five grouping functions are used as follows:

$$\begin{aligned} G_1(x, y) &= x + y - xy \\ G_2(x, y) &= \frac{2 - (1-x)^2(1-y) - (1-x)(1-y)^2}{2} \\ G_3(x, y) &= \sqrt{x^2 + y^2 - x^2y^2} \\ G_4(x, y) &= 1 - (1-x)(1-y) \end{aligned}$$

$$G_5(x, y) = 1 - \frac{2(1-x)(1-y)}{1+(1-x)(1-y)}.$$

The five pseudo grouping functions are selected as below:

$$PG_1(x, y) = G_1(x, y)$$

$$PG_2(x, y) = G_2(x, y)$$

$$PG_3(x, y) = G_3(x, y)$$

$$PG_4(x, y) = \begin{cases} 1, & 1.1(1-x) + (1-y) = 0 \\ 1 - \frac{(1.1+1)(1-x)(1-y)}{1.1(1-x)+(1-y)} & \text{otherwise} \end{cases}$$

$$PG_5(x, y) = G_5(x, y).$$

Through the convex combination of the above grouping functions and pseudo grouping functions, the results of threshold segmentations of 10 graphs are as follows (see Figures 13 and 14).




Figure 13. Obtained segmentations by the convex association of grouping functions.




Figure 14. Obtained segmentations from the convex association of pseudo grouping functions.

We show the experimental results through specific data. In Table 13, we reveal the threshold obtained from every picture as well as the percentage of good sorted picture elements. The second column presents the outcomes gained from the convex association of grouping functions, and the third column presents the outcomes gained from the convex association of PGFs. As we can see from the experiment, our method has higher accuracy, that is, the use of convex association of PGFs significantly improves the segmentation accuracy of each image.

Table 13. Thresholds and percentages obtained from convex association of grouping functions and convex association of pseudo grouping functions.

Image	Convex Combination of Grouping Functions		Convex Combination of Pseudo Grouping Functions	
	th	%	th	%
1	185	95.88	195	98.53
2	179	96.73	195	99.04
3	163	94.26	178	99.27
4	183	96.05	192	97.86
5	176	97.77	187	98.80
6	173	96.90	181	98.33
7	181	95.37	193	97.87
8	182	95.65	195	98.87
9	181	95.48	199	99.08
10	162	95.15	177	99.13

Then, in contrast with Otsu's algorithm ([41]), the thresholding method on the basis of pseudo grouping functions has greater advantages. In Table 14, we show the thresholds obtained by the two algorithms and the percentage of pixels with good classification. We can find that our method obtains the better results for each picture in the test case, hence it mends Otsu's thresholding technique with regard to the group of pictures.

Table 14. Thresholds and percentages obtained from presented algorithm as well as Otsu's algorithm.

Image	Convex Combination of Pseudo Grouping Functions		Otsu's	
	th	%	th	%
1	195	98.53	185	95.88
2	195	99.04	179	96.73
3	178	99.27	160	94.26
4	192	97.86	184	96.25
5	187	98.80	171	96.72
6	181	98.33	171	96.42
7	193	97.87	184	96.05
8	195	98.87	182	95.65
9	199	99.08	182	95.76
10	177	99.13	159	95.15

6. Conclusions

Starting from many practical application fields, we expand the theory of OFs as well as grouping functions, introduce new concepts of POFs and PGFs for the first time, and investigate their basic properties, construction methods, and induced residuated implication operators and residuated co-implication operators. Moreover, we discuss some applications of pseudo overlap (grouping) functions in multiple attributes group decision-making, fuzzy mathematical morphology and image processing. The experimental results show that: (1) pseudo overlap functions can aggregate the decision information of attributes or experts with different importance; (2) pseudo overlap functions and their convex combination can be used to enrich the operators in fuzzy mathematical morphology, and expand the contrast between foreground and background in gray images, so as to reduce the noise in edge extraction; (3) pseudo grouping functions can be used to improve the accuracy of thresholding image segmentation in image processing. Therefore, POFs and PGFs have a wide adaptability and application value. There are still many theoretical and applied problems to be studied in terms of pseudo overlap functions and fuzzy logic (see [42–47]).

Author Contributions: Writing—original draft preparation, R.L., M.L. and Q.O.; writing—review and editing, X.Z., H.B., B.B. and J.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by National Natural Science Foundation of China (No.12271319) and research project No.PID2019-108392GB-I00 (AEI/10.13039/501100011033). The Major Program of the National Social Science Foundation of China under Grant No.20&ZD047.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Grabisch, M.; Marichal, J.-L.; Mesiar, R.; Pap, E. Aggregation functions. In *Encyclopedia of Mathematics and Its Applications*; Cambridge University Press: Cambridge, UK, 2009; pp. 56–130.
- Rudas, I.J.; Pap, E.; Fodor, J. Information aggregation in intelligent systems: An application oriented approach. *Knowl.-Based Syst.* **2013**, *38*, 3–13. [[CrossRef](#)]
- Bustince, H.; Fernandez, J.; Mesiar, R.; Montero, J.; Orduna, R. Overlap functions. *Nonlinear Anal.* **2010**, *72*, 1488–1499. [[CrossRef](#)]
- Dimuro, G.P.; Bedregal, B. On residual implications derived from overlap functions. *Inf. Sci.* **2015**, *312*, 78–88. [[CrossRef](#)]

5. Qiao, J. On binary relations induced from overlap and grouping functions. *Int. J. Approx. Reason.* **2019**, *106*, 155–171. [\[CrossRef\]](#)
6. Dai, S. Comparison of overlap and grouping functions. *Axioms* **2022**, *11*, 420. [\[CrossRef\]](#)
7. Qiao, J. On (IO, O)-fuzzy rough sets based on overlap functions. *Int. J. Approx. Reason.* **2021**, *132*, 26–48. [\[CrossRef\]](#)
8. Cao, M.; Hu, B. On the ordinal sum of fuzzy implications: New results and the distributivity over a class of overlap and grouping functions. *Fuzzy Sets Syst.* **2022**, *446*, 93–123. [\[CrossRef\]](#)
9. Qiao, J. Overlap and grouping functions on complete lattices. *Inf. Sci.* **2021**, *542*, 406–424. [\[CrossRef\]](#)
10. Qiao, J.; Hu, B. On homogeneous, quasi-homogeneous and pseudo-homogeneous overlap and grouping functions. *Fuzzy Sets Syst.* **2019**, *357*, 58–90. [\[CrossRef\]](#)
11. Jurio, A.; Bustince, H.; Pagola, M.; Pradera, A.; Yager, R.R. Some properties of overlap and grouping functions and their application to image thresholding. *Fuzzy Sets Syst.* **2013**, *229*, 69–90. [\[CrossRef\]](#)
12. Jurio, A.; Paternain, D.; Pagola, M.; Bustince, H. Image thresholding by grouping functions: Application to MRI images. In *Recent Developments and New Directions in Soft Computing*; Springer: Cham, Switzerland, 2014; pp. 195–208.
13. Da Silva, I.A.; Bedregal, B.; Bustince, H. Weighted average operators generated by n-dimensional overlaps and an application in decision. In Proceedings of the 2015 Conference of the International Fuzzy Systems Association and the European Society for Fuzzy Logic and Technology, Gijon, Spain, 30 June–3 July 2015.
14. Qiao, J.; Gong, Z. On \tilde{r} -(quasi)-overlap functions. *IEEE Trans. Fuzzy Syst.* **2020**, *29*, 3178–3185. [\[CrossRef\]](#)
15. Lucca, G.; Antonio Sanz, J.; Pereira Dimuro, G.; Bedregal, B.; Jose Asiain, M.; Elkano, M.; Bustince, H. CC-integrals: Choquet-like copula-based aggregation functions and its application in fuzzy rule-based classification systems. *Knowl.-Based Syst.* **2017**, *119*, 32–43. [\[CrossRef\]](#)
16. Flondor, P.; Georgescu, G.; Iorgulescu, A. Pseudo-t-norms and pseudo-BL algebras. *Soft Comput.* **2001**, *5*, 355–371. [\[CrossRef\]](#)
17. Georgescu, G.; Popescu, A. Non-commutative fuzzy structures and pairs of weak negations. *Fuzzy Sets Syst.* **2003**, *143*, 129–155. [\[CrossRef\]](#)
18. Liu, H. Two classes of pseudo-triangular norms and fuzzy implications. *Comput. Math. Appl.* **2011**, *61*, 783–789. [\[CrossRef\]](#)
19. Bloch, I.; Maitre, H. Fuzzy mathematical morphologies: A comparative study. *Pattern Recognit.* **1995**, *28*, 1341–1387. [\[CrossRef\]](#)
20. De Baets, B. A fuzzy morphology: A logical approach. In *Uncertainty Analysis in Engineering and Sciences*; Springer: Berlin/Heidelberg, Germany, 1998; pp. 53–67.
21. Bloch, I. Duality vs. adjunction for fuzzy mathematical morphology and general form of fuzzy erosions and dilations. *Fuzzy Sets Syst.* **2009**, *160*, 1858–1867. [\[CrossRef\]](#)
22. Nachtgael, M.; Sussner, P.; Melange, T.; Kerre, E.E. On the role of complete lattices in mathematical morphology: From tool to uncertainty model. *Inf. Sci.* **2011**, *181*, 1971–1988. [\[CrossRef\]](#)
23. Gonzalez-Hidalgo, M.; Massanet, S.; Mir, A.; Ruiz-Aguilera, D. On the choice of the pair conjunction-implication into the fuzzy morphological edge detector. *IEEE Trans. Fuzzy Syst.* **2015**, *23*, 872–884. [\[CrossRef\]](#)
24. Roerdink, J.B.T.M. Mathematical morphology with noncommutative symmetry groups. *Math. Morphol. Image Process.* **1993**, *34*, 205–254.
25. Madrid, N.; Ojeda-Aciego, M.; Medina, J.; Perfilieva, I. L-fuzzy relational mathematical morphology based on adjoint triples. *Inf. Sci.* **2019**, *474*, 75–89. [\[CrossRef\]](#)
26. Gottwald, S. *Fuzzy Sets and Logic Fuzzy*; Friedrich, Vieweg & Sohn Verlag: Wiesbaden, Germany, 1993.
27. Mesiar, R.; Kolesarova, A. Copulas and fuzzy implications. *Int. J. Approx. Reason.* **2020**, *117*, 52–59. [\[CrossRef\]](#)
28. Ward, M.; Dilworth, R.P. Residuated lattices. *Trans. Am. Math. Soc.* **1939**, *45*, 335–354. [\[CrossRef\]](#)
29. De Baets, B.; Fodor, J. Residual operators of uninorms. *Soft Comput.* **1999**, *3*, 89–100. [\[CrossRef\]](#)
30. Ruiz, D.; Torrens, J. Residual implications and co-implications from idempotent uninorms. *Kybernetika* **2004**, *40*, 21–38.
31. Hartonas, C. Duality results for (co)residuated lattices. *Log. Universalis* **2019**, *13*, 77–99. [\[CrossRef\]](#)
32. Mesiarova, A. A note on two open problems of Alsina, Frank and Schweizer. *Aequationes Math.* **2006**, *72*, 41–46. [\[CrossRef\]](#)
33. Ouyang, Y. A note on metrics induced by copulas. *Fuzzy Sets Syst.* **2012**, *191*, 122–125. [\[CrossRef\]](#)
34. Sun, F.; Wang, X.; Qu, X.; Shu, Q.; Zhang, X. Residual operations of monotone binary operations over complete lattices. *Int. J. Approx. Reason.* **2019**, *110*, 127–144. [\[CrossRef\]](#)
35. Su, Y.; Wang, Z. Deresiduums of implications on a complete lattice. *Inf. Sci.* **2015**, *325*, 504–520.
36. Bustince, H.; Bedregal, B.; Campion, M.J.; da Silva, I.A.; Fernandez, J.; Indurain, E.; Raventos-Pujol, A.; Santiago, R.H.N. Aggregation of individual rankings through fusion functions: Criticism and optimality analysis. *IEEE Trans. Fuzzy Syst.* **2022**, *30*, 638–648. [\[CrossRef\]](#)
37. Nachtgael, M.; Kerre, E.E. Classical and fuzzy approaches towards mathematical morphology. In *Fuzzy Techniques in Image Processing*; Springer: Berlin/Heidelberg, Germany, 2000; pp. 1–57.
38. Bustince, H.; Barrenechea, E.; Pagola, M. Image thresholding using restricted equivalence functions and maximizing the measure of similarity. *Fuzzy Sets Syst.* **2007**, *158*, 496–516. [\[CrossRef\]](#)
39. Caldaïrou, B.; Passat, N.; Habas, P.; Studholme, C.; Rousseau, F. A non-local fuzzy segmentation method: Application to brain MRI. *Pattern Recognit.* **2011**, *44*, 1916–1927. [\[CrossRef\]](#)
40. Mayer, A.; Greenspan, H. An adaptive mean-shift framework for MRI brain segmentation. *IEEE Trans. Med. Imaging* **2009**, *28*, 105–118. [\[CrossRef\]](#) [\[PubMed\]](#)
41. Otsu, N. A threshold selection method from gray level histograms. *IEEE Trans. Syst. Man Cybern.* **2007**, *9*, 62–66. [\[CrossRef\]](#)

-
42. Liang, R.; Zhang, X. Interval-valued pseudo overlap functions and application. *Axioms* **2022**, *11*, 216. [[CrossRef](#)]
 43. Zhang, X.; Liang, R.; Bedregal, B. Weak inflationary BL-algebras and filters of inflationary (pseudo) general residuated lattices. *Mathematics* **2022**, *10*, 3394. [[CrossRef](#)]
 44. Sheng, N.; Zhang, X. Regular partial residuated lattices and their filters. *Mathematics* **2022**, *10*, 2429. [[CrossRef](#)]
 45. Hu, M.; Wang, C.; Yang, J.; Wu, Y.; Fan, J.; Jing, B. Rain Rendering and Construction of Rain Vehicle Color-24 Dataset. *Mathematics* **2022**, *10*, 3210. [[CrossRef](#)]
 46. Liu, H.; Zhao, B. New results on the distributive laws of uninorms over overlap functions. *IEEE Transactions on Fuzzy Systems* **2021**, *29*, 1927–1941. [[CrossRef](#)]
 47. Liang, R.; Zhang, X. Pseudo general overlap functions and weak inflationary pseudo BL-algebras. *Mathematics* **2022**, *10*, 3007. [[CrossRef](#)]