

Correction

Correction: Ullah et al. Using Halothermal Time Model to Describe Barley (*Hordeum vulgare* L.) Seed Germination Response to Water Potential and Temperature. *Life* 2022, 12, 209

Abd Ullah ^{1,2,3,4,†} , Sadaf Sadaf ^{5,†}, Sami Ullah ^{5,*}, Huda Alshaya ⁶, Mohammad K. Okla ⁷, Yasmeen A. Alwasel ⁷  and Akash Tariq ^{1,2,3,4,*} 

¹ Xinjiang Key Laboratory of Desert Plant Roots Ecology and Vegetation Restoration, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi 830011, China

² State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi 830011, China

³ Cele National Station of Observation and Research for Desert-Grassland Ecosystems, Cele 848300, China

⁴ University of Chinese Academy of Sciences, Beijing 100049, China

⁵ Department of Botany, University of Peshawar, Peshawar 25120, Pakistan

⁶ Cell and Molecular Biology, University of Arkansas, Fayetteville, NC 72701, USA

⁷ Botany and Microbiology Department, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia; yasmeen@ksu.edu.sa (Y.A.A.)

* Correspondence: samibotany@uop.edu.pk (S.U.); akash.malik786@mails.uca.ac.cn (A.T.)

† These authors contributed equally to this work.

Error in Title

In the original publication [1], there was an error in the title regarding the species name. The correct title should be:

“Using Halothermal Time Model to Describe Barley (*Hordeum vulgare* L.) Seed Germination Response to Water Potential and Temperature”.

Text Correction

There was an error in the original publication [1]. The Equations (15) and (18) of Timson germination index (TGI) and the time for 50% germination are both the same, and also the remark provided for Equation (15) does not follow the equation. A correction has been made to Section 2.5.7, “Timson Germination Index (TGI)” and Section 2.5.10, “Seed Vigor Index-II (SVI-II)”:

2.5.7. Timson Germination Index (TGI)

The Timson Germination Index (TGI) quantifies germination speed using the following formula [52]:

$$\text{Timson Germination Index (TGI)} = \Sigma G/t \quad (15)$$

where G is the number of seeds germinated on a given day and t is the corresponding day number (e.g., 1, 2, 3...).

2.5.10. Seed Vigor Index-II (SVI-II)

The dry weight of three seedlings from each pot was measured with a precision balance. These measures were used to calculate the Seed Vigor Index according to the following formula [55]:

$$\text{Seed Vigor Index} = \text{Seed dry weight (mg)} \times \text{Seed Germination \%} \quad (18)$$



Received: 12 November 2025

Revised: 2 December 2025

Accepted: 9 December 2025

Published: 4 February 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Ullah, A.; Sadaf, S.; Ullah, S.; Alshaya, H.; Okla, M.K.; Alwasel, Y.A.; Tariq, A. Using Halothermal Time Model to Describe Barley (*Hordeum vulgare* L.) Seed Germination Response to Water Potential and Temperature. *Life* **2022**, *12*, 209. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.