

Table S1. Impact of histoprognostic characteristics on the occurrence of discordance between ceCT Vs ¹⁸F-FDG PET/CT for treatment response classification

		<6 months		6 to 10 months		10 to 16 months		>16 months	
	Variables	Numbers of observation/ discordances	<i>p</i> value	Numbers of observation/ discordances	<i>p</i> value	Numbers of observation/ discordances	<i>p</i> value	Numbers of observation/ discordances	<i>p</i> value
Location	Head/Neck/Trunk								
	Lower limbs	55 / 18	0.500	56 / 20	0.600	48 / 13	0.970	34 / 15	0.688
	Upper limbs								
	Others								
Clinical stage	I/II	42 / 14	1.000	40 / 12	0.720	34 / 7	1.000	22 / 8	0.649
	III/IV								
Histological subtypes	NMM								
	SSM	47 / 15	0.499	48 / 16	0.592	48 / 13	0.727	30 / 13	0.318
	Others								
BRAF status	Mutated	55 / 18	0.394	56 / 20	0.762	41 / 9	0.360	34 / 15	0.715
	Non-mutated								
Breslow	≤1mm	38 / 12	1.000	37 / 12	0.659	33 / 9	1.000	22 / 11	0.611
	>1mm								
Ulceration	Yes	33 / 11	0.721	31 / 8	1.000	27 / 7	1.000	17 / 8	0.347
	No								
Regression	Yes	31 / 10	0.967	27 / 7	1.000	25 / 6	1.000	16 / 8	0.200
	No								