

Correction

Correction: Stämmler et al. Estimating Glomerular Filtration Rate from Serum Myo-Inositol, Valine, Creatinine and Cystatin C. *Diagnostics* 2021, 11, 2291

Frank Stämmler ¹, Marcello Grassi ¹, Jeffrey W. Meeusen ², John C. Lieske ^{2,3}, Surendra Dasari ⁴, Laurence Dubourg ⁵, Sandrine Lemoine ⁵, Jochen Ehrich ⁶ and Eric Schiffer ^{1,*}

- ¹ Department of Research and Development, Numares AG, 93053 Regensburg, Germany; frank.staemmler@numares.com (F.S.); marcello.grassi@numares.com (M.G.)
- ² Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN 55905, USA; meeusen.jeffrey@mayo.edu (J.W.M.); lieske.john@mayo.edu (J.C.L.)
- ³ Division of Nephrology and Hypertension, Mayo Clinic, Rochester, MN 55905, USA
- ⁴ Division of Endocrinology, Diabetes, Metabolism, and Nutrition, Mayo Clinic, Rochester, MN 55905, USA; dasari.surendra@mayo.edu
- ⁵ Service d'Explorations Fonctionnelles Rénales et Métaboliques, Hôpital Edouard Herriot, 69437 Lyon, France; laurence.dubourg@chu-lyon.fr (L.D.); sandrine.lemoine@chu-lyon.fr (S.L.)
- ⁶ Children's Hospital, Hannover Medical School, 30625 Hannover, Germany; ehrich.jochen@mh-hannover.de
- * Correspondence: eric.schiffer@numares.com; Tel.: +49-941-280-949-00

In the original publication [1], there is an error regarding the intercepts in Table 2. The corrected Table 2 appears below. The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table 2. GFR_{NMR} equation according to serum cystatin C cutoff levels and sex.

Sex	Serum Cystatin C [mg/L]	GFR _{NMR} Equation ¹
Female	<1.02	$238 \times \text{cystatin C}^{-0.4114} \times \text{creatinine}^{-0.3798} \times \text{valine}^{0.1628} \times 0.9979^{\text{myo_inositol}} \times 0.9963^{\text{age}}$
	≥1.02	$239 \times \text{cystatin C}^{-0.6443} \times \text{creatinine}^{-0.3798} \times \text{valine}^{0.1628} \times 0.9979^{\text{myo_inositol}} \times 0.9963^{\text{age}}$
Male	<1.22	$266 \times \text{cystatin C}^{-0.5867} \times \text{creatinine}^{-0.3798} \times \text{valine}^{0.1628} \times 0.9979^{\text{myo_inositol}} \times 0.9963^{\text{age}}$
	≥1.22	$269 \times \text{cystatin C}^{-0.6419} \times \text{creatinine}^{-0.3798} \times \text{valine}^{0.1628} \times 0.9979^{\text{myo_inositol}} \times 0.9963^{\text{age}}$

¹ Coefficients were rounded to four digits; the sex- and cutoff-dependent intercept was rounded to 0 digits; coefficients are brought to the scale of the respective metabolite serum concentration (μmol for valine, creatinine and myo-inositol; mg/L for cystatin C).

Reference

1. Stämmler, F.; Grassi, M.; Meeusen, J.W.; Lieske, J.C.; Dasari, S.; Dubourg, L.; Lemoine, S.; Ehrich, J.; Schiffer, E. Estimating Glomerular Filtration Rate from Serum Myo-Inositol, Valine, Creatinine and Cystatin C. *Diagnostics* **2021**, *11*, 2291. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.



Citation: Stämmler, F.; Grassi, M.; Meeusen, J.W.; Lieske, J.C.; Dasari, S.; Dubourg, L.; Lemoine, S.; Ehrich, J.; Schiffer, E. Correction: Stämmler et al. Estimating Glomerular Filtration Rate from Serum Myo-Inositol, Valine, Creatinine and Cystatin C. *Diagnostics* **2021**, *11*, 2291. *Diagnostics* **2024**, *14*, 1949. <https://doi.org/10.3390/diagnostics14171949>

Received: 13 August 2024

Accepted: 20 August 2024

Published: 3 September 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).