

Reply

# Reply to Comment: Evaluation of the Association of Omentin 1 rs2274907 A > T and rs2274908 G > A Gene Polymorphisms with Coronary Artery Disease in Indian Population: A Case–Control Study

Chandan K. Jha <sup>1</sup> , Rashid Mir <sup>2</sup>, Imadeldin Elfaki <sup>3</sup>, Jamsheed Javid <sup>2</sup>, Abdullatif Taha Babakr <sup>4</sup>, Shaheena Banu <sup>5</sup> and S. M. S. Chahal <sup>1,\*</sup>

<sup>1</sup> Department of Human Genetics, Punjabi University, Punjab 147002, India; chandujha58@gmail.com

<sup>2</sup> Prince Fahd Bin Sultan Research Chair, Department of Medical Lab Technology, Faculty of Applied Medical Sciences, University of Tabuk, Tabuk 71491, Saudi Arabia; rashidmirtabuk@gmail.com (R.M.); jamsheedjavidssj@gmail.com (J.J.)

<sup>3</sup> Department of Biochemistry, Faculty of Science, University of Tabuk, Tabuk 71491, Saudi Arabia; ielfaki@ut.edu.sa

<sup>4</sup> Department of Medical Biochemistry, Faculty of Medicine, Umm Al-Qura University, Makkah 57039, Saudi Arabia; abdullatifaha@yahoo.com

<sup>5</sup> Sri Jayadeva Institute of Cardiovascular science and Research, Bangalore 560069, India; shanbanubanu@gmail.com

\* Correspondence: smschahal1@gmail.com

Received: 19 October 2020; Accepted: 23 October 2020; Published: 26 October 2020



**Abstract:** Coronary artery disease (CAD) is a major cause of death all over the world. CAD is caused by atherosclerosis which is induced by the interaction of genetic factors and environmental factors. Genome-wide association studies have revealed the association of certain gene polymorphisms with susceptibility to CAD. Omentin 1 is an adipokine secreted by the visceral adipose tissues and has been reported to have anti-inflammatory, cardioprotective, and enhances insulin sensitivity. In this study, we examined the role of omentin-1 common single nucleotide polymorphisms (SNPs) (rs2274907 A > T and rs2274908 G > A) in CAD. We conclude that the AT genotype and the T allele of the rs2274907 A > T is associated with Cad in the south Indian population. Our results indicated that the rs2274907 SNP may be associated with CAD in this population. This finding needs further validation in well-designed and large-sample size studies before being introduced in clinical settings.

**Keywords:** Coronary Artery Disease (CAD); HWE; ARMS-PCR; Omentin 1

Thank you very much for your interest in our study by Jha et al., 2019 [1]. We acknowledge that no genotyping method is 100% efficient and that genotyping errors can occur. However, we used the amplification refractory mutation system PCR (ARMS-PCR) and allele-specific PCR to genotype omentin 1 single nucleotide polymorphisms (SNPs) rs2274907 A > T and rs2274908 G > A. These methods are standard, fast, simple, and reliable for genotyping [2,3]. Results were repeated several times and once perfection was achieved, the remaining CAD samples were screened. Results were validated by different methods. Some CAD cases were screened by RFLP using restriction enzyme AccI (XmiI) or by Sanger sequencing. No difference in results from the ARMS-PCR was observed.

Our genotyping result deviated from HWE and this was probably due to many factors including population structure and/or sample size, purifying selection, copy number variation, inbreeding, or the substructure of this population [4,5]. The genotyping result was not very different from previous results in different populations (Table 1). Deviation from HWE was associated with loss or gain of

heterozygosity [4]. In our study, it is likely that there was no loss or gain of the heterozygote genotype (AT) because the AT genotype percentage was 51% (Table 1). This percentage is more or less comparable to the AT genotype percentage obtained in some other populations (Table 1). We suggested that the deviation from HWE observed in our study occurred due to the substructure of this population. In support of this assumption is the recent study by Rathwa et al. [6] (Table 1), in which they report 0% AA genotype, 17.6% AT genotype, and more than 80% for the mutant genotype (TT). This indicates a complete shift to the mutant genotype (TT) in this population. Deviation from HWE observed in our study can also be due to the relatively small sample size used or other factors stated above [4,5]. We could not find the rs2274907 genotype distribution of an Indian population in other studies apart from the study by Rathwa et al. [6].

In Jha et al., 2019 [1], we report a potential association of the omentin SNP rs2274907 with CAD in a sample from a South Indian population. This result is consistent with previous studies conducted in Pakistani and Turkish populations [7,8], and to some extent in Iranian population [9]. Moreover, other studies reported an association of rs2274907 with type 2 diabetes [10,11], which is a risk factor for CAD.

In the database, the A allele is the major allele (<https://www.ncbi.nlm.nih.gov/snp/rs2274907>). Moreover, previous studies conducted in Pakistan and Iranian populations reported that the nucleotide change is from A to T [7,11,12]. The codon GAC is changed to GTC replacing the amino acid residue aspartate to valine at position 109 [13]. However, omentin 1 rs2274907 A > T is an SNP rather than a mutation that causes diseases [13].

**Table 1.** Genotype distribution of omentin gene polymorphism rs2274907 in different populations.

Reference	AA	AT	TT	Population
1—Kohan et al., 2016 [12]	57.5	38.8	3.7	Iran
2—Khoshi et al., 2019 [11]	70.6	27.1	2.3	Iran
3—Rakowska, et al., 2017 [10]	48.50	41.92	6.59	Polish
4—Rathwa et al., 2019 [6]	0	17.6	82.4	Indian
5—Shazia Nazar et al., 2017 [7]	63	30	7	Pakistan
6—Ümit Yörük et al., 2014 [8]	6.66	48	45.33	Turkish
7—Zhang et al., 2015 [14]	9.8	43.5	46.7	Chinese
8—Jha et al., 2019 [1]	49	51	0	Indian

In summary, we used a standard method for omentin SNPs genotyping. Our results indicated that the rs2274907 SNP may be associated with CAD in this population. This finding needs further validation in well-designed and large-sample size studies before being introduced in clinical settings.

**Author Contributions:** Conceptualization, all authors; methodology, C.K.J., R.M. and I.E.; sample collection, C.K.J. and S.B.; analyses, R.M. and I.E.; validation, all authors; supervision, S.M.S.C.; visualization, C.K.J., R.M., I.E., J.J., A.T.B., S.B., S.M.S.C.; writing, R.M. and I.E. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Jha, C.K.; Mir, R.; Elfaki, I.; Javid, J.; Babakr, A.T.; Banum, S.; Chahal, S.M.S. Evaluation of the Association of Omentin 1 rs2274907 A > T and rs2274908 G > A Gene Polymorphisms with Coronary Artery Disease in Indian Population: A Case Control Study. *J. Pers. Med.* **2019**, *9*, 30. [[CrossRef](#)] [[PubMed](#)]
- Medrano, R.F.V.; de Oliveira, C.A. Guidelines for the Tetra-Primer ARMS-PCR Technique Development. *Mol. Biotechnol.* **2014**, *56*, 599–608. [[CrossRef](#)] [[PubMed](#)]

3. Darawi, M.N.; Ai-Vyrn, C.; Ramasamy, K.; Hua, P.P.J.; Pin, T.M.; Kamaruzzaman, S.B.; Majeed, A.B.A. Allele-specific polymerase chain reaction for the detection of Alzheimer's disease-related single nucleotide polymorphisms. *BMC Med Genet.* **2013**, *14*, 27. [[CrossRef](#)] [[PubMed](#)]
4. Chen, B.; Cole, J.W.; Grond-Ginsbach, C. Departure from Hardy Weinberg Equilibrium and Genotyping Error. *Front. Genet.* **2017**, *8*, 167. [[CrossRef](#)] [[PubMed](#)]
5. Wittke-Thompson, J.K.; Pluzhnikov, A.; Cox, N.J. Rational Inferences about Departures from Hardy-Weinberg Equilibrium. *Am. J. Hum. Genet.* **2005**, *76*, 967–986. [[CrossRef](#)] [[PubMed](#)]
6. Rathwa, N.; Patel, R.; Palit, S.P.; Jadeja, S.D.; Narwaria, M.; Ramachandran, A.; Begum, R. Circulatory Omentin-1 levels but not genetic variants influence the pathophysiology of Type 2 diabetes. *Cytokine* **2019**, *119*, 144–151. [[CrossRef](#)]
7. Nazar, S.; Zehra, S.; Azhar, A. Association of single Nucleotide Missence Polymorphism Val109Asp of Omentin-1 gene and coronary artery disease in Pakistani population: Multicenter study. *Pak. J. Med. Sci.* **2017**, *33*, 1128–1133.
8. Yoruk, U.; Yaykasli, K.O.; Ozhan, H.; Memisogullari, R.; Karabacak, A.; Bulur, S.; Aslantas, Y.; Başar, C.; Kaya, E. Association of omentin Val109Asp polymorphism with coronary artery disease. *Anatol. J. Cardiol.* **2014**, *14*, 511–514. [[CrossRef](#)]
9. Jamshidi, J.; Ghanbari, M.; Asnaashari, A.; Jafari, N.; Valizadeh, G.A. Omentin Val109Asp polymorphism and risk of coronary artery disease. *Asian Cardiovasc. Thorac. Ann.* **2017**, *25*, 199–203. [[CrossRef](#)] [[PubMed](#)]
10. Mrozikiewicz-Rakowska, B.; Sobczyk-Kopciol, A.; Szymanski, K.; Nehring, P.; Szatkowski, P.; Bartkowiak-Wieczorek, J.; Bogacz, A.; Aniszczyk, A.; Drygas, W.; Ploski, R.; et al. Role of the rs2274907 allelic variant of the ITLN1 gene in patients with diabetic foot. *Pol. Arch. Intern. Med.* **2017**, *127*, 319–327. [[PubMed](#)]
11. Khoshi, A.; Bajestani, M.K.; Shakeri, H.; Goodarzi, G.; Azizi, F. Association of Omentin rs2274907 and FTO rs9939609 gene polymorphisms with insulin resistance in Iranian individuals with newly diagnosed type 2 diabetes. *Lipids Health Dis.* **2019**, *18*, 142. [[CrossRef](#)] [[PubMed](#)]
12. Kohan, L.; Safarpur, M.; Abdollahi, H. Omentin-1 rs2274907 and resistin rs1862513 polymorphisms influence genetic susceptibility to nonalcoholic fatty liver disease. *Mol. Biol. Res. Commun.* **2016**, *5*, 11–17. [[PubMed](#)]
13. Schäffler, A.; Zeitoun, M.; Wobser, H.; Buechler, C.; Aslanidis, C.; Herfarth, H. Frequency and significance of the novel single nucleotide missense polymorphism Val109Asp in the human gene encoding omentin in Caucasian patients with type 2 diabetes mellitus or chronic inflammatory bowel diseases. *Cardiovasc. Diabetol.* **2007**, *6*, 3. [[CrossRef](#)] [[PubMed](#)]
14. Zhang, C.; Zhu, K.J.; Liu, J.L.; Xu, G.X.; Liu, W.; Jiang, F.X.; Zheng, H.F.; Quan, C. Omentin-1 plasma levels and omentin-1 expression are decreased in psoriatic lesions of psoriasis patients. *Arch. Dermatol. Res.* **2015**, *307*, 455–459. [[CrossRef](#)] [[PubMed](#)]

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).