

Supplementary Table S2. Comparison between groups according to IL-17, IL-23 and standard inflammatory biomarkers levels

Variables (Median, IQR)	CD			UC		
	Mild or moderate (n=16)	Severe (n = 15)	p value	Mild or moderate (n = 14)	Severe (n = 17)	p value
Albumin (g/dL)	4 (3.82; 4.4)	3.7 (3.6; 4.2)	0.02	4 (3.97; 4.3)	3.9 (3.5; 4)	0.008
ESR	34.5 (17.25; 85.5)	65 (20; 102)	0.1	31 (17.5; 60)	48 (25; 106)	0.1
CRP (mg/L)	0.39 (0.32; 0.72)	3.1 (0.46; 7.9)	0.008	0.47 (0.37; 0.65)	1.2 (0.39; 5.29)	0.1
Calprotectin (µg/g)	50 (45; 370)	640 (400; 1000)	<0.001	260 (50; 812.5)	1200 (905; 1700)	0.001
IL-17 (pg/mL)	764.52 (528.43; 1222.05)	1695.00 (665.35; 2535.75)	0.1	976.47 (528.43; 1222.05)	2304.60 (1011.30; 2815.42)	0.004
IL-23 (pg/mL)	777.63 (744.15; 800.31)	1477.49 (1434.29; 1537.97)	<0.001	745.23 (731.19; 946.11)	1468.85 (1451.57; 1598.46)	< 0.001

¹CD, Crohn’s disease; UC, ulcerative colitis. IQR, Interquartile range. ESR, Erythrocyte sedimentation rate. CRP, C-reactive protein. IL, Interleukin.