Eucalyptus forests	Oak forests	Olive groves	Pine forests	Scrublands	Vineyards	Prays oleae	Chrysoperla carnea	SHDI	Grasslands
	Corr: 0.0663	-0.418	Corr: 0.027	Corr: 0.136	Corr: 0.176	Corr: -0.0845	Corr: 0.462	Corr: 0.448	Corr: 0.0171
	0.0005	-0.418	Corr. 0 000137	Corr: -0.318	Corr: 0.0732	-0.149	Corr: -0.0262	Corr: 0.143	Corr: -0.0608
ŀ	N	\frown	Corr. -0.0959	Corr: -0.343	Corr: -0.17	Corr: 0.412	Corr: 0.0159	Corr: -0.777	Corr: -0.466
			La	Corr: -0.105	Corr: -0.157	Corr: 0.197	Corr: -0.199	Corr: 0.286	Corr: 0.0896
i		6	21	L	Corr: -0.153	Corr: 0.053	Corr: 0.205	Corr: -0.00191	Corr: -0.399
	<u>.</u>	. <u>.</u>		<u>.</u>	\square	Corr: -0.376	Corr: -0.0586	Corr: 0.373	Corr: 0.249
1	÷		ř.: ·	č.	ŀ	\frown	Corr: 0.192	Corr: -0.375	Corr: -0.459
i. · ·	à		i ∙: .	÷	i		$ \land $	Corr: -0.044	Corr: -0.248
1	۴ <u>۰</u> .	. ³⁷ .\ ₂	ŀ' ' ·	Ę÷ .	ŀ	*.		\frown	Corr: 0.503
.		S	ļ.• · · .	b	. · ·	6			\wedge

Figure S1. Correlations of the different predictor variables included in the models.

	Estimate	Pr (< z)	
Intercept	-3.1380	< 0.001	
Surrounding olive groves (%)	0.0372	< 0.001	
C. carnea abundance	0.0246	< 0.001	
Interaction (Olive groves × <i>C. carnea</i>)	-0.0007	< 0.001	

Table S1. Estimated parameters of the selected model for the *Prays oleae* infestation. Pr (<|z|) shows the level of support of these parameters.