



Article Utilization of Pomelo (*Citrus maxima*) Peel Waste into Bioactive Essential Oils: Chemical Composition and Insecticidal Properties

Naduvilthara U. Visakh ¹, Berin Pathrose ^{1,*}, Arunaksharan Narayanankutty ^{2,*}, Ahmed Alfarhan ³ and Varsha Ramesh ⁴

- ¹ Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Thrissur 680656, India; visakhnu17palazhi@gmail.com
- ² Division of Cell and Molecular Biology, PG & Research Department of Zoology, St. Joseph's College (Autonomous), Devagiri, Calicut 673008, India
- ³ Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia; alfarhan@ksu.edu.sa
- ⁴ Department of Biotechnology, Deakin University, Burwood, VIC 3125, Australia; vramesh@deakin.edu.au
- * Correspondence: berin.pathrose@kau.in (B.P.); arunaksharann@devagiricollege.org (A.N.)

Simple Summary: The disposal of agricultural waste products is an emerging concern and an alternative to this is the development of value-added products from these wastes. Here we extracted the essential oil from *Citrus maxima* (CMEO) and examined its larvicidal and pest control potentials. Results pointed out that CMEO can be effective biopesticides against two major insect pests of stored grains. Furthermore, CMEO had a significant larvicidal action against different mosquito species. This study provided useful information on the compositional aspects and insecticidal properties of CMEO.

Abstract: The wastes generated during the post-harvest handling of various agricultural commodities is rather under-utilized. The peels of citrus fruits are often discarded as waste. Citrus peels are rich in essential oils and exhibit toxicity towards various insect species. The essential oils are also an eco-friendly option for insect pest management. The *Citrus maxima* peel essential oil (CMEO), a waste product, characterized it, and evaluated its potential for insect pest management. The major terpenoids present in CMEO are Limonene and α -Pinene. The CMEO displayed potentials in controlling the insect pests via contact and fumigant toxicity. Moreover, CMEO showed significant larvicidal activities against *Culex tritaeniorhynchus* and *Aedes aegypti* species of mosquitoes; however, *Armigeres subalbatus* was more resistant. The biological safety of the essential oil was also tested against the stored seeds, where no significant inhibition of seed germination was noticed compared to the control. Utilizing a waste product such as citrus peel for pest management can achieve the dual objective of waste utilization and eco-friendly pest management. Overall, the CMEO is therefore found to be a bioactive essential oil extracted from the wastes of pomelo (*C. maxima*).

Keywords: essential oil; larvicidal toxicity; insecticidal toxicity; fumigant toxicity; repellent activity; limonene; *Tribolium castaneum; Callosobruchus maculatus; Armigeres subalbatus; Culex tritaeniorhynchus; Aedes aegypti*

1. Introduction

The pollution and environmental issues created by agriculture and food wastes are an important and emerging concern [1]. However, the production of bioactive by-products and secondary metabolites is an emerging area of agri-food waste management. Among these, the essential oils isolated from food wastes can have significant economic value.

Essential oils are important plant-derived compounds with potent biological activities that can be utilizable in agriculture, medicine, and biotechnology. Essential oils are also



Citation: Visakh, N.U.; Pathrose, B.; Narayanankutty, A.; Alfarhan, A.; Ramesh, V. Utilization of Pomelo (*Citrus maxima*) Peel Waste into Bioactive Essential Oils: Chemical Composition and Insecticidal Properties. *Insects* 2022, *13*, 480. https://doi.org/10.3390/ insects13050480

Academic Editors: Adriano Pinter and Karin Kirchgatter

Received: 8 March 2022 Accepted: 10 May 2022 Published: 20 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). known to control the population kinetics of microbes, including bacteria [2], various viruses [3], infectious fungi [4] and so on. It is also evident that these essential oils inhibit the biofilm formation effects of various bacterial colonies [5,6]. Essential oils can also alleviate antibiotic resistance in multiple pathogenic microorganisms [7].

Besides antimicrobial properties, essential oils are also known for their potential effects on economically destructive pest species. The essential oils are emerging tools in the control of pest populations in warehoused grains. Various essential oils are effective against insect pests such as *Sitophilus* sp., *Callosobruchus* sp., *Tribolium* sp., etc. via their multi-organ toxic insults [8,9]. Apart from these, the essential oils are also known for their anticancer properties [10], antidiabetic effects [11], hypolipidemic effects [12], and several other pharmacological effects. Citrus peels are one of the major agriculture waste as well as an worthy source of essential oils that can be commercially exploited as an eco-friendly pest management [13]. Though there are several beneficial effects for the fruits of various citrus plants, their peels are always emerging as the predominant waste products.

Citrus plants are well-known for its nutritional and pharmacological effects [14,15]; further these fruits are important sources of essential oils with wide application [16]. The essential oils of citrus fruits are currently being used as botanical pesticides [13]. The terpenoids present in citrus essential oils exhibit insecticidal [17], fumigant [18], and repellent [19] activity against various insects pests present in warehouse grains. Essential oils derived from citrus fruits also possess significant activity against mosquitoes [20–22]. Among the various species of citrus fruits, the largest one and the least studied is *Citrus maxima* (L.) Osbeck (synonym *Citrus grandis*). Studies have indicated the possible uses of *C. maxima* peels. The extracts of *C. maxima* peels exhibit radical inhibitiory and hypoglycemic properties [23,24]. Besides, the extracts of the plant peels also inhibit neuro-inflammation and subsequent cognitive impairment [25,26]. Furthermore, significant radical reducing and antimicrobial benefits of the CMEO is reported [16,27]. Further, the CMEO has been effective in the suppression of hyphal growth in *Aspergillus flavus* [28].

However, studies on the utilization of CMEO extracted from its peel (an agro-food waste) for its insecticidal and pharmacological applications are lacking when compared to other species of citrus. Therefore, here we extracted the peel essential oil of *C. maxima* (CMEO) and characterized its chemical configuration through GC-MS to identify the bioactive components. The efficacy of CMEO against the insect pests of warehoused grains and mosquitoe larvae were analyzed. The biosafety aspects of the CMEO were determined using seed germination and non-targetted organism models.

2. Materials and Methods

2.1. Collection of Citrus Peel and Essential Oil Extraction

Pomelo (*Citrus maxima*) fruits were collected from the fruit orchard at Kerala Agricultural University, Thrissur, India (10.5449° N, 76.2864° E) during December 2021. After washing these pomelo fruits with distilled water, peels were removed by using knife. Peels were then kept in sterile ziplock bags at 4 °C until they were used for essential oil extraction. Hydro-distillation of fresh peels (300 g) was carried out using modified Clevenger method for the extraction of essential oil [29]. The hydro-distillation process lasted for 5 h (100 °C). The CMEO obtained was passed through sodium sulphate (AR) to remove the content of water. On a fresh weight basis, the oil yield was determined by using the following formula yield (%, v/w):

Yield (%,
$$v/w$$
) = V_{EO}/W_F × 100

where V_{EO} is the dry essential oil volume, and W_F is the weight of pomelo peels used for essential oil extraction. The CMEO thus obtained was put in storage using dark ambercoloured glass bottle at 4 °C inside the refrigerator until required for experiments [30].

2.2. Test Insects and Larval Culture

Red flour beetles (*Tribolium castaneum*) were reared using wheat flour. In order to eradicate prior insect infestation, wheat grains were heated for six hours at 50 °C and

then it was milled into wheat flour. Twenty adults of *T. castaneum* were released into plastic containers (18 cm × 10 cm) containing 250 g of sterilized wheat flour fortified with 5% (w/w) brewer's yeast. Ten such containers were prepared and kept in the culture room. After five days of oviposition, the adults were then sieved out and transferred to fresh rearing containers. Rearing containers were regularly examined to obtain uniformly aged adults (temperature 30 ± 2.5 °C; relative humidity of 80 ± 4.2%). All bioassays were conducted on adult insects (17 ± 2 days old) by sieving from wheat flour.

The culture of pulse beetle (*Callosobruchus maculatus*) was maintained on green gram grains procured from a local market. The grains were de-infestated by washing and drying 60 °C for one hour. Twenty adult male and female beetles were released into plastic containers (1 L volume) containing 100 g of sterilized green gram grains. Containers were covered with muslin cloth and kept for incubation for five days. After five days, adult insects were removed, and culture maintained (temperature 30 ± 2.5 °C; relative humidity of $80 \pm 4.2\%$). The adult insects emerged from this culture was used for sub-culturing. For bioassays, 5 ± 2 -day-old adult insects were used. All the experiments were conducted at the Pesticide Residue Testing Laboratory, Department of Agricultural Entomology, Kerala Agricultural University, Kerala, India.

The different mosquito species (*Ar. subalbatus, Ae. aegypti* and *Cx. Tritaeniorhynchus*) were reared in cages (450 mm \times 300 mm \times 300 mm) under standard conditions (28 \pm 2.5 °C; 75% humidity; 16 light: 8 h dark cycle). The larvae were reared in plastic trays and those in the third instar stage were collected for study.

2.3. Analaysis of Chemical Configuration Using GC-MS

The instrument used was TSQ 8000 Evo (Thermo scientific, Waltham, MA, USA) instrument equipped with an autosampler and TG-1MS capillary column (30 mm × 0.25 mm × 0.25 µm). The T_{OV} (temperature of oven) was sustained at 50 °C for 1 min and a temperature ramping of at 10 °C min⁻¹ to 120 °C, then to 270 °C for 5 min at 5 °C min⁻¹. The temperature of sample injector was retained at 250 °C. With a split ratio of 1:200, the samples (0.1 µL) were injected. Helium served as the carrier gas, with a flow rate of 1.0 mL min⁻¹. The GCMS spectra (35 to 500 *m*/*z*) with a dwell time of 0.2 ms. Xcalibur 1.1 software was recorded and analyse the mass spectra data (Thermo scientific). Spectral comparison was done with NIST library (library similarity and reverse similarity index > 800), the essential oil components were identified. Calculations were made for individual chemical component peak areas to determine relative percentage amounts.

2.4. Contact Toxicity

The adulticidal assay was conducted by using the residual film method with few modifications [31]. Initially, the concentrations of essential oils required to obtain insect mortality were assessed over a broad range of doses. Further, a narrow range of concentrations (40–120 mg/cm² for *T. castaneum* and 2–10 mg/cm² for *C. maculatus*) were bioassayed to obtain LC₅₀ and LC₉₀. The CMEO was dissolved in acetone (HPLC grade) to obtain the desired concentration and applied to 90 mm (mg/cm²). A uniform film of essential oils was formed over the Petri dish by gently rotating the Petri plate. For solvent evaporation, the treated Petri dishes were air-dried for 15 min, followed by the release of 10 adult insects and then incubating the plates at 28 ± 2 °C and 80 ± 5 %. Petri dishes were enclosed with perforated plastic lids to avoid any fumigant toxicity. Acetone (1 mL) was applied inside the Petri dish for control experiments. Three replications were employed for both treatments as well as the control. The percent mortality for *C. maculatus* and *T. castaneum* was measured at the end of 24 and 48 h and Mortality data was corrected using Abbott's formula [32].

2.5. Fumigant Toxicity

A modified method was used to determine the fumigant toxicity of essential oils derived from *C. maxima* against *T. castaneum* and *C. maculatus* adults [33]. Initially, preliminary bioassays were done with a wide range of concentrations to fix the desired concentrations for the experiment. In brief, fumigant toxicity was accomplished by placing filter paper discs (2.5 cm diameter) impregnated with the CMEO (1, 3, 5, 7, 10 mg/L air for both

discs (2.5 cm diameter) impregnated with the CMEO (1, 3, 5, 7, 10 mg/L air for both *T. castaneum* and *C. maculatus*) of plant essential oils (mg/L air) hanging down by a thread in the polyacrylamide plastic containers (80 mL) containing 10 adults of *T. castaneum* and *C. maculatus*, and sealing them tightly. Another airtight polyacrylamide plastic container (80 mL) was used for the control without essential oils on the filter paper. For both treatments and controls, three replicates were employed. Mortality was recorded after 24 and 48 h from the start of essential oil exposure. The treated sets were kept at a constant temperature of 28 ± 2 °C and relative humidity of $80 \pm 5\%$. Abbott's equation was used to correct mortality data as mentioned earlier.

2.6. Repellent Activity Assay

The method of area preference was utilized to estimate the repellent activity [34]. We evaluated CMEO's repellent activity against adult insects of *T. castaneum* and *C. maculatus*. Briefly, the test area was a 9 cm filter paper disc cut in to half. Varying concentrations of CMEO (0.5, 1.5, 2.5, 3.5 and 5 mg/cm²) were diluted in 0.5 mL acetone (HPLC grade) to prepare various solutions. The Petri plate was enclosed with a perforated plastic lid, and on each filter paper disc, ten adult insects were introduced to each concentration of CMEO. In each time interval (2, 4, 6, 12 and 24 h) the insects present on each half of disc was numerated. Percent repellence was calculated by aforementioned method [34].

Categorization of repellent effects by essential oils was discussed and estimated [35]. On the basis of percentage repellence (PR), five classes could be sorted: Class 0 = PR of 0 to 0.1%; Class I (0.2–10%); Class II (20.1–40%); Class III (40.1–60%); Class IV (60.1–80%), and Class V (80.1–100%).

2.7. Effect of CMEO on Wheat Germination

The impact of CMEO on the sprouting of wheat grains (stored products) is carried out to analyze the phytotoxic effects; briefly, the grains (25 nos) were kept in a moisturized container with access to light and water. The grains were exposed to varying concentrations (50, 100, and 250 μ g/mL) of CMEO (in 0.01% Tween 80). The relative humidity of the test samples were set at 75% and the incubator was maintained at 27 ± 2.5 °C in the dark (no photoperiod). Development of a radicle of length 1 cm was considered as a true germination and noted. A control involving 0.01% Tween 80 was used to dissolve the CMEO, and therefore a vehicle control was maintained to normalize the data. All assays were repeated six times, with each conducted in triplicate.

2.8. Screening of Mosquito Larvicidal Activity by CMEO

The larvae of *Cx. tritaeniorhynchus, Ar. Subalbatus* and *Ae. aegypti* were collected in the third instar stage, and approximately 50 of them were taken to individual chambers. The CMEO (dissolved in Tween 80 at 0.01%) was added to each of the beakers (0–200 μ g/mL) and observed for 24 h. Tween 80 (0.01%) was the control and the values were normalized using this group. The larvicidal assays were carried out four different times, each conducted in triplicate. The average mortality was plotted against concentration, and LC₅₀ value was determined by probit analysis.

2.9. Toxicity on Non-Targeted Organism (Poecilia reticulata)

The toxicity of pest and vector control agents can also harm non-targeted organisms including fishes. Hence, we selected *Poecilia reticulata* as a model organism for non-targeted species toxicity. Briefly, the male and female fishes of length 3.25 ± 0.08 cm and weighing 1.15 ± 0.12 g were treated with different doses of CMEO (0–250 µg/mL) over 48 h and observed for any kind of toxicity symptoms inlcuding body color change, difficulty in swimming, and mortality [36].

2.10. Data Analysis

The percent mortality in different tests was analysed using a one-way ANOVA test at 5% significance with Grapes version 1.1.0 software [37]. Using Polo Plus 2.0 software, the LC_{50} , LC_{90} , slope and 95% confidence levels were calculated for both contact and fumigant toxicity according to Finney's analysis. In the repellency test, to correct for heterogeneity of treatment variance, mean PR values were first transformed with an arcsine function before being assessed with one-way analysis of varience, and Tukey's HSD analysis was used to separate means. The phytotoxicity study was represented as mean \pm SD of germinating wheat grain, and tested by one-way ANOVA followed by Tukey's HSD test.

3. Results

3.1. Chemical Configuration of CMEO by GC-MS

CMEO obtained through hydrodistillation had a yield of $0.58\% \pm 0.016$ (v/w). Retention times (RTs) of each compound are listed. In total, we identified 25 different chemical compounds (Figure 1) from the oil (Table 1). The predominant components in *C. maxima* oil were D-limonene (33.61%), β -sitosterol (17.99%), α -sitosterol (12.19%), stigmasterol (5.22%) and α -pinene (4.32%).



Figure 1. The GC-MS chromatograms of Citrus maxima essential oil (CMEO).

3.2. Contact Toxicity

The results of contact toxicity bioassay indicated that CMEO at various concentrations had substantial contact toxicity via the residual film method to both *T. castaneum* and *C. maculatus* adult insects. CMEO caused more toxicity to adults of *C. maculatus* at 24 and 48 h after exposure, even at low concentrations $(2-10 \text{ mg/cm}^2)$ when compared to *T. castaneum* $(40-120 \text{ mg/cm}^2)$ (Table S1). The LC₅₀ and LC₉₀ values for CMEO against *T. castaneum* adults were 63.31 and 121.44 mg/cm², respectively, after 24 h exposure (Table 2). Similarly, at 48 h exposure, LC₅₀ and LC₉₀ values for CMEO against *T. castaneum* adults were 37.15 and 109.38 mg/cm², respectively. LC₅₀ and LC₉₀ values for *C. maculatus* adults were 7.12 and 16.73 mg/cm², respectively, at 24 h. Similarly, at 48 h exposure, LC₅₀ and LC₉₀ values for *C. maculatus* adults were 5.06 and 13.44 mg/cm², respectively. Increasing exposure times and concentrations resulted in a rise in mean mortality (%) with both stored grain insect pests. CMEO exhibited the highest efficacy against *C. maculatus* adults at lower concentration, followed by *T. castaneum* adults (Table 2).

Peak No.	Retention Time	Component	RSI ^a	%RA ^b
1	5.51	3-carene	934	1.49
2	6.12	Cyclohexene, 4-methylene-1-(1-methylethyl)	921	0.40
3	6.36	α-Pinene	895	4.32
4	7.02	D-Limonene	881	33.61
5	7.69	trans-Linalool oxide (furanoid)	878	1.17
6	7.93	cis-Linaloloxide	901	0.62
7	8.08	1,6-Octadien-3-ol, 3,7-dimethyl	921	0.76
8	9.66	α-Terpineol	923	0.57
9	10.48	2,6-Octadienal, 3,7-dimethyl-, (Z)	876	0.31
10	11.02	Citral	921	0.37
11	12.42	2-Carene	883	0.26
12	13.21	Geranyl acetate	917	0.64
13	14.20	Caryophyllene	921	0.41
14	15.49	α-Cubebene	882	3.18
15	15.81	α-Guaiene	904	0.90
16	29.50	4,4-dimethyl Cholesta-22,24-dien-5-ol	895	0.82
17	29.82	Stigmasterol 9		5.22
18	30.05	Desmosterol	864	3.79
19	30.12	(3á,22E) 3-methoxy-Stigmasta-5,22-diene,	873	1.95
20	35.29	campesterol	895	4.31
21	35.43	α-Sitosterol	838	12.19
22	35.59	β-Sitosterol	928	17.99
23	36.36	24-propylidene-, (3á) Cholest-5-en-3-ol,	874	1.90
24	36.49	Allopregnane-3á,7à, 11à-triol-20-one	829	0.39
25	38.91	9,19-Cyclolanost-24-en-3-ol, (3á)	898	2.41
		Total (%)		100.00

Table 1. Chemical configuration of Citrus maxima essential oil (CMEO).

^a Reverse similarity index; ^b Relative area.

3.3. Fumigant Toxicity

The CMEO exhibited fumigant potential against both stored grain insect pests. At various concentrations (1-10 mg/L air) for 24 and 48 h (Table S2). The CMEO showed high fumigant activity against *C. maculatus* adults compared to *T. castanaeum* adults. After fumigant toxicity bioassay, the LC₅₀ and LC₉₀ of the CMEO was calculated along with their respective confidence intervals (95%) (Table 3). The LC₅₀ and LC₉₀ for each exposure period were determined. Based on probit analysis, lethal concentrations of 4.95 mg/L and 4.13 mg/L resulted in 50% mortality (24 h and 48 h) in *T. castanaeum* adults. Similarly, lethal concentrations of 3.38 mg/L air and 1.34 mg/L air observed 50% mortality for *C. maculatus* adults (for 24 and 48 h), respectively. The effectiveness of CMEO was time-dependent. The results indicated that the lethal concentration decreased with increasing exposure periods. There was a greater vulnerability in *C. maculatus* adults compared with *T. castaneum* adults to CMEO (Table 3).

Test Insects	Exposure (h)	LC ₅₀ ^a (mg/cm ²)	LC ₉₀ ^a (mg/cm ²)	Slope \pm SEM ^b	χ^2 (d.f)
T. castaneum	24	63.31 (49.4–73.2)	121.44 (106.2–151.5)	0.022 ± 0.004	0.335 (3)
	48	37.15 (18.8–48.2)	109.38 (86.6–192.8)	2.733 ± 0.708	0.264 (3)
C. maculatus	24	7.12 (5.1–12.7)	16.73 (10.4–153.2)	3.458 ± 0.601	4.58 (3)
	48	5.06 (3.3–7.1)	13.44 (8.8–52.2)	3.023 ± 0.501	3.64 (3)

Table 2. Lethal concentrations on contact activity of CMEO against various insect pests.

 a Values in parenthesis represent lower and upper confidence limit; b SEM: Standard error of mean; χ^{2} : Chi-square.

Test Insects	Exposure Time (h)	LC ₅₀ (mg/L Air)	LC ₉₀ (mg/L Air)	Slope \pm SEM ^a	χ^2 (d.f)
Teachanan	24	4.95 (4.1–5.9)	12.68 (9.7–20.1)	3.138 ± 0.503	1.396 (3)
1. castaneum	48	4.13 (3.3–5.0)	11.82 (8.9–18.8)	2.806 ± 0.431	1.262 (3)
C. maculatus	24	3.38 (2.1–4.8)	29.61 (14.8–170.5)	1.360 ± 0.318	1.136 (3)
	48	1.34 (0.35–2.2)	17.80 (9.1–129.6)	1.144 ± 0.316	2.535 (3)

Table 3. Lethality of CMEO by fumigation against *T. castaneum* and *C. maculatus*.

^a SEM: Standard error of mean; χ^2 : Chi-square.

3.4. Repellent Activity

Both test insects, *T. castaneum* and *C. maculatus*, were strongly repelled by the CMEO (Table 4). According to the results of ANOVA, CMEO showed strong repellency to *T. castaneum* and *C. maculatus* adults (p > 0.05). At the lowest concentration (0.5 mg/cm^2 dosage), mean PR values were over 50 % (Class III) at 2 h to 24 h post-exposure for *T. castaneum*. Notably, for *C. maculatus* adults, CMEO at the lowest concentration (0.5 mg/cm^2) was able to cause repellence of only 23.90% (Class II) at 2 h to 24 h post-exposure. At higher concentration (5 mg/cm^2), CMEO repelled *T. castaneum* with the mean PR of 78.7% (Class IV). The mean PR value was 69.30% (Class IV) at the highest applied concentration (5 mg/cm^2) with *C. maculatus* adults. From the overall data, CMEO's were moderately repellent (Class II–IV) to *C. maculatus* adults at 2 h to 24 h; whereas, the CMEO repelled *T. castaneum* adults, (Class III–V). The repellence activities of CMEO was significantly different against *T. castaneum* and *C. maculatus* (Table 4).

3.5. Larvicidal Potential of CMEO

The *Ar. Subalbatus, Ae. aegypti* and *Cx. tritaeniorhynchus* species of mosquito larvae were opted for larvicidal screening; the essential oil of *C. maxima* was least effective against the *Ar. subalbatus* species (76.24 \pm 3.2 μ g/mL). Conversely, the essential oil was effective against *Ae. Aegypti* and *Cx. tritaeniorhynchus* species (47.07 \pm 2.4 and 58.04 \pm 2.8 μ g/mL) (Table 5).

3.6. Phytotoxicity Analysis of the CMEO

The phytotoxicity analysis of the essential oil was determined in terms of grain germination potentials. The Table 6 indicates no significant variation between any treated doses of CMEO on grain germination, and therefore was found to be non-phytotoxic.

Test	Concentration	Repellence Percentage of Treatments After				% Repellency	Repellent	
Insects	(mg/cm ²)	2 h	4 h	6 h	12 h	24 h	(Mean± ^b SEM)	Class
	0.5	$33.3\pm13.3~^{a}$	$46.7\pm6.6~^a$	60.0 ± 11.5 $^{\rm a}$	$66.7\pm6.6~^{a}$	60.0 ± 11.5 $^{\rm a}$	$53.3\pm5.8~^{\rm b}$	III
	1.5	$33.3\pm17.6~^{\rm a}$	66.7 ± 6.6 $^{\rm a}$	$60.0\pm11.5~^{\rm a}$	$66.6\pm6.6~^{a}$	$66.7\pm13.3~^{\rm a}$	$58.7\pm6.5~^{ab}$	III
	2.5	60.0 ± 11.5 a	$53.3\pm6.6~^{a}$	$66.7\pm13.3~^{\rm a}$	$73.3\pm6.6~^{a}$	73.3 ± 17.6 $^{\rm a}$	$65.3\pm3.8~^{ab}$	IV
T. castaneum	3.5	$66.7\pm13.3~^{a}$	$60.0\pm11.5~^{\rm a}$	$73.3\pm17.6~^{a}$	80.0 ± 11.5 $^{\rm a}$	$86.7\pm13.3~^{\rm a}$	$73.3\pm4.7~^{ab}$	IV
	5	$73.3\pm6.6~^{a}$	$66.7\pm13.3~^{\rm a}$	$73.3\pm13.3~^{\rm a}$	$86.7\pm13.3~^{\rm a}$	$93.3\pm6.6~^{a}$	78.7 ± 4.9 $^{\rm a}$	IV
-	F value	2.07	0.85	0.29	1.02	1.29	3.86	
	<i>p</i> value	0.16	0.53	0.87	0.44	0.33	0.02	
	^a d.f	4	4	4	4	4	20	
-	0.5	$6.7\pm6.6~^a$	13.3 ± 6.6 a	$33.3\pm17.6~^{a}$	$33.3\pm13.3~^{\rm a}$	$33.3\pm6.6~^{b}$	$23.9\pm8.9^{\text{ c}}$	Π
	1.5	$20.0\pm11.5~^{\rm a}$	$26.7\pm6.6~^a$	$40.0\pm11.5~^{\rm a}$	$53.3\pm6.6~^{a}$	$53.3\pm17.6~^{ab}$	$38.7\pm6.7^{\text{ bc}}$	II
C. maculatus	2.5	40.0 ± 11.5 $^{\rm a}$	$33.3\pm13.3~^{a}$	$53.3\pm17.6~^{\rm a}$	60.0 ± 11.5 $^{\rm a}$	$60.0\pm11.5~^{\rm ab}$	$49.3\pm5.4~^{\rm abc}$	III
-	3.5	$46.7\pm6.6~^{\rm a}$	$53.3\pm13.3~^{\rm a}$	$60.0\pm11.5~^{\rm a}$	80.0 ± 11.5 $^{\rm a}$	$86.7\pm6.6~^{a}$	65.3 ± 7.7 $^{\mathrm{ab}}$	IV
	5	40.0 ± 11.5 a	$60.0\pm19.9~^{\rm a}$	$80.0\pm11.5~^{\rm a}$	73.3 ± 13.3 $^{\rm a}$	$93.3\pm6.6~^{a}$	$69.3\pm9.1^{\rm a}$	IV
	F value	3.13	2.30	1.69	2.50	5.18	7.01	
	<i>p</i> value	0.06	0.12	0.22	0.10	0.01	0.001	
	^a d.f	4	4	4	4	4	20	

Table 4. Repellecy of CMEO against pests of stored grains at different exposure times.

^{a–c} Means having the same superscript show no significant variation (p < 0.05). ^ad.f: Degree of freedom, ^bSEM: Standard error of mean.

Table 5. Efficacy of the essential oil from *C. maxima* peels as a potential mosquito larvicidal agent (LC_{50} value expressed in $\mu g/mL$).

Mosquito	LC ₅₀ (µg/mL)	$\mathbf{Slope} \pm \mathbf{SEM}$	χ^2 (d.f)
Ar. subalbatus	76.24 ± 3.2	0.6566 ± 0.027	1.439 (3)
Ae. aegypti	47.07 ± 2.4	1.0643 ± 0.052	5.855 (3)
Cx. tritaeniorhynchus	58.04 ± 2.8	0.8630 ± 0.042	2.099 (3)

 Table 6. Effect of CMEO on the ability of wheat germination (% germination).

Duration (Hours)	% Germination in	C. maxima Essential Oil (µg/mL)			
Duration (mours)	Untreated Grains	50	100	250	
48	12.4 ± 1.0	13.1 ± 0.7 ⁿ	12.2 ± 0.9 ⁿ	11.9 ± 1.2 ⁿ	
72	30.7 ± 0.9	31.3 ± 0.6 ⁿ	30.8 ± 1.2 ⁿ	31.4 ± 1.2 ⁿ	
96	58.5 ± 1.5	59.2 ± 1.4 ⁿ	59.9± 1.1 ⁿ	58.4 ± 2.1 ⁿ	
120	80.6 ± 1.7	82.1 ± 1.6 ⁿ	81.4 ± 1.4 ⁿ	80.9 ± 2.1 ⁿ	
144	93.4 ± 2.1	$92.5\pm1.6\ ^{\rm n}$	$93.1\pm1.5^{\text{ n}}$	$92.4\pm2.5^{\text{ n}}$	

(ⁿ indicate no significant variation with untreated grains at p < 0.05).

3.7. Non-Targeted Organism Toxicity of CMEO

_

As indicated by the Table 7, no significant toxicity signs were observed in *P. reticulata* in terms of mortality, time spent on the top of water, or changes in body coloration. It is therefore confirmed that the CMEO has no toxic effects on the tested dose on non-targeted organisms.

Table 7. Effect of CMEO on the non-targeted organism (*P. reticulata*) and changes in the signs and symptoms of toxicity, as well as the mortality rate.

CMEO Dose (µg/mL)	% Mortality	Number of Fish Having Swimming Difficulty	Fishes Having Any Changes in Color	Total Time Spent on Top of Wate (Seconds)
0	0	0	0	27.6 ± 3.0
50	0	0	0	27.4 ± 3.0
100	0	0	0	29.7 ± 2.0
200	0	0	0	26.8 ± 3.0
250	0	0	0	29.5 ± 3.0

4. Discussion

Many studies have reported that essential oils exhibit repellent and insecticidal effects against storage pests, and therefore may be considered an substitute method for managing stored grain insect pests [38,39]. In traditional medicine, essential oils were used to prevent pest infestations by extracting the oils from aromatic plants [40]. Due to their efficacy, economic value, and storage capabilities, these essential oils might be more effective than chemical pesticides for grain protection [41]. The FDA (Food and Drug Administration) recognizes the safety of botanicals over synthetic pesticides [13]. Since aromatic essential oils are highly volatile, they can control stored-product pests through contact, fumigant, and repellent actions [40,42]. In this respect, using essential oils offers an environmentally friendly and food-safe alternative to chemical insecticides.

The results indicated that essential oils of *C. maxima* peels were rich in D-limonene (33.61%), β -sitosterol (17.99%), α -sitosterol (12.19%), stigmasterol (5.22%) and α -pinene (4.32%). Furthermore, several reports showed that *C. maxima* peels are rich in D-limonene and α -pinene as their major chemical constituents [43–45]. The difference in oil yields and chemical composition could be explained by genotypic variations, environmental factors, and external factors such as climate, collection time, extraction methods and soil composition [46,47].

Essential oils derived from citrus wastes possess immense potential in the management of various insect pests and vectors; *C. limon* and *C. sinensis* were effective against *Aedes albopictus* [48], whereas, de Andrade Dutra, et al. [49] and Ribeiro, et al. [50] reported that *C. latifolia*, *C. reticulata*, and *Citrus sinensis* controlled the growth of *C. maculatus* and *Tetranychus urticae*. *C. limetta* was capable of killing the larvae of *Anopheles stephensi* [51], and *C. aurantifolia* essential oil was effective against *Aedes aegypti* [52]. Almost half of the orange material utilized for extraction of orange juice is discarded as waste [53]. Essential oils from citrus peels can be commercially formulated by utilizing nanotechnology [54]. Based on the essential oil yield from our study, one tonne of *C. maxima* peels can yield 5.8 litres of essential oil. The yield of CMEO reinforces its potential for exploitation as a commercial biopesticide. Essential oil yield obtained from *C. maxima* peels in the various studies ranges from 0.29–14.25% [55–57]. This variation can be due to the difference in extraction methods employed and genotypic variations, as well as abiotic factors.

Contact toxicity of analysis of *C. maxima* peels revealed a higher toxicity towards C. maculatus adults. Similar studies have also indicated the potential of C. medica essential oil as a contact toxic biopesticide against on *T. castaneum* adult insects [58]. Furthermore, some studies have shown that the essential oils of *Citrus* spp. had high contact toxicity against stored grain pests such as C. maculatus [49] and Sitophilus zeamais [59]. The CMEO revealed higher contact lethality to warehouse grain products than essential oils of Pelargonium graveolens [60], Artemisia annua [61] and Melissa officinalis [62]. In stored product insects, different constituents of essential oils may cause different toxic reactions due to the variance in the chemical configuration of CMEO and their weight and cuticle thickness. Limonene and α -pinene were major monoterpenes presented in *C. maxima* peels essential oil. Both these secondary metabolites are reported to have insecticidal [63], fumigant [64] and repellent properties from other citrus species [65]. For stored grain insects, cuticle damage caused by hydrocarbon absorption and abrasion can also result in death due to desiccation [66]. This study found that both T. castaneum and C. maculatus adults experienced toxicity from the bioactive molecules in the tested essential oils. This is because they may interact with target sites in the nervous system, respiratory system, and cutaneous system of insects [66].

The fumigant toxicity analysis indicated that CMEO showed significant mortality against pests of warehouse grains. However, CMEO's were more toxic to *C. maculatus* adults than to *T. castaneum* adults. It has already reported that the essenitla oil of *C. sinensis* exerts higher sensitivity towards *C. maculatus* than *T. confusum* [67]. The results were not surprising as several studies have demonstrated *Citrus* spp. essential oil's fumigant properties against *T. castaneum* [19]. *C. maculatus* [49,68], *Oryzaephilus surinamensis* and

S. zeamais [69]. This result is in line with the fumigant effect of *Citrus reticulata* against *Cryptolestes ferrugineus* adults. Similarly, de Andrade Dutra, de Oliveira, Navarro and Santos [49] indicated the fumigant toxicity of different *Citrus* spp. against *C. maculatus* adults. We found that the LC_{50} value of *C. maxima* was 3.38 mg/L air against *C. maculatus* adults, which was better than that of the fumigant toxicity of essential oil from other *Citrus* spp.

Besides, the repellency of CMEO was tested against warehouse pests, and both test insects were repelled from essential oils from *C. maxima* peels. The essential oils extracted from *C. maxima* peels proved to be effective repellants against insect pests. Both *T. castaneum* and *C. maculatus* adults were repellent to the essential oils in a concentration-dependent manner. *Citrus* spp. have also been shown to repel storage insect pests such as the *T. castaneum* [58], *S. zeamais*, and *C. maculatus* [68]. Accordingly, the repellency of CMEO can be endorsed to the existence of D-limonene and α -pinene, which have been reported to be repellents [58,70]. The bioactivity of individual essential oil components of *C. maxima*, as well as their interactions with storage pests, should be further evaluated. In addition, the level of repellency has a direct effect on the reduction of oviposition, as well as the development of adults [71]. Hence, the essential oil of *C. maxima* can be used as a repellent biopesticide against pests of warehouse grains.

The majority of the bioactive chemicals found in essential oils had significant effects on non-target fishes, including fishes, or on human health or the environment. These essential oils have a higher efficiency and could serve as biopesticides in the future, as well as serving as better tools for integrated pest management because of their eco-friendliness [72].

5. Conclusions

The study concludes that the peel essential oil extracted from the *Citrus maxima* is highly useful as a biologically safe pesticide against the common stored grain insect pests *T. castaneum* and *C. maculatus*. Further, potent larvicidal toxicity against different mosquitoe species was observed, which may increase the economic value of the essential oil. Considering also the non-toxic nature on germinating grains and non-targeted organisms like fishes extends the possible application of the essential oil as a biologically safe green pesticide, even in the storage of food and seed grains.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/insects13050480/s1. Table S1: Efficacy of contact toxicity of CMEO against *T. castaneum* and *C. maculatus* at different exposure times; Table S2: Efficacy of fumigant toxicity of CMEO against *T. castaneum* and *C. maculatus*.

Author Contributions: Conceptualization, B.P. and A.N.; methodology, B.P. and A.A.; software, N.U.V.; validation, N.U.V. and A.N.; formal analysis, N.U.V., V.R. and A.A.; investigation, N.U.V.; resources, B.P., A.A. and A.N.; data curation, B.P., A.A., V.R. and A.N.; writing—original draft preparation, N.U.V. and A.N.; writing—review and editing, B.P., A.N., V.R. and A.A.; visualization, B.P.; supervision, B.P.; project administration, B.P.; funding acquisition, A.A. All authors have read and agreed to the published version of the manuscript.

Funding: The study was funded by King Saud University, Riyadh, Saudi Arabia through Researchers Supporting Project No: RSP 2022/11.

Institutional Review Board Statement: Not applicable.

Data Availability Statement: Data may be made available on valid request.

Acknowledgments: The authors acknowledge King Saud University, Riyadh, Saudi Arabia for funding this research through the Researchers Supporting Project No: RSP 2022/11.

Conflicts of Interest: There is no conflict of interest in the present study.

References

- Gomes-Araújo, R.; Martínez-Vázquez, D.G.; Charles-Rodríguez, A.V.; Rangel-Ortega, S.; Robledo-Olivo, A. Bioactive Compounds from Agricultural Residues, Their Obtaining Techniques, and the Antimicrobial Effect as Postharvest Additives. *Int. J. Food Sci.* 2021, 2021, 9936722. [CrossRef] [PubMed]
- Vasconcelos, N.G.; Silva, K.E.; Croda, J.; Simionatto, S. Antibacterial activity of *Cinnamomum cassia* L. essential oil in a carbapenemand polymyxin-resistant Klebsiella aerogenes strain. *Rev. Soc. Bras. Med. Trop.* 2020, 53, e20200032. [CrossRef] [PubMed]
- 3. Silva, J.K.R.D.; Figueiredo, P.L.B.; Byler, K.G.; Setzer, W.N. Essential Oils as Antiviral Agents. Potential of Essential Oils to Treat SARS-CoV-2 Infection: An In-Silico Investigation. *Int. J. Mol. Sci.* 2020, *21*, 3426. [CrossRef] [PubMed]
- 4. Chaudhari, A.K.; Singh, V.K.; Dwivedy, A.K.; Das, S.; Upadhyay, N.; Singh, A.; Dkhar, M.S.; Kayang, H.; Prakash, B.; Dubey, N.K. Chemically characterised *Pimenta dioica* (L.) Merr. essential oil as a novel plant based antimicrobial against fungal and aflatoxin B1 contamination of stored maize and its possible mode of action. *Nat. Prod. Res.* 2020, 34, 745–749. [CrossRef]
- 5. Wang, L.; Zhang, K.; Zhang, J.; Fu, J.; Li, J.; Wang, G.; Qiu, Z.; Wang, X. Antibacterial Activity of Cinnamomum camphora Essential Oil on Escherichia coli During Planktonic Growth and Biofilm Formation. *Front. Microbiol.* **2020**, *11*, 561002. [CrossRef]
- 6. Wijesinghe, G.K.; Maia, F.C.; de Oliveira, T.R.; de Feiria, S.N.B.; Joia, F.; Barbosa, J.P.; Boni, G.C.; Sardi, J.C.O.; Rosalen, P.L.; Hofling, J.F. Effect of Cinnamomum verum leaf essential oil on virulence factors of *Candida* species and determination of the in-vivo toxicity with *Galleria mellonella* model. *Mem. Do Inst. Oswaldo Cruz* 2020, 115, e200349. [CrossRef]
- Yap, P.S.; Lim, S.H.; Hu, C.P.; Yiap, B.C. Combination of essential oils and antibiotics reduce antibiotic resistance in plasmidconferred multidrug resistant bacteria. *Phytomed. Int. J. Phytother. Phytopharm.* 2013, 20, 710–713. [CrossRef]
- 8. Gaire, S.; O'Connell, M.; Holguin, F.O.; Amatya, A.; Bundy, S.; Romero, A. Insecticidal Properties of Essential Oils and Some of Their Constituents on the Turkestan Cockroach (Blattodea: Blattidae). *J. Econ. Entomol.* **2017**, *110*, 584–592. [CrossRef]
- 9. Francikowski, J.; Baran, B.; Cup, M.; Janiec, J.; Krzyżowski, M. Commercially Available Essential Oil Formulas as Repellents Against the Stored-Product Pest *Alphitobius diaperinus*. *Insects* **2019**, *10*, 96. [CrossRef]
- 10. Aburjai, T.; Mansi, K.; Azzam, H.; Alqudah, D.A.; Alshaer, W.; Abuirjei, M. Chemical Compositions and Anticancer Potential ofEssential Oil from Greenhouse-cultivated *Ocimum basilicum* Leaves. *Indian J. Pharm. Sci.* **2020**, *82*, 179–184. [CrossRef]
- Lee, S.C.; Xu, W.X.; Lin, L.Y.; Yang, J.J.; Liu, C.T. Chemical composition and hypoglycemic and pancreas-protective effect of leaf essential oil from indigenous cinnamon (*Cinnamomum osmophloeum* Kanehira). J. Agric. Food Chem. 2013, 61, 4905–4913. [CrossRef] [PubMed]
- 12. Lin, L.Y.; Chuang, C.H.; Chen, H.C.; Yang, K.M. Lime (*Citrus aurantifolia* (Christm.) Swingle) Essential Oils: Volatile Compounds, Antioxidant Capacity, and Hypolipidemic Effect. *Foods* **2019**, *8*, 398. [CrossRef] [PubMed]
- 13. Regnault-Roger, C.; Vincent, C.; Arnason, J.T. Essential oils in insect control: Low-risk products in a high-stakes world. *Annu. Rev. Entomol.* 2012, 57, 405–424. [CrossRef] [PubMed]
- Deng, M.; Jia, X.; Dong, L.; Liu, L.; Huang, F.; Chi, J.; Ma, Q.; Zhao, D.; Zhang, M.; Zhang, R. Structural elucidation of flavonoids from Shatianyu (*Citrus grandis* L. Osbeck) pulp and screening of key antioxidant components. *Food Chem.* 2022, 366, 130605. [CrossRef]
- 15. Somanathan Karthiga, R.; Sukhdeo, S.V.; Madhugiri Lakshminarayan, S.; Mysuru Nanjarajurs, S. Efficacy of Citrus maxima fruit segment supplemented paranthas in STZ induced diabetic rats. *J. Food Sci.* **2021**, *86*, 2091–2102. [CrossRef]
- 16. Tsai, M.L.; Lin, C.D.; Khoo, K.A.; Wang, M.Y.; Kuan, T.K.; Lin, W.C.; Zhang, Y.N.; Wang, Y.Y. Composition and Bioactivity of Essential Oil from *Citrus grandis* (L.) Osbeck 'Mato Peiyu' Leaf. *Molecules* **2017**, *22*, 2154. [CrossRef]
- 17. Oboh, G.; Ademosun, A.O.; Olumuyiwa, T.A.; Olasehinde, T.A.; Ademiluyi, A.O.; Adeyemo, A.C. Insecticidal activity of essential oil from orange peels (*Citrus sinensis*) against Tribolium confusum, Callosobruchus maculatus and Sitophilus oryzae and its inhibitory effects on acetylcholinesterase and Na+/K+-ATPase activities. *Phytoparasitica* **2017**, *45*, 501–508. [CrossRef]
- Campolo, O.; Malacrinò, A.; Zappalà, L.; Laudani, F.; Chiera, E.; Serra, D.; Russo, M.; Palmeri, V. Fumigant bioactivity of fiveCitrusessential oils againstTribolium confusum. *Phytoparasitica* 2014, 42, 223–233. [CrossRef]
- 19. Lou, H.; Huang, Y.; Ouyang, Y.; Zhang, Y.; Xi, L.; Chu, X.; Wang, Y.; Wang, C.; Zhang, L. Artemisia annua-sublingual immunotherapy for seasonal allergic rhinitis: A randomized controlled trial. *Allergy* **2020**, *75*, 2026–2036. [CrossRef]
- Giatropoulos, A.; Papachristos, D.P.; Kimbaris, A.; Koliopoulos, G.; Polissiou, M.G.; Emmanouel, N.; Michaelakis, A. Evaluation
 of bioefficacy of three Citrus essential oils against the dengue vector *Aedes albopictus* (Diptera: Culicidae) in correlation to their
 components enantiomeric distribution. *Parasitol. Res.* 2012, 111, 2253–2263. [CrossRef]
- 21. Michaelakis, A.; Papachristos, D.; Kimbaris, A.; Koliopoulos, G.; Giatropoulos, A.; Polissiou, M.G. Citrus essential oils and four enantiomeric pinenes against *Culex pipiens* (Diptera: Culicidae). *Parasitol. Res.* **2009**, *105*, 769–773. [CrossRef] [PubMed]
- 22. Adhikari, K.; Sarma, R.; Rabha, B.; Khanikor, B. Repellent Activity of Citrus Essential Oils and Two Constituent Compounds Against *Aedes aegypti. Proc. Natl. Acad. Sci. USA* 2022, *in press.* [CrossRef]
- Jiang, J.; Shan, L.; Chen, Z.; Xu, H.; Wang, J.; Liu, Y.; Xiong, Y. Evaluation of antioxidant-associated efficacy of flavonoid extracts from a traditional Chinese medicine Hua Ju Hong (peels of *Citrus grandis* (L.) Osbeck). *J. Ethnopharmacol.* 2014, 158 Pt A, 325–330. [CrossRef] [PubMed]
- Oboh, G.; Ademosun, A.O. Shaddock peels (*Citrus maxima*) phenolic extracts inhibit alpha-amylase, alpha-glucosidase and angiotensin I-converting enzyme activities: A nutraceutical approach to diabetes management. *Diabetes Metab. Syndr.* 2011, 5, 148–152. [CrossRef] [PubMed]

- Ademosun, A.O.; Adebayo, A.A.; Popoola, T.V.; Oboh, G. Shaddock (*Citrus maxima*) peels extract restores cognitive function, cholinergic and purinergic enzyme systems in scopolamine-induced amnesic rats. *Drug Chem. Toxicol.* 2020, 45, 1073–1080. [CrossRef]
- 26. Kuo, P.C.; Liao, Y.R.; Hung, H.Y.; Chuang, C.W.; Hwang, T.L.; Huang, S.C.; Shiao, Y.J.; Kuo, D.H.; Wu, T.S. Anti-Inflammatory and Neuroprotective Constituents from the Peels of *Citrus grandis*. *Molecules* **2017**, *22*, 967. [CrossRef]
- Ou, M.C.; Liu, Y.H.; Sun, Y.W.; Chan, C.F. The Composition, Antioxidant and Antibacterial Activities of Cold-Pressed and Distilled Essential Oils of Citrus paradisi and *Citrus grandis* (L.) Osbeck. *Evid.-Based Complement. Altern. Med. Ecam* 2015, 2015, 804091. [CrossRef]
- Singh, P.; Shukla, R.; Prakash, B.; Kumar, A.; Singh, S.; Mishra, P.K.; Dubey, N.K. Chemical profile, antifungal, antiaflatoxigenic and antioxidant activity of Citrus maxima Burm. and *Citrus sinensis* (L.) Osbeck essential oils and their cyclic monoterpene, DL-limonene. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* 2010, *48*, 1734–1740. [CrossRef]
- 29. Clevenger, J.F. Apparatus for the Determination of Volatile Oil. J. Am. Pharm. Assoc. 1928, 17, 345–349. [CrossRef]
- Patiño-Bayona, W.R.; Nagles Galeano, L.J.; Bustos Cortes, J.J.; Delgado Ávila, W.A.; Herrera Daza, E.; Suárez, L.E.C.; Prieto-Rodríguez, J.A.; Patiño-Ladino, O.J. Effects of Essential Oils from 24 Plant Species on *Sitophilus zeamais* Motsch (Coleoptera, Curculionidae). *Insects* 2021, 12, 532. [CrossRef]
- 31. Bagade, R.P.; Jadhav, A.D.; Chavan, R.V. Toxicity and repellency of four plant essential oils against *Tribolium castaneum* (Herbst)(Coleoptera: Tenebrionidae). *Int. J. Trop. Insect Sci.* 2021, *41*, 1505–1512. [CrossRef]
- 32. Culver, J.J. The Value of the Dry Substitutes for Liquid Lime Sulphur as a Control for San Jose Scale. J. Econ. Entomol. **1925**, 18, 265. [CrossRef]
- Denloye, A.; Makanjuola, W.; Teslim, O.; Alafia, O.; Kasali, A.; Eshilokun, A. Toxicity of *Chenopodium ambrosioides* L.(Chenopodiaceae) products from Nigeria against three storage insects. *J. Plant Prot. Res.* 2010, 50, 379–384. [CrossRef]
- Bai, L.; Jiao, M.L.; Zang, H.Y.; Guo, S.S.; Wang, Y.; Sang, Y.L.; Du, S.S. Chemical composition of essential oils from four Rhododendron species and their repellent activity against three stored-product insects. *Environ. Sci. Pollut. Res. Int.* 2019, 26, 23198–23205. [CrossRef]
- Ebrahimifar, J.; Jamshidnia, A.; Sadeghi, R.; Ebadollahi, A. Repellency of *Ferulago angulata* (Schlecht.) Boiss essential oil on two major stored-product insect pests without effect on wheat germination. *Int. J. Trop. Insect Sci.* 2021, 41, 217–223. [CrossRef]
- Narayanankutty, A.; Kuttithodi, A.M.; Alfarhan, A.; Rajagopal, R.; Barcelo, D. Chemical Composition, Insecticidal and Mosquito Larvicidal Activities of Allspice (*Pimenta dioica*) Essential Oil. *Molecules* 2021, 26, 6698. [CrossRef]
- 37. Gopinath, P.P.; Parsad, R.; Joseph, B.; Adarsh, V. grapesAgri1: Collection of Shiny Apps for Data Analysis in Agriculture. *J. Open Source Softw.* **2021**, *6*, 3437. [CrossRef]
- Kłyś, M.; Malejky, N.; Nowak-Chmura, M. The repellent effect of plants and their active substances against the beetle storage pests. J. Stored Prod. Res. 2017, 74, 66–77. [CrossRef]
- Rajendran, S.; Sriranjini, V. Plant products as fumigants for stored-product insect control. J. Stored Prod. Res. 2008, 44, 126–135. [CrossRef]
- 40. Isman, M.B. Bioinsecticides based on plant essential oils: A short overview. Z. Naturforsch. C. 2020, 75, 179–182. [CrossRef]
- Gueye, M.T.; Cissokho, P.S.; Goergen, G.; Ndiaye, S.; Seck, D.; Gueye, G.; Wathelet, J.-P.; Lognay, G. Efficacy of powdered maize cobs against the maize weevil *Sitophilus zeamais* (Coleoptera: Curculionidae) in stored maize in Senegal. *Int. J. Trop. Insect Sci.* 2012, 32, 94–100. [CrossRef]
- Nerio, L.S.; Olivero-Verbel, J.; Stashenko, E. Repellent activity of essential oils: A review. *Bioresour. Technol.* 2010, 101, 372–378. [CrossRef] [PubMed]
- 43. Dao, P.T.; Tran, N.Y.T.; Tran, Q.N.; Bach, G.L.; Lam, T.V. Kinetics of pilot-scale essential oil extraction from pomelo (Citrus maxima) peels: Comparison between linear and nonlinear models. *Alex. Eng. J.* **2022**, *61*, 2564–2572. [CrossRef]
- Chen, L.; Teng, H.; Xie, Z.; Cao, H.; Cheang, W.S.; Skalicka-Woniak, K.; Georgiev, M.I.; Xiao, J. Modifications of dietary flavonoids towards improved bioactivity: An update on structure-activity relationship. *Crit. Rev. Food Sci. Nutr.* 2018, 58, 513–527. [CrossRef] [PubMed]
- 45. Chen, G.-W.; Lin, Y.-H.; Lin, C.-H.; Jen, H.-C. Antibacterial Activity of Emulsified Pomelo (*Citrus grandis* Osbeck) Peel Oil and Water-Soluble Chitosan on *Staphylococcus aureus* and Escherichia coli. *Molecules* **2018**, *23*, 840. [CrossRef] [PubMed]
- Ocheng, F.; Bwanga, F.; Joloba, M.; Softrata, A.; Azeem, M.; Pütsep, K.; Borg-Karlson, A.-K.; Obua, C.; Gustafsson, A. Essential Oils from Ugandan Aromatic Medicinal Plants: Chemical Composition and Growth Inhibitory Effects on Oral Pathogens. *Evid.-Based Complement. Altern. Med.* 2015, 2015, 230832. [CrossRef] [PubMed]
- Pinto, Z.T.; Sánchez, F.F.; Santos, A.R.D.; Amaral, A.C.F.; Ferreira, J.L.P.; Escalona-Arranz, J.C.; Queiroz, M.M.D.C. Chemical composition and insecticidal activity of *Cymbopogon citratus* essential oil from Cuba and Brazil against housefly. *Rev. Bras. De Parasitol. Veterinária* 2015, 24, 36–44. [CrossRef] [PubMed]
- Akram, W.; Khan, H.A.A.; Hussain, A.; Hafeez, F. Citrus Waste-Derived Essential Oils: Alternative Larvicides for Dengue Fever Mosquito, *Aedes albopictus* (Skuse)(Culicidae: Diptera). *Pak. J. Zool.* 2011, 43, 367–372.
- de Andrade Dutra, K.; de Oliveira, J.V.; Navarro, D.M.D.A.F.; Santos, J.P.O. Control of *Callosobruchus maculatus* (FABR.)(Coleoptera: Chrysomelidae: Bruchinae) in *Vigna unguiculata* (L.) WALP. with essential oils from four *Citrus* spp. plants. *J. Stored Prod. Res.* 2016, 68, 25–32. [CrossRef]

- 50. Ribeiro, N.C.; da Camara, C.A.G.; Melo, J.P.R.; de Moraes, M.M. Acaricidal properties of essential oils from agro-industrial waste products from citric fruit against *Tetranychus urticae*. J. Appl. Entomol. 2019, 143, 731–743. [CrossRef]
- Kumar, S.; Warikoo, R.; Mishra, M.; Seth, A.; Wahab, N. Larvicidal efficacy of the Citrus limetta peel extracts against Indian strains of Anopheles stephensi Liston and Aedes aegypti L. Parasitol. Res. 2012, 111, 173–178. [CrossRef]
- Sarma, R.; Adhikari, K.; Mahanta, S.; Khanikor, B. Insecticidal activities of *Citrus aurantifolia* essential oil against *Aedes aegypti* (Diptera: Culicidae). *Toxicol. Rep.* 2019, 6, 1091–1096. [CrossRef] [PubMed]
- 53. Chavan, P.; Singh, A.K.; Kaur, G. Recent progress in the utilization of industrial waste and by-products of citrus fruits: A review. *J. Food Process Eng.* **2018**, *41*, e12895. [CrossRef]
- 54. Azmy, R.M. Nanoformulated Materials from Citrus Wastes. In *Waste Recycling Technologies for Nanomaterials Manufacturing;* Springer: Berlin/Heidelberg, Germany, 2021; pp. 649–669.
- 55. Baccati, C.; Gibernau, M.; Paoli, M.; Ollitrault, P.; Tomi, F.; Luro, F. Chemical Variability of Peel and Leaf Essential Oils in the Citrus Subgenus Papeda (Swingle) and Few Relatives. *Plants* **2021**, *10*, 1117. [CrossRef] [PubMed]
- 56. Thavanapong, N.; Wetwitayaklung, P.; Charoenteeraboon, J. Comparison of Essential Oils Compositions of *Citrus maxima* Merr. Peel Obtained by Cold Press and Vacuum Stream Distillation Methods and of Its Peel and Flower Extract Obtained by Supercritical Carbon Dioxide Extraction Method and Their Antimicrobial Activity. J. Essent. Oil Res. 2010, 22, 71–77.
- Ngan, T.; Muoi, N.; Quan, P.; Cang, M. Evaluation of physical and chemical properties of pomelo (*Citrus grandis* L.) essential oil using steam distillation process. *Asian J. Chem.* 2020, *6*, 1433–1436. [CrossRef]
- Luo, C.; Li, D.; Wang, Y.; Guo, S.; Zhang, D.; Du, S. Chemical Composition and Insecticide Efficacy of Essential Oils from *Citrus medica* L. var. sarcodactylis Swingle Against Tribolium castaneum Herbst in Stored Medicinal Materials. *J. Essent. Oil Bear. Plants* 2019, 22, 1182–1194. [CrossRef]
- Oyedeji, A.O.; Okunowo, W.O.; Osuntoki, A.A.; Olabode, T.B.; Ayo-Folorunso, F. Insecticidal and biochemical activity of essential oil from *Citrus sinensis* peel and constituents on *Callosobrunchus maculatus* and *Sitophilus zeamais*. *Pestic. Biochem. Physiol.* 2020, 168, 24. [CrossRef]
- Abouelatta, A.M.; Keratum, A.Y.; Ahmed, S.I.; El-Zun, H.M. Repellent, contact and fumigant activities of geranium (*Pelargonium graveolens* L.'Hér) essential oils against *Tribolium castaneum* (Herbst) and *Rhyzopertha dominica* (F.). *Int. J. Trop. Insect Sci.* 2020, 40, 1021–1030. [CrossRef]
- 61. Deb, M.; Kumar, D. Bioactivity and efficacy of essential oils extracted from *Artemisia annua* against *Tribolium casteneum* (Herbst. 1797) (Coleoptera: Tenebrionidae): An eco-friendly approach. *Ecotoxicol Env. Saf* **2020**, *189*, 22. [CrossRef]
- Upadhyay, N.; Singh, V.K.; Dwivedy, A.K.; Das, S.; Chaudhari, A.K.; Dubey, N.K. Assessment of *Melissa officinalis* L. essential oil as an eco-friendly approach against biodeterioration of wheat flour caused by *Tribolium castaneum* Herbst. *Environ. Sci. Pollut. Res. Int.* 2019, 26, 14036–14049. [CrossRef] [PubMed]
- 63. Papachristos, D.P.; Kimbaris, A.C.; Papadopoulos, N.T.; Polissiou, M.G. Toxicity of citrus essential oils against *Ceratitis capitata* (Diptera: Tephritidae) larvae. *Ann. Appl. Biol.* **2009**, *155*, 381–389. [CrossRef]
- 64. Rossi, Y.E.; Palacios, S.M. Fumigant toxicity of *Citrus sinensis* essential oil on *Musca domestica* L. adults in the absence and presence of a P450 inhibitor. *Acta Trop.* **2013**, 127, 33–37. [CrossRef]
- 65. Yoon, C.; Kang, S.-H.; Yang, J.-O.; Noh, D.-J.; Indiragandhi, P.; Kim, G.-H. Repellent activity of citrus oils against the cockroaches *Blattella germanica*, *Periplaneta americana* and *P. fuliginosa*. J. Pestic. Sci. 2009, 34, 77–88. [CrossRef]
- Devi, M.A.; Sahoo, D.; Singh, T.B.; Rajashekar, Y. Toxicity, repellency and chemical composition of essential oils from *Cymbopogon* species against red flour beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *J. Consum. Prot. Food Saf.* 2020, 15, 181–191. [CrossRef]
- 67. Tandorost, R.; Karimpour, Y. Evaluation of fumigant toxicity of orange peel *Citrus sinensis* (L.) essential oil against three stored product insects in laboratory condition. *Munis Entomol. Zool.* **2012**, *7*, 352–358.
- 68. Oyedeji, A.O.; Afolayan, A.J.; Hutchings, A. Compositional variation of the essential oils of *Artemisia afra* Jacq. from three provinces in South Africa—A case study of its safety. *Nat. Prod. Commun.* **2009**, *4*, 849–852. [CrossRef]
- Hernandez-Lambraño, R.; Pajaro-Castro, N.; Caballero-Gallardo, K.; Stashenko, E.; Olivero-Verbel, J. Essential oils from plants of the genus *Cymbopogon* as natural insecticides to control stored product pests. J. Stored Prod. Res. 2015, 62, 81–83. [CrossRef]
- Fouad, H.A.; da Camara, C.A. Chemical composition and bioactivity of peel oils from *Citrus aurantiifolia* and *Citrus reticulata* and enantiomers of their major constituent against *Sitophilus zeamais* (Coleoptera: Curculionidae). J. Stored Prod. Res. 2017, 73, 30–36. [CrossRef]
- 71. França, S.M.D.; Oliveira, J.V.D.; Esteves Filho, A.B.; Oliveira, C.M.D. Toxicity and repellency of essential oils to *Zabrotes subfasciatus* (Boheman)(Coleoptera, Chrysomelidae, Bruchinae) in *Phaseolus vulgaris* L. *Acta Amaz.* **2012**, *42*, 381–386. [CrossRef]
- 72. Chaubey, M.K. Essential oils as green pesticides of stored grain insects. Eur. J. Biol. Res. 2019, 9, 202–244.