

Article

Laboratory Evidence of 2-Isobutyl-3-methoxypyrazine as a Male-Released Aggregative Cue in *Labidostomis lusitanica* (Germar) (Coleoptera: Chrysomelidae)

Sergio López ^{1,*}, Sara Rodrigo-Gómez ², Enrique Fernández-Carrillo ², Clàudia Corbella-Martorell ¹ and Carmen Quero ^{1,*}

¹ Department of Biological Chemistry, Institute for Advanced Chemistry of Catalonia (IQAC-CSIC), Jordi Girona 18-26, 08034 Barcelona, Spain

² Instituto Regional de Investigación y Desarrollo Agroalimentario y Forestal (IRIAF)-Centro de Investigación Agroambiental "El Chaparrillo", 13071 Ciudad Real, Spain

* Correspondence: sergio.lopez@iqac.csic.es (S.L.); carme.quero@iqac.csic.es (C.Q.)

Simple Summary: The leaf beetle *Labidostomis lusitanica* (Germar) (Coleoptera: Chrysomelidae) is a voracious defoliator beetle regarded as a threat for pistachio (*Pistacia vera*) crops in Spain. During late April–early June, field aggregates containing both sexes are commonly found in pistachio leaves. Males collected from these aggregates release a sex-specific compound, namely 2-isobutyl-3-methoxypyrazine, that elicits both a strong electrophysiological response and positive chemotactic behavior in males and females. Altogether, our findings suggest that 2-isobutyl-3-methoxypyrazine may act as an aggregative cue, although further field assays are required to address the true role of the compound in natural conditions.

Abstract: In spite of its incidence on pistachio trees, the chemical ecology of *Labidostomis lusitanica* (Germar) (Coleoptera: Chrysomelidae) has been neglected so far. In this work, we provide the first evidence of a biologically active male-specific compound that may be promoting field aggregation. Headspace collections through solid-phase microextraction from feral males and females reported the presence of 2-isobutyl-3-methoxypyrazine exclusively in males. Electroantennographic recordings revealed that males and females responded in a dose-dependent manner to increasing stimuli of 2-isobutyl-3-methoxypyrazine, with females overall displaying a higher response than males. In dual-choice tests, both males and females showed a significant preference for the compound in comparison to a pure air stimulus. In light of these results, the possible role of 2-isobutyl-3-methoxypyrazine as an aggregation cue in *L. lusitanica* is discussed.

Keywords: *Labidostomis lusitanica*; Chrysomelidae; *Pistacia vera*; 2-isobutyl-3-methoxypyrazine; electroantennography; behavior

Citation: López, S.; Rodrigo-Gómez, S.; Fernández-Carrillo, E.; Corbella-Martorell, C.; Quero, C. Laboratory Evidence of 2-Isobutyl-3-methoxypyrazine as a Male-Released Aggregative Cue in *Labidostomis lusitanica* (Germar) (Coleoptera: Chrysomelidae). *Insects* **2023**, *14*, 107. <https://doi.org/10.3390/insects14020107>

Academic Editor: Aijun Zhang

Received: 16 December 2022

Revised: 13 January 2023

Accepted: 17 January 2023

Published: 19 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Pistachio (*Pistacia vera* L.) is a species native to central and southwestern Asia that is cultivated in some Mediterranean countries (Spain, Italy, Greece, Turkey, Tunisia), the Middle East (Iran, Syria), the United States (California and Arizona) and Australia [1–3]. Due to the quality of the edible nut, it is a species with a high commercial value worldwide. By the year 2003, pistachio nut production was ranked in sixth place in world tree nut production behind almond, walnut, cashew, hazelnut and chestnut [4], and currently only almond and walnut production surpass that of pistachio, with the United States (47%) as the major producer country, followed by Turkey (30%) and Iran (19%) [5]. Such relevance of *P. vera* as a highly exploited resource stresses the need of an exhaustive

knowledge and awareness about those biotic agents (i.e., pathogens and arthropod pests) that may affect cultivated plantations.

Diverse works have focused on describing the arthropods pests of pistachio in different countries [6–10]. Among the most damaging species are, for instance, the pistachio seed wasps *Eurytoma plotnikovi* Nikol'skaya (Hymenoptera: Eurytomidae) and *Megastigmus pistaciae* Walker (Hymenoptera: Torymidae) [11,12], the bark beetle *Chaetoptelius vestitus* (Mulsant and Rey) (Coleoptera: Curculionidae) [13], the pistachio psylla *Agonoscena pistaciae* Burckhardt and Lauterer (Hemiptera: Psyllidae) [14] and the lepidopteran species *Choristoneura rosaceana* (Harris) (Lepidoptera: Tortricidae), *Amyelois transitiella* (Walker) (Lepidoptera: Pyralidae) and *Kermania pistaciella* Amsel (Lepidoptera: Tineidae) [7,15].

In Spain, the European country with the greatest pistachio crop area, there is still a lack of knowledge on the insect pests that may threaten pistachio production [8,16]. Recently, a comprehensive study conducted by Gómez and coworkers highlighted the leaf beetle *Labidostomis lusitanica* (Germar) (Coleoptera: Chrysomelidae) as a potential serious threat for pistachio production in the Iberian Peninsula [17]. *L. lusitanica* is a polyphagous leaf beetle that feeds not only on pistachio, but also on *Quercus* L., *Salix* L. and *Populus* L., and on the herbaceous genera *Rumex* L. and *Polygonum* L. In pistachio plantations, it is considered a voracious species that may defoliate young trees in a few hours [17]. The species is found mostly in the eastern and southern part of the Iberian Peninsula, overlapping with pistachio crops [17]. During late April–early June, evenly distributed aggregates containing both sexes are commonly found, in which they feed, mate and lay eggs for a period of 4–5 weeks, and, subsequently, the individuals disperse across the crop. This behavior led us to suspect the existence of a possible aggregation pheromone in the species.

Hence, the aim of the current work was to isolate and identify the intraspecific chemical cues mediating this aggregation. For this purpose, we (a) characterized the volatile profile of both aggregative males and females, (b) measured the peripheral olfaction of a male-specific compound by means of electroantennographic recordings and, finally, (c) assessed the behavioral response of both sexes to the compound under laboratory conditions. Determining those chemical cues involved in the field aggregation of the insect would provide valuable knowledge for the further development of pheromone-based monitoring or mass trapping tools, in order to prevent the noxious effect of *L. lusitanica* on pistachio crops.

2. Materials and Methods

2.1. Insects

Feral aggregated *L. lusitanica* males and females were collected from 6th to 24th May 2021 in an organic farming-based pistachio orchard (0.36 ha) located in “El Chaparrillo” (39.0040, -3.9629; Ciudad Real, Spain). Immediately after collection, they were segregated by sex and sent to the facilities of the Institute for Advanced Chemistry of Catalonia, where each sex was kept separately inside plastic cubic cages (Bugdorm[®], 30 × 30 × 30 cm; Entomopraxis, Barcelona, Spain) at 25 ± 1 °C, 55 ± 5% RH and 16:8 L:D. Insects were fed on fresh pistachio leaves (var. Kerman), and leaves were renewed every two days.

2.2. Chemicals

2-isobutyl-3-methoxypyrazine (97%) was obtained from Tokyo Chemical Industry-Europe (Zwijndrecht, Belgium). A commercial mixture of straight-chain n-alkanes (C₁₂-C₆₀, Qualitative Retention Time Mix, Supelco) for calculating Kovats retention indices (KI) was acquired from Merck-Sigma Aldrich (Madrid, Spain). For the preparation of the corresponding dilutions of n-alkanes and 2-isobutyl-3-methoxypyrazine, n-hexane (GC purity, SupraSolv[®], Merck, Darmstadt, Germany) was used as solvent.

2.3. Headspace Solid-Phase Microextraction (HS-SPME)

The volatile profile of *L. lusitanica* males and females were analyzed separately through solid-phase microextraction (SPME), using a divinylbenzene/carboxen/polydimethylsiloxane (DVB/CAR/PDMS)-coated fiber (50/30 μm ; Supelco, Merck-Sigma Aldrich, Madrid, Spain). In each volatile collection, 20 individuals, either males or females, were exposed for 6 h to the SPME fiber inside a 40 mL screw-cap vial (Supelco, Merck Sigma Aldrich,). All collections ($n = 10$ for both sexes) were conducted from 10:00 to 18:00 at room temperature and 24–72 h after the arrival of the insects.

Samples were further analyzed in a Thermo Finnigan Trace 2000 gas chromatograph system coupled to a Trace MS quadrupole mass spectrometer (Thermo Fisher Scientific, Madrid, Spain). The injection port temperature was set at 270 $^{\circ}\text{C}$, and samples were run in splitless mode (5 min) in a non-polar TR-5MS column (30 m \times 0.25 mm I.D. \times 0.25 μm ; Thermo Fisher Scientific). The following temperature program was set up: an initial temperature of 50 $^{\circ}\text{C}$ held for 5 min, followed by an increase of 5 $^{\circ}\text{C}/\text{min}$ to 150 $^{\circ}\text{C}$ and, finally, an increase to 310 $^{\circ}\text{C}$ at 15 $^{\circ}\text{C}/\text{min}$, with a hold time of 10 min. The MS system operated in electron impact mode (70 eV), and scan mode (40–500 amu), at 1.0 scan/s. Compound identification was achieved by comparison of the experimental mass spectrum with those of the synthetic standard and a commercial mass spectral library (NIST Registry of Mass Spectral Data, 2005), and by comparison of the KI values of both the male-released compound and the synthetic standard. For determining the KI values, first, a 50 ng/ μL hexane dilution of the n-alkane mixture was prepared and, subsequently, 1 μL of this dilution was injected under the same conditions as those set for analyzing the HS-SPME samples.

2.4. Electroantennographic (EAG) Response

The olfactory response of *L. lusitanica* males ($n = 8$ –10) and females ($n = 8$ –10) to increasing quantities of 2-isobutyl-3-methoxy-pyrazine (1, 10, and 100 μg) was determined using electrophysiological recordings from severed antennae. Briefly, one antenna of a cold-anesthetized adult was excised with the aid of a microscalpel, and once excised, last two antennomeres were removed. The antenna was then fixed to a forked microelectrode holder (Syntech, Kirchzarten, Germany) with a drop of conductive gel (Spectra 360, Parker Lab. Inc., Hellendoorn, The Netherlands), placing the distal part of the antenna on the recording microelectrode, and the proximal part on the reference microelectrode. The holder was connected to an EAG Combi-Probe (Syntech, Kirchzarten, Germany) connected to a MP-5 micromanipulator (Syntech). Stimuli were delivered to the insect antennae by applying pure air puffs (ca. 300 mL/min) for 100 ms with a disposable glass Pasteur pipette (150 mm long) that contained a filter paper disc (2.5 cm diameter, Whatman[®], Merck-Sigma Aldrich, Madrid, Spain) loaded with 10 μL of the corresponding hexane dilution of 2-isobutyl-3-methoxy-pyrazine (0.1, 1.0, or 10 $\mu\text{g}/\mu\text{L}$). Two puffs for each amount of compound were applied to each antennal preparation in increasing order of magnitude, with an interval of 60 s between puffs. Control puffs with 10 μL of hexane were intercalated between two consecutive 2-isobutyl-3-methoxy-pyrazine stimuli to determine the baseline depolarization of the antenna. A continuous humidified pure air flow (ca. 650 mL/min) passed over the antenna through the open end of a T-shaped glass tube (7 cm long \times 5 mm diameter) positioned 1 cm over the sample, in order to prolong the life of the antennal preparation. The net EAG response evoked by each amount of 2-isobutyl-3-methoxy-pyrazine was calculated by subtracting the mean EAG amplitudes of the hexane puffs before and after the test compound. The EAG signals were amplified (10x) with the EAG Combi-Probe, filtered (DC to 1 kHz) with the aid of an IDAC-2 interface (Syntech, Kirchzarten, Germany), digitized on a PC and further analyzed with EAG Pro software (version 2.0, Syntech, Kirchzarten, Germany).

2.5. Behavioral Bioassays

A vertically placed dual-choice glass olfactometer (main arm 10 cm long \times 18 mm I.D., arms 8 cm long \times 18 mm I.D., angle between arms 90 $^{\circ}$) was used to determine the

walking preference of *L. lusitanica* males ($n = 31\text{--}40$) and females ($n = 36\text{--}49$) in response to 2-isobutyl-3-methoxy pyrazine (0.1, 1, and 10 μg). A Whatman® filter paper disc loaded with 10 μL of the corresponding hexane dilution of 2-isobutyl-3-methoxy pyrazine was placed in one of the olfactometer arms, whereas the control arm held a filter paper loaded with 10 μL of hexane, as a control. Filter papers were replaced every five insects, and the position of the arms were also switched to avoid any directionality. An incoming charcoal-filtered air flow at 350 mL/min was set for both arms. The whole system was surrounded by a white filter paper screen (45 cm height) to avoid the interference of visual stimuli [18]. Homogenous illumination (light intensity ca. 500 lux) was provided with a bulb placed 30 cm above the junction of the olfactometer. A Y-shaped copper wire was introduced to facilitate insect movement along the olfactometer. Prior to the beginning of each trial, beetles were individually isolated in polystyrene cell culture dishes (Corning Inc, NY, USA), and left for 1 h to acclimate to room conditions. A positive response was considered if the insect entered any arm, walked at least 2 cm beyond the junction and remained there or continued walking until the end of the arm. Beetles that did not make a choice were designated as non-responders and, therefore, discarded for further statistical analysis. All the behavioral assays were conducted from 10:00 to 18:00 at room temperature.

2.6. Statistical Analysis

Prior to any statistical analysis, the EAG net amplitudes in response to 2-isobutyl-3-methoxy pyrazine were subjected to the Shapiro–Wilk and Levene tests and, when necessary, data were log-transformed to fulfill the assumptions of normality and/or homogeneity of variance. Differences in absolute EAG responses within a sex were analyzed using one-way analysis of variance (ANOVA). A subsequent Tukey’s HSD post hoc test was applied for pairwise comparisons when ANOVA was significant. Differences between sexes in response to a concrete amount of stimulus were analyzed with Student’s *t*-test. The walking response in the double-choice olfactometer was subjected to a chi-square goodness-of-fit test to address if the arm preference displayed by each sex differed from a 50:50 distribution. All the statistical tests were performed using SPSS Statistics 17.0 software (SPSS, Chicago, IL, USA), at a significance level of $\alpha = 0.05$.

3. Results

3.1. HS-SPME Collections

All the volatile collections from males ($n = 10$) revealed the presence of a sex-specific compound (KI = 1185), whereas no traces were detected in any of the samples from females ($n = 10$) (Figure 1a). The base peaks at m/z 124, resulting from a McLafferty rearrangement and the loss of C_3H_6 , at m/z 151, produced by the loss of a methyl group from the isobutyl chain, and a possible molecular ion at m/z 166 tentatively suggested its identification as 2-isobutyl-3-methoxy pyrazine [19]. Further comparison of the mass spectrum with those of the commercial library and the synthetic standard (KI = 1185) corroborated the identification of the eluting compound as 2-isobutyl-3-methoxy pyrazine (Figure 1b).




Figure 1. Identification of 2-isobutyl-3-methoxy pyrazine from HS-SPME collections of feral *L. lusitanica*. (a) Zoomed-in region of the elution time of the male-specific peak (black arrow); (b) mass spectra of male-released compound (upper panel) and the synthetic standard of 2-isobutyl-3-methoxy pyrazine (lower panel). Calculated Kovats index (KI) is detailed for each mass spectrum.

3.2. EAG Response

The EAG response of both sexes followed a dose-dependent response pattern (males, $F_{2,24} = 5.304$, $p = 0.012$; females, $F_{2,23} = 10.249$, $p = 0.001$) (Figure 2). In terms of differences between sexes, overall, females showed a higher sensitivity to increasing amounts of 2-isobutyl-3-methoxy pyrazine than males (Figure 2). A stimulus of 10 μg elicited a significantly higher EAG response in females than in males ($t = -3.571$, $df = 11.526$, $p = 0.04$), while the difference in the response to 100 μg was close to significance ($t = -2.023$, $df = 14$, $p = 0.063$) (Figure 2).




Figure 2. Mean EAG response (mV \pm SEM) from excised antennae of feral *L. lusitanica* males ($n = 8-10$) and females ($n = 8-10$) to puffed stimuli of 2-isobutyl-3-methoxy pyrazine (1, 10 and 100 μg). Different letters within each sex denote significant differences in the EAG response among quantities (one-way ANOVA followed by Tukey’s HSD post hoc test, at $\alpha = 0.05$). Asterisk indicates significant differences between sexes in the response to a concrete amount of 2-isobutyl-3-methoxy pyrazine (Student’s t -test, at $\alpha = 0.05$; π , $p = 0.067$).

3.3. Behavioral Bioassays

Both males and females responded positively to 2-isobutyl-3-methoxypyrazine when they were released individually in a double-choice olfactometer (Figure 3). Specifically, 68% of tested males made a choice for the arm containing 0.1 μg of the compound ($X^2 = 4.235$, $df = 1$, $p = 0.04$) (Figure 3a), while high amounts of 2-isobutyl-3-methoxypyrazine yielded non-significant attraction percentages. Indeed, a significant aversive effect was detected when males were exposed to 10 μg of the compound ($X^2 = 3.846$, $df = 1$, $p = 0.05$) (Figure 3a). A similar response pattern was observed in females, with 73% of these displaying a significant preference for 1 μg of the compound ($X^2 = 5.538$, $df = 1$, $p = 0.019$) (Figure 3b). Conversely, no significant trend towards 2-isobutyl-3-methoxypyrazine was detected at either 0.1 ($X^2 = 1.485$, $df = 1$, $p = 0.223$) or 10 μg ($X^2 = 0.059$, $df = 1$, $p = 0.808$) (Figure 3b).




Figure 3. Behavioral response of feral *L. lusitanica* males (a) and females (b) to 2-isobutyl-3-methoxypyrazine (IBMP, at 0.1, 1 and 10 μg) and pure air (blank) in a double-choice olfactometer. Numbers beside each bar indicate the total number of insects tested, and those which made a choice for IBMP and the blank arm (B) are indicated within parentheses. Asterisks denote a significant preference towards the compound, while the dagger (†) indicates an aversive effect of the compound (chi-square goodness-of-fit test, at $\alpha = 0.05$).

4. Discussion

Pyrazines are nitrogen-containing heterocyclic compounds with relevance in insect chemical signaling at different levels (i.e., intraspecific and interspecific interactions). For instance, some plant-related pyrazines have been described as synomones [20–23]. Especially remarkable is the sexual deception displayed by some orchid species, which mimic the sex pheromone blend of female pollinator wasps (Hymenoptera: Thynnidae) [20,21]. Diverse 2-alkyl-methoxypyrazines released from the flowers of palm species of the genera *Acrocomia* and *Attalea* attract two florivorous scarab species (Coleoptera: Melolonthidae) [22], while the cones from the African cycad *Encephalartos villosus* Lem. emit 2-isopropyl-

3-methoxypyrazine (hereafter referred to as IPMP), making them attractive to pollinator beetles [23]. Additionally, 2-alkyl-methoxypyrazines act as allomones in aposematic insects of different orders [24–27]. 2-Alkyl-methoxypyrazines are common warning cues with a distinctive odor that, along with additional signals (e.g., visual [28,29]), constitute a multimodal defense mechanism to avoid predation. To cite an example, the wood tiger moth *Arctia plantaginis* (L.) (Lepidoptera: Arctiidae) produces 2-isobutyl-3-methoxypyrazine (IBMP) and 2-*sec*-butyl-3-methoxypyrazine (SBMP), which deter birds and ants [30], although they are ineffective against spiders [31]. In *Oncopeltus fasciatus* (Dallas) (Hemiptera: Lygaeidae), its chemical defense mechanism relies on the expelling of a fluid that contains IBMP [27]. Particularly remarkable is the role of 2-alkyl-methoxypyrazines in some ladybug species (Coleoptera: Coccinellidae), since they have been coadapted as allomones and aggregation pheromones in diapausing individuals [32,33].

In this sense, pheromone-mediated aggregation is a common behavior in insects, including leaf beetles (Coleoptera: Chrysomelidae) [34], in which chemical cues clearly play a pivotal, although not exclusive, role. Indeed, the elucidation of aggregation pheromones has been widely documented in chrysomelid species, with males as the sex responsible for their emission [35–44]. Here, we present the first empirical evidence of IBMP as a male-specific volatile compound that may be mediating aggregation in the leaf beetle *L. lusitanica*. This hypothesis is partially supported by our findings, with males being the sex responsible for the IBMP emission, as reported in other leaf beetle species, and both sexes showing a significant preference towards the compound. To the best of our knowledge, no pyrazine has been identified as a pheromone in leaf beetles so far. Indeed, most of the research on the pyrazine motif as pheromones has been conducted in ants, in which they act as alarm or trail pheromones [45–47], in true fruit flies (Diptera: Tephritidae) [48–50], in thynnine wasps [20] and in ladybugs [32,33,51]. As stated above, in some coccinellid species, an aggregative effect is induced by 2-alkyl-methoxypyrazines in diapausing individuals, aside from their roles as allomones. Al Abassi and coworkers reported that IPMP is attractive for both sexes of *Coccinella septempunctata* (L.) (Coleoptera: Coccinellidae) under laboratory conditions, inducing an attractive and arrestant effect attributable to the effect of an aggregation pheromone [32]. Similarly, IBMP alone or mixed with IPMP results in attraction for diapausing individuals in *Adalia bipunctata* (L.) (Coleoptera: Coccinellidae) [51]. In other ladybug species, such as *Harmonia axyridis* Pallas and *Hippodamia convergens* Guérin-Ménéville (Coleoptera: Coccinellidae), IPMP, IBMP and SBMP have been identified [52] and demonstrated to be behaviorally active. In the case of *H. convergens* adults, SBMP, and especially IBMP and the ternary blend at its natural ratio, induce significant aggregative effects in laboratory and field assays, whereas high amounts of IPMP and SBMP were shown to be repellent [33]. Likewise, an aversive behavior in *L. lusitanica* males was evident when they were confronted with the highest amount of IBMP in olfactometer trials. Given the dual activity of 2-alkyl-methoxypyrazines in ladybugs, it seems probable that IBMP may play a similar role in *L. lusitanica*, not only resulting in attraction for both sexes, but also conferring a defense mechanism against predation.

According to field observations, *L. lusitanica* aggregates with a high number of individuals (more than 50) are found only in a few pistachio trees, while most of the trees harbor smaller numbers (less than 15) [17]. In a similar vein, the response of males and females of *Aphthona nigriscutis* Foudras (Coleoptera: Chrysomelidae) to aggregation cues decreases according to the density of conspecifics on the plant [38]. In the latter, it is not acknowledged if this density-dependent response may be conditioned by deterrent plant volatile organic compounds released upon feeding by conspecifics [53–55]. For example, *Diorhabda carinulata* Desbrochers (Coleoptera: Chrysomelidae) avoids previously defoliated saltcedar plants, and this avoidance has been correlated with the release of 4-oxo-(*E*)-2-hexenal from the host, inducing a repellent effect on reproductive males and females [53]. Nonetheless, since *L. lusitanica* colonizes healthy and non-consumed pistachio leaves from young trees, this hypothesis should be a priori discarded. Alternatively, we suggest that chemical cues from pistachio trees may be signaling the suitability of the host to some

extent, partially explaining, therefore, the patchy spatial pattern of aggregates of variable numbers of individuals across the crop. Indeed, plant volatiles are demonstrated to be relevant in the aggregation processes of Chrysomelidae, either by luring in the pioneer individuals [43,56] or by enhancing/synergizing the effect of the aggregation pheromone [38,57–60], as it occurs in other coleopteran families [61,62]. Furthermore, being in contact with the host plant and/or feeding seem to be sine qua non conditions for pheromone release in leaf beetle species [41–44,56,63]. Examples highlighting this phenomenon are found in *Acalymma vittatum* (F.), whose feeding males are more attractive to conspecifics than non-feeding males [43], in *Phyllotreta* spp. [41,44] and in *Oulema melanopus* (L.), in which the release of aggregation pheromones is closely related to males feeding on the host plant [42]. In our case, IBMP was detected from males deprived of pistachio leaves during the volatile collection, although they were not subjected to starvation prior to the sampling. Thus, it remains unanswered whether the presence of plant volatiles and/or pistachio leaf consumption are crucial for the release of IBMP.

In summary, we have identified a male-specific compound that results in the attraction of both sexes under laboratory conditions. Future field research is ongoing to gain a broader understanding of the role of the compound in the chemical ecology of *L. lusitanica*, in order to determine its potential activity under natural conditions. Last, but not least, a thorough understanding of their population dynamics, and whether the production and release of 2-isobutyl-3-methoxypyrazine remain constant during the whole life cycle, would also be paramount for deciphering the biological implications of the compound.

Author Contributions: Conceptualization, S.L., S.R.-G., E.F.-C., C.C.-M. and C.Q.; methodology, S.L.; investigation, S.L. and C.C.-M.; data analysis, S.L.; writing—original draft preparation, S.L.; writing—review and editing, S.L. and C.Q.; supervision, S.L., S.R.-G., E.F.-C., C.C.-M. and C.Q. All authors have read and agreed to the published version of the manuscript.

Funding: Sara Rodrigo-Gómez acknowledges an innovation pilot project in plant health area promoted by 16.2.2. operation under the Rural Development Programme of Castilla-La Mancha, for the 2014–2020 period, cofounded by the European Agricultural Fund for Rural Development (EAFRD), Ministerio de Agricultura, Pesca y Alimentación (MAPA) and Junta de Comunidades de Castilla—La Mancha (JCCM) (SV-2019-02).

Data Availability Statement: All the data generated from the current work are available upon reasonable request to the corresponding authors.

Acknowledgments: We are very thankful to the personnel of the Centro de Investigación Agroambiental “El Chaparrillo” (IRIAF, Ciudad Real), and all those farmers involved in the abovementioned project for their invaluable support. Authors declare that all the people included in this section have consented to be acknowledged.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Whitehouse, W.E. The pistachio nut—a new crop for the Western United States. *Econ. Bot.* **1957**, *11*, 281–321, doi:10.1007/BF02903809.
2. Lemaistre, J.C. Le pistachier (étude bibliographique). *Fruits* **1959**, *14*, 57–77.
3. Maggs, D.H. *An introduction to pistachio growing in Australia*; Commonwealth Scientific and Industrial Research Organization: Adelaide, Australia, 1982; ISBN 0643029907.
4. Mehlenbacher, S.A. Progress and prospects in nut breeding. *Acta Hort.* **2003**, *622*, 57–79, doi:10.17660/ActaHortic.2003.622.4.
5. Mandalari, G.; Barreca, D.; Gervasi, T.; Rousell, M.A.; Klein, B.; Feeney, M.J.; Carughi, A. Pistachio nuts (*Pistacia vera* L.): production, nutrients, bioactives and novel health effects. *Plants* **2021**, *11*, 18, doi:10.3390/plants11010018.
6. Mourikis, P.A.; Tsourgianni, A.; Chitzanidis, A. Pistachio nut insect pests and means of control in Greece. *Acta Hort.* **1998**, *470*, 604–611, doi:10.17660/ActaHortic.1998.470.85.
7. Mehrnejad, M.R. Arthropod pests of pistachios, their natural enemies and management. *Plant Prot. Sci.* **2020**, *56*, 231–260, doi:10.17221/63/2019-PPS.
8. MAPAMA. *Guía de gestión integrada de plagas - Pistacho*; MAPAMA: Madrid, España, 2018; ISBN 9788449114984.
9. Davatchi, G.A. Biology of the insect fauna of cultivated and wild *Pistacia*. *Rev. Pathol. Veg.* **1958**, *37*, 3–166.
10. Bolu, H. Investigations on the fauna of insects and mites in pistachio areas in south Eastern Anatolia region of Turkey. *Turkish*

- J. Entomol.* **2002**, *26*, 197–208.
11. Doğanlar, M.; Karadağ, S.; Mendel, Z. Notes on pistachio seed wasps from two locations in the east Mediterranean. *Phytoparasitica* **2009**, *37*, 147–151, doi:10.1007/s12600-009-0022-y.
 12. Rice, R.E.; Michailides, T.J. Pistachio seed chalcid, *Megastigmus pistaciae* Walker (Hymenoptera: Torymidae), in California. *J. Econ. Entomol.* **1988**, *81*, 1446–1449, doi:10.1093/jee/81.5.1446.
 13. Braham, M.; Jardak, T. Contribution a l'étude de la bio-écologie du scolyte du pistachier *Chaetoptelius vestitus* Muls & Rey (Coleoptera, Scolytidae) dans les régions du centre et du sud tunisiens. *Rev. Ezzaitouna* **2012**, *13*, 1–17.
 14. Mehrnejad, M.R. Investigation into the overwintering and winter-management of the common pistachio psyllid, *Agonoscena pistaciae* (Hemiptera: Aphalaridae), a major pest in pistachio plantations. *Zool. Ecol.* **2018**, *28*, 384–388, doi:10.1080/21658005.2018.1516339.
 15. Haviland, D.R.; Bentley, W.J.; Siegel, J.P.; Holtz, B.A.; Daane, K.M.; Higbee, B.S. Navel orangeworm and obliquebanded leafroller. In *Pistachio Production Manual, Publication 3545*; Ferguson, L., Haviland, D.R., Eds.; University of California, Agriculture and Natural Resources: Oakland, United States, 2016; pp. 197–210.
 16. Fernández-Carrillo, E.; Fernández-Carrillo, J.L.; Rodrigo-Gómez, S.; Velázquez-de-Castro, A.J. Weevils (Coleoptera, Curculionidae) associated with pistachio tree (*Pistacia vera* L.) crops in Castilla-La Mancha (Central Spain). *Graellsia* **2022**, *78*, e168, doi:10.3989/graellsia.2022.v78.334.
 17. Gómez, S.R.; Gil-Tapetado, D.; García-Gila, J.; Blasco-Aróstegui, J.; Polidori, C. The leaf beetle *Labidostomis lusitanica* (Coleoptera: Chrysomelidae) as an Iberian pistachio pest: projecting risky areas. *Pest Manag. Sci.* **2022**, *78*, 217–229, doi:10.1002/ps.6624.
 18. Heisswolf, A.; Gabler, D.; Obermaier, E.; Müller, C. Olfactory versus contact cues in host plant recognition of a monophagous chrysomelid beetle. *J. Insect Behav.* **2007**, *20*, 247–266, doi:10.1007/s10905-007-9078-z.
 19. Gerritsma, D.A.; Brindle, I.D.; Jones, T.R.B.; Capretta, A. Preparation of labelled 2-methoxy-3-alkylpyrazines: synthesis and characterization of deuterated 2-methoxy-3-isopropylpyrazine and 2-methoxy-3-isobutylpyrazine. *J. Label. Compd. Radiopharm.* **2003**, *46*, 243–253, doi:10.1002/jlcr.664.
 20. Bohman, B.; Phillips, R.D.; Menz, M.H.M.; Berntsson, B.W.; Flematti, G.R.; Barrow, R.A.; Dixon, K.W.; Peakall, R. Discovery of pyrazines as pollinator sex pheromones and orchid semiochemicals: implications for the evolution of sexual deception. *New Phytol.* **2014**, *203*, 939–952, doi:10.1111/nph.12800.
 21. Bohman, B.; Peakall, R. Pyrazines attract *Catocheilus* thynnine wasps. *Insects* **2014**, *5*, 474–487, doi:10.3390/insects5020474.
 22. Maia, A.C.D.; Santos, G.K.N.; Gonçalves, E.G.; Navarro, D.M. do A.F.; Nuñez-Avellaneda, L.A. 2-Alkyl-3-methoxypyrazines are potent attractants of florivorous scarabs (Melolonthidae, Cyclocephalini) associated with economically exploitable Neotropical palms (Arecaceae). *Pest Manag. Sci.* **2018**, *74*, 2053–2058, doi:10.1002/ps.4895.
 23. Suinyuy, T.N.; Donaldson, J.S.; Johnson, S.D. Geographical matching of volatile signals and pollinator olfactory responses in a cycad brood-site mutualism. *Proc. R. Soc. B Biol. Sci.* **2015**, *282*, 20152053, doi:10.1098/rspb.2015.2053.
 24. Moore, B.P.; Brown, W.V.; Rothschild, M. Methylalkylpyrazines in aposematic insects, their hostplants and mimics. *Chemoecology* **1990**, *1*, 43–51, doi:10.1007/BF01325227.
 25. Burdfield-Steel, E.; Pakkanen, H.; Rojas, B.; Galarza, J.A.; Mappes, J. De novo synthesis of chemical defenses in an aposematic moth. *J. Insect Sci.* **2018**, *18*, 28, doi:10.1093/jisesa/iey020.
 26. Rothschild, M.; Moore, B.P.; Brown, W.V. Pyrazines as warning odour components in the Monarch butterfly, *Danaus plexippus*, and in moths of the genera *Zygaena* and *Amata* (Lepidoptera). *Biol. J. Linn. Soc.* **1984**, *23*, 375–380, doi:10.1111/j.1095-8312.1984.tb00153.x.
 27. Aldrich, J.R.; Avery, J.W.; Lee, C.-J.; Graf, J.C.; Harrison, D.J.; Bin, F. Semiochemistry of cabbage bugs (Heteroptera: Pentatomidae: *Eurydema* and *Murgantia*). *J. Entomol. Sci.* **1996**, *31*, 172–182, doi:10.18474/0749-8004-31.2.172.
 28. Rowe, C.; Guilford, T. Hidden colour aversions in domestic chicks triggered by pyrazine odours of insect warning displays. *Nature* **1996**, *383*, 520–522, doi:10.1038/383520a0.
 29. Lindström, L.; Rowe, C.; Guilford, T. Pyrazine odour makes visually conspicuous prey aversive. *Proc. Biol. Sci.* **2001**, *268*, 159–162, doi:10.1098/rspb.2000.1344.
 30. Rojas, B.; Burdfield-Steel, E.; Pakkanen, H.; Suisto, K.; Maczka, M.; Schulz, S.; Mappes, J. How to fight multiple enemies: target-specific chemical defences in an aposematic moth. *Proc. R. Soc. B Biol. Sci.* **2017**, *284*, 20171424, doi:10.1098/rspb.2017.1424.
 31. Burdfield-Steel, E.R.; Schneider, J.M.; Mappes, J.; Dobler, S. Testing the effectiveness of pyrazine defences against spiders. *Chemoecology* **2020**, *30*, 139–146, doi:10.1007/s00049-020-00305-5.
 32. Al Abassi, S.; Birkett, M.A.; Pettersson, J.; Pickett, J.A.; Woodcock, C.M. Ladybird beetle odour identified and found to be responsible. *Cell. Mol. Life Sci.* **1998**, *54*, 876–879, doi:10.1007/s000180050215.
 33. Wheeler, C.A.; Cardé, R.T. Defensive allomones function as aggregation pheromones in diapausing ladybird beetles, *Hippodamia convergens*. *J. Chem. Ecol.* **2013**, *39*, 723–732, doi:10.1007/s10886-013-0293-0.
 34. Santiago-Blay, J.; Jolivet, P.; Verma, K. A natural history of conspecific aggregations in terrestrial arthropods, with emphasis on cycloaexy in leaf beetles (Coleoptera: Chrysomelidae). *Terr. Arthropod Rev.* **2012**, *5*, 289–355, doi:10.1163/18749836-05031054.
 35. Dickens, J.C.; Oliver, J.E.; Hollister, B.; Davis, J.C.; Klun, J.A. Breaking a paradigm: male-produced aggregation pheromone for the Colorado potato beetle. *J. Exp. Biol.* **2002**, *205*, 1925–1933, doi:10.1242/jeb.205.13.1925.
 36. Beran, F.; Jiménez-Alemán, G.H.; Lin, M.-Y.; Hsu, Y.-C.; Mewis, I.; Srinivasan, R.; Ulrichs, C.; Boland, W.; Hansson, B.S.; Reinecke, A. The aggregation pheromone of *Phyllotreta striolata* (Coleoptera: Chrysomelidae) revisited. *J. Chem. Ecol.* **2016**, *42*, 748–755, doi:10.1007/s10886-016-0743-6.

37. Cossé, A.A.; Bartelt, R.J.; Zilkowski, B.W.; Bean, D.W.; Petroski, R.J. The aggregation pheromone of *Diorhabda elongata*, a biological control agent of saltcedar (*Tamarix* spp.): identification of two behaviorally active components. *J. Chem. Ecol.* **2005**, *31*, 657–670, doi:10.1007/s10886-005-2053-2.
38. Tansey, J.A.; McClay, A.S.; Cole, D.E.; Keddie, B.A. Evidence for the influence of conspecific chemical cues on *Aphthona nigricutis* (Coleoptera: Chrysomelidae) behaviour and distribution. *BioControl* **2005**, *50*, 343–358, doi:10.1007/s10526-004-1232-3.
39. Zilkowski, B.W.; Bartelt, R.J.; Cossé, A.A.; Petroski, R.J. Male-produced aggregation pheromone compounds from the eggplant flea beetle (*Epitrix fuscula*): identification, synthesis, and field bioassays. *J. Chem. Ecol.* **2006**, *32*, 2543–2558, doi:10.1007/s10886-006-9163-3.
40. Rao, S.; Cossé, A.A.; Zilkowski, B.W.; Bartelt, R.J. Aggregation pheromone of the cereal leaf beetle: field evaluation and emission from males in the laboratory. *J. Chem. Ecol.* **2003**, *29*, 2165–2175, doi:10.1023/a:1025698821635.
41. Beran, F.; Mewis, I.; Srinivasan, R.; Svoboda, J.; Vial, C.; Mosimann, H.; Boland, W.; Büttner, C.; Ulrichs, C.; Hansson, B.S.; et al. Male *Phyllotreta striolata* (F.) produce an aggregation pheromone: identification of male-specific compounds and interaction with host plant volatiles. *J. Chem. Ecol.* **2011**, *37*, 85–97, doi:10.1007/s10886-010-9899-7.
42. Cossé, A.A.; Bartelt, R.J.; Zilkowski, B.W. Identification and electrophysiological activity of a novel hydroxy ketone emitted by male cereal leaf beetles. *J. Nat. Prod.* **2002**, *65*, 1156–1160, doi:10.1021/np020063q.
43. Smyth, R.R.; Hoffmann, M.P. A male-produced aggregation pheromone facilitating *Acalymma vittatum* [F.] (Coleoptera: Chrysomelidae) early-season host plant colonization. *J. Insect Behav.* **2003**, *16*, 347–359, doi:10.1023/A:1024824025210.
44. Peng, C.; Bartelt, R.J.; Weiss, M.J. Male crucifer flea beetles produce an aggregation pheromone. *Physiol. Entomol.* **1999**, *24*, 98–99, doi:10.1046/j.1365-3032.1999.00113.x.
45. Vander Meer, R.K.; Preston, C.A.; Choi, M.-Y. Isolation of a pyrazine alarm pheromone component from the fire ant, *Solenopsis invicta*. *J. Chem. Ecol.* **2010**, *36*, 163–170, doi:10.1007/s10886-010-9743-0.
46. Hölldobler, B.; Morgan, E.D.; Oldham, N.J.; Liebig, J. Recruitment pheromone in the harvester ant genus *Pogonomyrmex*. *J. Insect Physiol.* **2001**, *47*, 369–374, doi:10.1016/S0022-1910(00)00143-8.
47. Tentschert, J.; Bestmann, H.-J.; Hölldobler, B.; Heinze, J. 2,3-Dimethyl-5-(2-methylpropyl)pyrazine, a trail pheromone component of *Eutetramorium mocquersyi* Emery (1899) (Hymenoptera: Formicidae). *Naturwissenschaften* **2000**, *87*, 377–380, doi:10.1007/s001140050745.
48. Chuman, T.; Landolt, P.J.; Heath, R.R.; Tumlinson, J.H. Isolation, identification, and synthesis of male-produced sex pheromone of papaya fruit fly, *Toxotrypana curvicauda* Gerstaecker (Diptera: Tephritidae). *J. Chem. Ecol.* **1987**, *13*, 1979–1992, doi:10.1007/BF01014680.
49. Robledo, N.; Vega, M.; Escalante, J.; Arzuffi, R. A new component of the male papaya fruit fly (Diptera: Tephritidae) sex pheromone. *Fla. Entomol.* **2014**, *97*, 1260–1262, doi:10.1653/024.097.0338.
50. Robacker, D.; Aluja, M.; Bartelt, R.J.; Patt, J. Identification of chemicals emitted by calling males of the sapote fruit fly, *Anastrepha serpentina*. *J. Chem. Ecol.* **2009**, *35*, 601–609, doi:10.1007/s10886-009-9631-7.
51. Susset, E.C.; Ramon-Portugal, F.; Hemptinne, J.-L.; Dewhurst, S.Y.; Birkett, M.A.; Magro, A. The role of semiochemicals in short-range location of aggregation sites in *Adalia bipunctata* (Coleoptera, Coccinellidae). *J. Chem. Ecol.* **2013**, *39*, 591–601, doi:10.1007/s10886-013-0285-0.
52. Cudjoe, E.; Wiederkehr, T.B.; Brindle, I.D. Headspace gas chromatography-mass spectrometry: a fast approach to the identification and determination of 2-alkyl-3-methoxypyrazine pheromones in ladybugs. *Analyst* **2005**, *130*, 152–155, doi:10.1039/b410659h.
53. Gaffke, A.M.; Sing, S.E.; Millar, J.G.; Dudley, T.L.; Bean, D.W.; Peterson, R.K.D.; Weaver, D.K. An herbivore-induced plant volatile from saltcedar (*Tamarix* spp.) is repellent to *Diorhabda carinulata* (Coleoptera: Chrysomelidae). *Environ. Entomol.* **2020**, *49*, 1063–1070, doi:10.1093/ee/nvaa079.
54. Piesik, D.; Wenda-Piesik, A.; Kotwica, K.; Łyszczarz, A.; Delaney, K.J. *Gastrophysa polygoni* herbivory on *Rumex confertus*: single leaf VOC induction and dose dependent herbivore attraction/repellence to individual compounds. *J. Plant Physiol.* **2011**, *168*, 2134–2138, doi:10.1016/j.jplph.2011.06.012.
55. Piesik, D.; Wenda-Piesik, A.; Krasieńska, A.; Wrzesieńska, D.; Delaney, K.J. Volatile organic compounds released by *Rumex confertus* following *Hypera rumicis* herbivory and weevil responses to volatiles. *J. Appl. Entomol.* **2016**, *140*, 308–316, doi:10.1111/jen.12238.
56. Bartelt, R.J.; Cossé, A.A.; Zilkowski, B.W.; Wiedenmann, R.N.; Raghu, S. Early-summer pheromone biology of *Galerucella californiensis* and relationship to dispersal and colonization. *Biol. Control* **2008**, *46*, 409–416, doi:10.1016/j.biocontrol.2008.05.010.
57. Weber, D.C.; Konstantinov, A.S.; Khimian, A.; Bier, A.D.; Lubenow, L.A.; Knodel, J.J.; Haber, A.I.; Wallingford, A.K.; Mason, J.A.C.; Kuhar, T.P. Trapping of crucifer-feeding flea beetles (*Phyllotreta* spp.) (Coleoptera: Chrysomelidae) with pheromones and plant kairomones. *J. Econ. Entomol.* **2022**, *115*, 748–756, doi:10.1093/jee/toac042.
58. Soroka, J.J.; Bartelt, R.J.; Zilkowski, B.W.; Cossé, A.A. Responses of flea beetle *Phyllotreta cruciferae* to synthetic aggregation pheromone components and host plant volatiles in field trials. *J. Chem. Ecol.* **2005**, *31*, 1829–1843, doi:10.1007/s10886-005-5929-2.
59. Li, Y.; Zhao, P.; Yin, C.; Liu, X.; Zhang, Q. Attraction of *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae) by several plant volatiles and aggregation pheromone. *Acta Entomol. Sin.* **2010**, *53*, 734–740.
60. Cossé, A.A.; Bartelt, R.J.; Zilkowski, B.W.; Bean, D.W.; Andress, E.R. Behaviorally active green leaf volatiles for monitoring the leaf beetle, *Diorhabda elongata*, a biocontrol agent of saltcedar, *Tamarix* spp. *J. Chem. Ecol.* **2006**, *32*, 2695–2708, doi:10.1007/s10886-006-9193-x.

61. Giblin-Davis, R.M.; Weissling, T.J.; Oehlschlager, A.C.; Gonzalez, L.M. Field response of *Rhynchophorus cruentatus* (Coleoptera: Curculionidae) to its aggregation pheromone and fermenting plant volatiles. *Fla. Entomol.* **1994**, 164–177, doi:10.2307/3495883.
62. Wibe, A.; Borg-Karlson, A.-K.; Cross, J.; Bichão, H.; Fountain, M.; Liblikas, I.; Sigsgaard, L. Combining 1,4-dimethoxybenzene, the major flower volatile of wild strawberry *Fragaria vesca*, with the aggregation pheromone of the strawberry blossom weevil *Anthonomus rubi* improves attraction. *Crop Prot.* **2014**, 64, 122–128, doi:10.1016/j.cropro.2014.06.016.
63. Hitchner, E.M.; Kuhar, T.P.; Dickens, J.C.; Youngman, R.R.; Schultz, P.B.; Pfeiffer, D.G. Host plant choice experiments of Colorado potato beetle (Coleoptera: Chrysomelidae) in Virginia. *J. Econ. Entomol.* **2008**, 101, 859–865, doi:10.1093/jee/101.3.859.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.