

Supplementary Materials

Diverse epidemiology of *Leptospira* serovars notified in New Zealand, 1999-2017

Shahista Nisa ¹, David A. Wilkinson ², Olivia Angelin-Bonnet ³, Shevaun Paine ⁴, Karen Cullen ⁴, Jackie Wight ⁴, Michael G. Baker ⁵ and Jackie Benschop ^{1,6,*}

¹ Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, School of Veterinary Science, Massey University, Palmerston North, New Zealand; s.nisa@massey.ac.nz

² New Zealand Food Safety Science and Research Centre, Massey University, Palmerston North, New Zealand; david.wilkinson@univ-reunion.fr

³ School of Fundamental Sciences, College of Sciences, Massey University, Palmerston North, New Zealand; Angelin-bonnet@massey.ac.nz

⁴ Institute of Environmental Science and Research, Wellington, New Zealand; Shevaun.Paine@esr.cri.nz; karen.cullen@esr.cri.nz; Jackie.Wright@esr.cri.nz

⁵ Department of Public Health, University of Otago, Wellington, New Zealand; michael.baker@otago.ac.nz

⁶ Global Leptospirosis Environmental Action Network, World Health Organization, Geneva, Switzerland; J.Benschop@massey.ac.nz


* Correspondence: j.benschop@massey.ac.nz; Tel.: +64 6 359 9099 ext: 83994

Received: date; Accepted: date; Published: date




Figure S1: Location boundaries for the 20 District Health Board boundaries in New Zealand. Source: Ministry of Health, New Zealand.

28



30

31 **Figure S2:** Choropleth maps of early and late study period. Average annual incidence per 100 000 of
 32 notified leptospirosis cases per District Health Boards in New Zealand for the early (1999-2007) and
 33 the late (2008-2017) study period for all cases, Europeans cases and Maori cases. Grey indicates DHB
 34 with no incidences.



35

36 **Figure S3:** Type of laboratory tests used to diagnose notified confirmed and probable leptospirosis
37 cases in New Zealand, 1999 to 2017.