

Article

MDPI

Identification of Clinically Relevant *Streptococcus* and *Enterococcus* Species Based on Biochemical Methods and 16S rRNA, *sodA*, *tuf*, *rpoB*, and *recA* Gene Sequencing

Maja Kosecka-Strojek ^{1,*}, Mariola Wolska ¹, Dorota Żabicka ², Ewa Sadowy ³ and Jacek Międzobrodzki ¹

- ¹ Department of Microbiology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, 30-387 Krakow, Poland; mariola.wolska@doctoral.uj.edu.pl (M.W.); jacek.miedzobrodzki@uj.edu.pl (J.M.)
- ² Department of Molecular Microbiology, National Medicines Institute, 00-725 Warsaw, Poland; d.zabicka@nil.gov.pl
- ³ Department of Epidemiology and Clinical Microbiology, National Medicines Institute, 00-725 Warsaw, Poland; e.sadowy@nil.gov.pl
- * Correspondence: maja.kosecka-strojek@uj.edu.pl; Tel.: +48-12-664-6365

Received: 13 October 2020; Accepted: 9 November 2020; Published: 11 November 2020

Abstract: Streptococci and enterococci are significant opportunistic pathogens in epidemiology and infectious medicine. High genetic and taxonomic similarities and several reclassifications within genera are the most challenging in species identification. The aim of this study was to identify *Streptococcus* and *Enterococcus* species using genetic and phenotypic methods and to determine the most discriminatory identification method. Thirty strains recovered from clinical samples representing 15 streptococcal species, five enterococcal species, and four nonstreptococcal species were subjected to bacterial identification by the Vitek[®] 2 system and Sanger-based sequencing methods targeting the 16S rRNA, *sodA*, *tuf*, *rpoB*, and *recA* genes. Phenotypic methods allowed the identification of 10 streptococcal strains, five enterococcal strains, and four nonstreptococcal strains (*Leuconostoc*, *Granulicatella*, and *Globicatella* genera). The combination of sequencing methods allowed the identification of 21 streptococcal strains, five enterococcal strains, or sequencing methods allowed the identification of several molecular methods was sufficient for unambiguous confirmation of species identity. This study will be useful for comparison of several identification methods, both those used as a first choice in routine microbiology and those used for final confirmation.

Keywords: streptococci; enterococci; molecular diagnostics; genetic methods; Sanger sequencing

1. Introduction

Gram-positive bacteria of the Streptococcus and Enterococcus genera are of great clinical and epidemiological importance, and most species are components of the natural human microbiota [1]. The genus Streptococcus includes а large number (at least 135) [2,3] [https://www.bacterio.net/genus/streptococcus] of species that colonize human and animal mucous membranes. Species such as Streptococcus pyogenes, Streptococcus pneumoniae, and Streptococcus agalactiae are highly virulent and cause infections and diseases such as scarlet and rheumatic fevers, pneumonia or neonatal sepsis [4-6]. Streptococci are classified based on colony morphology, hemolysis type, and serological specificity. The serological specificity is based on antigenic differences in cell wall carbohydrates, in cell wall pilus-associated proteins, and in the polysaccharide capsule in group B streptococci [7]. The classification and nomenclature of streptococci are based on group antigens (Lancefield serotyping system) as follows: group A *Streptococcus* (GAS); group B *Streptococcus* (GBS); group C *Streptococcus*; group G *Streptococcus*; the viridans group, with the subgroups anginosus, mitis, mutans, and salivarius; and the bovis group [8–10].

The members of the genus now known as *Enterococcus* were formerly considered to be group D Streptococcus until 1984 [11]. Isolates from the Enterococcus genus are commensals of the gastrointestinal tracts of humans and animals and include 64 species [12,13] [https://bacterio.net/genus/enterococcus]. All Enterococcus species are classified into the antigen D group by the Lancefield system [11] and exhibit gamma-hemolysis on blood agar, although some strains are alpha-hemolytic or beta-hemolytic [14,15]. Enterococcus faecalis and Enterococcus faecium can cause a variety of infections, including endocarditis and urinary tract infections [16,17].

The addition of new species, changing taxonomy and modification of the systematic names of streptococci and enterococci, poses a challenge to proper identification of species. Therefore, precise identification of these species is laborious. Clinical laboratories use phenotypic biochemical methods such as Vitek® 2 (bioMérieux, La Balme Les Grottes, France) and BD Phoenix (BD Diagnostic Systems, Sparks, MD, USA), commercial rapid test kits such as API® Strep (bioMérieux, La Balme Les Grottes, France) and matrix-assisted laser desorption ionization-time-of-flight mass spectrometry (MALDI-TOF MS). In routine diagnostics, especially the Vitek® 2 system is used. This system is based on kinetic analysis detecting metabolic changes and by additional continuous monitoring of reactions, provides much faster species identifications [18]. Nevertheless, the technique so far has failed at differentiating between mitis, bovis groups, and other closely relative species [19,20]. On the other hand, commercially available MALDI-TOF MS systems provide accurate identification of many clinically relevant streptococcal species. However, MALDI-TOF spectra databases are limited to only some species, and further improvements of Streptococcus and Enterococcus spectra databases seem necessary. The phenotypic trait variability within strains and species using this method compared to methods based on genetic discrimination causes limited differentiation capacity; consequently, more than 50% of these bacteria are incorrectly identified [21,22].

The development of molecular biological techniques has made it possible to rapidly and reliably diagnose infections caused by bacteria of the *Streptococcus* and *Enterococcus* genera. Genetic methods are based on PCR or sequencing, and identification is based on selected molecular target amplification, sequencing, and comparison to a reference sequence deposited in a nucleotide database [13]. 16S rRNA gene sequencing has proven to be one of the most powerful tools for the classification of microorganisms, including streptococci and enterococci [1,23]. However, due to low specificity, the correct identification of bacterial species should not be based on the nucleotide sequence of a single gene. For unambiguous species confirmation, it is necessary to use additional molecular markers. For the identification of *Streptococcus* and *Enterococcus* isolates, several gene targets, such as genes encoding manganese-dependent superoxide dismutase (*sodA*) [24], the elongation factor Tu (*tuf*) [25], and beta-subunit of RNA polymerase (*rpoB*) [26], have been used. Furthermore, for species included in the mitis (currently includes about 20 different species [27,28]) and bovis (*Streptococcus bovis, Streptococcus equinus, Streptococcus gallolyticus, Streptococcus lutetiensis, Streptococcus alactolyticus* [29]) groups, which are closely related, other conserved molecular targets, such as the subunit of the bacterial recombinase (*recA*) gene, may be used [30,31].

The aim of this study was to identify clinically relevant *Streptococcus* and *Enterococcus* species using genetic and phenotypic methods and to determine the most discriminatory identification method. In our study, the Vitek[®] 2 system and Sanger sequencing of five genes, namely, the 16S rRNA, *sodA*, *tuf*, *rpoB*, and *recA* genes, were used.

2. Results

After recovering the isolates from clinical samples, the hospital laboratories identified all of the isolates at the genus level. All isolates were identified as *Streptococcus* and *Enterococcus* with routine diagnostic methods. Afterwards, serotyping and identification at the species level were performed in our laboratory. The Lancefield serotype groups were assigned: 57% streptococci, 60% enterococci, and 50% other nonstreptococci. No visible agglutination of latex or autoagglutination with more than one reagent with antibody particles was interpreted as ambiguous. Briefly, in the streptococcal serotype identification performed with the Pastorex[™] Strep Test Kit (Bio-Rad, Hercules, CA, USA), a positive reaction is indicated by red clumps on a green background, visible to the naked eye. Agglutination with only one of the six latex suspensions convincingly establishes the group of the strain tested. A negative reaction is indicated by a homogenous brown suspension, without clumps, after one minute of agitation. A reaction is un-interpretable if small clumps appear on a brown background, or if agglutination appears with more than one latex reagent in the kit [32].

The Vitek[®] 2 system allowed for identification of 10 of the 21 *Streptococcus* strains, all five *Enterococcus* strains, and three nonstreptococcal strains (*Globicatella sanguinis, Leuconostoc lactis,* and *Leuconostoc citreum*) (Table 1). The Vitek[®] 2 procedure and serotyping were performed for all isolates, and MALDI-TOF MS was performed for ambiguous and untypable isolates. Most streptococci and enterococci species were identified at excellent (67% *Streptococcus;* 40% *Enterococcus*) and very good (14% *Streptococcus;* 60% *Enterococcus*) discrimination levels. For 29% of the streptococcal strains (PL427, S63, 1816/15, 1226/14, PL431 1374/11), the Vitek[®] 2 system did not allow identification at the species level, and only the indistinguishable *S. mitis* or *S. oralis* group was assigned. The strains 6922/09 and 1860/08 were assigned as *Streptococcus anginosus/Streptococcus gordonii* and *Streptococcus agalactiae/Streptococcus dysgalactiae,* respectively. The PL434 strain was identified as *Kocuria rosea,* and p41 was not identified at all.

For isolates which were not identified as *Streptococcus* or *Enterococcus* by the Vitek[®] 2 system (PL434, 1113/11, 3696/08, p41, and 1375/11), MALDI-TOF MS was used. All isolates were identified with a high degree of confidence (\geq 2.00). For the strains 1113/11, 3696/08, and 1375/11, MALDI-TOF MS showed the same identification results as the Vitek[®] 2 system. In the case of PL434 (identified as *Kocuria rosea* by Vitek[®] 2), MALDI-TOF MS identified this isolate as *Granulicatella adiacens*, and p41 was identified as *S. pneumoniae* (Table 1).

			Vitek® 2 Sy	stem Results	
Isolate No.	Serotype	Species	Probability of	Discrimination	MALDI-TOF MS
			Identification	Level	Species (Score Values)
E15	G	Enterococcus avium	95%	Very good	not performed
E2	D	Enterococcus casseliflavus	98%	Excellent	not performed
E5	AGL	Enterococcus durans	94%	Very good	not performed
E28	AGL	Enterococcus faecalis	99%	Excellent	not performed
E10	D	Enterococcus raffinosus	94%	Very good	not performed
1375/11	-	Globicatella sanguinis	96%	Excellent	Globicatella sanguinis (2.39)
PL434	D	Kocuria rosea	90%	Good	Granulicatella adiacens (2.22)
3696/08	D	Leuconostoc citreum	97%	Excellent	Leuconostoc citreum (2.10)
1113/11	AGL	Leuconostoc lactis	97%	Excellent	Leuconostoc lactis (2.20)
S19	В	Streptococcus agalactiae	97%, 99%	Excellent	not performed
4734/08	С	Streptococcus anginosus	97%	Excellent	not performed
5898/10	AGL	Streptococcus anginosus	95%	Very good	not performed
1816/15	-	Streptococcus mitis/oralis	99%	Excellent	not performed
1226/14	-	Streptococcus mitis/oralis	95%	Very good	not performed
1107/09	С	Streptococcus gordonii	97%	Excellent	not performed

Table 1. Performance of the serotyping of the Vitek[®] 2 system and MALDI-TOF MS identification of *Streptococcus* spp. and *Enterococcus* spp.

Pathogens 2020, 9, 939

6922/09	С	Streptococcus anginosus/gordonii	96%	Low	not performed
1860/08	G	Streptococcus agalactiae/ dysgalactiae	98%	Low	not performed
8190/10	G	Streptococcus dysgalactiae	96%	Excellent	not performed
5010/12	AGL	Streptococcus gallolyticus	99%	Excellent	not performed
S18	AGL	Streptococcus gallolyticus	97%	Excellent	not performed
PL427	-	Streptococcus mitis/oralis	99%	Excellent	not performed
PL428	-	Streptococcus lutetiensis	97%	Excellent	not performed
PL431	D	Streptococcus mitis/oralis	90%	Good	not performed
S16	-	Streptococcus parasanguinis	99%	Excellent	not performed
S63	С	Streptococcus mitis/oralis	98%	Excellent	not performed
1374/11	-	Streptococcus mitis/oralis	99%	Excellent	not performed
p63	С	Streptococcus pneumoniae	99%	Excellent	not performed
p41	D	×	×	×	Streptococcus pneumoniae (2.08)
S47	A	Streptococcus pyogenes	95%	Very good	not performed
PL433	G	Streptococcus salivarius	96%	Excellent	not performed

AGL—agglutination; All ambiguous *Streptococcus*, *Enterococcus* and nonstreptococcus species are indicated in a dark red color. ×—lack of identification.

2.2. Sanger Sequencing of the 16S rRNA Gene

Sanger sequencing of the 16S rRNA gene allowed identification of 19 *Streptococcus* strains (90% of all streptococcal strains), four *Enterococcus* strains (80% of all enterococcal strains), and three of the four nonstreptococcal strains (*G. adiacens, G. sanguinis, L. citreum*). Identification of the following pairs of enterococcal, streptococcal, and nonstreptococcal species was impossible because the 16S rRNA gene sequences were identical or almost identical (\geq 99.8% identity): *Enterococcus raffinosus/Enterococcus gilvus; Streptococcus australis/Streptococcus sanguinis; S. pneumoniae/S. mitis; L. lactis/Leuconostoc garlicum* (Table 2).

2.3. Sanger Sequencing of the sodA Gene

Sanger sequencing of the *sodA* gene allowed identification of 12 *Streptococcus* strains (57% of all streptococcal strains), four *Enterococcus* strains (80% of all enterococcal strains), and two of the four nonstreptococcal strains (*G. adiacens, G. sanguinis*). Identification of the following pairs of enterococcal, streptococcal, and nonstreptococcal species was impossible because the *sodA* gene sequences were identical or almost identical (\geq 99.8% identity): *E. faecalis/E. faecium; S. anginosus/Streptococcus milleri; Streptococcus lutetiensis/Streptococcus infantarius; Streptococcus parasanguinis/S. oralis; S. mitis/Streptococcus cristatus; S. pyogenes/S. dysgalactiae; L. citreum/S. parasanguinis. For strain 1113/11 (<i>L. lactis*), there was no *sodA* gene reference sequence in any database (Table 2).

2.4. Sanger Sequencing of the tuf Gene

Sanger sequencing of the *tuf* gene allowed identification of 13 *Streptococcus* strains (62% of all streptococcal strains), five *Enterococcus* strains (100% of all enterococcal strains), and three of the four nonstreptococcal strains (*G. adiacens, G. sanguinis, L. citreum*). Identification of the following pairs of enterococcal, streptococcal, and nonstreptococcal species was impossible because the *tuf* gene sequences were identical or almost identical (≥99.8% identity): *S. anginosus/S. milleri; S. infantis/S. oralis; S. lutetiensis/S. infantarius; S. oralis/S. infantarius; S. pneumoniae/S. mitis; L. lactis/L. garlicum* (Table 2).

2.5. Sanger Sequencing of the rpoB Gene

Sanger sequencing of the *rpoB* gene allowed identification of 18 *Streptococcus* strains (86% of all streptococcal strains), five *Enterococcus* strains (100% of all enterococcal strains), and all four nonstreptococcal strains (*G. adiacens*, *G. sanguinis*, *L. lactis*, *L. citreum*). Identification of the following

pairs of enterococcal, streptococcal, and nonstreptococcal species was impossible because the *rpoB* gene sequences were identical or almost identical (≥99.8% identity): *S. anginosus/Streptococcus intermedius; S australis/S. infantis; S. pseudopneumoniae/S. mitis* (Table 2).

Identified Species	Isolate No.	16S rRNA Gene	sodA Gene	<i>tuf</i> Gene	rpoB Gene
Enterococcus avium	E15	Enterococcus avium	Enterococcus avium	Enterococcus avium	Enterococcus avium
Enterococcus	F2	Enterococcus	Enterococcus	Enterococcus	Enterococcus
casseliflavus	62	casseliflavus	casseliflavus	casseliflavus	casseliflavus
Enterococcus durans	E5	Enterococcus durans	Enterococcus durans	Enterococcus durans	Enterococcus durans
Enterococcus faecalis	E28	Enterococcus faecalis	Enterococcus faecium	Enterococcus faecalis	Enterococcus faecalis
Enterococcus raffinosus	E10	Enterococcus raffinosus/gilvus	Enterococcus raffinosus	Enterococcus raffinosus	Enterococcus raffinosus
Globicatella sanguinis	1375/11	Globicatella sanguinis	Globicatella sanguinis	Globicatella sanguinis	Globicatella sanguinis
Granulicatella adiacens	PL434	Granulicatella adiacens	Granulicatella adiacens	Granulicatella adiacens	Granulicatella adiacens
Leuconostoc citreum	3696/08	Leuconostoc citreum	no amplification product	Leuconostoc citreum	Leuconostoc citreum
Leuconostoc lactis	1113/11	Leuconostoc lactis /garlicum	×	Leuconostoc lactis /garlicum	Leuconostoc lactis
Streptococcus agalactiae	S19	Streptococcus agalactiae	Streptococcus agalactiae	Streptococcus agalactiae	Streptococcus agalactiae
Streptococcus anginosus	4734/08	Streptococcus anginosus	Streptococcus anginosus	Streptococcus anginosus	Streptococcus anginosus
Streptococcus	E909/10	Streptococcus	Streptococcus	Streptococcus	Streptococcus
anginosus	5696/10	anginosus	anginosus/milleri	anginosus/milleri	anginosus/ intermedius
Streptococcus australis	1816/15	Streptococcus australis	Streptococcus australis	Streptococcus australis	Streptococcus australis
Streptococcus australis	1226/14	Streptococcus australis/sanguinis	Streptococcus australis	Streptococcus australis	Streptococcus australis
Streptococcus constellatus	1107/09	Streptococcus constellatus	Streptococcus anginosus/milleri	Streptococcus anginosus/milleri	Streptococcus constellatus
Streptococcus	6022/00	Streptococcus	Streptococcus	Streptococcus	Streptococcus
constellatus	0922/09	constellatus	anginosus/milleri	anginosus/milleri	constellatus
Streptococcus	1860/08	Streptococcus	Streptococcus	Streptococcus	Streptococcus
dysgalactiae	1000/00	dysgalactiae	dysgalactiae	dysgalactiae	dysgalactiae
Streptococcus	8190/10	Streptococcus	Streptococcus	Streptococcus	Streptococcus
dysgalactiae	0 - 2 0 0 - 2 0	dysgalactiae	dysgalactiae	dysgalactiae	dysgalactiae
Streptococcus	5010/12	Streptococcus	Streptococcus	Streptococcus	Streptococcus
gallolyticus		gallolyticus	gallolyticus	gallolyticus	gallolyticus
Streptococcus	S18	Streptococcus	Streptococcus	Streptococcus	Streptococcus
gallolyticus		gallolyticus	gallolyticus	gallolyticus	gallolyticus
Streptococcus	PL427	infantic	Streptococcus	Streptococcus oralis	Streptococcus
injuntis		injunits	Strentococcue		<i>uustrutisritijuttis</i>
Streptococcus lutetiensis	PL428	Streptococcus lutetiensis	lutetiensis/ infantarius	Streptococcus lutetiensis/ infantarius	Streptococcus lutetiensis
Streptococcus mitis	PL431	Streptococcus mitis	Streptococcus mitis	Streptococcus mitis	Streptococcus mitis
Streptococcus oralis	S16	Streptococcus oralis	Streptococcus oralis	Streptococcus oralis/infantis	Streptococcus oralis
Streptococcus oralis	S63	Streptococcus oralis	Streptococcus oralis	Streptococcus oralis/infantis	Streptococcus oralis
Streptococcus parasanguinis	1374/11	Streptococcus parasanguinis	Streptococcus parasanguinis/orali s	Streptococcus parasanguinis	Streptococcus parasanguinis
Streptococcus pneumoniae	p63	Streptococcus pneumoniae	Streptococcus pneumoniae	Streptococcus pneumoniae	Streptococcus pneumoniae

Table 2. Summary of the species identification based on 16S rRNA, *sodA*, *tuf*, and *rpoB* genes.

Streptococcus pseudopneumoniae	p41	Streptococcus pneumoniae/mitis	Streptococcus mitis/cristatus	Streptococcus pneumoniae/mitis	Streptococcus pseudopneumoniae/miti s	
Streptococcus pyogenes	S47	Streptococcus pyogenes	Streptococcus pyogenes/dysgalacti ae	Streptococcus pyogenes	Streptococcus pyogenes	
Streptococcus salivarius	PL433	Streptococcus salivarius	Streptococcus salivarius	Streptococcus salivarius	Streptococcus salivarius	

All ambiguous *Streptococcus, Enterococcus* and nonstreptococcal species are indicated in dark red color. ×—lack of reference sequences.

2.6. Analysis of the recA Gene for the Streptococcal mitis Group

The streptococcal species that belong to the mitis group (*S. pneumoniae, S. pseudopneumoniae, S. mitis, S. oralis, S. gordonii, S. sanguinis,* and *S. parasanguinis*) are closely related phylogenetically.

For precise differentiation of species within this group, sequencing of the *recA* gene was used. The specific nucleotide signatures of the 313-bp fragment of the *recA* gene sequence were compared to reference sequences in GenBank (HM572273–HM572277). Sanger sequencing of the *recA* gene allowed precise identification of strains from the mitis group, namely, *S. pneumoniae*, *S. pseudopneumoniae*, *S. mitis*, *S. oralis*, and *S. infantis*. The alignment showed six specific nucleotides at positions 97, 160, 199, 247, 250, and 280 (Figure 1). The nucleotide signature is based on homology analyses of *recA* gene sequences from reference strains of the aforementioned species and our strains. The *recA* gene sequence of the p41 strain was almost identical to the reference sequence (*S. pseudopneumoniae*), with a one-nucleotide difference at position 280. For PL427, differences at two nucleotide positions were observed in comparison to *S. infantis*. The only method that allowed unambiguous identification of *S. pseudopneumoniae* was Sanger sequencing of the *recA* gene.

	Specific nucleotide position					
	313-bp fragment of <i>recA</i> /gene				ne	
	Position					
Species	97	160	199	247	250	280
Streptococcus pseudopneumoniae						
Ref. strain: Streptococcus pseudopneumoniae ATCC BAA-960(HM572274)	А	A	C	Т	Т	С
Our strain: p41 (MT560905)	А	A	С	Т	Т	Т
Streptococcus pneumoniae						
Ref. strain: Streptococcus pneumoniae NCTC 7465 (HM572273)	G	Т	Т	С	C	Т
Our strain: p63 (MT560906)	G	Т	Т	С	C	Т
Streptococcus infantis						
Ref. strain: Streptococcus infantis ATCC 700779 (HM572277)	А	A	С	Т	Т	С
Our strain: PL427 (MT560907)	А	Т	С	Т	Т	Т
Streptococcus mitis						
Ref. strain: Streptococcus mitis NCTC 12261 (HM572275)	А	C	C	Т	Т	С
Our strain: PL431 (MT560908)	А	С	С	Т	Т	С
Streptococcus oralis						
Ref. strain: Streptococcus oralis NCTC 11427 (HM572276)	Т	A	C	А	Т	С
Our strain: S63 (MT560909)	Т	А	С	А	Т	С

Figure 1. Alignment of specific nucleotides for the streptococcal mitis complex observed in the 313bp *recA* fragment.

2.7. Comparison of the Sequencing Methods

The combination of sequencing methods based on the 16S rRNA, *sodA*, *tuf*, *rpoB*, and *recA* genes allowed the identification of 21 streptococcal strains, five enterococcal strains, two *Leuconostoc* strains, one *Globicatella sanguinis* strain, and one *Granulicatella adiacens* strain. Due to high (or identical) similarity or a lack of similarity with the reference sequences in GenBank and leBIBIQBPP, it was not possible to identify all the strains at the species level by using the targets separately (Table 2).

For *Streptococcus*, Sanger sequencing of the 16S rRNA gene had the highest identification potential, allowing the identification of 19 (90%) strains. Additionally, *rpoB* gene sequencing had high discriminative potential, allowing the identification of 18 (86%) *Streptococcus* strains. Sanger sequencing of the *tuf* gene had moderate identification potential and identified 13 (62%) streptococcal strains. Sanger sequencing of the *sodA* gene had the lowest discriminatory potential, allowing the identification.

Sanger sequencing of *rpoB* and *tuf* allowed the identification of five (100%) analyzed enterococcal strains. Sequencing of the 16S rRNA and *sodA* genes had moderate identification potential and allowed the identification of four (80%) enterococcal strains (Table 3).

Streptoco	<i>ccus</i> spp. (<i>n</i> = 21 Str	Unambiguous	No Identification at	
<i>Enterococcus</i> spp. (<i>n</i> = 5 Strains)			Species Identification	the Species Level
	16S rPNA gong -		4 (80%)	1 (20%)
	165 IKINA gene	Streptococcus	19 (90%)	2(10%)
	and A mama	Enterococcus	4 (80%)	1 (20%)
Company Conversion	souA gene	Streptococcus	12 (57%)	9 (43%)
Sanger Sequencing	tul cono	Enterococcus	5 (100%)	0 (0%)
	<i>tuj</i> gene	Streptococcus	13 (62%)	8 (38%)
	ma P como	Enterococcus	5 (100%)	0 (0%)
	rpob gene	Streptococcus	18 (86%)	3 (14%)

Table 3. Summary of the performance of 16S rRNA, *sodA*, *tuf*, and *rpoB* genes sequencing used for differentiation of *Streptococcus* and *Enterococcus* genera.

2.8. Phylogenetic Analysis of Streptococcus and Enterococcus

To show the relationships among the species, phylogenetic trees were constructed. The evolutionary distances were computed using the Jukes–Cantor method and are shown in units of the number of base substitutions per site. The computed overall means for the 16S rRNA, *rpoB*, *soda*, and *tuf* genes were 0.098, 0.225, 0.348, and 0.176, respectively. In the phylogenetic tree constructed for the *tuf* gene, the *Leuconostoc* species sequences are shorter because sequences of the same length as those of other species could not be obtained. Both streptococci and enterococci are grouped into separate clusters. Moreover, the *Streptococcus* strains are divided into mitis, bovis, and anginosus complexes. Sequencing of the 16S rRNA, *rpoB*, and *tuf* genes showed that *L. lactis*, *L. citreum*, *G. sanguinis*, and *G. adiacens* were distantly related to the other species (Figures 2–5).

16S rRNA gene



Figure 2. The phylogenetic tree of streptococcal and enterococcal species evolutionary relationships based on the 16S rRNA gene. The phylogenetic trees were constructed using the neighbor-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Jukes–Cantor method and are in the units of the number of base substitutions per site. The strains which are placed in boxes have grouped together in all methods used. The length of the compared sequences was 1296 bp.



Figure 3. The phylogenetic tree of streptococcal and enterococcal species evolutionary relationships based on the *sodA* gene. The phylogenetic trees were constructed using the neighbor-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Jukes–Cantor method and are in the units of the number of base substitutions per site. The strains which are placed in boxes have grouped together in all methods used. The length of the compared sequences was 418 bp.





Figure 4. The phylogenetic tree of streptococcal and enterococcal species evolutionary relationships based on the *tuf* gene. The phylogenetic trees were constructed using the neighbor-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Jukes–Cantor method and are in the units of the number of base substitutions per site. The strains which are placed in boxes have grouped together in all methods used. The length of the compared sequences was 770 bp.

rpoB gene



Figure 5. The phylogenetic tree of streptococcal and enterococcal species evolutionary relationships based on the *rpoB* gene. The phylogenetic trees were constructed using the neighbor-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Jukes–Cantor method and are in the units of the number of base substitutions per site. The strains which are placed in boxes have grouped together in all methods used. The length of the compared sequences was 666 bp.

3. Discussion

Because of the variability of strains and challenging taxonomic changes of *Streptococcus* and *Enterococcus* species, it is necessary to use a reliable identification method to better understand the pathogenic potential of various streptococcal and enterococcal species. The currently used phenotypic identification methods based on morphological and biochemical characteristics appear to be unreliable and are characterized by low discriminatory potential [33–35].

12 of 21

In this study, we applied biochemical methods and genetic sequencing-based methods to identify clinically relevant *Streptococcus* and *Enterococcus* species. We showed that the Vitek® 2 system and MALDI-TOF MS did not correctly identify particular closely related species, such as *S. mitis, S. oralis,* and other species of the mitis group. Overall, the phenotypic methods allowed the identification of 48% of streptococcal and 100% of enterococcal strains. These data are consistent with previous data in the literature [19,36–39]. Therefore, applying genetic methods in standard microbiological diagnostics can lead to unambiguous confirmation at the species level. Genotypic methods utilizing Sanger sequencing of targeted genes were shown to be useful for both *Streptococcus* and *Enterococcus* identification [13,25]. 16S rRNA is mostly used to identify unknown organisms because of the availability of universal primers [23,40]. However, most reports show that the discriminatory power of 16S rRNA gene sequencing is very low for closely related streptococcal and enterococcal species [13,41,42]. Analysis based on only one gene target is not recommended because duplication, gene transfer, and gene loss can affect the reliability of the results [43,44].

In this study, we used a combination of four gene targets (16S rRNA, *sodA*, *tuf*, *rpoB*) to unambiguously confirm the identity at the species level for 21 streptococci and five enterococcal strains. None of the individual sequencing-based methods allowed the identification of all species. In our study, Sanger sequencing of the 16S rRNA gene had the highest discriminatory power, allowing unambiguous identification of 19 (90%) of the analyzed streptococcal strains, but the *rpoB* gene had almost identification potential, allowing the identification of 18 (86%) *Streptococcus* strains. For *Enterococcus* strains, Sanger sequencing of the *tuf* and *rpoB* genes allowed the identification of five (100%) strains. The 16S rRNA and *sodA* genes did not allow identification of all *Enterococcus* strains, but in our study, this group was very small (only five strains).

Over the years, the taxonomy of bacteria has changed, and streptococcal groups, i.e., mitis and bovis, have undergone several reclassifications. Moreover, incorrect systematic names of bacteria have been deposited in publicly available databases [45]. In our study, several problematic situations occurred. First, *Streptococcus tigurinus* was classified as *S. oralis* subsp. *tigurinus*, but in 2012, this species was separated into two different species. Finally, in 2016, it was again proposed that this species be classified as *S. oralis* subsp. *tigurinus* [27,46]. Our sequence was aligned to the sequence of *S. oralis*, but the next closest species was *S. tigurinus*. Incorrect taxonomic annotations of DNA sequences are often present in databases [45]. A similar situation was found for *S. lutetiensis* (PL428 strain), which was described as *S. infantarius* subsp. *coli* based on the *sodA* and *tuf* genes. In 2005, the International Committee on Systematics of Prokaryotes (Status of strains that contravene Rules 27 (3) and 30 of the International Code of Nomenclature of Bacteria, Opinion 81) accepted *S. lutetiensis* as the correct systematic name [47], but in databases, double taxonomic annotation for one organism can be found.

The *Enterococcus* strain E28 (*E. faecalis*) *sodA* gene sequence matched *E. faecium* (412/412 nucleotide identity). In our study, such a situation did not occur for other gene targets, yet it has been reported in the literature [48,49]. On the other hand, for strain E10 (*E. raffinosus*), the 16S rRNA gene sequence matched two enterococcal species, namely, *E. gilvus* (1289/1290 nucleotide identity) and *E. raffinosus* (1289/1291 nucleotide identity).

For *Streptococcus*, there were also some problematic cases in the anginosus group (also known as the *S. milleri* group). Strain 5898/10 was identified as *S. anginosus* by 16S rRNA gene sequencing, but other molecular methods showed ambiguous identification among the *S. anginosus-S. milleri-S. intermedius* species. Such a situation was observed by others [50,51]. A similar problem was observed in the identification of the 1107/08 and 6922/09 strains. Only 16S rRNA and *rpoB* allowed *Streptococcus constellatus* identification, while for the *sodA* and *tuf* genes, our strain sequences shared high nucleotide similarities with both the *S. anginosus* and *S. milleri* sequences. The *Streptococcus milleri* group proved to be challenging to identify in previous studies [51,52].

Both phenotypic and genetic methods correctly identified the nonstreptococcal species as *Globicatella sanguinis, Granulicatella adiacens, Leuconostoc citreum,* and *Leuconostoc lactis. Globicatella sanguinis* was initially described as *Streptococcus uberis* and *Aerococcus viridans* due to similar phenotypic properties. The advanced methods allowed the distinguishing and classification of *G*.

sanguinis into a new species [9,53–55]. In our study, this species was identified by all four gene targets (16S rRNA, *sodA*, *rpoB*, and *tuf*).

Granulicatella adiacens was first described as *Streptococcus adjacens* and then as belonging to the *Abiotrophia* genus due to distant relations with streptococci. Collins and Lawson proposed a new genus, *Granulicatella*, due to significant differences [56,57]. In our study, strain PL434 was identified as *G. adiacens* by all sequencing methods.

The *Leuconostoc* genus is often identified as *Streptococcus* spp. Because similar biochemical properties and serotypes of the D group are observed, *Leuconostoc* species are difficult to detect with routine diagnostic methods [9]. It has been suggested that *Leuconostoc* is a pathogen that colonizes the gastrointestinal tract and is present in neutropenic patients [58,59]. For the *Leuconostoc* genus, strain 1113/11 was correctly identified by the Vitek[®] 2 system and based on the *rpoB* gene, but the 16S rRNA and *tuf* genes were ambiguous between *L. lactis* and *L. garlicum*. For the *sodA* gene, there was no *L. lactis* reference sequence available in databases, but the sequence was identical to *S. parasanguinis*. Such results were not observed by other research groups, but our study showed that in some cases the distinction between two bacterial genera is not possible by only one molecular target. For both *Leuconostoc* strains (1113/11 and 3696/08), the other set of primers for *tuf* gene amplification had to be used [60].

Strain 3696/08 was correctly identified as *L. citreum* by 16S rRNA, *tuf*, and *rpoB* gene sequencing, but amplification of the *sodA* gene was problematic. The primers d1 and d2 [24] used for the *sodA* gene in other *Streptococcus* strains did not result in PCR product amplification.

In our study, *S. pseudopneumoniae* was not identified by any of the four Sanger sequencing-based or phenotypic methods. Arbique et al. and Harf-Monteil et al. observed similarity between the isolates identified as *S. pseudopneumoniae* and *S. pneumoniae*, which demonstrated a high degree of homology and shared phenotypic characteristics [61,62]. However, in 2011, Zbinden et al. suggested that sequencing of the *recA* gene could differentiate between *S. pneumoniae* and *S. pseudopneumoniae* [31]. In our study, in addition to identification of the *Streptococcus mitis* group, we used Sanger sequencing of the *recA* gene, which successfully confirmed the identities of the *S. pseudopneumoniae*, *S. pneumoniae*, *S. mitis*, *S. oralis*, and *S. infantis* species. Moreover, it was the only method that correctly identified the p41 strain as *S. pseudopneumoniae*.

In *Streptococcus* species genetic diagnostics, other molecular target such as sequencing of the *ddl* or *gdh* genes could also be used [63,64]. However, these targets are not commonly used and are usually used for identification of specific species groups [65,66]. The advanced molecular diagnostics precision should definitely be strengthened with methods based on next-generation sequencing, but the costs and challenging data analysis are the pitfalls of these methods to be used in routine diagnostic laboratories [67].

To conclude, phenotypic methods such as the Vitek[®] 2 system and MALDI-TOF MS constitute basic methods because the results are received after approximately 8 h and are characterized by lower costs than those of genetic methods. However, Sanger sequencing and PCR-based approaches proved to be excellent tools for identification at the species level for both *Streptococcus* and *Enterococcus* strains. We also proved that the use of only one method is often not enough for appropriate identification at the species level.

4. Materials and Methods

4.1. Ethical Approval

This article does not contain any studies with human participants or animals performed by any of the authors.

4.2. Bacterial Isolates

The collection of bacterial isolates included 30 isolates of 15 *Streptococcus* species, five *Enterococcus* species, two *Leuconostoc* species, and one isolate each from *Globicatella* and *Granulicatella*

species recovered from clinical origin (Table 4). Most isolates were recovered from the National Medicines Institute in Warsaw (n = 13), with five isolates from the University Medical Center Groningen and 12 from Pescara Local Hospital. The isolates were cultured on blood agar medium with 5% sheep blood (bioMérieux, La Balme Les Grottes, France) and incubated at 37 °C in an atmosphere of 5% CO₂ for 20 h.

	Species	Isolate Number
	Enterococcus avium	E15 c
Futanacas	Enterococcus casseliflavus	E2 °
Enterococcus	Enterococcus durans	E5 °
n - 5	Enterococcus faecalis	E28 c
	Enterococcus raffinosus	E10 c
	Globicatella sanguinis	1375/11 ª
Other species	Granulicatella adiacens	PL434 ^b
n = 4	Leuconostoc citreum	3696/08 ª
	Leuconostoc lactis	1113/11 ª
	Streptococcus agalactiae	S19 c
	Churche and an a in and	4734/08 a
	Streptococcus unginosus	5898/10 ª
		1816/15 ª
	Streptococcus australis	1226/14ª
		1107/09 a
	Streptococcus constellatus	6922/09 a
		1860/08 ª
	Streptococcus aysgalactiae	8190/10 ª
		5010/12 ª
Streptococcus	Streptococcus gallolyticus	S18 c
n = 21	Streptococcus infantis	PL427 ^b
	Streptococcus lutetiensis	PL428 b
	Streptococcus mitis	PL431 ь
	Chumphococcus ouglis	S16 c
	Streptococcus oruns	S63 c
	Streptococcus parasanguinis	1374/11 ª
	Streptococcus pneumoniae	р63 ^с
	Streptococcus pseudopneumoniae	p41 °
	Streptococcus pyogenes	- S47 ۲
	Streptococcus salivarius	PL433 b

Table 4. Streptococcus and	Enterococcus species	used for analyses.
----------------------------	----------------------	--------------------

^a National Medicines Institute in Warsaw; ^b University Medical Center Groningen; ^c Pescara Local Hospital–Italy.

4.3. Phenotypic Identification Tests

All isolates were identified using two phenotypic tests. The Vitek[®] 2 system (bioMérieux, La Balme Les Grottes, France) was used to identify isolates at the genus and species levels. The suspension used in the Vitek[®] 2 system was adjusted to a McFarland standard of 0.5 by using a densitometer and interpreted according to the manufacturer's instructions. A score of \geq 96% indicated excellent species identification; 91–95% indicated very good species identification. A score of 89–92% indicated good species identification. For streptococcal serotype identification, the PastorexTM Strep Test Kit (Bio-Rad, Hercules, CA, USA) was used. The bacterial cells were suspended in 300 µL of enzymatic extract and incubated at 37 °C for 15 min. After incubation, the reagent with antibodies

and bacterial suspension was applied to identification cards and mixed. The results were read after 30 s.

4.4. MALDI-TOF MS Identification

The MicroFlex MALDI-TOF mass spectrometer with MALDI Biotyper software 2.0 (Bruker Daltonics, Bremen, Germany) was used for isolate identification. Identification of isolates PL434, 1113/11, 3696/08, p41, and 1375/11 using MALDI-TOF MS was performed by The Microbiological Laboratory of the Jagiellonian Center of Innovation (Krakow, Poland). Sample extraction and strain identification were performed following the manufacturer's instructions. A score of >2 indicated correct genus and probable species identification.

4.5. Genomic DNA Extraction

The Qiagen DNeasy Blood & Tissue Kit (Qiagen, Germantown, MD, USA) was used for genomic DNA extraction. Bacteria were homogenized with a TissueLyser II (Qiagen, Germantown, MD, USA) for five minutes at a frequency of 50 Hz. After homogenization, the tubes were centrifuged for 10 min at 13 200 rpm. The subsequent steps were performed according to the manufacturer's instructions.

4.6. PCR Amplification of the 16S rRNA, sodA, rpoB tuf, and recA Genes

Both bacterial DNA and the negative control (nuclease-free H₂O (EurX—Molecular Biology Products, Gdansk, Poland)) were amplified with primers for a given locus. As shown in Tables 5 and 6, primers specific for the targeted locus were used as described previously [21,24–26,31,60,68]. Based on our previous studies, the PCR programs were modified slightly to obtain increased product quality [13].

All PCR products were resolved by electrophoresis in 1% agarose with 1× TAE and then purified using the DNA Clean & Concentrator[™] Kit (Zymo Research, Irvine, CA, USA; A&A Biotechnology, Gdynia, Poland). Concentrations and purity were measured using a NanoDrop ND-1000. Sanger sequencing was performed at GATC Eurofins Genomics (Ebersberg, Germany) and Genomed S.A. (Warsaw, Poland) with the same primers as those used for PCR (Tables 5 and 6).

Target Gene		PCR Program	Cycles (Steps 2–4)	Amplification Primers (5'→3')	Amount of Sequenced PCR Product	Reference
16S	1. 2.	94 °C for 2 min 94 °C for 30 s 58 °C for 20 s	25	LPW57 AGTTTGATCCTGGCTCAG	250 ng	[69]
rKNA 3. (1284-bp) 4. 5.	3. 4. 5.	72 °C for 1 min 72 °C for 5 min	25 -	LPW58 AGGCCCGGGAACGTATTCAC	- 250 lig	[66]
sodA	1. 2. 3	95 °C for 3 min 95 °C for 30 s 43 °C for 1 min A	35 -	d1 CCITAYICITAYGAYGCIYTIGARCC	- 100 ng	[24]
(430-bp)	3. 4. 5.	72 °C for 1 min 30 s 72 °C for 10 min	55	d2 ARRTARTAIGCRTGYTCCCAIACRTC	- 100 llg	
tuf	1. 2. 3	95 °C for 2 min 94 °C for 30 s 50 °C for 30 s B	30 -	tuf-F CCAATGCCACAAACTCGT	- 200 ng	[25]
(830-bp)	5. 5.	72 °C for 1 min 30 s 72 °C for 10 min	50	tuf-R CCTGAACCAACAGTACGT	- 200 Hg	[23]
<i>гроВ</i> (860-bp)	1. 2. 3	95 °C for 2 min 94 °C for 30 s 52 °C for 30 s	35	Strepto-F AARYTIGGMCCTGAAGAAAT	200	[0/]
	3. 4. 5.	72 °C for 1 min 72 °C for 5 min		Strepto-R TGIARTTTRTCATCAACCATGTG	- 200 lig	[20]

Table 5. Oligonucleotide sequencing primers and PCR program used for amplification of 16S rRNA, *sodA*, *tuf*, *rpoB* and *recA* genes.

5

72 °C for 10 min

recA (850-bp)

1.	95 °C for 2 min		RStrGseq81			
2.	94 °C for 30 s		GAAAWWIATYGARAAAGAITTTGGTA			
3.	54 °C for 30 s	30	А	150 ng	[31]	
4.	72 °C for 1 min 30 s		RStrGseq937			

TTYTCAGAWCCTTGICCAATYTTYTC

^A **40** °C (strains: S18, S19, 6922/08, 1226/14, 5898/10, PL434, PL434, E10) **45** °C (strain: 1860/08) **50** °C (strains: S16, 1816/15, 8190/10) **52** °C (strain: p41) ^B **53** °C (strains: 1226/14, PL427) **59** °C (strains: E10, E15).

Table 6. Alternative oligonucleotide sequencing primers and PCR program used for amplification of *sodA* and *tuf* genes.

$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Target Gene		PCR Program	Cycles (Steps 2–4)	Amplification Primers (5'→3')	Amount of Sequenced PCR Product	Reference
$\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} $	sodA	1. 2.	94 °C for 5 min 94 °C for 30 s		sodA-F TRCAYCATGAYAARCACCAT		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	(430-bp)	3. 4. 5.	50 °C for 1 min ^c 72 °C for 30 s 72 °C for 5 min	30	sodA-R ARRTARTAMGCRTGYTCCCARACRTC	100 ng	[21]
$\begin{array}{c} 3. & 55 \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	<i>tuf</i> (830-bp)	1. 2.	95 °C for 3 min 95 °C for 30 s	25	U1 AAYATGATIACIGGIGCIGCICARATGGA	200	[60]
5. 72°C for 7 min		3. 4. 5.	55 °C for 30 s b 72 °C for 1 min 72 °C for 7 min	35	U2 AYRTTITCICCIGGCATIACCAT	200 ng	

^c 50 °C (strain: 3696/08) ^D 55 °C (strains: 1113/11, 3696/08).

4.7. Sanger Sequencing Analysis of the 16S rRNA, sodA, rpoB, and tuf Genes

The Sanger sequencing results were analyzed using Chromas software (version: 2.6.6). Nucleotide BLAST (Basic Local Alignment Search Tool http://www.ncbi.nlm.nih.gov/BLAST/) was used to analyze the obtained sequences and align them to the reference sequences deposited in the GenBank (https://www.ncbi.nlm.nih.gov/nucleotide/) and leBIBIQBPP (leBIBI-Quick BioInformatic Phylogeny of Prokaryotes) (https://umr5558-bibiserv.univ-lyon1.fr/lebibi/lebibi.cgi) databases. The first and second best species alignments were analyzed. To identify the selected strain at the species level, the criterion of \geq 99% first best match with the reference database and a difference of at least two nucleotides between the first and second best matches was used [13,69]. All sequences were aligned in ClustalW. The phylogenetic trees were constructed using the neighbor-joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) and evolutionary distances were computed using the Jukes–Cantor method (MEGA, version 7.0.26, Pennsylvania State University, State College, PA, USA). Pairwise comparison of each pair of sequences was performed using CLC Genomics Workbench (version 8.1, Qiagen, USA).

4.8. Sanger Sequencing Analysis of the recA Gene

The obtained *recA* gene sequences (313 bp) for the S63, PL427, PL431, p63, and p41 strains were analyzed at six specific nucleotide positions (97, 160, 199, 247, 250, and 280). For precise differentiation of species within the mitis complex, the reference sequences of the *recA* genes from *S. pneumoniae* NCTC 7465, *Streptococcus mitis* NCTC 12261, *Streptococcus oralis* NCTC 11427, *Streptococcus pseudopneumoniae* ATCC BAA-960, and *Streptococcus infantis* ATCC 700779 (reference numbers in GenBank: HM572273, HM572275, HM572276, HM572274, HM572277, respectively) were used [25].

4.9. Nucleotide Sequence Accession Numbers

The 124 sequences for 21 *Streptococcus*, five *Enterococcus*, and four other species were annotated using the NCBI BankIt tool and deposited in the GenBank database (https://www.ncbi.nlm.nih.gov/genbank/) under the following accession numbers: for the 16S rRNA gene, MT535599-MT535603, MT535764 and MT535859-MT535882; for the *sodA* gene, MT560910-

MT560938; for the *tuf* gene, MT560846-MT560874 and MT707819; for the *rpoB* gene, MT560875-MT560904; and for the *recA* gene, MT560905-MT560909.

Author Contributions: Conceptualization, M.K.-S.; methodology, M.K.-S.; validation, M.K.-S., M.W. and J.M.; formal analysis, M.K.-S.; investigation, M.K.-S. and M.W.; resources, D.Ż., E.S. and J.M.; data curation, M.K.-S.; writing—original draft preparation, M.K.-S. and M.W.; writing—review and editing, M.K.-S., M.W., D.Ż., E.S. and J.M.; visualization, M.K.S. and M.W.; supervision, M.K.-S.; project administration, M.K.-S.; funding acquisition, M.K.-S. All authors have read and agreed to the published version of the manuscript.

Funding: This project was partly financed by funds granted by the National Science Centre (NCN, Poland) on the basis of the decision no. UMO-2016/21/N/NZ6/00981 (for M.K.S.).

Acknowledgments: The authors are thankful to Artur Sabat from the Department of Medical Microbiology, University Medical Center Groningen, University of Groningen, The Netherlands, and Vincenzo Savini from Clinical Microbiology and Virology, Spirito Santo Hospital, Pescara, Italy, for providing some *Streptococcus* and *Enterococcus* strains.

Conflicts of interest: The authors declare that they have no conflict of interest.

References

- Lal, D.; Verma, M.; Lal, R. Exploring internal features of 16S rRNA gene for identification of clinically relevant species of the genus *Streptococcus. Ann. Clin. Microbiol. Antimicrob.* 2011, 10, 28, doi:10.1186/1476-0711-10-28.
- 2. Kohler, W. The present state of species within the genera *Streptococcus* and *Enterococcus*. *Int. J. Med. Microbiol.* **2007**, 297, 133–150, doi:10.1016/j.ijmm.2006.11.008.
- 3. Gao, X.Y.; Zhi, X.Y.; Li, H.W.; Klenk, H.P.; Li, W.J. Comparative genomics of the bacterial genus *Streptococcus* illuminates evolutionary implications of species groups. *PLoS ONE* **2014**, *9*, e101229, doi:10.1371/journal.pone.0101229.
- 4. Farley, M.M.; Strasbaugh, L.J. Group B streptococcal disease in nonpregnant adults. *Clin. Infect. Dis.* **2001**, 33, 556–561, doi:10.1086/322696.
- 5. Henriques-Normark, B.; Tuomanen, E.I. The pneumococcus: Epidemiology, microbiology, and pathogenesis. *Cold Spring Harb. Perspect. Med.* **2013**, *3*, a010215, doi:10.1101/cshperspect.a010215.
- Pekuz, S.; Soysal, A.; Akkoc, G.; Atıcı, S.; Yakut, N.; Gelmez, G.A.; Kadayifci, E.K.; Güneser, D.; Demir, S.O.; Söyletir, G.; et al. Prevalence of nasopharyngeal carriage, serotype distribution, and antimicrobial resistance of *Streptococcus pneumoniae* among children with chronic diseases. *Jpn. J. Infect. Dis.* 2019, 72, 7– 13, doi:10.7883/yoken.JJID.2017.410.
- Gerhardt, P.; Murray, R.G.E.; Willis, A.; Krieg, N.R. Methods for General and Molecular Bacteriology, 2nd ed.; Am. Soc. Microbiol.: Washington, DC, USA, 1994; pp. 619–647.
- Kawamura, Y.; Hou, X.G.; Sultana, F.; Miura, H.; Ezaki, T. Determination of 16S rRNA sequences of Streptococcus mitis and Streptococcus gordonii and phylogenetic relationships among members of the genus Streptococcus. Int. J. Syst. Bacteriol. 1995, 45, 406–408, doi:10.1099/00207713-45-2-406.
- 9. Facklam, R.; Elliott, J.A. Identification, classification, and clinical relevance of catalase-negative, grampositive cocci, excluding the streptococci and enterococci. *Clin. Microbiol. Rev.* **1995**, *8*, 479–495.
- 10. Parks, T.; Barrett, L.; Jones, N. Invasive streptococcal disease: A review for clinicians. *Br. Med. Bull.* **2015**, *115*, 77–89, doi:10.1093/bmb/ldv027.
- 11. Schleifer, K.H.; Kilpper-Balz, R. Transfer of *Streptococcus faecalis* and *Streptococcus faecium* to the genus *Enterococcus* nom. rev. as *Enterococcus faecalis* comb. nov. and *Enterococcus faecium* comb. nov. *Int. J. Syst. Evol.* **1984**, 34, 31–34, doi:10.1099/00207713-34-1-31.
- 12. Devriese, L.; Baele, M.; Butaye, P. The Genus Enterococcus. In *The Prokaryotes*; Springer: Berlin/Heidelberg, Germany, 2006; pp. 163–174, doi:10.1007/0-387-30744-3_5.
- Kosecka-Strojek, M.; Sabat, A.J.; Akkerboom, V.; Kooistra-Smid, A.M.D.M.; Miedzobrodzki, J.; Friedrich, A.W. Development of a reference data set for assigning *Streptococcus* and *Enterococcus* species based on next generation sequencing of the 16S-23S rRNA region. *Antimicrob Resist. Infect. Control.* 2019, *8*, 178, doi:10.1186/s13756-019-0622-3.
- 14. Savini, V.; Franco, A.; Gherardi, G.; Marrollo, R.; Argentieri, A.V.; De Araujo, F.P.; Amoruso, R.; Battisti, A.; Fazii, P.; Carretto, E. Beta-hemolytic, multi-lancefield antigen-agglutinating *Enterococcus durans* from a

pregnant woman, mimicking *Streptococcus agalactiae*. J. Clin. Microbiol. **2014**, 52, 2181–2182, doi:10.1128/JCM.00296-14.

- 15. Ferede, Z.T.; Tullu, K.D.; Derese, S.G.; Yeshanew, A.G. Prevalence and antimicrobial susceptibility pattern of *Enterococcus* species isolated from different clinical samples at Black Lion Specialized Teaching Hospital, Addis Ababa, Ethiopia. *BMC Res. Notes* **2018**, *11*, 793, doi:10.1186/s13104-018-3898-0.
- Top, J.; Willems, R.; Bonten, M. Emergence of CC17 *Enterococcus faecium*: From commensal to hospitaladapted pathogen. *FEMS Immunol. Med. Microbiol.* 2008, 52, 297–308, doi:10.1111/j.1574-695X.2008.00383.x.
- 17. Fernández-Hidalgo, N.; Escolà-Vergé, L. *Enterococcus faecalis* Bacteremia: Consider an echocardiography, but consult an infectious diseases specialist. *J. Am. Coll. Cardiol.* **2019**, *74*, 202–204, doi:10.1016/j.jacc.2019.03.526.
- Ligozzi, M.; Bernini, C.; Bonora, M.G.; De Fatima, M.; Zuliani, J.; Fontana, R. Evaluation of the VITEK 2 system for identification and antimicrobial susceptibility testing of medically relevant gram-positive cocci. *J. Clin. Microbiol.* 2002, 40, 1681–1686, doi:10.1128/jcm.40.5.1681-1686.2002.
- Chen, J.H.; She, K.K.; Wong, O.Y.; Teng, J.L.; Yam, W.C.; Lau, S.K.; Woo, P.C.; Cheng, V.C.; Yuen, K.Y. Use of MALDI Biotyper plus ClinProTools mass spectra analysis for correct identification of *Streptococcus pneumoniae* and *Streptococcus mitis/oralis. J. Clin. Pathol.* 2015, *68*, 652–656, doi:10.1136/jclinpath-2014-202818.
- 20. Singhal, N.; Kumar, M.; Kanaujia, P.K.; Virdi, J.S. MALDI-TOF mass spectrometry: An emerging technology for microbial identification and diagnosis. *Front. Microbiol.* **2015**, *6*, 791.
- 21. Hoshino, T.; Fujiwara, T.; Kilian, M. Use of phylogenetic and phenotypic analyses to identify nonhemolytic streptococci isolated from bacteremic patients. *J. Clin. Microbiol.* **2005**, *43*, 6073–6085, doi:10.1128/JCM.43.12.6073-6085.2005.
- Isaksson, J.; Rasmussen, M.; Nilson, B.; Stadler, L.S.; Kurland, S.; Olaison, L.; Ek, E.; Herrmann, B. Comparison of species identification of endocarditis associated viridans streptococci using *rnpB* genotyping and 2 MALDI-TOF systems. *Diagn. Microbiol. Infect. Dis.* 2015, *81*, 240–245, doi:10.1016/j.diagmicrobio.2014.12.007.
- 23. Clarridge, J.E. Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. *Clin. Microbiol. Rev.* **2004**, *17*, 840–862, doi:10.1128/CMR.17.4.840-862.2004.
- 24. Poyart, C.; Quesne, G.; Coulon, S.; Berche, P.; Trieu-Cuot, P. Identification of streptococci to species level by sequencing the gene encoding the manganese-dependent superoxide dismutase. *J. Clin. Microbiol.* **1998**, *36*, 41–47.
- 25. Li, X.; Xing, J.; Li, B.; Wang, P.; Liu, J. Use of *tuf* as a target for sequence-based identification of Grampositive cocci of the genus *Enterococcus*, *Streptococcus*, coagulase-negative *Staphylococcus*, and *Lactococcus*, *Ann. Clin. Microbiol. Antimicrob.* **2012**, *11*, 31, doi:10.1186/1476-0711-11-31.
- 26. Drancourt, M.; Roux, V.; Fournier, P.E.; Raoult, D. *rpoB* gene sequence-based identification of aerobic Gram-positive cocci of the genera *Streptococcus*, *Enterococcus*, *Gemella*, *Abiotrophia*, and *Granulicatella*. J. Clin. *Microbiol*. **2004**, *42*, 497–504, doi:10.1128/JCM.42.2.497-504.2004.
- 27. Jensen, A.; Scholz, C.F.P.; Kilian, M. Re-evaluation of the taxonomy of the Mitis group of the genus *Streptococcus* based on whole genome phylogenetic analyses, and proposed reclassification of *Streptococcus alis subsp. dentisani* as *Streptococcus oralis* subsp. *dentisani* comb. nov., *Streptococcus tigurinus* as Streptococcus oralis subsp. *tigurinus* comb. nov., and *Streptococcus oligofermentans* as a later synonym of *Streptococcus cristatus*. *Int. J. Syst. Evol. Microbiol.* **2016**, *66*, 4803–4820, doi:10.1099/ijsem.0.001433.
- 28. Sadowy, E.; Hryniewicz, W. Identification of *Streptococcus pneumoniae* and other Mitis streptococci: Importance of molecular methods. *Eur. J. Clin. Microbiol. Infect. Dis.* **2020**, doi:10.1007/s10096-020-03991-9.
- 29. Dekker, J.P.; Lau, A.F. An Update on the *Streptococcus bovis* Group: Classification, Identification, and Disease Associations. *J. Clin. Microbiol.* **2016**, *54*, 1694–1699, doi:10.1128/JCM.02977-15.
- Werno, A.M.; Christner, M.; Anderson, T.P.; Murdoch, D.R. Differentiation of *Streptococcus pneumoniae* from nonpneumococcal streptococci of the *Streptococcus mitis* group by Matrix-Assisted Laser Desorption Ionization–Time of Flight Mass Spectrometry. *J. Clin. Microbiol.* 2012, *50*, 2863–2867, doi:10.1128/JCM.00508-12.
- 31. Zbinden, A.; Köhler, N.; Bloemberg, G.V. *recA*-based PCR assay for accurate differentiation of *Streptococcus pneumoniae* from other viridans streptococci. *J. Clin. Microbiol.* **2011**, *49*, 523–527, doi:10.1128/JCM.01450-10.
- 32. BIO-RAD. 2017. Manufacturer's Instructions: Agglutination Test for the Grouping Streptococci Belonging to Groups A, B, C, D, F, G. PASTOREXTM STREP. Available online:

https://www.biorad.com/webroot/web/pdf/inserts/CDG/en/61721_880999_EN.pdf (accessed on 6 November 2020).

- 33. Gherardi, G.; Angeletti, S.; Panitti, M.; Pompilio, A.; Di Bonaventura, G.; Crea, F.; Avola, A.; Fico, L.; Palazzo, C.; Sapia, G.F.; et al. Comparative evaluation of the Vitek-2 Compact and Phoenix systems for rapid identification and antibiotic susceptibility testing directly from blood cultures of Gram-negative and Gram-positive isolates. *Diagn. Microbiol. Infect. Dis.* 2012, *72*, 20–31, doi:10.1016/j.diagmicrobio.2011.09.015.
- 34. de Cueto, M.; Ceballos, E.; Martinez-Martinez, L.; Perea, E.J.; Pascual, A. Use of positive blood cultures for direct identification and susceptibility testing with the Vitek 2 system. *J. Clin. Microbiol.* **2004**, *42*, 3734–3738, doi:10.1128/JCM.42.8.3734-3738.2004.
- 35. Chen, J.; Lee, S.; Yang, B.; Lu, J. Rapid identification and susceptibility testing using the VITEK 2 system using culture fluids from positive BacT/ALERT blood cultures. *J. Microbiol. Immunol. Infect.* **2008**, *41*, 259–264.
- Romero-Gómez, M.P.; Gómez-Gil, R.; Paño-Pardo, J.R.; Mingorance, J. Identification and susceptibility testing of microorganism by direct inoculation from positive blood culture bottles by combining MALDI-TOF and Vitek-2 Compact is rapid and effective. J. Infect. 2012, 65, 513–520, doi:10.1016/j.jinf.2012.08.013.
- Dubois, D.; Segonds, C.; Prere, M.F.; Marty, N.; Oswald, E. Identification of clinical *Streptococcus pneumoniae* isolates among other alpha and nonhemolytic streptococci by use of the Vitek MS matrix-assisted laser desorption ionization-time of flight mass spectrometry system. *J. Clin. Microbiol.* 2013, *51*, 1861–1867, doi:10.1128/JCM.03069-12.
- 38. Monteiro, J.; Inoue, F.M.; Lobo, A.P.; Sugawara, E.K.; Boaretti, F.M.; Tufik, S. Fast and reliable bacterial identification direct from positive blood culture using a new TFA sample preparation protocol and the Vitek[®] MS system. *J. Microbiol. Methods* **2015**, *109*, 157–159, doi:10.1016/j.mimet.2014.12.009.
- 39. Febbraro, F.; Rodio, D.M.; Puggioni, G.; Antonelli, G.; Pietropaolo, V.; Trancassini, M. MALDI-TOF MS Versus VITEK[®]2: Comparison of systems for the identification of microorganisms responsible for bacteremia. *Curr. Microbiol.* **2016**, *73*, 843–850, doi:10.1007/s00284-016-1121-x.
- Srinivasan, R.; Karaoz, U.; Volegova, M.; MacKichan, J.; Kato-Maeda, M.; Miller, S.; Nadarajan, R.; Brodie, E.L.; Lynch, S.V. Use of 16S rRNA Gene for identification of a broad range of clinically relevant bacterial pathogens. *PLoS ONE* 2015, *10*, e0117617, doi:10.1371/journal.pone.0117617.
- 41. Janda, M.J.; Sharon, L.A. 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: Pluses, Perils, and Pitfalls. *J. Clin. Microbiol.* **2007**, *45*, 2761–2764, doi:10.1128/JCM.01228-07.
- 42. Summanen, P.H.; Rowlinson, M.; Wooton, J.; Finegold, S.M. Evaluation of genotypic and phenotypic methods for differentiation of the members of the Anginosus group streptococci. *Eur. J. Clin. Microbiol. Infect. Dis.* **2009**, *28*, 1123–1128, doi:10.1007/s10096-009-0758-9.
- Stackebrandt, E.; Frederiksen, W.; Garrity, G.M.; Grimont, P.; Kämpfer, P.A.D.; Maiden, M.C.J.; Nesme, X.; Rosselló-Mora, R.; Swings, J.; Trüper, H.G.; et al. Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *Int. J. Syst. Evol. Microbiol.* 2002, *52*, 1043–1047, doi:10.1099/00207713-52-3-1043.
- 44. Lerat, E.; Daubin, V.; Ochman, H.; Moran, N.A. Evolutionary origins of genomic repertoires in bacteria. *PLoS Biol.* **2005**, *3*, e130, doi:10.1371/journal.pbio.0030130.
- 45. Leray, M.; Knowlton, N.; Ho, S.L.; Nguyen, B.N.; Machida, R.J. GenBank is a reliable resource for 21st century biodiversity research. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 22651–22656, doi:10.1073/pnas.1911714116.
- 46. Zbinden, A.; Mueller, N.J.; Tarr, P.E.; Spröer, C.; Keller, P.M.; Bloemberg, G.V. *Streptococcus tigurinus* sp. nov., isolated from blood of patients with endocarditis, meningitis and spondylodiscitis. *Int. J. Syst. Evol. Microbiol.* **2012**, *62*, 2941–2945, doi:10.1099/ijs.0.038299-0.
- Poyart, C.; Quesne, G.; Trieu-Cuot, P. Taxonomic dissection of the *Streptococcus bovis* group by analysis of manganese-dependent superoxide dismutase gene (*sodA*) sequences: Reclassification of '*Streptococcus infantarius* subsp. *coli*' as *Streptococcus lutetiensis* sp. nov. and of *Streptococcus bovis* biotype 11.2 as *Streptococcus pasteurianus* sp. nov. *Int. J. Syst. Evol. Microbiol.* 2002, *52*, 1247–1255, doi:10.1099/00207713-52-4-1247.
- 48. Jackson, C.R.; Fedorka-Cray, P.J.; Barrett, J.B. Use of a genus- and species-specific multiplex PCR for identification of enterococci. *J. Clin. Microbiol.* **2004**, *42*, 3558–3565, doi:10.1128/JCM.42.8.3558-3565.2004.
- 49. Castillo-Rojas, G.; Mazari-Hiríart, M.; de León, S.P.; Amieva-Fernández, R.I.; Agis-Juárez, R.A.; Huebner, J.; López-Vidal, Y. Comparison of *Enterococcus faecium* and *Enterococcus faecalis* strains isolated from water

and clinical samples: Antimicrobial susceptibility and genetic relationships. *PLoS ONE* 2013, *8*, e59491, doi:10.1371/journal.pone.0059491.

- 50. Clarridge, J.E.; Osting, C.; Jalali, M.; Osborne, J.; Waddington, M. Genotypic and phenotypic characterization of *"Streptococcus milleri"* group isolates from a Veterans Administration hospital population. *J. Clin. Microbiol.* **1999**, *37*, 3681–3687.
- Arinto-Garcia, R.; Pinho, M.D.; Carriço, J.A.; Melo-Cristino, J.; Ramirez, M. Comparing Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry and phenotypic and molecular methods for identification of species within the *Streptococcus anginosus* Group. *J. Clin. Microbiol.* 2015, *53*, 3580–3588, doi:10.1128/JCM.01892-15.
- 52. Limia, A.; Alarcón, T.; Jiménez, M.L.; López-Brea, M. Comparison of three methods for identification of *Streptococcus milleri* group isolates to species level. *Eur. J. Clin. Microbiol. Infect. Dis.* 2000, 19, 128–131, doi:10.1007/s100960050444.
- 53. Collins, M.D.; Aguirre, M.; Facklam, R.R.; Shallcross, J.; Williams, A.M. *Globicatella sanguis* gen.nov., sp.nov., a new gram-positive catalase-negative bacterium from human sources. *J. Appl. Bacteriol.* **1992**, *73*, 433–437, doi:10.1111/j.1365-2672.1992.tb05000.x.
- 54. Shewmaker, P.L.; Steigerwalt, A.G.; Shealey, L.; Weyant, R.; Facklam, R. DNA relatedness, phenotypic characteristics, and antimicrobial susceptibilities of *Globicatella sanguinis* strains. *J. Clin. Microbiol.* **2001**, *39*, 4052–4057, doi:10.1128/JCM.39.11.4052-4057.2001.
- 55. Miller, A.O.; Buckwalter, S.P.; Henry, M.W.; Wu, F.; Maloney, K.F.; Abraham, B.K.; Hartman, B.J.; Brause, B.D.; Whittier, S.; Walsh, T.J.; et al. *Globicatella sanguinis* Osteomyelitis and Bacteremia: Review of an emerging human pathogen with an expanding spectrum of disease. *Open Forum Infect. Dis.* **2017**, *4*, ofw277, doi:10.1093/ofid/ofw277.
- Kawamura, Y.; Hou, X.G.; Sultana, F.; Liu, S.; Yamamoto, H.; Ezaki, T. Transfer of *Streptococcus adjacens* and *Streptococcus defectivus* to *Abiotrophia* gen. nov. as *Abiotrophia adiacens* comb. nov. and *Abiotrophia defectiva* comb. nov., respectively. *Int. J. Syst. Bacteriol.* 1995, 45, 798–803, doi:10.1099/00207713-45-4-798.
- Collins, M.D.; Lawson, P.A. The genus *Abiotrophia* (Kawamura et al.) is not monophyletic: Proposal of *Granulicatella* gen. nov., *Granulicatella adiacens* comb. nov., *Granulicatella elegans* comb. nov. and *Granulicatella balaenopterae* comb. nov. *Int. J. Syst. Evol. Microbiol.* 2000, *50*, 365–369, doi:10.1099/00207713-50-1-365.
- 58. Patel, T.; Molloy, A.; Smith, R.; Balakrishnan, I. Successful treatment of *Leuconostoc* bacteremia in a neutropenic patient with tigecycline. *Infect. Dis. Rep.* **2012**, *4*, e31, doi:10.4081/idr.2012.e31.
- Albanese, A.; Spanu, T.; Sali, M.; Novegno, F.; D'Inzeo, T.; Santangelo, R. Molecular identification of *Leuconostoc mesenteroides* as a cause of brain abscess in an immunocompromised patient. *J. Clin. Microbiol.* 2006, 44, 3044–3045, doi:10.1128/JCM.00448-06.
- 60. Ke, D.; Picard, F.J.; Martineau, F.; Ménard, C.; Roy, P.H.; Ouellette, M.; Bergeron, M.G. Development of a PCR assay for rapid detection of enterococci. *J. Clin. Microbiol.* **1999**, *37*, 3497–3503.
- Arbique, J.C.; Poyart, C.; Trieu-Cuot, G.; Quesne, M.; Carvalho, G.S.; Steigerwalt, A.G.; Morey, R.E.; Jackson, D.; Davidson, R.J.; Facklam, R.R. Accuracy of phenotypic and genotypic testing for identification of *Streptococcus pneumoniae* and description of *Streptococcus pseudopneumoniae* sp. nov. *J. Clin. Microbiol.* 2004, 42, 4686–4696, doi:10.1128/JCM.42.10.4686-4696.2004.
- 62. Harf-Monteil, C.; Granello, C.; Le Brun, C.; Monteil, H.; Riegel, P. Incidence and pathogenic effect of *Streptococcus pseudopneumoniae*. J. Clin. Microbiol. **2006**, 44, 2240–2241, doi:10.1128/JCM.02643-05.
- 63. Nielsen, X.C.; Justesen, U.S.; Dargis, R.; Kemp, M.; Christensen, J.J. Identification of clinically relevant nonhemolytic Streptococci on the basis of sequence analysis of 16S-23S intergenic spacer region and partial *gdh* gene. *J. Clin. Microbiol.* **2009**, *47*, 932–939, doi:10.1128/JCM.01449-08.
- 64. Garnier, F.; Gerbaud, G.; Courvalin, P.; Galimand, M. Identification of clinically relevant viridans group streptococci to the species level by PCR. *J. Clin. Microbiol.* **1997**, *35*, 2337–2341, doi:10.1128/JCM.35.9.2337-2341.1997.
- 65. Harju, I.; Lange, C.; Kostrzewa, M.; Maier, T.; Rantakokko-Jalava, K.; Haanperä, M. Improved differentiation of *Streptococcus pneumoniae* and other *S. mitis* group streptococci by MALDI Biotyper using an improved MALDI Biotyper database content and a novel result interpretation algorithm. *J. Clin. Microbiol.* 2017, 55, 914–922, doi:10.1128/JCM.01990-16.

- 66. Scott, J.E.; Li, K.; Filkins, L.M.; Zhu, B.; Kuchma, S.L.; Schwartzman, J.D.; O'Toole, G.A. *Pseudomonas aeruginosa* can inhibit growth of streptococcal species via siderophore production. *J. Bacteriol.* **2019**, 201, e00014-19, doi:10.1128/JB.00014-19.
- Besser, J.; Carleton, H.A.; Gerner-Smidt, P.; Lindsey, R.L.; Trees, E. Next-generation sequencing technologies and their application to the study and control of bacterial infections. *Clin. Microbiol. Infect.* 2018, 24, 335–341, doi:10.1016/j.cmi.2017.10.013.
- 68. Woo, P.C.; Leung, A.S.; Leung, K.W.; Yuen, K.Y. Identification of slide coagulase positive, tube coagulase negative *Staphylococcus aureus* by 16S ribosomal RNA gene sequencing. *Mol. Pathol.* **2001**, *54*, 244–247, doi:10.1136/mp.54.4.244.
- Sabat, A.J.; Van Zanten, E.; Akkerboom, V.; Wisselink, G.; van Slochteren, K.; de Boer, R.F.; Hendrix, R.; Friedrich, A.W.; Rossen, J.W.A.; Kooistra-Smid, A.M.D. Targeted next-generation sequencing of the 16S-23S rRNA region for culture-independent bacterial identification-increased discrimination of closely related species. *Sci. Rep.* 2017, 7, 1–12, doi:10.1038/s41598-017-03458-6.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).