

Supporting Information

Metabolic Engineering of *Escherichia coli* for Hyperoside Biosynthesis

Guosi Li ¹, Fucheng Zhu ¹, Peipei Wei ¹, Hailong Xue ², Naidong Chen ¹, Baowei Lu ¹, Hui Deng ¹, Cunwu Chen ^{1,*}, Xinjian Yin ^{3,*}

¹ Anhui Engineering Laboratory for Conservation and Sustainable Utilization of Traditional Chinese Medicine Resources, Department of Biological and Pharmaceutical Engineering, West Anhui University, Lu'an, 237012, Anhui, China; 02000159@wxc.edu.cn (G.L.); fucheng323@163.com (F.Z.); 1209091181@qq.com (P.W.); 2004cnd@163.com; lu_baowei123@126.com; dhup@qq.com

² Key Laboratory of Biomass Chemical Engineering of Ministry of Education, College of Chemical and Biological Engineering, Zhejiang University, Hangzhou, 310027, China; 11428043@zju.edu.cn

³ School of Marine Science, Sun Yat-sen University, Zhuhai, 519080, China

* Correspondence: chengcw@wxc.edu.cn (C.C.); yinxj5@mail.sysu.edu.cn (X.Y.)

List of Entries

Figure S1 Multiple sequence alignment of PhUGT, MdP2'GT, and GmSGT2.....	3
Figure S2 ^{13}C NMR spectra of purified hyperoside.	3
Figure S3 ^1H NMR spectra of purified hyperoside.	4




Figure S3. ^1H NMR spectra of purified hyperoside.