



## Article

# Phylogenomic Analysis of *Salmonella enterica* subsp. *enterica* Serovar Bovismorbificans from Clinical and Food Samples Using Whole Genome Wide Core Genes and *kmer* Binning Methods to Identify Two Distinct Polyphyletic Genome Pathotypes

Gopal R. Gopinath <sup>1,\*</sup>, Hyein Jang <sup>1,†</sup>, Junia Jean-Gilles Beaubrun <sup>1,2,†</sup>, Jayanthi Gangiredla <sup>1</sup>, Mark K. Mammel <sup>1</sup>, Andrea Müller <sup>3</sup>, Sandeep Tamber <sup>4,‡</sup>, Isha R. Patel <sup>1</sup>, Laura Ewing <sup>1</sup>, Leah M. Weinstein <sup>1</sup>, Caroline Z. Wang <sup>1</sup>, Samantha Finkelstein <sup>1</sup>, Flavia Negrete <sup>1</sup>, Tim Muruvanda <sup>5</sup>, Marc Allard <sup>5</sup>, Donald C. Sockett <sup>6</sup>, Franco Pagotto <sup>4</sup>, Ben D. Tall <sup>1</sup> and Roger Stephan <sup>3,‡</sup>

**Citation:** Gopinath, G.R.; Jang, H.; Beaubrun, J.J.-G.; Gangiredla, J.; Mammel, M.K.; Müller, A.; Tamber, S.; Patel, I.R.; Ewing, L.; Weinstein, L.M.; et al. Phylogenomic Analysis of *Salmonella enterica* subsp. *enterica* Serovar Bovismorbificans from Clinical and Food Samples Using Whole Genome Wide Core Genes and *kmer* Binning Methods to Identify Two Distinct Polyphyletic Genome Pathotypes. *Microorganisms* **2022**, *10*, 1199. <https://doi.org/10.3390/microorganisms10061199>

Academic Editor: Lisa Gorski

Received: 12 May 2022

Accepted: 1 June 2022

Published: 11 June 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

<sup>1</sup> Center of Food Safety and Applied Nutrition, U. S. Food and Drug Administration, Laurel, MD 20708, USA; hyein.jang@fda.hhs.gov (H.J.); junia.jean-gillesbeaubrun.civ@mail.mil (J.J.-G.B.); jayanthi.gangiredla@fda.hhs.gov (J.G.); mark.mammel@fda.hhs.gov (M.K.M.); isha.patel@fda.hhs.gov (I.R.P.); laura.ewing-peoples@fda.hhs.gov (L.E.); lmweinstein1@gmail.com (L.M.W.); caroline.z.wang@duke.edu (C.Z.W.); sfinkel6@terpmail.umd.edu (S.F.); flavianegrete@yahoo.com (F.N.); shakee2326@aol.com (B.D.T.)

<sup>2</sup> Biological Analysis Division, Public Health Command Europe Laboratory Sciences, Room 102, Bldg 3810, Kirchberg Kaserne, RP 66849 Landstuhl, Germany

<sup>3</sup> Institute for Food Safety and Hygiene, University of Zurich, CH-8057 Zurich, Switzerland; andreamonika.mueller@uzh.ch (A.M.); stephanr@fsafety.uzh.ch (R.S.)

<sup>4</sup> Food Directorate, Bureau of Microbial Hazards/Health Canada, Ottawa, ON K1A 0K9, Canada; sandeep.tamber@canada.ca (S.T.); franco.pagotto@canada.ca (F.P.)

<sup>5</sup> Center of Food Safety and Applied Nutrition, U. S. Food and Drug Administration, College Park, MD 20740, USA; tim.muruvanda@fda.hhs.gov (T.M.); marc.allard@fda.hhs.gov (M.A.)

<sup>6</sup> Wisconsin Veterinary Diagnostic Laboratory, University of Wisconsin-Madison, Madison, WI 53706, USA; donald.sockett@wvdl.wisc.edu

\* Correspondence: gopal.gopinathrao@fda.hhs.gov; Tel.: +1-240-402-3612

† These authors contributed equally to this work.

‡ Contacts for strains used in this study: Roger Stephan (Swiss collection; stephanr@fsafety.uzh.ch), Sandeep Tamber (Canadian collection; sandeep.tamber@canada.ca) and Gopal Gopinath for FDA collection.

**Abstract:** *Salmonella enterica* subsp. *enterica* serovar Bovismorbificans has caused multiple outbreaks involving the consumption of produce, hummus, and processed meat products worldwide. To elucidate the intra-serovar genomic structure of *S. Bovismorbificans*, a core-genome analysis with 2690 loci (based on 150 complete genomes representing *Salmonella enterica* serovars developed as part of this study) and a *k-mer*-binning based strategy were carried out on 95 whole genome sequencing (WGS) assemblies from Swiss, Canadian, and USA collections of *S. Bovismorbificans* strains from foodborne infections. Data mining of a digital DNA tiling array of legacy SARA and SARB strains was conducted to identify near-neighbors of *S. Bovismorbificans*. The core genome analysis and the *k-mer*-binning methods identified two polyphyletic clusters, each with emerging evolutionary properties. Four STs (2640, 142, 1499, and 377), which constituted the majority of the publicly available WGS datasets from >260 strains analyzed by *k-mer*-binning based strategy, contained a conserved core genome backbone with a different evolutionary lineage as compared to strains comprising the other cluster (ST150). In addition, the assortment of genotypic features contributing to pathogenesis and persistence, such as antimicrobial resistance, prophage, plasmid, and virulence factor genes, were assessed to understand the emerging characteristics of this serovar that are relevant clinically and for food safety concerns. The phylogenomic profiling of polyphyletic *S. Bovismorbificans* in this study corresponds to intra-serovar variations observed in *S. Napoli* and *S. Newport* serovars using similar high-resolution genomic profiling approaches and contributes to the understanding

of the evolution and sequence divergence of foodborne *Salmonellae*. These intra-serovar differences may have to be thoroughly understood for the accurate classification of foodborne *Salmonella* strains needed for the uniform development of future food safety mitigation strategies.

**Keywords:** *Salmonella* Bovismorbificans; phylogenomics; plasmids; phages; virulence factors

## 1. Introduction

*Salmonella* species are one of the leading causes of foodborne outbreaks and systemic infections worldwide [1] and in the USA [2], leading to thousands of deaths every year. Human infections with *Salmonella enterica* subspecies *enterica* serovar Bovismorbificans are rare compared with other *Salmonella* serovars such as Typhimurium or Enteritidis. However, *S. Bovismorbificans* have increasingly been reported as an emerging human pathogen causing foodborne illnesses in Asia [3] and in the western hemisphere. The molecular epidemiology of a sprout-borne outbreak of *S. Bovismorbificans* [4,5] in Finland had been reported earlier. Most foodborne outbreaks associated with *S. Bovismorbificans* reported in Europe, the USA, and Canada has been traced to pork products, lettuce, hummus, and sprouts [6–8]. The first genomes of *S. Bovismorbificans* were generated from a Malawian bacteremia case and UK veterinary samples belonging to Sequence Type (ST)142 (ST142) including that of a virulence plasmid pVirBov from clinical strain 3114 [9]. More ST142 virulent strains were isolated from outbreaks associated with uncooked ham products in the Netherlands [10]. Interestingly, these strains possessed a 5.1 kb col156 plasmid. The draft genomes from food and clinical strains from a 2011 outbreak of *S. Bovismorbificans* in Washington, DC, USA isolated from contaminated hummus samples were made available [11–13]. These strains were typed as ST377 and were phylogenetically distinct from other USA food and environmental strains reported independently [14] belonging to ST150. These reports implicated ST142 and ST377 as the predominant sequence types of *S. Bovismorbificans* contaminating the food supply and causing foodborne illnesses in the USA and Europe. In 2018, the Wisconsin Veterinary Diagnostic Laboratory (WVDL) isolated *S. Bovismorbificans* strains from environmental samples that were collected from a Minnesota dairy calf production facility. During this time, the dairy industry experienced an increased mortality event in 25% of dairy calves at 1–3 weeks of age caused by *S. Bovismorbificans* (personal communication, DCS). Environmental sampling also showed the presence of *S. Bovismorbificans*. This finding prompted the hypothesis that *S. Bovismorbificans* may be persistent in this dairy production environment. A study conducted [15] on the impact of *S. Bovismorbificans* along the food supply chain in Hungary investigated contaminated food production environments, animals, foods of animal origin, and humans and provided evidence that *S. Bovismorbificans* is less invasive to host animals than *S. Enteritidis* but may colonize and persist in several animal species leading to contamination of meat.

There is considerable interest in identifying underlying bacterial genomic attributes related to the increased transmission of minor serovars such as *S. Bovismorbificans* in humans and within the animal food supply chains. The diversity of virulence mechanisms, differences in the genomic features, and emergent sources/niches are increasingly recognized as major contributors to the success of *Salmonella* as a major foodborne pathogenic group [16]. Genome-wide variations of different groups within the serovars of *S. enterica* have been extensively studied using whole genome sequencing (WGS) datasets. A high number of polyphyletic lineages in many serovars have been predicted using MLST and genome-wide SNP profiling methods [17–22]. For example, phylogenetic analysis of 156 WGS datasets from 78 serovars using about 120,000 whole genome SNPs present in at least 95% of the strains identified the presence of polyphyletic lineages in a handful of serovars [19]. Similar WGS-based methods were applied to analyze *S. Newport* [20] and *S. Napoli* [21] strains. In the present study, 95 strains of *S. Bovismorbificans* were

collected from the USA, Switzerland, and Canada from various types of sources and years. Phylogenomic sequence analysis of the strains was conducted using a core genome schema developed using 150 complete genomes. This resulted in a scalable ad hoc bioinformatic workflow to identify the core genome among *S. Bovismorbificans*, which resulted in the identification of two distinct polyphyletic groups within the serovar with significant divergence in their core gene loci. In addition, a *k-mer*-binning method and data mining of digital DNA tiling array profiles were applied in parallel to illustrate this evolutionary relationship among the sub-groups of this important emerging *Salmonella* pathogen. We suggest that this approach can be applied to predict and annotate the emerging virulence properties of under-surveyed minor serovars of *Salmonella* with the potential to cause sporadic foodborne outbreaks.

## 2. Materials and Methods

### 2.1. Bacterial Strains

A total of 95 strains of *S. Bovismorbificans* isolated from Switzerland, the USA, and Canada were analyzed in this study. The collection represents clinical, food, feed, animal, and environmental strains, and metadata for these strains including country, year of isolation, and genomic attributes. *S. Bovismorbificans* strains collected from Switzerland, the USA, and Canada were provided by University of Zurich, University of Wisconsin-Madison, and Health Canada, respectively. In addition to the newly sequenced 81 strains, genomes from 14 *S. Bovismorbificans* genomes, which were associated with several human USA outbreaks, were included [7,11,14]. For comparative analysis, a scaffolded genome from the clinical strain (3114) from Malawi [9], an environmental strain (CIES13) from Mexico, and 25 more *S. Bovismorbificans* genomes were downloaded from NCBI. All strains were stored at  $-80^{\circ}\text{C}$  in Trypticase soy broth (TSB; BBL, Cockeysville, MD) supplemented with 50% glycerol. All strains were serotyped and identified according to the White–Kauffmann–Le Minor scheme by slide agglutination [23], USA strains were also serotyped by PCR analysis, as described earlier [24]. Genome assemblies from the strains were used to confirm each strain's serotype using the SeqSero2 v1.0.2 [25] application on CFSAN's GalaxyTrakr [26] at <https://galaxytrakr.org/> (accessed on 5 January 2022).

### 2.2. Genomic DNA Preparation

Frozen stocks of each strain were streaked onto Xylose-lysine-tergitol 4 (XLT4; BBL, 63Cockeysville, MD, USA) agar plates and cultured at  $37^{\circ}\text{C}$  for overnight. A single typical colony (black or black-centered) of each strain on XLT4 was inoculated into 5 mL of TSB supplemented with 1% NaCl (TSBS), and then incubated at  $37^{\circ}\text{C}$  for 20 h with shaking at 150 rpm. Genomic DNA was extracted from the overnight cultures using a Qiagen QIACube instrument and its automated technology (QIAGEN Sciences, Germantown, MD, USA) according to the manufacturer's instructions. Typical yields of the purified genomic DNA are 5–50  $\mu\text{g}$  from a final elution volume of 200  $\mu\text{L}$ , and duplicated DNA samples were prepared for WGS and microarray experiments. Each strain's DNA was quantified using a Qubit dsDNA BR assay kit (Invitrogen, Thermo Fisher Scientific, Wilmington, DE, USA) and Qubit 2.0 fluorometer (Life Technologies, Grand Island, NY, USA).

### 2.3. Whole Genome Sequencing (WGS), Assembly, and Annotation

Eighty-one strains were obtained from Switzerland, the USA, and Canada, and representing clinical (69), food (9), feed (1), animal (1), and environment (1) strains (isolated during 1984–1989 and 2011–2018) were sequenced in this study. Each DNA sample prepared as described above was diluted in nuclease-free deionized water (molecular biology grade, Thermo Fisher Scientific, Waltham, MA, USA) to achieve a final concentration of 0.2 ng/ $\mu\text{L}$ . WGS libraries of these strains (50 $\times$  coverage) were constructed using the Nextera XT DNA sample preparation kit (Illumina, San Diego, CA, USA). Genome sequencing was performed on a Miseq platform using either 500 or 600 cycles of paired-end reads

(Illumina). FastQ datasets (raw reads) were trimmed for removal of adaptor sequences and for quality control purposes, and de novo assembled using CLC Genomics Workbench version 9.0 (CLC Bio, Aarhus, Denmark). The genomes were independently annotated using the Rapid Annotation Subsystems Technology (RAST) annotation server [27] for quality control and accuracy. The genome sizes, the number of coding sequences (CDS), ST assignments, NCBI BioSample ID, and accession numbers of these assemblies are shown in Table 1. PacBio RSII platform was used following the manufacturer's protocols to generate a complete genome of a 93kb plasmid from Sal610. All the assemblies (Table 1) along with the prokaryotic genome annotation pipeline (PGAP) annotations [28] were deposited into NCBI's GenBank and used in the subsequent analyses as needed. The datasets were released to the public through submission to NCBI under the FDA-CFSAN's GenomeTrakr *Salmonella enterica* BioProject PRJNA378379, which is part of the CFSAN's foodborne pathogen research umbrella project PRJNA186875 at NCBI [29].

**Table 1.** Strain information (Strain name, Source, Country, Year of isolation, and Reference), genomic characterization (Genome size, No. of CDSs, and Sequence Type), NCBI Biosample, and GenBank accession numbers of 95 *S. Bovismorbificans* isolates used in this study.

| Strain    | Source          | Country     | Year    | Genome Size (kb) | No. of CDSs | ST <sup>a</sup> | NCBI Biosample | NCBI Accession No. | Reference  |
|-----------|-----------------|-------------|---------|------------------|-------------|-----------------|----------------|--------------------|------------|
| M18_12182 | Environment     | USA         | 2018    | 4679             | 4399        | 150             | SAMN12657258   | WSCV000000000      | This study |
| Bovis_38  | Beef            | Canada      | 1989    | 4579             | 4321        | 142             | SAMN12657217   | WSCW000000000      | This study |
| Bovis_120 | Seafood, squid  | Canada      | missing | 4706             | 4474        | 1499            | SAMN12657218   | WSCX000000000      | This study |
| Bovis_187 | Seafood, shrimp | Canada      | 1989    | 4739             | 4498        | 1499            | SAMN12657204   | WSCY000000000      | This study |
| Bovis_277 | Clinical        | Canada      | 1984    | 4832             | 4638        | 1499            | SAMN12657210   | WSCZ000000000      | This study |
| Bovis_278 | Clinical        | Canada      | 1984    | 4738             | 4498        | 142             | SAMN12657232   | WSDA000000000      | This study |
| N14_0147  | Clinical        | Switzerland | 2014    | 4690             | 4471        | 142             | SAMN12657231   | WSDB000000000      | This study |
| N14_0646  | Clinical        | Switzerland | 2014    | 4702             | 4449        | 1499            | SAMN12657228   | WSDC000000000      | This study |
| N14_1189  | Clinical        | Switzerland | 2014    | 4790             | 4575        | 142             | SAMN12657257   | WSDD000000000      | This study |
| N14_1190  | Clinical        | Switzerland | 2014    | 4784             | 4566        | 142             | SAMN12657229   | WSD E000000000     | This study |
| N14_1194  | Clinical        | Switzerland | 2014    | 4789             | 4580        | 142             | SAMN12657219   | WSD F000000000     | This study |
| N14_1212  | Clinical        | Switzerland | 2014    | 4790             | 4577        | 142             | SAMN12657254   | WSD G000000000     | This study |
| N14_1217  | Clinical        | Switzerland | 2014    | 4854             | 4688        | 142             | SAMN12657212   | WSD H000000000     | This study |
| N14_1222  | Clinical        | Switzerland | 2014    | 4791             | 4577        | 142             | SAMN12657214   | WSD I000000000     | This study |
| N14_1232  | Clinical        | Switzerland | 2014    | 4778             | 4560        | 142             | SAMN12657205   | WSD J000000000     | This study |
| N14_1233  | Clinical        | Switzerland | 2014    | 4790             | 4568        | 142             | SAMN12657209   | WSD K000000000     | This study |
| N14_1237  | Clinical        | Switzerland | 2014    | 4781             | 4565        | 142             | SAMN12657230   | WSD L000000000     | This study |
| N14_1238  | Clinical        | Switzerland | 2014    | 4781             | 4563        | 142             | SAMN12657235   | WSD M000000000     | This study |
| N14_1239  | Clinical        | Switzerland | 2014    | 4778             | 4562        | 142             | SAMN12657165   | WSD N000000000     | This study |
| N14_1252  | Clinical        | Switzerland | 2014    | 4781             | 4562        | 142             | SAMN12657181   | WSD O000000000     | This study |
| N14_1255  | Clinical        | Switzerland | 2014    | 4786             | 4576        | 142             | SAMN12657154   | WSD P000000000     | This study |
| N14_1266  | Clinical        | Switzerland | 2014    | 4779             | 4561        | 142             | SAMN12657194   | WSD Q000000000     | This study |
| N14_1267  | Clinical        | Switzerland | 2014    | 4775             | 4602        | 142             | SAMN12657197   | WSD R000000000     | This study |
| N14_1268  | Clinical        | Switzerland | 2014    | 4783             | 4567        | 142             | SAMN12657192   | WSD S000000000     | This study |
| N14_1285  | Clinical        | Switzerland | 2014    | 4780             | 4561        | 142             | SAMN12657190   | WSD T000000000     | This study |
| N14_1527  | Clinical        | Switzerland | 2014    | 4813             | 4590        | 1499            | SAMN12657150   | WSD U000000000     | This study |
| N14_1673  | Clinical        | Switzerland | 2014    | 4819             | 4609        | 142             | SAMN12657188   | WSD V000000000     | This study |
| N14_1674  | Clinical, blood | Switzerland | 2014    | 4783             | 4571        | 142             | SAMN12657156   | WSD W000000000     | This study |
| N14_1962  | Clinical        | Switzerland | 2014    | 4715             | 4468        | 1499            | SAMN12657164   | WSD X000000000     | This study |
| N14_2376  | Food, onion     | Switzerland | 2014    | 4723             | 4486        | 377             | SAMN12657147   | WSD Y000000000     | This study |
| N15_0039  | Food, grain     | Switzerland | 2015    | 4726             | 4493        | 377             | SAMN12657185   | WSD Z000000000     | This study |
| N15_0111  | Clinical        | Switzerland | 2015    | 5024             | 4860        | 1499            | SAMN12657157   | WSE A000000000     | This study |
| N15_0387  | Clinical        | Switzerland | 2015    | 4643             | 4404        | 142             | SAMN12657198   | WSE B000000000     | This study |
| N15_0584  | Clinical        | Switzerland | 2015    | 4670             | 4468        | 142             | SAMN12657182   | WSE C000000000     | This study |
| N15_1010  | Food            | Switzerland | 2015    | 4725             | 4483        | 377             | SAMN12657196   | WSE D000000000     | This study |
| N15_1120  | Food            | Switzerland | 2015    | 4724             | 4486        | 377             | SAMN12657193   | WSE E000000000     | This study |
| N15_1199  | Food            | Switzerland | 2015    | 4720             | 4485        | 377             | SAMN12657191   | WSE F000000000     | This study |
| N15_1496  | Clinical        | Switzerland | 2015    | 4738             | 4530        | 142             | SAMN12657189   | WSE G000000000     | This study |
| N15_2048  | Clinical        | Switzerland | 2015    | 4847             | 4671        | 1499            | SAMN12657202   | WSE H000000000     | This study |
| N15_2285  | Clinical        | Switzerland | 2015    | 4806             | 4598        | 377             | SAMN12657199   | WSE I000000000     | This study |
| N15_2358  | Clinical        | Switzerland | 2015    | 4893             | 4724        | 377             | SAMN12657233   | WSE J000000000     | This study |
| N16_0208  | Clinical        | Switzerland | 2016    | 4798             | 4571        | 1499            | SAMN12657155   | WSE B000000000     | This study |

|                        |                 |             |         |                 |      |                      |              |                        |            |
|------------------------|-----------------|-------------|---------|-----------------|------|----------------------|--------------|------------------------|------------|
| N16_0447               | Clinical, urine | Switzerland | 2016    | 4636            | 4399 | 142                  | SAMN12657160 | WSBI00000000           | This study |
| N16_0656               | Clinical        | Switzerland | 2016    | 4838            | 4628 | 2640                 | SAMN12657256 | WSBJ00000000           | This study |
| N16_1158               | Clinical        | Switzerland | 2016    | 4781            | 4566 | 142                  | SAMN12657253 | WSBK00000000           | This study |
| N16_1250               | Clinical        | Switzerland | 2016    | 4788            | 4571 | 142                  | SAMN12657211 | WSBL00000000           | This study |
| N16_1558               | Clinical        | Switzerland | 2016    | 4571            | 4290 | 150                  | SAMN12657220 | WSBM00000000           | This study |
| N16_1675               | Clinical        | Switzerland | 2016    | 4782            | 4573 | 142                  | SAMN12657207 | WSBN00000000           | This study |
| N16_1715               | Clinical        | Switzerland | 2016    | 4784            | 4570 | 142                  | SAMN12657200 | WSBO00000000           | This study |
| N16_1722               | Clinical        | Switzerland | 2016    | 4785            | 4569 | 142                  | SAMN12657206 | WSBP00000000           | This study |
| N16_1740               | Clinical        | Switzerland | 2016    | 4778            | 4556 | 142                  | SAMN12657234 | WSBQ00000000           | This study |
| N16_1757               | Clinical        | Switzerland | 2016    | 4784            | 4568 | 142                  | SAMN12657203 | WSBR00000000           | This study |
| N16_1786               | Clinical        | Switzerland | 2016    | 4781            | 4567 | 142                  | SAMN12657259 | WSBS00000000           | This study |
| N16_1788               | Clinical        | Switzerland | 2016    | 4784            | 4584 | 142                  | SAMN12657260 | WSBT00000000           | This study |
| N16_1794               | Clinical        | Switzerland | 2016    | 4794            | 4573 | 142                  | SAMN12657221 | WSBU00000000           | This study |
| N16_1829               | Clinical        | Switzerland | 2016    | 4785            | 4571 | 142                  | SAMN12657255 | WSBV00000000           | This study |
| N16_2042               | Clinical        | Switzerland | 2016    | 4778            | 4561 | 142                  | SAMN12657208 | WSBW00000000           | This study |
| N16_2132               | Clinical        | Switzerland | 2016    | 4787            | 4563 | 142                  | SAMN12657216 | WSBX00000000           | This study |
| N16_2207               | Animal, cat     | Switzerland | 2016    | 4778            | 4573 | 142                  | SAMN12657146 | WSBY00000000           | This study |
| N16_2424               | Clinical        | Switzerland | 2016    | 4775            | 4556 | 142                  | SAMN12657163 | WSBZ00000000           | This study |
| N16_2574               | Clinical        | Switzerland | 2016    | 4947            | 4754 | 142                  | SAMN12657148 | WSCA00000000           | This study |
| N16_2598               | Clinical        | Switzerland | 2016    | 4706            | 4481 | 142                  | SAMN12657161 | WSCB00000000           | This study |
| N16_2682               | Clinical        | Switzerland | 2016    | 4777            | 4574 | 142                  | SAMN12657159 | WSCC00000000           | This study |
| N16_2712               | Clinical        | Switzerland | 2016    | 4703            | 4489 | 142                  | SAMN12657184 | WSCD00000000           | This study |
| N16_2718               | Clinical        | Switzerland | 2016    | 4723            | 4482 | 377                  | SAMN12657144 | WSEC00000000           | This study |
| N16_2802               | Clinical        | Switzerland | 2016    | 4724            | 4508 | 142                  | SAMN12657195 | WSEF00000000           | This study |
| N16_2849               | Clinical        | Switzerland | 2016    | 4725            | 4489 | 377                  | SAMN12657142 | WSEG00000000           | This study |
| N16_2955               | Clinical        | Switzerland | 2016    | 4811            | 4599 | 1499                 | SAMN12657187 | WSEH00000000           | This study |
| N17_0002               | Clinical        | Switzerland | 2017    | 4706            | 4492 | 142                  | SAMN12657149 | WSEI00000000           | This study |
| N17_0502               | Clinical        | Switzerland | 2017    | 4873            | 4696 | 1499                 | SAMN12657152 | WSEJ00000000           | This study |
| N17_0607               | Clinical        | Switzerland | 2017    | 4640            | 4401 | 142                  | SAMN12657145 | WSEK00000000           | This study |
| N17_0620               | Food            | Switzerland | 2017    | 4731            | 4493 | 377                  | SAMN12657151 | WSEL00000000           | This study |
| N17_0857               | Feed            | Switzerland | 2017    | 4724            | 4507 | 377                  | SAMN12657158 | WSEM00000000           | This study |
| N17_1212               | Clinical        | Switzerland | 2017    | 4773            | 4576 | 2640                 | SAMN12657162 | WSEN00000000           | This study |
| N17_1364               | Clinical        | Switzerland | 2017    | 4869            | 4680 | 2640                 | SAMN12657186 | WSEO00000000           | This study |
| N17_1393               | Clinical        | Switzerland | 2017    | 4779            | 4564 | 2640                 | SAMN12657153 | WSEP00000000           | This study |
| N17_2111               | Clinical        | Switzerland | 2017    | 4775            | 4561 | 2640                 | SAMN12657143 | WSEQ00000000           | This study |
| N18_0002               | Clinical        | Switzerland | 2018    | 4805            | 4605 | 142                  | SAMN12657183 | WSEK00000000           | This study |
| N18_0969               | Clinical        | Switzerland | 2018    | 4640            | 4402 | 142                  | SAMN12657213 | WSES00000000           | This study |
| N18_1092               | Clinical        | Switzerland | 2018    | 4728            | 4492 | 377                  | SAMN12657215 | WSET00000000           | This study |
| N18_1144               | Clinical        | Switzerland | 2018    | 4722            | 4544 | 377                  | SAMN12657201 | WSEU00000000           | This study |
| Sal609 <sup>b</sup>    | Clinical        | USA         | 2011    | 4896            | 4925 | 377                  | SAMN02422699 | AZKW00000000           | [11]       |
| Sal610                 | Clinical        | USA         | 2011    | 4857            | 4870 | 377                  | SAMN02422700 | AZKX00000000           | [11]       |
| Sal615                 | Clinical        | USA         | 2011    | 4845            | 4891 | 377                  | SAMN02422701 | AZKY00000000           | [11]       |
| Sal616                 | Food            | USA         | 2011    | 4865            | 4893 | 377                  | SAMN02422702 | AZKZ00000000           | [11]       |
| Sal617                 | Food            | USA         | 2011    | 4872            | 4887 | 377                  | SAMN02422703 | AZLA00000000           | [11]       |
| Sal644                 | Clinical        | USA         | 2001    | 4769            | 4769 | 1499                 | SAMN02422688 | AZLC00000000           | [11]       |
| Sal676                 | Clinical        | USA         | 2012    | 4569            | 4476 | 150                  | SAMN02422698 | AZKV00000000           | [11]       |
| Sal677                 | Clinical        | USA         | 2012    | 4663            | 4614 | 150                  | SAMN02422693 | AZKR00000000           | [14]       |
| Sal678                 | Clinical        | USA         | 2012    | 4579            | 4629 | 150                  | SAMN02422694 | AZKS00000000           | [14]       |
| Sal679                 | Clinical        | USA         | 2012    | 4567            | 4471 | 150                  | SAMN02422695 | AZKT00000000           | [14]       |
| Sal680                 | Clinical        | USA         | 2012    | 4596            | 4506 | 150                  | SAMN02422696 | AZKU00000000           | [14]       |
| Sal681                 | Clinical        | USA         | 2012    | 4575            | 4492 | 150                  | SAMN02422697 | AZLB00000000           | [14]       |
| Sal682                 | Clinical        | USA         | 2012    | 4926            | 4944 | 377                  | SAMN02422690 | AZKQ00000000           | [11]       |
| Sal683                 | Clinical        | USA         | 2012    | 4574            | 4586 | 150                  | SAMN02422689 | AZKP00000000           | [14]       |
| bovis3114 <sup>b</sup> | Clinical        | Malawi      | 1997    | 4680            | 4599 | 142                  | SAMEA3138815 | HF969015               | [9]        |
| bovispt13              | Unknown         | Unknown     | Unknown | NA <sup>c</sup> | NA   | Unknown <sup>d</sup> | SAMN01081634 | SRS347148 <sup>e</sup> | NCBI SRA   |
| boviscies13            | Water           | Mexico      | 2013    | NA              | NA   | 150                  | SAMN02335370 | SRS476367 <sup>f</sup> | NCBI SRA   |
| pSal610                | Clinical        | USA         | 2011    | 93.8            | 111  | 377                  | SAMN02422700 | CP076746               | This study |

<sup>a</sup>Sequence type (ST) was determined by uploading genome assemblies to <https://cge.cbs.dtu.dk/services/MLST/> (accessed on 1 June 2020). <sup>b</sup>Genome size and number of CDSs of strains named with ‘Sal’ and ‘bovis’ were determined by the SEED Viewer of RAST annotation. <sup>c</sup>NA represents ‘not available’. <sup>d</sup>Nearest ST is 142. <sup>e,f</sup>Public sequence reads were downloaded from NCBI SRA (Sequence Read Archive) and locally assembled. Note: External genome sequences downloaded from NCBI were downloaded by entering the assembly accessions at the end of the URL:

<https://www.ncbi.nlm.nih.gov/nuccore?term=> (last accessed on 6 April 2022) QDND01, MXTS01, MZFY01, QAUR01, MXXQ01, MYFY01, WFIN01, MXTX01, MXUD01, JYUT01, MXFZ01, QAUR01, AUQE01, ARYX01, AAADAU000000000.1 (for AAADAU01), NPMA01, VCTV01, AAOBOO000000000.1 (for AAOBOO01), AAIBEG000000000.1 (for AAIBEG01). Four sequence reads datasets were downloaded from NCBI SRA database: ERR1755559, SRR1783167, SRR16148890 and SRR3473097 for generating WGS assemblies for this analysis.

#### 2.4. Identification of Whole-Genome Core Genes and High-Resolution Phylogenomic Analysis

A total of 645 complete genomes (Supplemental File S1) of *S. enterica* representing 150 serovars were downloaded from NCBI's Genome database (20 February 2020) to generate a large local BLAST database. Some *Salmonella* serovars such as *S. Typhimurium* and *S. Enteritidis* were overrepresented in this pool, while most others had few or single completed genomes. A shorter list of representative genomes of the 150 serovars was randomly chosen for subsequent analysis in this bioinformatic workflow. CDS (4606) annotations of *S. enterica* serovar Typhimurium LT2 genome (NC\_003197) were downloaded from GenBank and used as a query in the BLAST analysis at 50, 90, and 95% identity levels. In-house Perl and Python scripts (available upon request) were used to parse the BLAST outputs and to create a SNP-finding workflow. Manual curation of the resulting data matrix with homology search and alignment contributed to the generation of a whole genome core genes (wg-core) set of 2690 loci representing a conserved genomic backbone of representative genomes for 150 complete *Salmonella* serovars.

WGS assemblies from *S. Bovismorbificans* strains from this study and external sources (Table 1) were queried with the wg-core gene set to identify conserved backbone genes. A SNP data matrix consisting of homologs of wg-core genes in the evaluated *S. Bovismorbificans* strains with at least one allele in each of the 2512 out of 2690 loci was created (Supplemental File S2 also available from <https://github.com/gopal-gopinath/S.bovismorbificans-SNP-matrix1> (last accessed 6 May 2022)). The evolutionary distances were computed using the Maximum Composite Likelihood method [30] and the phylogenetic tree was built based on the neighbor-joining method [31] as implemented on the MEGA X phylogenetic suite [32] and UPGMA algorithm implemented in SplitsTree version 5 [33]. The SNP matrices were curated for quality to remove missing genes and partial gene sequences represented by gaps after multiple alignments as part of the routine quality control. The genome-wide *k*-mer-binning analysis was carried out by generating a Jaccard similarity matrix with *k*-mer content using in-house scripts. For each genome, a list of all *k*-mers ( $k = 30$ ) present in the sequences was stored. Each pair of genomes was compared for *k*-mer content to derive the Jaccard similarity score defined as the size of the intersection divided by the size of the union of the *k*-mer sets.

#### 2.5. Data Mining of a DNA Tiling Microarray Database for Genomic Comparisons with Legacy Strain Collections

The FDA *Salmonella* custom high-density Affymetrix DNA microarray platform was used, as previously described [34,35]. An 8 µg aliquot of purified genomic DNA was fragmented by incubation at 37 °C for 10 min in a 50 µL reaction containing 1× One-Phor-All Plus Buffer [Tris, Magnesium and Potassium acetate (Ratios 1:1:5)] and 0.1 units DNase I (GE Healthcare, Pittsburgh, PA, USA). Following fragmentation, the DNA was labeled at the 3'-end using 1 mM biotin-11-ddATP (PerkinElmer NEL508, Waltham, MA, USA), 5X terminal transferase buffer (Promega, Madison, WI, USA), and 60 units of terminal transferase enzyme (Promega), as described earlier. The genomic DNA samples were hybridized following the Affymetrix GeneChip Expression Analysis Technical Manual (Affymetrix, Santa Clara, CA, USA, 2014), washed in the Affymetrix FS-450 fluidics station (Affymetrix, Santa Clara, CA, USA), and scanned using software of the Affymetrix GeneChip Command Console (AGCC) Scanner 3000. Reagents used in hybridization, washing, and staining were prepared according to the Affymetrix GeneChip Expression Analysis Technical Manual [36]. For microarray data analysis, a probe set intensity for

each allele represented on the microarray chip were assessed using the Robust MultiArray Averaging (RMA) function in the Affymetrix package of R-Bioconductor [37].

## 2.6. Characterization of Multi-Locus Sequence Typing (MLST) and Antimicrobial Resistance Gene (AMR) Patterns

Sequence types of all *S. Bovismorbificans* genomes used in this study were determined using the MLST schema [17,38] implemented on the Center for Genomic Epidemiology (CGE) web servers (<http://cge.cbs.dtu.dk/services/MLST>, accessed on 6 August 2021). For serological confirmation and antimicrobial resistance gene analyses, genome assemblies (FASTA files) were uploaded to the CFSAN's GalaxyTrakr SeqSero and AMRFinderPlus tool, respectively (<https://galaxytrakr.org/root/login?redirect=%2F>, accessed on 6 August 2021). CFSAN's Galaxy GenomeTrakr AMRFinderPlus tool scans each genome against the accompanying database, which is designed to find acquired antimicrobial, biocide, heat, acid, and metal resistance genes in bacterial protein or assembled nucleotide sequences, as well as known point mutations for several taxa [39]. The classes of antimicrobials that the CFSAN Galaxy GenomeTrakr AMRFinderPlus tool identifies include aminoglycoside, beta-lactams, bleomycin, colistin, fosfomycin, fusidic acid, glycopeptide, nitroimidazole, oxazolidinone, phenicol, quaternary ammonium, quinolone, rifamycin, streptogramin, sulfonamide, tetracycline, and trimethoprim.

## 2.7. Identification of *Salmonella* Pathogenicity Islands (SPIs), Plasmids, and Prophage Genomic Regions and Virulence Factors

pVIRBov, a virulence plasmid from *S. Bovismorbificans* strains 3114 [9], was downloaded (Acc. No.: HF969016) from NCBI. pVIRBov sequences were used to identify homologous plasmid sequences in the WGS datasets generated from this study (Table 1). For this step, a local database of *S. Bovismorbificans* strains was first created for BLAST analysis as needed. The stringent parameters used for the BLAST analysis (percent identity: 50%, best hit score: 0.05, best hit overhang: 0.25, e-value:  $1 \times 10^{-10}$ ) were chosen to minimize both alignment errors and random mutations for screening the WGS database. In-house Python and Perl scripts were then used to parse the BLAST output and identify homologous sequences of 111 total pSal610/pVIRBov CDS in 95 genomes of *S. Bovismorbificans* strains. BRIG 0.95 software [40] was used for the visualization of plasmid sequence comparisons. Assembled genomes of 34 *S. Bovismorbificans* strains representative of the various STs were submitted to the CGE SPIFinder (<https://cge.cbs.dtu.dk/services/SPIFinder>, accessed on 6 August 2021) to detect *Salmonella* pathogenicity islands (SPIs). The default parameters with threshold for ID of 95% and minimum length at 60% for SPIFinder 1.0 tool settings were used to minimize noise and to eliminate any gene fragments, which enables the detection of genes in the start or end of contigs. The CGE PlasmidFinder tool [41] available at <https://cge.cbs.dtu.dk/services/PlasmidFinder> (accessed on 6 August 2021) was employed to detect replicon-associated plasmid sequences using the *Enterobacteriaceae* as the default settings: 95% minimal identity and 60% minimal coverage. Prophage genomic regions were identified using the PHASTER (Phage Search Tool Enhanced Release) web server (<https://phaster.ca>, accessed on 6 August 2021). Upon uploading to the PHASTER pipeline, putative phage and prophage regions (i.e., incomplete, questionable, and intact prophage regions) were identified through examination of prophage genes and comparison of predicted proteins against PHASTER's complete prophage databases [42]. The SEED server was routinely used for sequence comparisons and datamining functional annotations [43]. Prophage sequences from the *S. Bovismorbificans* WGS datasets were identified using the *Salmonella* phage RE-2010 (GenBank Accn#: HM770079). NCBI Genome Tree for *S. Bovismorbificans* at <https://www.ncbi.nlm.nih.gov/genome/tree/152/> (accessed on 6 September 2021) was used to identify nearest neighbors for *S. Bovismorbificans*.

### 3. Results and Discussion

#### 3.1. Genome Sequencing, Assembly and MLST Typing

The *S. Bovismorbificans* genomes in this study ranged from 4.5 to 5.0 MB. A complete genome of a 93 kb virulence plasmid from strain Sal610 was generated (Accession No. #: CP076746; Table 1). The shared *S. Bovismorbificans* and *S. Hindmarsh* immunogenic cluster signature from SeqSero was 8:r:1,5. The MLST tool on the CGE server grouped the *S. Bovismorbificans* strains into five different STs: 142, 2640, 377, 1499, and 150 (Supplemental Table S1). Nine out of 95 genomes were typed as ST150 and the rest were identified as ST142 or one of the highly similar STs: 2640, 377, and 1499 (Table 1) that differ from each other in one of the seven MLST loci. ST150 has a highly divergent allelic pattern (Supplemental File S1), where all seven alleles display significant differences from the ST142 strains and the other three STs. Comparative sequence analysis on the SEED server and local BLAST analysis of strain sal610 (ST377), Swiss strains (ST142, ST 1499, and ST2650), and strains sal676-681, N16\_1558 and M18\_12182 (ST150) consistently pointed to a high degree of genomic differences (data not shown) among these groups of strains. These preliminary analyses suggested extensive genome-wide sequence differences among the *S. Bovismorbificans* strains that were not fully captured by the conventional techniques consistent with the limitations of such approaches observed in other situations [44].

#### 3.2. Data Mining of DNA Tiling Digital Hybridization Profiles Identified Distinct Genomic Backbone Differences

Digital profiles from DNA tiling microarray hybridization experiments were created for representative strains of different STs and isolation sources using the FDA SEEC microarray platform. Initial data mining of an accompanying database made up of digital hybridization profiles of thousands of *Salmonella* strains sourced from past surveillance efforts and institutional collections suggested nucleotide diversity in the gene content of the queried genomes. For example, the sequence diversity of genes harvested from digital hybridization experiments of previously reported hummus-outbreak and surveillance *S. Bovismorbificans* strains [11,14] showed that the hummus strains were more closely related to *S. Typhimurium* (SARA2) and *S. Muenchen* consisting of SARA63-66 [38,45], while the surveillance strains (ST150) clustered as a distinct, independent group (Supplemental Figure S1A). Additionally, clinical Swiss *S. Bovismorbificans* strains belonging to ST142, ST2640, and ST1499 were more related to one another than to the ST150 strains. When the digital profiles of representative strains from different STs originally reported [8] and included in this study were queried against this database, they sorted between these two clusters (Supplemental Figure S1B), as expected. Previous reports [17,35] had pointed to significant sequence variations within STs using techniques ranging from PFGE to WGS to a limited number of genes used for clustering. As foodborne outbreaks of minor, zoonotic serovars such as *S. Bovismorbificans* are being reported to cause illness in humans [8,9] by entering the food-supply chain, a detailed understanding of the genomic differences between the lineages that constitute *S. Bovismorbificans* and other serovars not currently on the top list of public health agencies becomes important for risk assessment, import control, and the development of prevention measures for efficient food safety and public health outbreak management.

#### 3.3. Development of a Workflow for Core Genome Analysis Based on Complete Genomes from 150 *S. enterica* Serovars


We started with 645 complete and fully annotated genomes representing around 150 serovars that were obtained from the publicly available NCBI Genome database (Supplemental File S1). These sequences were combined with the new WGS assemblies of *S. Bovismorbificans* generated from this study (Table 1) to create a large BLAST database. When 4606 *S. Typhimurium* LT2 chromosomal genes were tested against this database at

50, 90, and 95% similarity levels, it yielded 1900 to 3400 chromosomal genes. After initial screening, 2830 genes at 90% similarity level were selected for manual curation to remove loci with alignment gaps, redundant annotations, and sequence quality errors. 2690 LT2 gene loci that contained at least one allelic difference in at least 90% of the genomes were selected as the 'wg-core gene' set for our phylogenomic studies. This list included the seven MLST housekeeping gene sequences and all the *S. Typhimurium* LT2 loci used as a reference, which were annotated and available in Supplemental File S6. The wg-core gene schema was tested for accuracy and reproducibility on genomes from an in-house collection of new and published strains (sub-sets of genomes from the data matrix presented in Supplemental File S2) prior to its application on the complete *S. Bovismorbificans* dataset from this study. It must be noted that this schema was tested only on the listed serovars of *S. enterica* subsp. *enterica*. Furthermore, extensive analysis is needed to evaluate the performance of this schema against the genome assemblies of other *S. enterica* sub-species. A broader comparison of the loci from this schema with genome-wide MLST markers in Enterobase [46], various bioinformatic methods used in genomic epidemiology [47] and SISTR database [48] is also necessary for developing a unified wg-core MLST marker dataset.

### 3.4. Genome Pathotyping of *S. Bovismorbificans* Strains Using Wg-Core Genes and *k*-Mer-Binning Methods

The 2690 wg-core genes clustered the 95 *S. Bovismorbificans* WGS assemblies into two high-level groups that exhibit distinct genomic differences (Figure 1). The two distinct clusters consisted of a larger cluster (Cluster 1) made up of Swiss, DC-hummus outbreak, and European strains, and a smaller cluster (Cluster 2) consisting of singletons from veterinary, food, and clinical sources from the USA, Switzerland, and Canada. More than 85 *S. Bovismorbificans* genomes from this study are represented in Cluster 1 and phylogenetically sort into the four highly related MLST groups identified earlier (ST142, ST2460, ST377, and ST1499). A ST142 sub-cluster included only Swiss strains from 2014–2016; the 2011 hummus outbreak sub-cluster strains belonging to ST377 grouped together; the ST1499 strains comprised a mixture of Canadian and Swiss strains, and the ST2460 cluster had five Swiss strains. The two polyphyletic genomic pathotypes observed in our study appear to be a serovar-wide phenomenon for *S. Bovismorbificans* when geographically and temporally different external genomes were also included for analysis. For example, seven of the external *S. Bovismorbificans* strains were sorted into Cluster 1 along with these genomes. A publicly available genome assembly, QAUR01 from NCBI, was a single ST1058 strain noted in this study. The five ST groups profiled in Cluster 1 had overlapping MLST allelic formulae with differences in up to three loci in some cases (Supplemental File S3). Our analyses demonstrated a conserved genomic backbone among these strains constituting the bulk of *S. Bovismorbificans* strains with publicly available genomic data (Figure 1). Cluster 2 consisted only of ST150 strains obtained from food, clinical ('sal' strains), and veterinary (M18\_12182) sources from the US and Canada ('Bovis' strains) along with a single Swiss strain, N16\_1558. The analysis identified CIES13, MXTS01, and QDND0, sharing a similar genomic backbone with the ST150 strains. The conserved core genomic backbone of the Cluster 1 strains was phylogenetically closer to the *S. Typhimurium* LT2 (reference strain) genome than that of Cluster 2 *S. Bovismorbificans* genomes as determined by this wg-core gene analysis. The conventional serotyping and sequence-based determination of the serovars of these two clusters were based on the immunogenic antigens and their coding sequences, respectively. Even if the *S. Bovismorbificans* strains of these two clusters appear to be sharing similar sequences for the serotyping antigens (based on sequence-based serotyping results from SeqSero2) resulting in the designation of the same serovar, these strains possess highly divergent core genome backbones and appear to be polyphyletic based on the 2690 wg-core gene schema developed as part of this study. These observations with the *S. Bovismorbificans* strains from different regions matched with our initial separation of hummus outbreak and surveillance strains into


clusters when compared with the digital DNA tiling microarray profiles of a legacy strain collection (Supplemental Figures S1A,B).



**Figure 1.** Wg-core gene analysis of 95 *S. Bovismorbificans* identifies two distinct clusters having different genomic backbones. *S. Bovismorbificans* strains were obtained from clinical, animal, feed or food and water or unknown sources isolated during 1984–1989 and 2011–2018 (from this study) and representative strains from NCBI. Cluster analysis was carried out using single nucleotide polymorphisms in 2690 core genes representing conserved backbone, and the phylogenetic tree was developed using the Maximum-Likelihood method [30] available on MEGAX's phylogenetic suite [32]. *S. Typhimurium* (LT2, single dot) was used as an outlier. Alleles in 48,344 positions were considered across 110 genomes spanning 2650+ out of 2690 core genes, which were considered tested over 500 bootstrapping iterations. The resulting circular tree is shown here. A browsable vertical layout of the tree is available in Supplemental Figure S1. Refer to Supplemental File S2 on the page for the comprehensive allelic data matrix from this analysis.

The 95 strains from the current study used for this analysis have a large proportion of food and clinical isolates (Table 1) from a Swiss collection. Nevertheless, the architecture of Cluster 1 strains displayed emerging sequence variations characteristic of robust microevolutionary processes, as observed from the emergence of different ST groups. When a sub-set of 35 Cluster 1 genomes was analyzed at higher resolution using wg-core genes, sub-groups of strains with significant SNP differences emerged (Figure 2). Partial quantitation of allelic differences in ST377, ST2640, and ST1499 strains individually in comparison with ST142 strains and micro-evolutionary profiling of the associated core-gene loci was carried out using annotations of reference strain LT2 (Supplemental File S3). For example, the ST1499 group consists of strains demonstrating significant sequence

divergence in hundreds of core gene loci when compared with ST142. The core gene loci associated with the ST1499 strains included kinases, transport and membrane proteins, fimbriae, and enzymes involved in various metabolic pathways, which appeared to be undergoing a faster rate of evolutionary change (Supplemental File S3, worksheet 2). The Cluster 2 ST150 strains also predictably exhibited sequence variations in the core-gene loci (Supplemental File S3, worksheet 4). Metadata analysis combined with these phylogenetic clusters also showed that in Switzerland from 2014–2018, *S. Bovismorbificans* strains belonging to four STs: 142, 377, 1499, or 2640 were prevalent at different time points, indicating long-term persistence of *S. Bovismorbificans* strains in the food supply chain. These results suggested that: (i) many Cluster 1 *S. Bovismorbificans* strains belonging to different conventional ST groups displayed a phylogenetically distinct, shared genomic backbone, and (ii) the emergent properties within these sub-groups of spatially and temporally discrete lineages were undergoing robust, quantifiable micro-evolutionary changes. However, robust biological experimentation is required to assess the phenotypic impact of these micro-evolutionary changes.



**Figure 2.** Phylogenetic analysis with wg-core gene SNPs in emerging ST lineages within genome type Cluster 1 of the *S. Bovismorbificans* serovar. The Cluster 1 consisted of 4 major STs and diverged significantly from ST150 strains of Cluster 2. Even among these related strains in Cluster 1, emerging polymorphisms exhibiting different rates of microevolution were observed using the SNP data matrix. UPGMA tree drawn on SPlitsTree 5.0. Refer to Supplemental File S3 for details of SNPs among the strains of different Cluster 1 STs.


Manual curation of NCBI Genome Tree for *S. Bovismorbificans* and data mining of the DNA Tiling database from the SEEC microarray platform (Supplemental File S2B) identified *Salmonella* serovars Muenchen and Hindmarsh as the closest neighbors to the strains in Cluster 1, and serovar *S. Takoradi* for strains in Cluster 2. To resolve the nearest neighbors to the *S. Bovismorbificans* clusters that would clarify the differences in the core-gene loci we had observed earlier, we carried out *k-mer*-binning analyses on a global collection of whole genomes from 265 *S. Bovismorbificans* (inclusive of the 95 from this study from Table 1), 60 *S. Muenchen* and nine *S. Hindmarsh* strains (Accession and serovars listed in columns A and B, respectively, in Supplemental File S4). The analysis generated a data-matrix with  $330 \times 330$  datapoints and quantified values of each binary genomic

comparison as a *k-mer* ratio were visualized as a heat-map color-coded in a teal blue (proximal) to orange (distal) values (Supplemental File S4, worksheet titled 'Jaccard Matrix'). Many *S. Hindmarsh* strains were similar to the *S. Bovismorbificans* strains found associated with Cluster 1, strains; however, two—AAOBOO01 (containing "8:r:" formula according to SeqSero) and SRR3710239 (containing the typical "8:r:1,5" formula on SeqSero)—appear to contain a Cluster 2 genomic backbone such as ST150 *S. Bovismorbificans* strains.

Other interesting findings were that a sample with assembly AAADAU01 was mistyped as *S. Hindmarsh* and appeared to be unrelated to any of the known *S. Hindmarsh* or *S. Bovismorbificans* strains. Furthermore, two ambiguously typed strains (AAOBOO01 and SRR3473907) with overlapping serotyping signatures were almost identical to the other ST150 *S. Bovismorbificans* strains. A subset of this data matrix was illustrated that depicts a snapshot of this whole genome-based *k-mer*-binning analysis (Figure 3) and the evolutionary distance between pair-wise genome comparisons is shown [represented by teal blue (highly similar) to sky blue (somewhat similar) to orange (highly dissimilar) values]. The evolutionary distance between pair-wise genome comparisons was shown in values from teal blue (highly similar) to sky blue (some similarity) to orange (highly dissimilar) interspersed with a gradient of tan values for decreasing similarity. Clusters 1 and 2 were placed apart and their individual comparisons shown in orange highlighted the evolutionary distance measured by the *k-mer*-binning method. Interestingly, *S. Typhimurium* LT2 and *S. Muenchen* AUQE01 shown in sky blue for comparison with Cluster 1 appeared to suggest a more similar genome backbone among these groups when compared with Cluster 2. Similarly, *S. Takoradi* strain (NPMA01) defined the boundary for Cluster 2 (shown in tan value) strains when compared with Cluster 1. A ST150 strain CIES13 appeared to have significant genomic divergence when compared with other Cluster 2 strains. When the WGS assemblies from these nearest neighbors of the two cluster groups were queried for wg-core SNP-based clustering (Figure 4), the patterns similar to the *k-mer* profiling, as observed in Figure 3, were observed. ST150 strains formed a separate cluster with two *S. Takoradi* strains as their nearest neighbors (Figure 4). AAADAU01, a mistyped *S. Hindmarsh* isolate, was identified as a strain of serovar *S. Weltevreden* based on its core-genome profile (data not shown). Two *S. Muenchen* strains and *S. Typhimurium* LT2 (reference genome) flanked the top Cluster 1 that contained *S. Bovismorbificans* genomes from this study and *S. Hindmarsh* strains from NCBI. The polyphyletic nature of *S. Hindmarsh* strains evident from our analysis needs further investigation. The wg-core gene method relied on identifying sequence variations (SNPs) in generating different clusters while the *k-mer* binning approach used the number of conserved *k-mers* between any two genomes. The concordance of the two different whole genome analytic methods we have applied to understand the phylogenetic relationship between the two clusters in this study was evident from these results illustrated in Figures 1–4 and associated Supplemental File S4.

|             | Bovis3114 | Bovis_277 | N14_0147 | N14_1267 | N15_0111 | N15_0584 | N15_2285 | N16_0447 | N16_0656 | N16_1158 | N16_2042 | N17_0002 | N17_2111 | QAUR01 | Sal610 | Sal616 | AAIBEG01 | ERR1755559 | JYUT01 | LT2Genome | AUQE01 | NPMA01 | AAADAU01 | M18_12182 | QDN001 | Sal676 | Sal678 | Sal681 | AAOBOO01 | Boviscies13 |       |
|-------------|-----------|-----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|--------|--------|--------|----------|------------|--------|-----------|--------|--------|----------|-----------|--------|--------|--------|--------|----------|-------------|-------|
| Bovis3114   | 1.000     | 0.897     | 0.949    | 0.992    | 0.864    | 0.951    | 0.945    | 0.959    | 0.976    | 0.993    | 0.994    | 0.942    | 0.993    | 0.865  | 0.932  | 0.929  | 0.893    | 0.963      | 0.950  | 0.565     | 0.568  | 0.503  | 0.478    | 0.477     | 0.476  | 0.479  | 0.478  | 0.478  | 0.471    | 0.399       |       |
| Bovis_277   | 0.897     | 1.000     | 0.899    | 0.897    | 0.865    | 0.899    | 0.890    | 0.906    | 0.885    | 0.898    | 0.899    | 0.891    | 0.900    | 0.856  | 0.871  | 0.870  | 0.898    | 0.906      | 0.898  | 0.580     | 0.571  | 0.503  | 0.494    | 0.475     | 0.474  | 0.476  | 0.475  | 0.476  | 0.469    | 0.397       |       |
| N14_0147    | 0.949     | 0.899     | 1.000    | 0.947    | 0.870    | 0.975    | 0.949    | 0.985    | 0.933    | 0.949    | 0.950    | 0.966    | 0.949    | 0.862  | 0.923  | 0.920  | 0.886    | 0.973      | 0.976  | 0.564     | 0.566  | 0.502  | 0.479    | 0.476     | 0.475  | 0.477  | 0.476  | 0.477  | 0.470    | 0.397       |       |
| N14_1267    | 0.992     | 0.897     | 0.947    | 1.000    | 0.863    | 0.949    | 0.944    | 0.957    | 0.975    | 0.992    | 0.993    | 0.941    | 0.993    | 0.863  | 0.931  | 0.928  | 0.891    | 0.961      | 0.948  | 0.565     | 0.567  | 0.503  | 0.478    | 0.477     | 0.476  | 0.478  | 0.477  | 0.478  | 0.471    | 0.398       |       |
| N15_0111    | 0.864     | 0.865     | 0.870    | 0.863    | 1.000    | 0.860    | 0.856    | 0.867    | 0.852    | 0.864    | 0.865    | 0.852    | 0.865    | 0.893  | 0.844  | 0.842  | 0.907    | 0.870      | 0.874  | 0.548     | 0.545  | 0.483  | 0.465    | 0.456     | 0.455  | 0.457  | 0.456  | 0.457  | 0.450    | 0.383       |       |
| N15_0584    | 0.951     | 0.899     | 0.975    | 0.949    | 0.860    | 1.000    | 0.956    | 0.985    | 0.934    | 0.950    | 0.951    | 0.987    | 0.951    | 0.864  | 0.925  | 0.922  | 0.894    | 0.975      | 0.962  | 0.565     | 0.567  | 0.503  | 0.479    | 0.477     | 0.476  | 0.478  | 0.477  | 0.478  | 0.471    | 0.398       |       |
| N15_2285    | 0.945     | 0.890     | 0.949    | 0.944    | 0.856    | 0.956    | 1.000    | 0.957    | 0.930    | 0.945    | 0.947    | 0.948    | 0.946    | 0.860  | 0.927  | 0.924  | 0.886    | 0.950      | 0.938  | 0.562     | 0.560  | 0.499  | 0.479    | 0.473     | 0.472  | 0.474  | 0.473  | 0.473  | 0.467    | 0.395       |       |
| N16_0447    | 0.959     | 0.906     | 0.985    | 0.957    | 0.867    | 0.985    | 0.957    | 1.000    | 0.943    | 0.959    | 0.959    | 0.976    | 0.959    | 0.871  | 0.933  | 0.931  | 0.895    | 0.984      | 0.971  | 0.568     | 0.571  | 0.507  | 0.482    | 0.480     | 0.479  | 0.482  | 0.481  | 0.481  | 0.474    | 0.400       |       |
| N16_0656    | 0.976     | 0.885     | 0.933    | 0.975    | 0.852    | 0.934    | 0.955    | 0.977    | 1.000    | 0.977    | 0.978    | 0.926    | 0.980    | 0.852  | 0.917  | 0.915  | 0.879    | 0.947      | 0.934  | 0.558     | 0.561  | 0.497  | 0.473    | 0.471     | 0.470  | 0.473  | 0.472  | 0.472  | 0.465    | 0.394       |       |
| N16_1158    | 0.993     | 0.898     | 0.949    | 0.992    | 0.864    | 0.950    | 0.955    | 0.977    | 1.000    | 0.996    | 0.942    | 0.994    | 0.864    | 0.932  | 0.929  | 0.892  | 0.963    | 0.950      | 0.565  | 0.568     | 0.503  | 0.478  | 0.477    | 0.476     | 0.478  | 0.477  | 0.478  | 0.471  | 0.398    |             |       |
| N16_2042    | 0.994     | 0.899     | 0.950    | 0.993    | 0.865    | 0.951    | 0.947    | 0.959    | 0.978    | 0.996    | 1.000    | 0.943    | 0.995    | 0.865  | 0.933  | 0.930  | 0.893    | 0.963      | 0.950  | 0.565     | 0.568  | 0.503  | 0.478    | 0.477     | 0.476  | 0.478  | 0.477  | 0.478  | 0.471    | 0.398       |       |
| N17_0002    | 0.942     | 0.891     | 0.966    | 0.941    | 0.852    | 0.987    | 0.948    | 0.976    | 0.926    | 0.942    | 0.943    | 1.000    | 0.942    | 0.867  | 0.917  | 0.914  | 0.887    | 0.966      | 0.953  | 0.561     | 0.563  | 0.500  | 0.477    | 0.474     | 0.473  | 0.475  | 0.474  | 0.474  | 0.468    | 0.396       |       |
| N17_2111    | 0.993     | 0.900     | 0.949    | 0.993    | 0.865    | 0.951    | 0.946    | 0.959    | 0.980    | 0.994    | 0.995    | 0.942    | 1.000    | 0.866  | 0.932  | 0.930  | 0.894    | 0.963      | 0.950  | 0.566     | 0.568  | 0.504  | 0.478    | 0.477     | 0.476  | 0.479  | 0.478  | 0.478  | 0.471    | 0.399       |       |
| QAUR01      | 0.865     | 0.856     | 0.862    | 0.863    | 0.893    | 0.864    | 0.860    | 0.871    | 0.852    | 0.864    | 0.865    | 0.867    | 0.866    | 1.000  | 0.845  | 0.843  | 0.873    | 0.874      | 0.862  | 0.542     | 0.542  | 0.488  | 0.469    | 0.460     | 0.459  | 0.461  | 0.460  | 0.461  | 0.455    | 0.386       |       |
| Sal610      | 0.932     | 0.871     | 0.923    | 0.931    | 0.844    | 0.925    | 0.927    | 0.933    | 0.917    | 0.932    | 0.933    | 0.917    | 0.932    | 0.845  | 1.000  | 0.990  | 0.871    | 0.937      | 0.924  | 0.556     | 0.554  | 0.494  | 0.467    | 0.468     | 0.467  | 0.469  | 0.468  | 0.469  | 0.462    | 0.391       |       |
| Sal616      | 0.929     | 0.870     | 0.920    | 0.928    | 0.842    | 0.922    | 0.924    | 0.931    | 0.915    | 0.929    | 0.930    | 0.914    | 0.930    | 0.843  | 0.990  | 1.000  | 0.870    | 0.935      | 0.922  | 0.556     | 0.553  | 0.493  | 0.467    | 0.467     | 0.467  | 0.469  | 0.468  | 0.468  | 0.462    | 0.391       |       |
| AAIBEG01    | 0.893     | 0.898     | 0.886    | 0.891    | 0.907    | 0.894    | 0.886    | 0.895    | 0.879    | 0.892    | 0.893    | 0.887    | 0.894    | 0.873  | 0.871  | 0.870  | 1.000    | 0.899      | 0.887  | 0.563     | 0.558  | 0.495  | 0.481    | 0.467     | 0.466  | 0.468  | 0.467  | 0.468  | 0.461    | 0.391       |       |
| ERR1755559  | 0.963     | 0.906     | 0.973    | 0.961    | 0.870    | 0.975    | 0.950    | 0.984    | 0.947    | 0.963    | 0.963    | 0.966    | 0.963    | 0.874  | 0.937  | 0.935  | 0.899    | 1.000      | 0.974  | 0.570     | 0.573  | 0.508  | 0.481    | 0.482     | 0.481  | 0.483  | 0.482  | 0.482  | 0.476    | 0.401       |       |
| JYUT01      | 0.950     | 0.898     | 0.976    | 0.948    | 0.874    | 0.962    | 0.938    | 0.971    | 0.934    | 0.950    | 0.950    | 0.953    | 0.950    | 0.862  | 0.924  | 0.922  | 0.887    | 0.974      | 1.000  | 0.564     | 0.566  | 0.504  | 0.476    | 0.478     | 0.477  | 0.479  | 0.478  | 0.479  | 0.472    | 0.399       |       |
| LT2Genome   | 0.565     | 0.580     | 0.564    | 0.565    | 0.548    | 0.562    | 0.565    | 0.562    | 0.568    | 0.558    | 0.565    | 0.565    | 0.561    | 0.566  | 0.542  | 0.556  | 0.556    | 0.563      | 0.570  | 0.564     | 1.000  | 0.557  | 0.491    | 0.490     | 0.465  | 0.464  | 0.466  | 0.465  | 0.466    | 0.459       | 0.389 |
| AUQE01      | 0.568     | 0.571     | 0.566    | 0.567    | 0.545    | 0.567    | 0.560    | 0.571    | 0.561    | 0.568    | 0.568    | 0.563    | 0.568    | 0.542  | 0.554  | 0.553  | 0.558    | 0.573      | 0.566  | 0.557     | 1.000  | 0.494  | 0.471    | 0.468     | 0.467  | 0.470  | 0.469  | 0.469  | 0.462    | 0.391       |       |
| NPMA01      | 0.503     | 0.503     | 0.502    | 0.503    | 0.483    | 0.503    | 0.499    | 0.507    | 0.497    | 0.503    | 0.503    | 0.500    | 0.504    | 0.488  | 0.494  | 0.493  | 0.495    | 0.508      | 0.504  | 0.491     | 0.494  | 1.000  | 0.514    | 0.559     | 0.558  | 0.561  | 0.560  | 0.560  | 0.552    | 0.463       |       |
| AAADAU01    | 0.478     | 0.494     | 0.479    | 0.478    | 0.465    | 0.479    | 0.479    | 0.482    | 0.473    | 0.478    | 0.478    | 0.477    | 0.478    | 0.469  | 0.467  | 0.467  | 0.481    | 0.481      | 0.476  | 0.490     | 0.471  | 0.514  | 1.000    | 0.482     | 0.481  | 0.484  | 0.483  | 0.483  | 0.476    | 0.403       |       |
| M18_12182   | 0.477     | 0.475     | 0.476    | 0.477    | 0.456    | 0.477    | 0.473    | 0.480    | 0.471    | 0.477    | 0.477    | 0.474    | 0.477    | 0.460  | 0.468  | 0.467  | 0.467    | 0.482      | 0.478  | 0.465     | 0.468  | 0.559  | 0.482    | 1.000     | 0.983  | 0.991  | 0.988  | 0.989  | 0.970    | 0.778       |       |
| QDN001      | 0.476     | 0.474     | 0.475    | 0.476    | 0.455    | 0.476    | 0.472    | 0.479    | 0.470    | 0.476    | 0.476    | 0.473    | 0.476    | 0.459  | 0.467  | 0.467  | 0.466    | 0.481      | 0.477  | 0.464     | 0.467  | 0.558  | 0.481    | 0.983     | 1.000  | 0.985  | 0.993  | 0.994  | 0.966    | 0.774       |       |
| Sal676      | 0.479     | 0.476     | 0.477    | 0.478    | 0.457    | 0.478    | 0.474    | 0.482    | 0.473    | 0.478    | 0.478    | 0.475    | 0.479    | 0.461  | 0.469  | 0.469  | 0.468    | 0.483      | 0.479  | 0.466     | 0.470  | 0.561  | 0.484    | 0.991     | 0.986  | 1.000  | 0.994  | 0.974  | 0.781    |             |       |
| Sal678      | 0.478     | 0.475     | 0.476    | 0.477    | 0.456    | 0.477    | 0.473    | 0.481    | 0.472    | 0.477    | 0.477    | 0.474    | 0.478    | 0.460  | 0.468  | 0.468  | 0.467    | 0.482      | 0.478  | 0.465     | 0.469  | 0.560  | 0.483    | 0.988     | 0.983  | 0.992  | 1.000  | 0.992  | 0.971    | 0.778       |       |
| Sal681      | 0.478     | 0.476     | 0.477    | 0.478    | 0.457    | 0.478    | 0.473    | 0.481    | 0.472    | 0.478    | 0.478    | 0.474    | 0.478    | 0.461  | 0.469  | 0.468  | 0.468    | 0.482      | 0.479  | 0.466     | 0.469  | 0.560  | 0.483    | 0.989     | 0.984  | 0.994  | 0.992  | 1.000  | 0.972    | 0.779       |       |
| AAOBOO01    | 0.471     | 0.469     | 0.470    | 0.471    | 0.450    | 0.471    | 0.467    | 0.474    | 0.465    | 0.471    | 0.471    | 0.468    | 0.471    | 0.455  | 0.462  | 0.462  | 0.461    | 0.476      | 0.472  | 0.459     | 0.462  | 0.552  | 0.476    | 0.970     | 0.966  | 0.974  | 0.971  | 0.972  | 1.000    | 0.767       |       |
| Boviscies13 | 0.399     | 0.397     | 0.397    | 0.398    | 0.383    | 0.398    | 0.395    | 0.400    | 0.394    | 0.398    | 0.398    | 0.396    | 0.399    | 0.386  | 0.391  | 0.391  | 0.391    | 0.401      | 0.399  | 0.389     | 0.391  | 0.463  | 0.403    | 0.778     | 0.774  | 0.781  | 0.778  | 0.779  | 0.767    | 1.000       |       |

**Figure 3.** *k*-mer-binning analysis of genomes from *S. Bovismorbificans* and closely related serovars. A snap shot of the Jaccard index from the *k*-mer-binning analysis ( $k = 30$ ) of 330 WGS assemblies dataset from five *Salmonella* serovars was illustrated (Heatmap orange to teal blue = most divergent to most similar). *S. Typhimurium* (LT2Genome) and *S. Muenchen* (AUQE01) are closer to Cluster 1 strains (shown in sky blue), and *S. Takoradi* (NPMA01) is the closest serovar to the ST150 Cluster 2 strains (shown in tan). Distinct genomic pathotypes of the two *S. Bovismorbificans* clusters suggest evolutionarily independent hinging of a single serotyping gene cluster (“8:r:1,5”) in two different lineages of SE with different genomic backbones. The complete phylogenomic analysis with 330 genomes is presented as Supplemental File S4.




**Figure 4.** Phylogenomic profiling of *S. Bovismorbificans* and closely related serovars. *S. Bovismorbificans* is made up of strains with two distinct genomic backbones, naturally grouped into two distant clusters. Cluster 1 strains are closer to *S. Typhimurium* (represented by the reference genome LT2) and *S. Muenchen* (AUQE01 and ARYX01) than to the Cluster 2 *S. Bovismorbificans* strains. WGS analysis from this study identified *S. Takoradi* (NPMA01 and VCTV01) as the closest serovar to the Cluster 2 *S. Bovismorbificans* strains. *S. Hindmarsh* strains sharing the antigen-cluster with *S. Bovismorbificans* in White–Kauffmann–Le Minor scheme also exhibit different genomic backbones as in the case of *S. Bovismorbificans*. Two of the Hindmarsh strains are aligned with Cluster 1 (SRR1783167 and AAIBEG01), while AAADAU01 and AAOBOO01 align within Cluster 2. For this analysis, Neighbor-Joining method implemented on MEGA X suite was used yielding a data matrix made up of 46,403 base positions across 35 genomes. MLST, DNA tiling Microarray (MA), whole genome sequence (WGS) based phylogenetic analysis, core-gene alleles spanning more than 2700 core genes, and k-mer binning based on NGS datasets and conventional assays all point to two

divergent genome types bearing a single serotyping cluster. SRR3473907 was not clearly serotyped either by its NCBI BioSample record or by SeqSero in this study.

Intra-serovar differences among lineages within some *Salmonella* serovars are being recognized from the growing volume of WGS data from outbreak and surveillance samples, geographically distributed and within legacy strain collections. Genome-wide and core genome analysis of *Salmonella* serovars using WGS datasets has been proven to be capable of profiling underlying evolutionary processes contributing to the genome backbone differences among different lineages constituting a conventionally typed serovar [19–22,46,47]. The application of WGS technology for sequence-based serotyping [25,48], characterization of outbreak isolates [49,50], source-tracking in the epidemiological investigation of foodborne outbreaks, and phylogenetic relationships [51–55] have been well documented. The robust bioinformatic workflow from this study utilizes an ad hoc set of complete genome assemblies for high-resolution phylogenetic analysis and adds to the existing *S. enterica* core-genes (cgMLST) and whole genome MLST (wgMLST) typing schema hosted on public resources such as Enterobase and SISTR [46,48]. Detailed genomic analysis in combination with sequence- and/or serology-based serotyping alone could provide a broader explanation for the complex intra-serovar differences that are not easily explained solely in terms of conventional serological methods.


### 3.5. Plasmids, Phage, AMR, and Virulence Profiling of *S. Bovismorbificans* Genome Types

Predictive analysis of mobilome and virulence factor elements in the WGS assemblies of *S. Bovismorbificans* from this study was carried out on PlasmidFinder, PHASTER, SPIFinder and GalaxyTrakr AMRFinderPlus tools as described earlier. A 94 kb putative virulence plasmid, pVirBov (HF969016), such as the *S. Typhimurium* LT2 plasmid pSLT, was reported from a Malawian clinical *S. Bovismorbificans* strain 3114 [9]. PlasmidFinder web tool identified IncFIB(s), IncFII(s) plasmids in 59 out of 60 Cluster 1 strains (Supplemental Figure S3), while other plasmid types such as IncI1, IncI2, and colI56 were noted in a few strains (Supplemental File S5). Previously, the presence of a putative 90 kb plasmid was predicted [7] in some of the hummus-outbreak strains included in this study (Table 1). We designed a local BLAST analysis to identify the homologous sequences of this plasmid in the WGS assemblies. Fifty-seven *S. Bovismorbificans* strains hosted the homologous plasmid sequences on a single contig. The alignment of these contigs illustrated a possible loss of complete annotation (Supplemental File S4, mauve alignment of putative plasmid-bearing contigs). We sequenced the complete genome of the pVirBov-like virulence plasmid from the clinical strain Sal610. The 93,477 bp closed assembly of pSal610 was then used to compare with pVirBov and representative genome contigs from the study strains (Figure 5 and Supplemental Figure S4). Low complexity regions, missing sequences, and ambiguous sequence fragments in pVirBov, Bovis\_277\_contig24 and Sal682\_contig35 were identified. The first complete sequence of *S. Bovismorbificans* virulence plasmid, pSal610 (CP076746), from clinical strain Sal610 isolated during a hummus outbreak in 2011, differed from the homologous pVirBov from strain 3114 by additional 360 bases. Further analysis is needed to understand the occurrence of this plasmid among other SE lineages, and its contribution to the virulence of the *S. Bovismorbificans* Cluster 1 strains.



**Figure 5.** Comparison of pVirBov 3114, plasmids from Swiss strains and the closed assembly pSal610 (humus outbreak). Closed genome of 93,777 bp long virulence plasmid from *S. Bovismorbificans* strain Sal610 (CP076746) was used as the reference genome to compare pVIRBov from strain 3114 [9] and a few selected Cluster 1 isolates (Table 1) from this study. BLAST+ based comparison and visualization were carried out on BRIG 0.95 [39]. White patches in a circle indicate any missing sequences in the query genome.

PHASTER tool results (Supplemental File S5) suggested a variety of homologous sequences only in Cluster 1 strains. A *S. Enteritidis* ELPhiS prophage RE2010-like sequence was predicted only in ST377 strains from the 2011 hummus outbreak in Washington DC and in various isolates from the Swiss collection. A complete RE2010-like phage sequence was identified from the ST377 strains by BLAST analysis (data not shown). Figure 6 illustrates the presence of the RE2010-like phage in Sal610 (contig1), N16\_2718 (contig8), and N14\_2376 (contig 17). The GalaxyTrakr AMRFinderPlus tool was used to evaluate the presence of antimicrobial resistance genes that may be present in the 95 *S. Bovismorbificans* strains. Previously, genes conferring resistance to Quinolone and other antimicrobials had been reported in some serovars including *S. Bovismorbificans* [56,57]. The results presented in Table 2 show that seven Cluster 1 strains were identified to contain multiple resistance genes against antimicrobials, such as Cephalosporin, Quinolone, Tetracycline, Beta-Lactam, Tetracycline, Gentamicin, Kanamycin, Bleomycin, Bleomycin, Sulfonamide, Streptomycin, Chloramphenicol, Streptomycin and Quaternary Ammonium. The results of this predictive analysis (Supplemental File S3) further highlighted extensive genotypic differences between the two polyphyletic clusters identified in this study for *S. Bovismorbificans* and mirror the established pathogenic properties of major serovars such as Typhimurium, Enteritidis, Heidelberg, and Newport [58].



**Figure 6.** ST377 strains from clinical and food sources contain a unique *S. Enteritidis* RE-2010-like phage. An ELPhiS prophage RE-2010 (Accession: HM770079) from *S. Enteritidis* LK5 [59] was identified in ST377 genomes. A comparison of representative genomes from three strains isolated from different sample sources are illustrated above: Sal610 contig 1 (AZKX01000024)—Clinical, N16\_2718\_contig\_8 (WSCE01000008)—Clinical and N14\_2376\_contig\_17 (WSDY01000017)—Food, Onion. The *S. Bovismorbificans* version of the phage appeared to be homologous with predicted additional ORFs in the structure protein core. Prevalence of other phage sequences in *S. Bovismorbificans* strains from this study are listed in the Supplemental File S5.

**Table 2.** Antimicrobial resistance genes identified in the *S. Bovismorbificans* genomes <sup>a</sup>.

| Strain   | Gene                          | Subclass                         | Sequence Name/Description                                   |
|----------|-------------------------------|----------------------------------|-------------------------------------------------------------|
| N14_0147 | <i>aph(3'')-Ib</i>            | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(3'')-Ib             |
|          | <i>sul2</i>                   | Sulfonamide                      | Sulfonamide-resistant dihydropteroate synthase Sul2         |
|          | <i>aph(6)-Id</i>              | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(6)-Id               |
| N15_0111 |                               |                                  | Class A extended-spectrum beta-lactamase CTX-M-55           |
|          | <i>bla<sub>CTX-M-55</sub></i> | Cephalosporin                    | Quinolone resistance pentapeptide repeat protein QnrS1      |
|          | <i>qnrS1</i>                  | Quinolone                        | Tetracycline efflux MFS transporter Tet(A)                  |
|          | <i>tet(A)</i>                 | Tetracycline                     | Class A broad-spectrum beta-lactamase TEM-1                 |
|          | <i>bla<sub>TEM-1</sub></i>    | Beta-lactam                      | Tetracycline resistance ribosomal protection protein Tet(M) |
|          | <i>tet(M)</i>                 | Tetracycline                     | Aminoglycoside N-acetyltransferase AAC(3)-IId               |
|          | <i>aac(3)-IId</i>             | Gentamicin                       | Aminoglycoside O-phosphotransferase APH(3')-IIa             |
|          | <i>aph(3')-IIa</i>            | Kanamycin                        | Bleomycin binding protein BLMT                              |
|          | <i>ble</i>                    | Bleomycin                        | Bleomycin binding protein                                   |
|          | <i>bleO</i>                   | Bleomycin                        | Sulfonamide-resistant dihydropteroate synthase Sul3         |
|          | <i>sul3</i>                   | Sulfonamide                      | ANT(3'')-Ia family aminoglycoside                           |
|          | <i>aadA2</i>                  | Streptomycin                     | Nucleotidyltransferase AadA2                                |
|          | <i>cmlA1</i>                  | Chloramphenicol                  | Chloramphenicol efflux MFS transporter CmlA1                |
|          | <i>aadA1</i>                  | Streptomycin                     | ANT(3'')-Ia family aminoglycoside                           |
|          | <i>qacL</i>                   | Quaternary ammonium <sup>b</sup> | Nucleotidyltransferase AadA1                                |
| N16_0208 |                               |                                  | Quaternary ammonium compound efflux SMR transporter QacL    |
|          | <i>aph(6)-Id</i>              | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(6)-Id               |
|          | <i>sul2</i>                   | Sulfonamide                      | Sulfonamide-resistant dihydropteroate synthase Sul2         |
|          | <i>aph(3'')-Ib</i>            | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(3'')-Ib             |
| N16_2574 | <i>tet(A)</i>                 | Tetracycline                     | Tetracycline efflux MFS transporter Tet(A)                  |
|          | <i>dfrA1</i>                  | Trimethoprim                     | Trimethoprim-resistant dihydrofolate reductase DfrA1        |
|          | <i>bla<sub>TEM-1</sub></i>    | Beta-lactam                      | Class A broad-spectrum beta-lactamase TEM-1                 |
|          | <i>sul2</i>                   | Sulfonamide                      | Sulfonamide-resistant dihydropteroate synthase Sul2         |
| N16_2598 | <i>bla<sub>CTX-M-1</sub></i>  | Cephalosporin                    | Class A extended-spectrum beta-lactamase CTX-M-1            |
|          | <i>tet(A)</i>                 | Tetracycline                     | Tetracycline efflux MFS transporter Tet(A)                  |
| N16_2955 | <i>aph(3'')-Ib</i>            | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(3'')-Ib             |
|          | <i>aph(6)-Id</i>              | Streptomycin                     | Aminoglycoside O-phosphotransferase APH(6)-Id               |

|          |                    |                             |                                                         |
|----------|--------------------|-----------------------------|---------------------------------------------------------|
| N17_0502 | <i>tet(A)</i>      | Tetracycline                | Tetracycline efflux MFS transporter Tet(A)              |
|          | <i>sul2</i>        | Sulfonamide                 | Sulfonamide-resistant dihydropteroate synthase Sul2     |
|          | <i>floR</i>        | Chloramphenicol/florfenicol | Chloramphenicol/florfenicol efflux MFS transporter FloR |
|          | <i>qnrB19</i>      | ol                          | Quinolone resistance pentapeptide repeat protein QnrB19 |
|          | <i>sul2</i>        | Quinolone                   | Sulfonamide-resistant dihydropteroate synthase Sul2     |
|          | <i>aph(3'')-Ib</i> | Sulfonamide                 | Aminoglycoside O-phosphotransferase APH(3'')-Ib         |
|          | <i>aph(6)-Id</i>   | Streptomycin                | Aminoglycoside O-phosphotransferase APH(6)-Id           |
|          | <i>tet(A)</i>      | Streptomycin                | Aminoglycoside O-phosphotransferase APH(6)-Id           |
|          |                    | Tetracycline                | Tetracycline efflux MFS transporter Tet(A)              |

<sup>a</sup> Galaxy GenomeTrakr AMRFinder tool used for identification of antimicrobial resistance (AMR). All FASTA genomes of 98 *S. Bovismorbificans* were scanned, and 7 strains captured the acquired antimicrobial resistance genes. <sup>b</sup> Element type: Stress; Element subtype: Biocide.

#### 4. Conclusions

The high resolution WGS analyses of *S. Bovismorbificans* reported in this study and other SE serovars by others clearly highlight the need for a broader sequence-based genome pathotyping framework that recognizes genome-wide genetic boundaries (vertical) and the emergent adaptive/incidental features (horizontal) of the genome pathotypes even within conventionally defined serovars [20,21]. Together, these results demonstrate the genomic diversity of *S. Bovismorbificans* strains and provide details of the sequence diversity separating the two polyphyletic lineages recognized and characterized in this study. The schema of 2690 wg-core genome loci used to identify the conserved genome backbone could be applied to determine the genome structure of other SE serovars having intra-serovar differences and elucidate the genomic structure within a population of *S. enterica* pathogens that persist and circulate for years, with different STs succeeding over time. As demonstrated for *S. Bovismorbificans*, wg-core genome analysis could be used to identify misidentified strains by the limitations of conventional serotyping methods and ameliorate the characterization of emerging sub-groups within a serovar. The characteristic mobilome features described in this study will facilitate proper classification of emerging genome pathotypes of *S. Bovismorbificans*. The application of WGS datasets and methods based on the genome sequences have become an integral part of not just understanding the source-tracking of strains, but also in understanding the genomic diversity of *S. enterica* strains in food-safety investigations. The results from this study of *S. Bovismorbificans* using phylogenomic analysis highlight the need to use the WGS data, including annotations to expand our understanding of the genomic and phenotypic diversity among the populations of strains within all *Salmonella* serovars that display the potential to infect humans and in contaminating the animal food supply chain. A comprehensive understanding of emerging pathotypes also of minor serovars of *S. enterica* in addition to those already on top of the list for surveillance would necessarily complement ongoing post-genomics efforts to apply NGS data and methods to prevent new sources of foodborne illnesses and additional serovars adding to the growing list of *S. enterica* pathogens.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/microorganisms10061199/s1>, Supplemental File S1 to S6 consisting of the list of serovars used to develop wg-core genome scheme, Jaccard matrix comparing 330 genomes using *k-mers*, mobilome data, 2650 core genes developed as part of this study, and a large SNP matrix consisting of 48,344 positions in 2512 of 2650 core genes; Supplemental Figures-S1A,S1B, S2, S3 and S4 with data mining of microarray analysis of *S. Bovismorbificans* from humus, other foods, clinical and unknown origins, and a snapshot of MAUVE comparison of the pSal610 virulence plasmid with many other representative strains; and Supplemental Table S1 with allelic profiles by 6 ST spanning all the known *S. Bovismorbificans* isolates.

**Author Contributions:** G.R.G., B.D.T., and R.S. designed the study, wrote the drafts, and finalized the submission; H.J. and J.J.-G.B. contributed to the initial and final drafts; R.S. and A.M. characterized and provided 76 Swiss strains; S.T. and F.P. provided the five Canadian assemblies; D.C.S. provided the MN calf production strain; J.J.-G.B. and L.E. carried out the characterization of hummus outbreak strains; J.G., I.R.P., and M.K.M. generated and analyzed microarray data, M.K.M. generated the matrices from *k-mer* analysis, and G.R.G. carried out the rest of the genomic analysis and illustrations; H.J. carried out genome sequencing, data submission, and supervised assembly and annotations by S.F. and C.Z.W.; L.M.W. and F.N. carried out phage and plasmid analysis; T.M. and M.A. closed the plasmid genome. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by through the USFDA appropriations.

**Institutional Review Board Statement:** The study did not require approval from an institutional Review Board.

**Informed Consent Statement:** Informed consent was not required.

**Data Availability Statement:** All WGS datasets have been submitted to the NCBI GenomeTrakr project and the details are provided in Table 1. Supplemental Files S1–S6, Supplemental Figures S1–S4, and Supplemental Table S1 contain additional information and illustrations relevant to the results presented in the main text.

**Acknowledgments:** We thank the University of Maryland, Joint Institute for Food Safety and Applied Nutrition (JIFSAN) for supporting JIFSAN interns, L.W., F.N., and S.F. We also thank the Oak Ridge Institute for Science and Education of Oak Ridge, Tennessee, USA, for sponsoring research fellow, H.J. We would also like to thank the FDA Pathways program for supporting intern, C.Z.W. This research was internally funded through USA FDA appropriations, and in part by the University of Maryland JIFSAN Program through a cooperative agreement with the FDA (Contract no. FDU001418). All authors read and approved the manuscript.

**Conflicts of Interest:** All authors declare no conflict of interest with regards to the work reported in this manuscript.

## References

1. Havelaar, A.H.; Kirk, M.D.; Torgerson, P.R.; Gibb, H.J.; Hald, T.; Lake, R.J.; Praet, N.; Bellinger, D.C.; de Silva, N.R.; Gargouri, N.; et al.; World Health Organization Foodborne Disease Burden Epidemiology Reference Group. World Health Organization Global Estimates and Regional Comparisons of the Burden of Foodborne Disease in 2010. *PLoS Med.* **2015**, *12*, e1001923.
2. Scallan, E.; Hoekstra, R.M.; Angulo, F.J.; Tauxe, R.V.; Widdowson, M.A.; Roy, S.L.; Jones, J.L.; Griffin, P.M. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* **2011**, *17*, 7–15.
3. Nguyen, D.T.; Kanki, M.; Nguyen, P.D.; Le, H.T.; Ngo, P.T.; Tran, D.N.; Le, N.H.; Dang, C.V.; Kawai, T.; Kawahara, R.; et al. Prevalence, antibiotic resistance, and extended-spectrum and AmpC  $\beta$ -lactamase productivity of *Salmonella* isolates from raw meat and seafood samples in Ho Chi Minh City, Vietnam. *Int. J. Food Microbiol.* **2016**, *236*, 115–122.
4. Pönkä, A.; Andersson, Y.; Siitonen, A.; de Yong, B.; Jahkola, M.; Haikala, O.; Kuhmonen, A.; Pakkala, P. *Salmonella* in alfalfa sprouts. *Lancet* **1995**, *345*, 462–463.
5. Puohiniemi, R.; Heiskanen, T.; Siitonen, A. Molecular epidemiology of two international sprout-borne *Salmonella* outbreaks. *J. Clin. Microbiol.* **1997**, *35*, 2487–2491.
6. Rimhanen-Finne, R.; Niskanen, R.T.; Lienemann, T.; Johansson, T.; Sjöman, M.; Korhonen, T.; Guedes, S.; Kuronen, H.; Virtanen, M.J.; Mäkinen, J.; et al. Nationwide outbreak of *Salmonella* Bovismorbificans associated with sprouted alfalfa seeds in Finland, 2009. *Zoonoses Public Health* **2011**, *58*, 589–596.
7. Blaylock, M.; Blackwell, R.; Merid, S.; Jackson, S.; Kotewicz, M.; Gopinath, G.; Ayers, S.L.; Abbott, J.; Sabo, J.; Ewing, L.; et al. Comparison of *Salmonella* enterica serovar Bovismorbificans 2011 hummus outbreak strains with non-outbreak strains. *Food Microbiol.* **2015**, *46*, 627–634.
8. Knoblauch, A.M.; Bratschi, M.W.; Zuske, M.K.; Althaus, D.; Stephan, R.; Hächler, H.; Baumgartner, A.; Prager, R.; Rabsch, W.; Altpeter, E.; et al. Cross-border outbreak of *Salmonella* enterica ssp. enterica serovar Bovismorbificans: Multiple approaches for an outbreak investigation in Germany and Switzerland. *Swiss Med. Wkly.* **2015**, *145*, w14182.
9. Bronowski, C.; Fookes, M.C.; Gilderthorp, R.; Ashelford, K.E.; Harris, S.R.; Phiri, A.; Hall, N.; Gordon, M.A.; Wain, J.; Hart, C.A.; et al. Genomic characterisation of invasive non-typhoidal *Salmonella* enterica Subspecies enterica Serovar Bovismorbificans isolates from Malawi. *PLoS Negl. Trop. Dis.* **2013**, *7*, e2557.
10. Brandwagt, D.; van den Wijngaard, C.; Tulen, A.D.; Mulder, A.C.; Hofhuis, A.; Jacobs, R.; Heck, M.; Verbruggen, A.; van den Kerkhof, H.; Slegers-Fitz-James, I.; et al. Outbreak of *Salmonella* Bovismorbificans associated with the consumption of uncooked ham products, the Netherlands, 2016 to 2017. *Eurosurveillance* **2018**, *23*, 17-00335.

11. Gopinath, G.; Jean-Gilles Beaubrun, J.; Grim, C.; Hanes, D. Draft Genome Sequences of Nine *Salmonella enterica* Serovar Bovismorbificans Isolates from Various Sources. *Genome Announc.* **2014**, *2*, e01249-13.
12. CDC. Multistate Outbreak of *Salmonella* serotype Bovismorbificans Infections Associated with Hummus and Tahini d United States. 2011. Available online: [www.cdc.gov/mmwr/preview/mmwrhtml/mm6146a3.htm](http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6146a3.htm) (accessed on 6.4.2022).
13. Centers for Disease Control and Prevention (CDC). Multistate outbreak of *Salmonella* serotype Bovismorbificans infections associated with hummus and tahini--United States, 2011. *MMWR Morb. Mortal. Wkly. Rep.* **2012**, *61*, 944–947.
14. Gopinath, G.; Jean-Gilles Beaubrun, J.; Grim, C.; Blaylock, M.; Blackwell, R.; Merid, S.; Diallo, A.; Hanes, D. Whole-Genome Sequences of Six *Salmonella enterica* Serovar Bovismorbificans Isolates Associated with a 2011 Multistate Hummus-Borne Outbreak. *Genome Announc.* **2014**, *2*, e01239-13.
15. Nógrády, N.; Imre, A.; Kostyák, A.; Tóth, A.; Nagy, B. Molecular and pathogenic characterization of *Salmonella enterica* serovar Bovismorbificans strains of animal, environmental, food, and human origin in Hungary. *Foodborne Pathog. Dis.* **2010**, *7*, 507–513. <https://doi.org/10.1089/fpd.2009.0420>.
16. Cheng, R.A.; Eade, C.R.; Wiedmann, M. Embracing Diversity: Differences in Virulence Mechanisms, Disease Severity, and Host Adaptations Contribute to the Success of Nontyphoidal *Salmonella* as a Foodborne Pathogen. *Front. Microbiol.* **2019**, *10*, 1368.
17. Achtman, M.; Wain, J.; Weill, F.X.; Nair, S.; Zhou, Z.; Sangal, V.; Krauland, M.G.; Hale, J.L.; Harbottle, H.; Uesbeck, A.; et al. Multilocus sequence typing as a replacement for serotyping in *Salmonella enterica*. *PLoS Pathog.* **2012**, *8*, e1002776. Erratum in: *PLoS Pathog.* **2020**, *16*, e1009040.
18. Den Bakker, H.C.; Moreno Switt, A.I.; Govoni, G.; Cummings, C.A.; Ranieri, M.L.; Degoricija, L.; Hoelzer, K.; Rodriguez-Rivera, L.D.; Brown, S.; Bolchacova, E.; et al. Genome sequencing reveals diversification of virulence factor content and possible host adaptation in distinct subpopulations of *Salmonella enterica*. *BMC Genom.* **2011**, *12*, 425.
19. Timme, R.E.; Pettengill, J.B.; Allard, M.W.; Strain, E.; Barrangou, R.; Wehnes, C.; VanKessel, J.S.; Karns, J.S.; Musser, S.M.; Brown, E.W. Phylogenetic diversity of the enteric pathogen *Salmonella enterica* subsp. *enterica* inferred from genome-wide reference-free SNP characters. *Genome Biol. Evol.* **2013**, *5*, 2109–2123.
20. Cao, G.; Meng, J.; Strain, E.; Stones, R.; Pettengill, J.; Zhao, S.; McDermott, P.; Brown, E.; Allard, M. Phylogenetics and differentiation of *Salmonella* Newport lineages by whole genome sequencing. *PLoS ONE*. **2013**, *8*, e55687.
21. Mastrorilli, E.; Petrin, S.; Orsini, M.; Longo, A.; Cozza, D.; Luzzi, I.; Ricci, A.; Barco, L.; Losasso, C. Comparative genomic analysis reveals high intra-serovar plasticity within *Salmonella* Napoli isolated in 2005–2017. *BMC Genom.* **2020**, *21*, 202.
22. Shariat, N.W.; Timme, R.E.; Walters, A.T. Phylogeny of *Salmonella enterica* subspecies *arizonae* by whole-genome sequencing reveals high incidence of polyphyly and low phase 1 H antigen variability. *Microb. Genom.* **2021**, *7*, 000522.
23. Grimont, P.A.D.; Weill, F.X. *Antigenic Formulae of the Salmonella Serovars*, 9th ed.; 2007; World Health Organization Collaborating Centre for Reference and Research on *Salmonella*, Institut Pasteur: Geneva, Switzerland.
24. Jean-Gilles Beaubrun, J.; Cheng, C.; Chen, K.; Ewing, L.; Wang, H.; Agpaoa, M.C.; Huang, M.J.; Dickey, E.; Du, J.M.; Williams-Hill, D.M.; et al. The use of a PCR-based method for identification of *Salmonella enterica* serotypes Newport, Typhimurium, Javiana and Saint Paul from various food matrices. *Food Microbiol.* **2012**, *31*, 199e209.
25. Zhang, S.; Yin, Y.; Jones, M.B.; Zhang, Z.; Deatherage Kaiser, B.L.; Dinsmore, B.A.; Fitzgerald, C.; Fields, P.I.; Deng, X. *Salmonella* serotype determination utilizing high-throughput genome sequencing data. *J. Clin. Microbiol.* **2015**, *53*, 1685–1692.
26. Gangiredla, J.; Rand, H.; Benisatto, D.; Payne, J.; Strittmatter, C.; Sanders, J.; Wolfgang, W.J.; Libuit, K.; Herrick, J.B.; Prarat, M.; et al. GalaxyTrakr: A distributed analysis tool for public health whole genome sequence data accessible to non-bioinformaticians. *BMC Genom.* **2021**, *22*, 114.
27. Aziz, R.K.; Bartels, D.; Best, A.A.; DeJongh, M.; Disz, T.; Edwards, R.A.; Formsma, K.; Gerdes, S.; Glass, E.M.; Kubal, M.; et al. The RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genom.* **2008**, *9*, 75.
28. Haft, D.H.; DiCuccio, M.; Badretdin, A.; Brover, V.; Chetvernin, V.; O'Neill, K.; Li, W.; Chitsaz, F.; Derbyshire, M.K.; Gonzales, N.R.; et al. RefSeq: An update on prokaryotic genome annotation and curation. *Nucleic Acids Res.* **2018**, *46*, D851–D860. <https://doi.org/10.1093/nar/gkx1068>. PubMed PMID: 29112715.
29. Allard, M.W.; Strain, E.; Melka, D.; Bunning, K.; Musser, S.M.; Brown, E.W.; Timme, R. Practical Value of Food Pathogen Traceability through Building a Whole-Genome Sequencing Network and Database. *J. Clin. Microbiol.* **2016**, *54*, 1975–1983.
30. Tamura, K.; Nei, M.; Kumar, S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 11030–11035.
31. Saitou, M.; Nei, N. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **1987**, *4*, 406–425.
32. Kumar, S.; Stecher, G.; Tamura, K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* **2016**, *33*, 1870–1874. <https://doi.org/10.1093/molbev/msw054>.
33. Huson, D.H.; Bryant, D. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* **2006**, *23*, 254–267. <https://doi.org/10.1093/molbev/msj030>. Epub 2005 Oct 12. PMID: 16221896.
34. Feldgarden, M.; Brover, V.; Haft, D.H.; Prasad, A.B.; Slotta, D.J.; Tolstoy, I.; Tyson, G.H.; Zhao, S.; Hsu, C.H.; McDermott, P.F.; et al. Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates. *Antimicrob. Agents Chemother.* **2019**, *63*, e00483-19.
34. Jackson, S.A.; Patel, I.R.; Barnaba, T.; LeClerc, J.E.; Cebula, T.A. Investigating the global diversity of *Escherichia coli* using a multi-genome DNA microarray platform with novel gene prediction strategies. *BMC Genom.* **2011**, *12*, 349. <https://doi.org/10.1186/1471-2164-12-349>.

35. Li, B.; Jackson, S.; Gangiredla, J.; Wang, W.; Liua, H.; Tall, B.D.; Jean-Gilles Beaubrun, J.; Jay-Russell, M.; Vellidis, G.; Elkins, C.A. Genomic Evidence Reveals Numerous Re-Introduction Events of *Salmonella enterica* serovar Newport in Irrigation Ponds in Suwannee Watershed. *Appl. Environ. Microbiol.* **2015**, *81*, 8243–8253.
36. Affymetrix. Expression Analysis Technical Manual, with Specific Protocols for Use with the Hybridization, Wash, and Stain Kit. 2014. Available online: [http://www.affymetrix.com/support/downloads/manuals/expression\\_analysis\\_technical\\_manual.pdf](http://www.affymetrix.com/support/downloads/manuals/expression_analysis_technical_manual.pdf) (accessed on 6 April 2022).
37. Bolstad, B.M.; Irizarry, R.A.; Astrand, M.; Speed, T.P. A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* **2003**, *19*, 185–193. <https://doi.org/10.1093/bioinformatics/19.2.185>.
38. Achtman, M.; Hale, J.; Murphy, R.A.; Boyd, E.F.; Porwollik, S. Population structures in the SARA and SARB reference collections of *Salmonella enterica* according to MLST, MLEE and microarray hybridization. *Infect. Genet. Evol.* **2013**, *16*, 314–325.
39. Feldgarden, M.; Brover, V.; Haft, D.H.; Prasad, A.B.; Slotta, D.J.; Tolstoy, I.; Tyson, G.H.; Zhao, S.; Hsu, C.H.; McDermott, P.F.; et al. Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates. *Antimicrob. Agents Chemother.* **2019**, *63*, e00483-19.
40. Alikhan, N.F.; Petty, N.K.; Ben Zakour, N.L.; Beatson, S.A. BLAST Ring Image Generator (BRIG): Simple prokaryote genome comparisons. *BMC Genom.* **2011**, *12*, 402.
41. Carattoli, A.; Zankari, E.; García-Fernández, A.; Voldby Larsen, M.; Lund, O.; Villa, L.; Møller Aarestrup, F.; Hasman, H. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* **2014**, *58*, 3895–3903.
42. Arndt, D.; Grant, J.; Marcu, A.; Sajed, T.; Pon, A.; Liang, Y.; Wishart, D.S. PHASTER: A better, faster version of the PHAST phage search tool. *Nucleic Acids Res.* **2016**, *44*, W16–W21.
43. Overbeek, R.; Olson, R.; Pusch, G.D.; Olsen, G.J.; Davis, J.J.; Disz, T.; Edwards, R.A.; Gerdes, S.; Parrello, B.; Shukla, M.; et al. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* **2014**, *42*, D206–D214.
44. Maiden, M.C.; Jansen van Rensburg, M.J.; Bray, J.E.; Earle, S.G.; Ford, S.A.; Jolley, K.A.; McCarthy, N.D. MLST revisited: The gene-by-gene approach to bacterial genomics. *Nat. Rev. Microbiol.* **2013**, *11*, 728–736.
45. Achtman, M.; Zhou, Z.; Alikhan, N.F.; Tyne, W.; Parkhill, J.; Cormican, M.; Chiou, C.S.; Torpdahl, M.; Litrup, E.; Prendergast, D.M.; et al. Genomic diversity of *Salmonella enterica*—The UoWUCC 10K genomes project. *Wellcome Open Res.* **2020**, *5*, 22.
46. Worley, J.; Meng, J.; Allard, M.W.; Brown, E.W.; Timme, R.E. *Salmonella enterica* Phylogeny Based on Whole-Genome Sequencing Reveals Two New Clades and Novel Patterns of Horizontally Acquired Genetic Elements. 2018. mBio. 9: Pii: e02303-18.
47. Deng, X.; den Bakker, H.C.; Hendriksen, R.S. Genomic Epidemiology: Whole-Genome-Sequencing-Powered Surveillance and Outbreak Investigation of Foodborne Bacterial Pathogens. *Annu Rev Food Sci Technol.* **2016**, *7*, 353–374.
48. Yoshida, C.E.; Kruczkiewicz, P.; Laing, C.R.; Lingohr, E.J.; Gannon, V.P.; Nash, J.H.; Taboada, E.N. The *Salmonella* In Silico Typing Resource (SISTR): An Open Web-Accessible Tool for Rapidly Typing and Subtyping Draft *Salmonella* Genome Assemblies. *PLoS ONE*. **2016**, *11*, e0147101.
49. Pearce, M.E.; Alikhan, N.F.; Dallman, T.J.; Zhou, Z.; Grant, K.; Maiden, M.C.J. Comparative analysis of core genome MLST and SNP typing within a European *Salmonella* serovar Enteritidis outbreak. *Int. J. Food Microbiol.* **2018**, *274*, 1–11.
50. Alikhan, N.F.; Zhou, Z.; Sergeant, M.J.; Achtman, M. A genomic overview of the population structure of *Salmonella*. *PLoS Genet.* **2018**, *14*, e1007261.
51. Timme, R.E.; Strain, E.; Baugher, J.D.; Davis, S.; Gonzalez-Escalona, N.; Sanchez Leon, M.; Allard, M.W.; Brown, E.W.; Tallent, S.; Rand, H. Phylogenomic Pipeline Validation for Foodborne Pathogen Disease Surveillance. *J. Clin. Microbiol.* **2019**, *57*, e01816-18.
52. Barretto, C.; Rincón, C.; Portmann, A.C.; Ngom-Bru, C. Whole Genome Sequencing Applied to Pathogen Source Tracking in Food Industry: Key Considerations for Robust Bioinformatics Data Analysis and Reliable Results Interpretation. *Genes* **2021**, *12*, 275.
53. Brown, E.; Dessai, U.; McGarry, S.; Gerner-Smidt, P. Use of Whole-Genome Sequencing for Food Safety and Public Health in the United States. *Foodborne Pathog Dis.* **2019**, *16*, 441–450.
54. Zankari, E.; Hasman, H.; Kaas, R.S.; Seyfarth, A.M.; Agersø, Y.; Lund, O.; Larsen, M.V.; Aarestrup, F.M. Genotyping using whole-genome sequencing is a realistic alternative to surveillance based on phenotypic antimicrobial susceptibility testing. *Antimicrob. Chemother.* **2013**, *68*, 771–777.
55. Branchu, P.; Bawn, M.; Kingsley, R.A. Genome Variation and Molecular Epidemiology of *Salmonella enterica* Serovar Typhimurium Pathovariants. *Infect. Immun.* **2018**, *86*, e00079-18.
56. Cavaco, L.M.; Hasman, H.; Xia, S.; Aarestrup, F.M. *qnrD*, a novel gene conferring transferable quinolone resistance in *Salmonella enterica* Serovar Kentucky and Bovismorbificans strains of human origin. *Antimicrob. Agents Chemother.* **2009**, *53*, 603–608.
57. Finley, R.; Reid-Smith, R.; Ribble, C.; Popa, M.; Vandermeer, M.; Aramini, J. The occurrence and anti-microbial susceptibility of *Salmonellae* isolated from commercially available pig ear pet treats. *Zoonoses Public Health*. **2008**, *55*, 455–461.
58. Jajere, S.M. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and antimicrobial resistance including multidrug resistance. *Vet. World* **2019**, *12*, 504–521.
59. Hanna, L.F.; Matthews, T.D.; Dinsdale, E.A.; Hasty, D.; Edwards, R.A. Characterization of the ELPhiS prophage from *Salmonella enterica* serovar Enteritidis strain LK5. *Appl. Environ. Microbiol.* **2012**, *78*, 1785–1793.