



Article

Antibiotic Susceptibility and Molecular Typing of Invasive *Haemophilus influenzae* Isolates, with Emergence of Ciprofloxacin Resistance, 2017–2021, Italy

Maria Giufrè ^{1,*}, Rita Cardines ¹, Manuela Marra ², Maria Carollo ², Marina Cerquetti ¹ and Paola Stefanelli ¹

¹ Department of Infectious Diseases, Istituto Superiore di Sanità, 00161 Rome, Italy

² Core Facilities Technical-Scientific Service (FAST), Istituto Superiore di Sanità, 00161 Rome, Italy

* Correspondence: maria.giufre@iss.it

Table S1. List of Sequence type by serotype, grouped by Clonal complex.

NTHi			Capsulated		
CC (n)	ST	n	CC (n)	ST	n
3 (20)	14	6	Hia		
	102	2	—	4	1
	136	2	23 (3)	23	2
	143	1		2053	1
	180	1	Hib		
	1034	4	6 (48)	6	38
	1524	3		78	1
	2031	1		95	1
11 (18)	103	15		101	1
	142	1		190	1
	145	1		227	3
	644	1		641	2
12 (6)	12	5		662	1
	786	1	222 (1)	116	1
34 (1)	34	2	Hie		
41 (1)	41	1	18 (3)	69	3
57 (1)	57	2	Hif		
107 (4)	107	2	105 (1)	105	1
	159	1	124 (30)	124	26
	1218	1		598	4
155 (1)	155	1			
165 (1)	165	1			
210 (1)	2129	1			
390 (1)	834	1			
393 (2)	1850	2			
395 (4)	388	4			
396 (1)	396	1			
422 (2)	422	1			
	1667	1			
425 (1)	425	1			
487 (3)	160	3			
513 (1)	513	1			
746 (1)	716	1			
836 (5)	836	5			

1025 (1)	1025	1	
–	184	1	
–	760	1	
–	919	1	
–	943	1	
–	946	1	
–	947	1	
–	949	1	
–	1390	1	
–	1412	1	
–	2027	1	
Total		88	87

CC, clonal complex; ST, sequence type.