microorganisms

MDPI

Article

Biocontrol Potential of Three Bacterial Strains, Isolated from Vermicompost, Against Pre-Emergence Damping-Off of Tomato Caused by *Rhizoctonia solani,* Under Laboratory and Greenhouse Conditions

Mohamed Baha Saeed [1,2,3,4,*], Abdulaziz M. Alnasser [2], Nasser I. Alaruk [2], Abdulrahmn A. Algrwai [2], Salman A. Aloudah [1,2], Sultan M. Al-Eid [2] and Mark D. Laing [4]

[1] Saudi Organic Farming Association, Riyadh 12482, Saudi Arabia
[2] National Organic Agriculture Center, Ministry of Environment, Water and Agriculture, Unaizah 56467, Saudi Arabia
[3] Department of Crop Protection, Faculty of Agriculture, University of Khartoum, Khartoum North 13314, Sudan
[4] Discipline of Plant Pathology, School of Agricultural, Earth and Environmental Science, University of KwaZulu-Natal, Durban 4041, South Africa
* Correspondence: mohamedbaha75@gmail.com; Tel.: +966-57-029-8915

Abstract

Synthetic chemicals, such as fertilizers and pesticides, are widely used in agriculture to improve soil fertility and to control weeds, pests and diseases. Numerous studies have highlighted the negative effects of these chemicals on the soil environment. In contrast, during vermicomposting, earthworms generate numerous beneficial outcomes. This study aimed to screen antagonistic bacteria found after vermicomposting for their potential to inhibit the pre-emergence damping-off of tomato seedlings caused by *Rhizoctonia solani*. Using a dual culture method, 85 bacterial isolates were screened, three of which demonstrated antagonistic activity against *R. solani*. Molecular characterization based on 16S ribosomal RNA identified the bacterial isolates as *Bacillus subtilis* (NOAC.B77), *Bacillus vallismortis* (NOAC.B42), and *Bacillus cereus* (NOAC.B17). The strains NOAC.B77 and NOAC.B42 exhibited the most significant inhibitory effects on the mycelial growth of *R. solani*, with inhibition levels of 80.8% and 79.2%, respectively. In greenhouse trials, only 13% of the Inoculated, Unprotected Control tomato seedlings emerged, i.e., the *R. solani* inoculum caused an 87% level of preemergence damping off. In contrast, after treatment with the bacterial strains NOAC.B77 and NOAC.B42, tomato seedling emergence was not significantly different from the Uninoculated Control. These results suggest that the bacterial strains NOAC.B77 and NOAC.B42 could be commercialized as biological agents to control damping-off of tomato seedlings under greenhouse conditions.

Keywords: *Rhizoctonia solani*; *Bacillus* spp.; tomato; seedling damping-off; biocontrol

Check for updates

Academic Editor: Nazrul Islam

Received: 17 December 2025
Revised: 26 January 2026
Accepted: 30 January 2026
Published: 4 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Tomato (*Solanum lycopersicum* L.) belongs to the Solanaceae family and is one of the most widely grown and consumed horticultural crops worldwide [1]. Global tomato production exceeds 5 million hectares annually, making it a key horticultural crop [2]. The tomato is a nutritious and valuable food containing various bioactive compounds,

including carotenoids (notably lycopene and ß-carotene), ascorbic acid (vitamin C), tocopherols (vitamin E), fibre, vitamin A, and polyphenolic compounds (such as phenols, flavonoids, and phenolic acids) [1]. In addition to its nutritional value, the tomato is an economically important crop, and a high level of productivity is essential for the economic viability of tomato farms. However, tomato crops are affected by numerous fungal pathogens that adversely affect their production globally, leading to significant economic losses [3]. These fungal pathogens include *Rhizoctonia solani* (Kuhn), *Fusarium* spp., and *Pythium* spp., all of which cause seed rot, pre-emergence damping-off, and stem rot in greenhouses and field crops [4].

Rhizoctonia solani is an aggressive, necrotrophic plant pathogen, with a wide host range, including solanaceous crops, cereals, vegetables, cotton, cucurbits, ornamental plants and forest trees [5]. Losses caused by this pathogen range from 20% to 40% worldwide [6]. The fungus produces highly resistant sclerotia, enabling *R. solani* to survive under harsh environmental conditions. *R. solani* is currently managed using fungicides, crop rotation, biological control agents, and resistant cultivars [7–9]. Biological control is considered safe, environmentally friendly, and effective [10]. The biological control of *R. solani* on various crops has been extensively studied [11–13]. Due to concerns related to chemical control and its sustainability, alternative approaches such as biological control are gaining attention.

Vermicompost is a product of the decomposition and stabilization of organic matter in soil, manure, or compost by earthworms such as *Eisenia fetida* (Savigny), *Eudrilus eugeniae* (Kinberg), and *Perionyx excavatus* (Perrier) [14]. Treating livestock and poultry manure with earthworms can result in effective bioresource utilization, reduced environmental protection costs, and may result in a significant decrease in the levels of pathogenic microorganisms in manure. Vermicompost is valuable because it usually contains microorganisms capable of controlling soil-borne pathogens [15–17]. In addition, microbes present in vermicompost can enhance crop production, and contribute to the stability of agricultural soils. Various bacterial genera have been reported in vermicompost, depending on environmental conditions and the raw materials used in the vermicomposting process [15,18,19]. The disease-suppressive potential of vermicompost is mainly attributed to these microbial communities.

Biocontrol strains of *Bacillus* spp. establish protective biofilm layers on roots or soil particles and then produce a range of antimicrobial secondary metabolites that inhibit plant pathogens. These include antimicrobial peptides (AMP), volatile compounds, and hydrolytic enzymes (such as chitinases, glucanases, and proteases). They may also suppress the pathogens through competition for nutrients and space [20–22]. However, information on antagonistic bacteria associated with vermicompost remains limited. This is the first study in Saudi Arabia to screen vermicompost for antagonistic bacteria with inhibitory effects on *R. solani*, to identify these bacteria, and to evaluate their potential to control *R. solani* infections of tomato plants, under laboratory and greenhouse conditions.

2. Materials and Methods

2.1. Isolation and Identification of Rhizoctonia solani Causing Root Rot

Tomato seedlings that exhibited damping-off symptoms were collected and processed for pathogen isolation. The seedlings were initially rinsed with tap water to remove soil debris from the roots. Small sections from infected tissues, usually from the stem near the soil surface, were excised. Plant tissues were surface disinfected using two successive treatments, each lasting one minute: first with a 1% sodium hypochlorite (NaOCl; Sigma-Aldrich, St. Louis, MO, USA) solution, followed by sterile distilled water. The disinfected samples were then placed on potato dextrose agar (PDA; Neogen, Ltd., Lansing, MI, USA) medium in Petri dishes (90 mm) (ISOLAB Laborgeräte GmbH, Eschen-

strasse 5, 22844 Norderstedt, Germany) after excess moisture was removed using sterile filter paper ((Whatman No. 1); Cytiva, Marlborough, MA, USA). The plates were then incubated at 25 ± 1 °C for 2 days. Mycelia growing from the plant tissues were transferred onto PDA slants using a sterile inoculation needle (HiMedia, Mumbai, India) and incubated at 25 ± 1 °C for one week [9,23]. Identification of the isolate of *R. solani* was based on its characteristic mycelial morphology [24]. Pathogenicity was confirmed by inoculation of the fungus onto 3-week-old tomato seedlings (cv. Nurdan F1). Ten seedlings were grown per pot (28 cm diameter), and each pot was filled with a peat-based artificial growing medium. Inoculation was with a 5 mm plug of the isolate taken from the margin of a 15-day-old culture grown on PDA. The causal agent of damping-off of these seedlings was reisolated, as per the initial isolation technique. The mycelium of the freshly isolated fungal colonies was examined microscopically to confirm the identity of the pathogen as *R. solani* [24].

2.2. Vermicompost Sample Collection

Fresh vermicompost samples were collected from the vermicomposting unit at the National Organic Agriculture Center (NOAC), Qassim, Saudi Arabia (25°48′23″ N 42°52′24″ E). Nine samples, each weighing 0.5 kg, were collected from nine different locations within the unit using an auger. The samples were sieved to remove insects and other debris and stored at 4 °C until they were used.

2.3. Isolation of Antagonistic Bacteria from Vermicompost, and Screening for Inhibitory Activity Against Rhizoctonia solani

A 10 g sample of vermicompost from each source was weighed and added to a flask (SIMAX, Kavalierglass a.s., Sázava, Czech Republic) containing 90 mL of sterile distilled water, then shaken for 3 min to obtain a vermicompost suspension stock. For a serial dilution series, 1 mL of the suspension stock was transferred into 9 mL of sterile distilled water. Dilutions of 10^{-3}, 10^{-4}, and 10^{-5} were prepared from the original suspension. From each dilution sample, 0.1 mL was spread onto nutrient agar (NA) medium (Neogen, Ltd., Lansing, MI, USA) in 90 mm plastic Petri dishes. After incubation at 30 °C for 48 h, single colonies were selected and a single pure culture of each isolate was established [25].

Bacterial isolates were screened for their antifungal activity against *R. solani* using an in vitro dual culture assay of the putative antagonists, and the pathogen, which was grown on PDA (Neogen, Ltd.). Each bacterial isolate was inoculated at four evenly distributed points, 2.5 cm from the edge of a Petri dish (90 mm in diameter), with three replicates per isolate. After inoculation for 48 h at 28 °C in the dark, a 5 mm plug taken from the margin of a 15-day-old *R. solani* culture grown on PDA was inoculated in the centre of each plate. Plates inoculated with the fungus but not the bacterial isolates served as controls. The plates were incubated at 28 °C for five days, after which the diameter of the fungal colony was measured. The level of inhibition of the colony diameter was calculated using the following formula.

$$\text{Inhibition rate (\%)} = [(C - T)/C] \times 100\%$$

where C is the colony diameter in the control plate, and T is the colony diameter in the treatment plate.

2.4. In Vitro Antagonistic Activity Against R. solani

To evaluate the effects of various incubation time, the three most effective bacterial strains (NOAC.B77, NOAC.B42, and NOAC.B17) were assessed for their antagonistic activity against *R. solani*. Each strain was inoculated onto NA plates for 0, 1, 2, and 3 days, as described in Section 2.3. Subsequently, a 5 mm plug of *R. solani* grown on PDA was inoculated in the centre of each plate and incubated at 28 °C. At 0, 1, 2, and 3 days after the

inoculation of each plate with *R. solani*, the level of inhibition of the colony diameter was calculated, as described in Section 2.3.

2.5. Morphological Changes in R. solani Mycelia Affected by Antagonistic Bacterial Strains

A thin layer of nutrient agar (NA) was poured into Petri dishes to facilitate microscopic observation. Three Petri dishes (90 mm in diameter) were prepared for each bacterial strain. Bacterial incubation was performed as described in Section 2.3. After three days of incubation, a 5 mm plug of *R. solani* from a 15-day-old culture was placed onto each plate. Control plates contained *R. solani* only. All plates were examined under a light microscope (Euromex, Duiven, The Netherlands) to look for any morphological changes in fungal mycelial growth caused by bacterial antagonism, in comparison with the normal growth of fungal mycelium observed in the control treatment. Control plates were grown with only the fungal isolate, to confirm that the observed effects on fungal growth were due to bacterial activity.

2.6. Plant Growth Promotion by Bacterial Strains

The potential plant growth-promoting activity of the most effective biocontrol strains (NOAC.B17, NOAC.B42 and NOAC.B77) was evaluated using a paper towel seedling technique. Un-treated tomato seeds (cv. Nurdan F1 Argeto Vegetable Seeds, Antalya, Turkey) were soaked in a bacterial suspension (1×10^8 CFU ml^{-1}) for 3 h at 25 °C. Three replicates were used for each strain, with ten seeds per replicate. Treated seeds were placed between two moistened germination papers, rolled, and incubated at 25 °C. Sterile distilled water was used to treat seeds in the control treatment. After 15 days of incubation, germination percentage, root length, and shoot length were recorded [26]. The seedling vigour index was calculated as follows: Vigour index = (mean root length + mean shoot length in cm) × percentage germination [27].

2.7. Root Colonization

To assess the colonization ability of the most effective bacterial strains, tomato seeds (cv. Nurdan F1) were surface-sterilized with 1% sodium hypochlorite for 1 min, followed with two rinses in sterile distilled water. The seeds were immersed in a bacterial suspension and shaken for 20 min. The coated seeds were placed on sterile filter paper in Petri dishes and incubated at 25 °C. After germination and radical emergence, when the primary root of each seedling reached approximately 50 mm in length, segments from the elongation zone were excised and transferred to NA medium. The Petri dishes were incubated at 29 °C for 3–5 days and examined daily for bacterial growth emerging from the surface-sterilized root segments.

2.8. Identification of the Selected Bacterial Isolates

2.8.1. DNA Extraction and PCR Conditions

DNA of the bacteria was extracted according to Cook and Myers [28], with minor modifications. The three isolates were each cultured in nutrient broth (Oxoid, Ltd., Cheshire, UK) at 30 °C for 24 h. Cells were harvested by centrifugation at 12,000 rpm for 3 min, then washed with Tris-EDTA (TE) buffer (pH 7.7). (Sigma-Aldrich, St. Louis, MO, USA) The cell pellets were re-suspended in TE buffer (300 µL), heated at 95 °C for 10 min in a hot water bath, cooled to room temperature, and centrifuged at 12,000 rpm for 8 min. The supernatant samples containing genomic DNA were transferred into sterile tubes and stored at 4 °C until PCR amplification. PCR amplification of the 16S rDNA region was performed in 30 µL reaction volumes using DreamTaq™ Green PCR Master Mix (Thermo Scientific, Waltham, MA, USA) and the universal primers 27F (5′-AGAGTTTGATCATGGCTCAG-3′) and 1492R (5′-TACGGTTACCTTGTTACGACTT-3′). Amplification products were verified

by electrophoresis on a 1% agarose gel, (Sigma-Aldrich, St. Louis, MO, USA) T purified, and sequenced in both directions (Macrogen, Seoul, Republic of Korea).

2.8.2. Phylogenetic Analysis

The Maximum Likelihood method was used to infer the phylogenetic relationships using the following parameters: the Maximum Composite Likelihood (MCL) approach, the Tamura–Nei substitution model, and the Neighbour-Joining algorithm. Analyses were performed using MEGA version 11 software (MEGA Software, Tempe, AZ, USA) [29]. Type strains are indicated in the phylogenetic tree by the letter T. The obtained sequences were deposited in the GenBank (NCBI) database under accession numbers PP961956, PP961957 and KY034383.

2.9. Greenhouse Experiment

2.9.1. Preparation of Fungal Inoculum and Bacterial Isolates

The three selected bacterial isolates were cultured in nutrient broth at 30 °C on a rotary shaker (JS Research Inc. (JSR), Gongju-City, Republic of Korea) for 48 h. A bacterial suspension containing 1×10^8 CFU ml^{-1} was prepared for experimental use. To prepare *R. solani* inoculum, 500 g of barley grains (local supplier, Riyadh, Saudi Arabia) were washed, soaked overnight, and autoclaved at 121 °C for 15 min, followed by a 24 h cooling period at 22–25 °C. Plugs of *R. solani* were cultured on PDA using the previously isolated strain (Section 2.5). Five agar plugs (5 mm × 5 mm) from a 15-day-old culture were added to sterilized barley grains. The inoculated grains were incubated at 25 °C for 15 days until fully colonized by the fungus.

2.9.2. Pot Experiment Design

A greenhouse experiment was conducted to evaluate the effect of the selected bacterial isolates on the seedling emergence of tomato seedlings inoculated with *R. solani*. Tomato seeds (cv. Nurdan F1) were grown in pots that were 28 cm in diameter, which were filled with a peat-based artificial growing medium. The experiment followed a randomized complete block design (RCBD) with three replicates and eight treatments: (1) *B. subtilis* (NOAC.B77), (2) *B. cereus* (NOAC.B17) (3) *B. vallismortis* (NOAC.B42), (4) *B. subtilis* (NOAC.B77) + *R. solani*, (5) *B. cereus* (NOAC.B17) + *R. solani*, (6) *B. vallismortis* (NOAC.B42) + *R. solani* (7) *R. solani* (inoculated, untreated control), and (8) tomato seeds only (uninoculated, untreated control).

2.9.3. Inoculation Technique and Data Collection

For Treatments 1–6, 50 mL of the appropriate bacterial suspension was applied to each pot as a soil drench. For Treatments 4–7, 10 g of barley grains colonized with *R. solani* were incorporated into the soil of each pot. Tomato seeds (cv. Nurdan F1) were sown at a rate of 15 seeds per pot, three days after bacterial application. Seedling emergence percentage was calculated once all seeds in the T8 pots (uninoculated, untreated Control) had fully germinated. Approximately five days after sowing, seedling emergence was evaluated. This was done by determining the number of sown seeds that emerged after each treatment, relative to T8, the uninoculated, untreated control. Seedling Emergence was expressed as the proportion of sown seeds that successfully germinated and grew in a healthy state. The effectiveness of bacterial treatments was quantified by calculating.

$$\text{Corrected Seedling Emergence (\%)} = 100 - ([(C\% - T\%)/C\%] \times 100]))$$

where C% is the percentage of seedlings that emerged in the Uninoculated, Untreated Control pot (T8 = maximum potential emergence), and T% is the percentage of seedlings that emerged from each of the treated pots (T1–T7).

2.10. Statistical Analysis

Data were subjected to analysis of variance (ANOVA). Means were compared using the Duncan's multiple range test at a 5% significance level. Statistical analyses were conducted using GenStat for Windows (23rd edition); VSN International, Hemel Hempstead, UK.

3. Results

3.1. Isolation of Antagonistic Bacteria from Vermicompost

Eighty-five bacterial isolates were obtained from the collected vermicompost samples and evaluated for their antagonistic activity against *R. solani*. Among the tested isolates, five showed antagonistic activity in the dual culture assay and three exhibited strong antifungal activity against *R. solani*.

3.2. In Vitro Antagonistic Activity Against R. solani

The selected bacterial isolates significantly reduced the growth of *R. solani* in vitro (Figure 1). Significant differences among the tested isolates were observed at all exposure times following bacterial inoculation, with inhibition percentages ranging from 56% to 65% at day zero and day one after inoculation (Table 1). A notable increase in inhibition was observed by day three, reaching 79.15% and 80.78% for *B. subtilis* (NOAC.B77) and *B. vallismortis* (NOAC.B42), respectively.




Figure 1. In vitro antagonistic activity of three bacterial isolates against *Rhizoctonia solani* in a dual culture assay after three days of bacterial incubation: (**a**) *Bacillus subtilis* (NOAC.B77), (**b**) *Bacillus vallismortis* (NOAC.B42), (**c**) *Bacillus cereus* (NOAC.B17), and (**d**) *R. solani* only, the untreated control.

Table 1. In vitro inhibition of the mycelial growth of *R. solani* in culture following exposure to three *Bacillus* isolates at 0, 1, 2, and 3 days after bacterial inoculation.

Isolate	Inhibition (%) of *R. solani* Time After Bacterial Inoculation (Days)			
	(0) Day	(1) Day	(2) Days	(3) Days
B. subtilis (NOAC.B77)	62.38 b	64.70 b	72.72 b	80.77 b
B. vallismortis (NOAC.B42)	64 b	65.30 b	74.05 b	79.15 b
B. cereus (NOAC.B17)	56.12 a	57.0 a	58.8 a	61.75 a
LSD	4.329	6.39	6.03	4.025
p value	0.015 *	0.034 *	<0.001 ***	<0.001 ***
F Value	9.14	6.28	30.81	82.21
CV%	4.5	5.9	5.5	3.1

Means followed by the same letter do not differ significantly at $p < 0.05$ according to the Duncan's multiple range test. Asterisks indicate significant differences at *** $p < 0.001$, * $p < 0.05$.

These results identified specific bacterial strains that produced inhibitory compounds over time, as evidenced by comparisons with untreated fungal colony measurements taken days zero, one, and three after inoculation (Table 1). A linear response in inhibition of *R. solani* colony growth as a result of increasing bacterial inoculation time was observed for all isolates, with $R^2 > 0.9$. (Figure 2).




Figure 2. Linear regression analysis showing the relationships between percentage inhibition of the mycelial growth of *R. solani*, and increasing time after inoculation, following the application of three bacterial isolates: *Bacillus subtilis* (NOAC.B77), *Bacillus vallismortis* (NOAC.B42) and *Bacillus cereus* (NOAC.B17).

3.3. Morphological Changes in the Mycelium of R. solani Induced by Three Antagonistic Bacterial Isolates

Light microscopic observation of the mycelium of *R. solani* in the inhibition zones of dual culture assays revealed structural changes, presumably caused by antifungal compounds produced by the bacterial isolates (Figure 3). Common abnormalities included hyphal shrivelling, twisting, and distortion when *R. solani* was co-cultivated with the bacterial strains. In contrast, the mycelium of the untreated control cultures exhibiting a smooth surface and normal branched, tubular morphology.




Figure 3. Light micrographs of *Rhizoctonia solani* hyphae in the inhibition zone of the dual culture assay, showing morphological abnormalities: (**a**) *R. solani* mycelium grown in the presence of *Bacillus subtilis* (NOAC.B77), (**b**) *R. solani* mycelium grown in the presence of *Bacillus vallismortis* (NOAC.B42), (**c**) *R. solani* mycelium grown in the presence of *Bacillus cereus* (NOAC.B17), and (**d**) *R. solani* only, the untreated control.

3.4. Plant Growth Promotion and Root Colonization by the Selected Bacterial Strains

No significant differences ($p < 0.05$) were observed in seed germination percentage, shoot length or root length between the control and rhizobacteria-treated tomato seedlings in the rolled paper towel assay (Table 2).

Table 2. Effects of seed treatments with antagonistic bacterial strains on seed germination and seedling vigour of tomato seedlings grown under laboratory conditions.

Strain	Germination (%)	Root Length (cm)	Shoot Length (cm)	Vigour Index
B. subtilis (NOAC. B77)	96.80 a	3.340 a	0.64 a	385.7 a
B. vallismortis (NOAC. B42)	96.80 a	3.540 a	0.80 a	384.2 a
B. cereus (NOAC. B17)	94.20 a	3.680 a	0.40 a	382.5 a
Control	95.20 a	3.600 a	0.18 a	352.2 a

Means followed by the same letter do not differ significantly at $p < 0.05$ according to the Duncan's multiple range test.

3.5. Root Colonization

The development of bacterial colonies following the sterile transfer of excised tomato root segments onto nutrient agar (NA) medium confirmed that the tested isolates successfully colonized the surface of tomato roots (Figure 4). Root fragments were surface

disinfected in sodium hypochlorite prior culturing, a procedure that was expected to eliminate vegetative bacterial cells.




Figure 4. Root colonization of tomato seedlings by three *Bacillus* strains following aseptic transfer of root segments onto nutrient agar medium: (**a**) *Bacillus subtilis* (NOAC.B77), (**b**) *Bacillus vallismortis* (NOAC.B42), (**c**) *Bacillus cereus* (NOAC.B17), and (**d**) untreated control.

3.6. Identification of the Selected Bacterial Isolates

The 16S rRNA gene sequences obtained from the selected isolates were compared with reference sequences in GenBank databases. Phylogenetic analysis revealed the relationships between the isolates and closely related *Bacillus* species (Figure 5). Strain NOAC.B17 (PP961956) showed a 99.75% sequence similarity to *Bacillus cereus* ATCC 14579 (KY034383) from Saudi Arabia. Strain NOAC.B42 (PP961957) displayed a relatedness of (99.65%) to *Bacillus vallismortis* Strain NRRL B-14890 from USA. Strain NOAC.B77 (PP961958) shared of 97.85% similarity with *Bacillus subtilis* strain NBRC 13719 from Japan and *Bacillus subtilis* strain DSM 10 from India.

3.7. Greenhouse Experiment

Significant differences were observed among the treatments when assessing the effects of bacterial strains on differences in % Emergence of tomato seedlings caused by pre-emergence damping that was caused by *R. solani*, under greenhouse conditions (F 111.17; $p < 0.001$; d.f. = 7). No adverse effects were observed when tomato seeds were treated with one of the three bacterial strains alone, and the seedling emergence level percentages were similar to those of the untreated control (>92%). In contrast, the percentage of seedlings that emerged in the *R. solani* infected control was only 13%. The isolates *B. subtilis* strain

NOAC.B77, *B. vallismortis* strain NOAC.B42 and *B. cereus* strain NOAC.B17 increased seedling emergence to 55.0%, 54.3% and 47.7%, respectively (Figure 6), by reducing the level of pre-emergence damping off caused by *R. solani*.




Figure 5. Molecular phylogenetic tree based on the 16S rRNA gene sequences of the three selected bacterial isolates. Type strains are indicated by the letter T. Two *Pseudomonas* species were included as out-group taxa to root the phylogenetic tree. The scale bar represents the number of nucleotide substitutions per site.




Figure 6. Effect of three bacterial isolates: *Bacillus subtilis* strain NOAC.B77, *Bacillus vallismortis* strain NOAC.B42 and *Bacillus cereus* strain NOAC.B17 on Seedling Emergence% under greenhouse conditions. The control inoculated with *Rhizoctonia solani* had an emergence of 13%. Error bars represent standard error (SE). Levels of seedling emergence (%) listed with the same latter are not significantly different at $p = 0.05$.

4. Discussion

Biological control represents an option for the management of soil-borne pathogens of vegetable crops, including those caused by *R. solani*. The advantages they provide include their environmentally friendly, cost-effective, and user-friendly nature. Many commercial biopesticide products are currently available worldwide for large-scale field applications [30]. However, the successful biocontrol of soil-borne pathogens relies on the competitiveness of the microbial biocontrol agents in soils that vary geographically. Therefore, the present study focused on the isolation and screening of indigenous bacterial strains with antifungal activity from vermicompost, which is widely used as a natural fertilizer in organic agriculture in Saudi Arabia.

Among the 85 bacterial isolates screened, three isolates (*Bacillus subtilis* (NOAC.B77), *Bacillus vallismortis* (NOAC.B42), and *Bacillus cereus* (NOAC.B17)) exhibited antagonistic activity against *R. solani* in in vitro dual culture assays, significantly reducing growth of the fungal colonies, without creating obvious inhibition zones or halos. The antagonistic efficacy of these strains increased over time, from approximately 56% on Day Zero to 79.2% and 80.8% by Day Three for *B. subtilis* (NOAC.B77) and *B. vallismortis* (NOAC.B42), respectively. This increase was assumed to be due to the accumulation of antifungal compounds produced by the bacterial strains in the agar medium. Applying these bacterial strains to tomato seeds or seedlings three days prior to planting may enhance their effectiveness against *R. solani* under field conditions by allowing them sufficient time to multiply, colonize, and to release antifungal compounds into the rhizosphere soil. This would lead to a reduction in the inoculum levels of *R. solani*, reducing pre-emergence seed and seedlings infections, and increasing seedling emergence. These findings are consistent with those of Zohora et al. [31], who reported that *B. subtilis* strain RB14 strongly inhibited *R. solani* growth in vitro. Similarly, Rashad et al. [32] found that *B. subtilis* strain SR22 inhibited *R. solani* growth in a dual culture assay by 51%. Variations in the levels of inhibition provided by different bacterial isolates in the present study may be attributed to differences in the quantity and toxicity of the antifungal compounds produced by each bacterial strain [26]. These inhibitory effects may be mediated by the production of enzymes (e.g., chitinases, glucanases, proteases), antibiotics, or other bioactive metabolites, although their precise nature was not determined in this study.

The antagonistic effects of the bacterial isolates were further supported by microscopic examination of *R. solani* hyphae growing within the inhibition zones of dual culture agar plates. Hyphae of *R. solani*, co-cultivated with antagonistic *Bacillus* strains, exhibited morphological abnormalities, including distortion, shrivelling, twisting, and irregular growth patterns. Similar morphological alterations were reported by Lee et al. [33], who observed hyphal damage when *B. velezensis* GH1-13 was co-cultivated with a strain of *R. solani*, a pathogen affecting isolated from cool-season turfgrasses. Irregular hyphal growth was also described by Wu et al. [34] when *B. subtilis* SL-44 was co-cultivated with *R. solani* associated with *Capsicum annuum* (L.). They suggested that antifungal metabolites produced by the tested *Bacillus* strain played a key role in its biocontrol activity.

Bacterial grew out from the surface sterilized root fragments from seedlings treated with the three antagonistic bacterial strains. No bacteria grew out of the roots of the untreated control plants. It is likely that endospores of the three *Bacillus* strains were present on the root tissues, survived the surface sterilization, and subsequently germinated and multiplied under culture conditions. Root colonization is a well-documented characteristic of *Bacillus* species [35], including numerous biocontrol strains [36,37].

In the in vitro plant growth promotion assay, the tested bacterial strains did not significantly enhance or diminish the levels of seed germination or seedling vigour of the tomato seed. This contrasts with the findings of other studies where growth suppression

was observed to follow seed bacterization. For example, Al-Hussini et al. [38] demonstrated that treatments of tomato seeds with *Klebsiella oxytoca* strain D1/3 led to a reduction in both shoot and root lengths, as well as overall seedling vigour, compared with untreated controls, indicating that this bacterium was releasing phytotoxic compounds. In other studies, *Bacillus* isolates enhanced plant growth significantly [39,40].

In the greenhouse pot experiment, *B. subtilis* (NOAC.B77), *B. vallismortis* (NOAC.B42) and *B. cereus* (NOAC.B17) increased tomato seedling emergence by 55.0%, 54.3% and 47.7%, respectively, by reduced damping-off caused by *R. solani*. In contrast, only 13% of tomato seeds emerged when the tomato seeds were exposed to *R. solani* in the absence of bacterial treatment. These findings align with those of Khedher et al. [41], who found that bacterization of potato tubers with *B. subtilis* strain V26 significantly reduced the incidence of black scurf disease caused by *R. solani*, and improved plant growth. These results confirm the antagonistic potential of the tested strains against *R. solani* under greenhouse conditions, without any observable effect on seedling vigour, indicating that their primary role was biocontrol protection of the seedlings rather than plant growth promotion.

Relatively few studies have investigated *B. vallismortis* as a biocontrol agent against *R. solani* on tomato [42]. In the present study, the *B. vallismortis* isolate demonstrated a level of biocontrol efficacy comparable to that of *B. subtilis* under greenhouse conditions. Previous studies have reported the effectiveness of *B. vallismortis* strains against several plant pathogens, including *Fusarium solani* ((Mart.) Sacc.) infecting *Panax ginseng* (Mey), [43], and *Fusarium oxysporum*, *Fusarium moniliforme* (Sheld.), *Fusarium proliferatum* (Matsush.) Nirenberg ex Gerlach & Nirenberg), and *F. solani* on apple (*Malus domestica* (Suckow) Borkh.) [44].

5. Conclusions

Vermicompost is a promising source of antagonistic bacteria with potential applications in the biocontrol of soil-borne plant diseases, such as those caused by *R. solani*. Among the bacterial strains evaluated in this study, *B. subtilis* (NOAC.B77) and *B. vallismortis* (NOAC.B42) were the most effective strains in reducing seedling damping-off of tomato seedlings caused by *R. solani*. These strains show potential for further development as commercial biocontrol agents.

Author Contributions: Conceptualization, M.B.S.; methodology M.B.S. and A.M.A.; software, M.B.S.; validation, N.I.A. and A.A.A.; formal analysis, M.B.S.; investigation, All authors.; resources, S.M.A.-E. and S.A.A.; data curation, M.B.S. and A.M.A.; writing—original draft preparation, M.B.S.; writing—review and editing, M.B.S. and M.D.L.; visualization, M.B.S. and A.A.A.; supervision, M.B.S. and S.M.A.-E.; project administration, S.A.A.; funding acquisition, S.A.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Ministry of Environment, Water, and Agriculture (Saudi Arabia) and the Saudi Organic Farming Association through the Scientific Research Development Project in Organic Farming (Grant No. 23060114771).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in this study are included in the article. Further inquiries can be directed to the corresponding author.

Acknowledgments: The authors acknowledge the Ministry of Environment, Water, and Agriculture (Saudi Arabia), the Organic Farming Administration, and the Saudi Organic Farming Association for overseeing and funding the current work through the Scientific Research Development Project in

Organic Farming (Grant No. 230601147751). The authors also acknowledge the National Center for Sustainable Agriculture Research and Development (Estidama) for overseeing the research project.

Conflicts of Interest: The authors declare that they have no competing interests.

Abbreviations

The following abbreviations are used in this manuscript:

NaOCl	Sodium hypochlorite
PDA	Potato Dextrose Agar
NOAC	National Organic Agriculture Center
NA	Nutrient Agar
PCR	Polymerase Chain Reaction
MCL	Maximum Composite Likelihood
MEGA 11	Molecular Evolutionary Genetics Analysis version 11
NCBI	National Center for Biotechnology Information
CFU	Colony-Forming Unit
RCBD	Randomized Complete Block Design
ANOVA	Analysis of Variance
LSD	Least Significant Difference
CV%	Coefficient of Variation

References

1. Collins, E.J.; Bowyer, C.; Tsouza, A.; Chopra, M. Tomatoes: An extensive review of the associated health impacts of tomatoes and factors that can affect their cultivation. *Biology* **2022**, *11*, 239. [CrossRef]
2. Food and Agriculture Organization of the United Nations (FAOSTAT). *World Food and Agriculture—Statistical Yearbook 2022*; FAO: Rome, Italy, 2022. [CrossRef]
3. Al-Askar, A.A.; Ghoneem, K.M.; Rashad, Y.M.; Abdulkhair, W.M.; Hafez, E.E.; Shabana, Y.M.; Baka, Z.A. Occurrence and distribution of tomato seed-borne mycoflora in Saudi Arabia and its correlation with the climatic variables. *Microb. Biotechnol.* **2014**, *7*, 556–569. [CrossRef]
4. Nefzi, A.; Jabnoun-Khiareddine, H.; Abdallah, R.A.B.; Ammar, N.; Medimagh-Saïdana, S.; Haouala, R.; Daami-Remadi, M. Suppressing *Fusarium* crown and root rot infections and enhancing the growth of tomato plants by *Lycium arabicum* extracts. *S. Afr. J. Bot.* **2017**, *113*, 288. [CrossRef]
5. Ghosh, S.; Kanwar, P.; Jha, G. Alterations in rice chloroplast integrity, photosynthesis and metabolome associated with pathogenesis of *Rhizoctonia solani*. *Sci. Rep.* **2017**, *7*, 41610. [CrossRef]
6. Srivastava, S.; Bist, V.; Srivastava, S.; Singh, P.C.; Trivedi, P.K.; Asif, M.H.; Chauhan, P.S.; Nautiyal, C.S. Unraveling aspects of *Bacillus amyloliquefaciens*-mediated enhanced production of rice under biotic stress of *R. solani*. *Front. Plant Sci.* **2016**, *7*, 587. [CrossRef] [PubMed]
7. Pandey, C.; Prabha, D.; Negi, Y.K.; Maheshwari, D.K.; Dheeman, S.; Gupta, M. Macrolactin A mediated biocontrol of *Fusarium oxysporum* and *R. solani* infestation on *Amaranthus hypochondriacus* by *Bacillus subtilis* BS-58. *Front. Microbiol.* **2023**, *14*, 1105849. [CrossRef]
8. Mogazy, A.M.; Abdallah, W.E.; Mohamed, H.I.; Omran, A.A. The efficacy of chemical inducers and fungicides in controlling tomato root rot disease caused by *R. solani*. *Plant Physiol. Biochem.* **2024**, *210*, 108669. [CrossRef] [PubMed]
9. Rafiq, M.; Shoaib, A.; Javaid, A.; Perveen, S.; Umer, M.; Arif, M.; Cheng, C. Exploration of resistance level against black scurf caused by *R. solani* in different cultivars of potato. *Plant Stress* **2024**, *12*, 100476. [CrossRef]
10. Aydın, M.H. *Rhizoctonia solani* and its biological control. *Turk. J. Agric. Res.* **2022**, *9*, 118–135. [CrossRef]
11. Akber, M.A.; Fang, X. Research progress on diseases caused by the soil-borne fungal pathogen *Rhizoctonia solani* in alfalfa. *Agronomy* **2024**, *14*, 1483. [CrossRef]
12. Intana, W.; Suwannarach, N.; Kumla, J.; Wonglom, P.; Sunpapao, A. Plant growth promotion and biological control against *Rhizoctonia solani* in Thai local rice variety "Chor Khing" using *Trichoderma breve* Z2-03. *J. Fungi* **2024**, *10*, 417. [CrossRef]
13. Wekesa, T.B.; Wekesa, V.W.; Onguso, J.M.; Kavesu, N.; Okanya, P.W. Study of pathogenicity test, antifungal activity, and secondary metabolites of *Bacillus* spp. from Lake Bogoria as biocontrol of *Rhizoctonia solani* Kühn in *Phaseolus vulgaris* L. *Int. J. Microbiol.* **2024**, *2024*, 6620490. [CrossRef]

14. Gebrehana, Z.G.; Gebremikael, M.T.; Beyene, S.; Sleutel, S.; Wesemael, W.M.L.; De Neve, S. Organic residue valorization for Ethiopian agriculture through vermicomposting with native (*Eudrilus eugeniae*) and exotic (*Eisenia fetida* and *Eisenia andrei*) earthworms. *Eur. J. Soil Biol.* **2023**, *116*, 103488. [CrossRef]
15. Soltan, H.; Dakhly, O.; Mahmoud, M.; Fayz, Y.F. Microbiological and genetical identification of some vermicompost beneficial associated bacteria. *SVU-Int. J. Agric. Sci.* **2022**, *4*, 21. [CrossRef]
16. Malakar, C.; Barman, D.; Kalita, M.C.; Deka, S. A biosurfactant-producing novel bacterial strain isolated from vermicompost having multiple plant growth-promoting traits. *J. Basic Microbiol.* **2023**, *63*, 746. [CrossRef]
17. Mediyawa, H.; Munidasa, W. Diversity and ecological significance of vermicomposting bacteria. In *Transformative Applied Research in Computing, Engineering, Science and Technology*; Damayanthi, D., Miruna, R., Eds.; CRC Press: Boca Raton, FL, USA, 2025; p. 141.
18. Kawicha, P.; Laopha, A.; Chamnansing, W.; Sopawed, W.; Wongcharone, A.; Sangdee, A. Biocontrol and plant growth-promoting properties of *Streptomyces* isolated from vermicompost soil. *Indian Phytopathol.* **2020**, *73*, 655–666. [CrossRef]
19. Jiang, L.; Yan, H.; Yin, Y.; Dong, R.; Wu, C. Use of *Bacillus velezensis* JNS-1 from vermicompost against strawberry root rot. *Front. Microbiol.* **2025**, *16*, 1566949. [CrossRef]
20. Sowmiya, K.; Aravindharajan, S.; Bhagyashree, K.; Manoj, M. Biofilm-forming capability of *Bacillus* and its related genera. In *Applications of Bacillus and Bacillus-Derived Genera in Agriculture, Biotechnology and Beyond*; Mageshwaran, V., Singh, U.B., Saxena, A.K., Singh, H.B., Eds.; Springer Nature: Singapore, 2024; p. 71.
21. Zhou, H.; Dong, K.; Du, Q.; Wei, Q.; Wu, J.; Deng, J.; Wang, F. Biofilm-forming of *Bacillus tequilensis* DZY 6715 enhanced suppression of the *Camellia oleifera* anthracnose caused by *Colletotrichum fructicola* and its mechanism. *Sci. Hortic.* **2024**, *338*, 113676. [CrossRef]
22. Liu, T.; Zheng, Y.; Wang, L.; Wang, X.; Wang, H.; Tian, Y. Optimizing surfactin yield in *Bacillus velezensis* BN to enhance biocontrol efficacy and rhizosphere colonization. *Front. Microbiol.* **2025**, *16*, 1551436. [CrossRef] [PubMed]
23. Parajuli, P.; Yadav, R.K.; Manandhar, H.K.; Parajulee, M.N. Management of Root Rot (*Rhizoctonia solani* Kühn) of Common Bean Using Host Resistance and Consortia of Chemicals and Biocontrol Agents. *Biology* **2025**, *14*, 235. [CrossRef]
24. Sharma, M.; Gupta, S.; Sharma, T. Characterization of variability in *Rhizoctonia solani* by using morphological and molecular markers. *J. Phytopathol.* **2005**, *153*, 449. [CrossRef]
25. Wang, X.; Yan, S.; Min, C.; Yang, Y. Isolation and antimicrobial activities of actinomycetes from vermicompost. *China J. Chin. Mater. Med.* **2015**, *40*, 614.
26. Al-Daghari, D.S.S.; Al-Sadi, A.M.; Al-Mahmooli, I.H.; Janke, R.; Velazhahan, R. Biological control efficacy of indigenous antagonistic bacteria isolated from the rhizosphere of cabbage grown in biofumigated soil against *Pythium aphanidermatum* damping-off of cucumber. *Agriculture* **2023**, *13*, 626. [CrossRef]
27. Abdul-Baki, A.A.; Anderson, J.D. Vigor determination in soybean seed by multiple criteria. *Crop Sci.* **1973**, *13*, 630–633. [CrossRef]
28. Cook, A.E.; Meyers, P.R. Rapid identification of filamentous actinomycetes to the genus level using genus-specific 16S rRNA gene restriction fragment patterns. *Int. J. Syst. Evol. Microbiol.* **2003**, *53*, 1907–1915. [CrossRef] [PubMed]
29. Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular evolutionary genetics analysis version 11. *Mol. Biol. Evol.* **2021**, *38*, 3022. [CrossRef]
30. Lahlali, R.; Ezrari, S.; Radouane, N.; Kenfaoui, J.; Esmaeel, Q.; El Hamss, H.; Belabess, Z.; Barka, E.A. Biological control of plant pathogens: A global perspective. *Microorganisms* **2022**, *10*, 596. [CrossRef] [PubMed]
31. Zohora, U.; Ano, T.; Rahman, M. Biocontrol of *Rhizoctonia solani* K1 by iturin A producer *Bacillus subtilis* RB14 seed treatment in tomato plants. *Adv. Microbiol.* **2016**, *6*, 424. [CrossRef]
32. Rashad, Y.M.; Abdalla, S.A.; Sleem, M.M. Endophytic *Bacillus subtilis* SR22 triggers defense responses in tomato against Rhizoctonia root rot. *Plants* **2022**, *11*, 2051. [CrossRef]
33. Lee, G.; Choi, H.; Liu, H.; Han, Y.H.; Paul, N.C.; Han, G.H.; Kim, H.; Kim, P.I.; Seo, S.I.; Song, J. Biocontrol of the causal brown patch pathogen *Rhizoctonia solani* by *Bacillus velezensis* GH1 13 and development of a bacterial strain specific detection method. *Front. Plant Sci.* **2023**, *13*, 1091030. [CrossRef]
34. Wu, Z.; Huang, Y.; Li, Y.; Dong, J.; Liu, X.; Li, C. Biocontrol of *Rhizoctonia solani* via induction of the defense mechanism and antimicrobial compounds produced by *Bacillus subtilis* SL 44 on pepper (*Capsicum annuum* L.). *Front. Microbiol.* **2019**, *10*, 2676. [CrossRef]
35. Allard-Massicotte, R.; Tessier, L.; Lécuyer, F.; Lakshmanan, V.; Lucier, J.-F.; Garneau, D.; Caudwell, L.; Vlamakis, H.; Bais, H.P.; Beauregard, P.B. *Bacillus subtilis* early colonization of *Arabidopsis thaliana* roots involves multiple chemotaxis receptors. *mBio* **2016**, *7*, e01664-16. [CrossRef]
36. Al-Ali, A.; Deravel, J.; Krier, F.; Béchet, M.; Ongena, M.; Jacques, P. Biofilm formation is determinant in tomato rhizosphere colonization by *Bacillus velezensis* FZB42. *Environ. Sci. Pollut. Res.* **2018**, *25*, 29910. [CrossRef] [PubMed]
37. Mendis, H.C.; Thomas, V.P.; Schwientek, P.; Salamzade, R.; Chien, J.-T.; Waidyarathne, P.; Kloepper, J.; De La Fuente, L. Strain-specific quantification of root colonization by plant growth promoting rhizobacteria *Bacillus firmus* I-1582 and *B. amyloliquefaciens* QST713 in non-sterile soil and field conditions. *PLoS ONE* **2018**, *13*, e0193119. [CrossRef] [PubMed]

38. Al-Hussini, H.S.; Al-Rawahi, A.Y.; Al-Marhoon, A.A.; Al-Abri, S.A.; Al-Mahmooli, I.H.; Al-Sadi, A.M.; Velazhahan, R. Biological control of damping-off of tomato caused by *Pythium aphanidermatum* by using native antagonistic rhizobacteria isolated from Omani soil. *J. Plant Pathol.* **2019**, *101*, 315. [CrossRef]
39. Satognon, F.; Watts, D.; Adesemoye, A.O.; Torbert, H.; Prasad, R.; Mullenix, M.K. Impact of Bacillus PGPR consortium on germination and seedling vigor of cool- and warm-season forage grasses under different temperature regimes. *Arid. Land Res. Manag.* **2025**, 2549838. [CrossRef]
40. Widawati, S.; Suliasih. The Effect of Plant Growth Promoting Rhizobacteria (PGPR) on Germination and Seedling Growth of Sorghum bicolor L. Moench. *IOP Conf. Ser. Earth Environ. Sci.* **2018**, *166*, 012022. [CrossRef]
41. Khedher, S.B.; Kilani-Feki, O.; Dammak, M.; Jabnoun-Khiareddine, H.; Daami-Remadi, M.; Tounsi, S. Efficacy of *Bacillus subtilis* V26 as a biological control agent against *Rhizoctonia solani* on potato. *Comptes Rendus Biol.* **2015**, *338*, 784. [CrossRef]
42. Chebotar, V.K.; Gancheva, M.S.; Chizhevskaya, E.P.; Erofeeva, A.V.; Khiutti, A.V.; Lazarev, A.M.; Zhang, X.; Xue, J.; Yang, C.; Tikhonovich, I.A. Endophyte *Bacillus vallismortis* BL01 to control fungal and bacterial phytopathogens of tomato (*Solanum lycopersicum* L.) plants. *Horticulturae* **2024**, *10*, 1095. [CrossRef]
43. Hu, Y.; Yang, L.-Y.; Lei, M.-Y.; Yang, Y.-X.; Sun, Z.; Wang, W.; Han, Z.-M.; Cheng, L.; Lv, Z.-L.; Han, M. *Bacillus vallismortis* acts against ginseng root rot by modifying the composition and microecological functions of ginseng root endophytes. *Front. Microbiol.* **2025**, *16*, 1561057. [CrossRef]
44. Duan, Y.; Chen, R.; Zhang, R.; Jiang, W.; Chen, X.; Yin, C.; Mao, Z. Isolation and identification of *Bacillus vallismortis* HSB-2 and its biocontrol potential against apple replant disease. *Biol. Control* **2022**, *170*, 104921. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.