

Supplement material of

Bioactive compounds produced by *Hypoxyylon fragiforme* against *Staphylococcus aureus* biofilms

Kamila Tomoko Yuyama¹, Clara Chepkirui², Lucile Wendt², Diana Fortkamp^{1,3}, Marc Stadler², Wolf-Rainer Abraham^{1,*}

¹ Helmholtz Centre for Infection Research (HZI), Chemical Microbiology, Inhoffenstraße 7, 38124 Braunschweig, Germany; kamila.yuyama@helmholtz-hzi.de, diana.fortkamp@helmholtz-hzi.de

² Helmholtz Centre for Infection Research (HZI), Microbial Drugs; clara.chepkirui@helmholtz-hzi.de, lucile.wendt@helmholtz-hzi.de, marc.stadler@helmholtz-hzi.de

³ Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ), Department of Exact Sciences, Piracicaba - SP, Brazil; dianafortkamp@usp.br

* Correspondence: wolf-rainer.abraham@helmholtz-hzi.de; Tel.: +49-531-6181-4300




Figure S1: ^1H NMR spectrum of sclerin (**2**) in CDCl_3




Figure S2: ¹H NMR of the diacid of sclerin (**3**) in d₆-methanol




Figure S3: ¹H NMR spectrum of methyl 1-(5-hydroxy-6-carboxylic-2,3,4-trimethylphenyl) propionate (**4**) in CDCl₃