



Article

# Modeling the Reduction and Cross-Contamination of *Salmonella* in Poultry Chilling Process in China

Xingning Xiao<sup>1,2</sup>, Wen Wang<sup>2,\*</sup> , Jianmin Zhang<sup>3</sup>, Ming Liao<sup>3</sup>, Hua Yang<sup>2</sup>, Weihuan Fang<sup>4</sup> and Yanbin Li<sup>1,5,\*</sup>

<sup>1</sup> College of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou 310058, China; xingningxiao@126.com

<sup>2</sup> State Key Laboratory for Quality and Safety of Agro-products, MOA Laboratory of Quality & Safety Risk Assessment for Agro-products (Hangzhou), Institute of Quality and Standard of Agricultural Products, Zhejiang Academy of Agricultural Sciences, Hangzhou 310021, China; yanghua@mail.zaas.ac.cn

<sup>3</sup> College of Veterinary Medicine, South China Agricultural University, Guangzhou 510642, China; junfeng-v@163.com (J.Z.); mliao@scau.edu.cn (M.L.)

<sup>4</sup> College of Animal Sciences, Zhejiang University, Hangzhou 310058, China; whfang@zju.edu.cn

<sup>5</sup> Department of Biological & Agricultural Engineering, University of Arkansas, Fayetteville, Arkansas, AR 72701, USA

\* Correspondence: wangwen@zaas.ac.cn (W.W.); yanbinli@zju.edu.cn (Y.L.); Tel.: +86-571-86419052 (W.W.); +86-571-88982536 (Y.L.); Fax: +86-571-86401834 (W.W.); +86-571-88982530 (Y.L.)

Received: 10 September 2019; Accepted: 11 October 2019; Published: 13 October 2019



**Abstract:** The study was to establish a predictive model for reduction and cross-contamination of *Salmonella* on chicken in chilling process. Reduction of *Salmonella* on chicken was  $0.75 \pm 0.04$ ,  $0.74 \pm 0.08$ , and  $0.79 \pm 0.07$  log CFU/g with 20, 50, and 100 mg/L of chlorine, respectively. No significant differences of bacterial reductions with 20–100 mg/L of chlorine were found and a Normal ( $-0.75, 0.1$ ) distribution could describe the uncertainty of bacterial reductions. Inoculated and non-inoculated chicken samples were washed together and bacterial transfer rates among them were 0.13%–0.004% with 20–100 mg/L of chlorine. No significant differences of transfer rates with 50–100 mg/L of chlorine were observed and a Triangle ( $-2.5, -1.5, -1.1$ ) distribution could describe the log transfer rate. Additionally, a 3-factor response surface model based on the central composite design was developed to evaluate the effects of initial contamination level (1–5 log CFU/g), pre-chill incidence (3%–40%) and chlorine concentration (0–100 mg/L) on post-chill incidence. The post-chill incidences in these treatments were within 30%–91.7%. The developed model showed a satisfactory performance to predict the post-chill incidence as evidenced by statistical indices (pseudo- $R^2 = 0.9$ ;  $p < 0.0001$ ; RMSE = 0.21) and external validation parameters ( $B_f = 1.02$ ;  $A_f = 1.11$ ).

**Keywords:** modeling; *Salmonella*; reduction; cross-contamination; poultry chilling

## 1. Introduction

Microbial contamination is a widespread public concern in the meat industry because it can shorten the shelf life and increase the safety risk of fresh meat and meat products [1,2]. Recently, contamination of *Salmonella* continues to be a major concern in poultry industry [3,4]. It was found that non-typhoid *Salmonella* annually caused 9.87 million gastroenteritis cases in China and more than half of the retail chicken carcasses were contaminated with *Salmonella* [4,5]. *Salmonella* from the skin, feather, spillage of intestinal, and water are disseminated from carcass to carcass as they are moved down the poultry slaughtering line [6]. Incoming animal-associated contamination of the abattoir environment as well as the slaughter of large numbers of animals on the same slaughter line contributes to direct contamination or cross-contamination of chicken carcasses during slaughtering.

Following poultry evisceration, the next step in poultry processing is chilling by adding chlorine into the water to reduce microbial contamination and cross-contamination [6–8]. Chlorine-based decontamination disinfectants, such as sodium hypochlorite, chlorine dioxide, acidified sodium chlorite, and monochloramine, are the most commonly used sanitizers in poultry slaughterhouses [9–11]. Chlorine concentrations used in poultry slaughterhouses are different from country to country. The U.S. Department of Agriculture Food Safety and Inspection Service (USDA-FSIS) requires that processors add 20 to 50 mg/L of chlorine to chilling water to reduce pathogens and prevent pathogen cross-contamination of poultry carcasses [12]. In China, 50 to 100 mg/L of chlorine is commonly used in poultry chilling process [13]. Although other antimicrobial agents are approved for poultry processing such as organic acids, chlorine still remains the most widely used antimicrobial chemical by the poultry industry due to its antimicrobial efficacy, convenience, and low price [8,14,15].

The chilling tank is a high-risk area where cross-contamination between contaminated and non-contaminated carcasses occurred via washing water, leading the changes of bacterial contamination level and incidence [6]. A higher incidence of *Salmonella* was observed in poultry carcasses from post-chill than those from pre-chill [16,17]. Several factors have been determined to be involved with bacterial incidence during chilling, including poultry contamination level, chlorine concentration and seasonal effect [6,16,18]. Predictive microbiology is an important tool for investigating the behavior of pathogens under prescribed environmental conditions through the development of mathematical models [19,20]. In previously studies, a Weibull model was used to describe bacterial reduction and a logistic model was developed to predict post-chill incidence in poultry chilling process with 0–50 mg/L of chlorine, while chlorine concentration with 50–100 mg/L is common used in China according to the national performance standard of the operation procedure for poultry slaughtering [6,21,22]. Very few predictive models are available for the description of *Salmonella* reduction and cross-contamination during chilling for quantitative microbial risk assessment (QMRA) of the poultry supply chain in China.

Therefore, the objectives of this study were: (i) To investigate the effect of chlorine concentration on the reduction and transfer of *Salmonella* in chilling process; (ii) to determine the post-chill incidence of *Salmonella* on chicken products with different initial contamination levels, pre-chill incidences and chlorine concentrations; and (iii) to develop probability distribution and predictive models for describing the bacterial reduction, bacterial transfer rate and bacterial post-chill incidence. The developed predictive models in this study were critical to provide reliable inputs to a QMRA model for the whole poultry supply chain in China.

## 2. Materials and Methods

### 2.1. Bacterial Inoculum

Five serovars of *Salmonella* (Stanley BYC12, Indiana HZC10, Typhimurium YXC1, Thompson LWC10, Kentucky CBC2) isolated from poultry slaughterhouses (Guangzhou, Guangdong, China) in China were used in this study. The bacterial strains were maintained in brain heart infusion broth (BHI, Becton Dickinson (BD), Franklin Lakes, NJ, USA) containing 20% (*v/v*) glycerol at –80 °C. Each strain was separately incubated in BHI at 37 °C for 24 h and cultured to approximately 9 log CFU/mL. Equal volumes of each culture suspension were mixed to obtain a five-strain mixture of *Salmonella*. Appropriate 10-fold dilutions in sterile phosphate buffered saline (PBS, Sigma, St. Louis, MO, USA) were made and plated on xylose lysine Tergitol 4 (XLT4, BD) agar to determine the cell number in the inoculums.

### 2.2. Preparation and Inoculation of Chicken Samples

Skin injuries will happen due to the weak skin is more susceptible to mechanical tears, leading to skinless chicken meat exposure to the chilling water, especially in small scale slaughterhouses in China. In the preliminary test, bacterial reduction and bacterial transfer rate on skin-on chicken wingettes and skinless chicken breasts were compared, which was not significantly different ( $p > 0.05$ )

as determined by analysis of variance (ANOVA). Therefore, chicken breast samples (weight:  $25 \pm 1$  g, size:  $5 \times 3 \times 2$  cm) were used in the tests, which were purchased from a local supermarket (Hangzhou, Zhejiang, China) and stored in a freezer at  $-20$  °C. After thawing overnight at  $4$  °C, the samples were exposed to UV light for 30 min in a biosafety cabinet (Thermo Fisher 1389, Waltham, MA, USA) to decontaminate initial microbial contamination. Controls in the test showed that no *Salmonella* was initially present on the chicken breast samples. Then the chicken breast samples were submerged into *Salmonella* suspensions for 30 min and transferred to plastic plates for another 30 min to allow bacterial full attachment, both of which were performed at ambient temperature ( $25 \pm 1$  °C).

### 2.3. Chilling Treatments

The chilling conditions were designed based on the operation of poultry slaughterhouses (Guangzhou, Guangdong) in China, the ratio of chicken samples to chilling water was 1:10 (*w/v*). Chilling treatments at  $4$  °C were carried out in a laboratory water bath (TX150, Grant, Royston, UK) equipped with a digital thermometer (34970A, Agilent, Santa Clara, CA, USA) to monitor temperatures of water. Sodium hypochlorite (NaClO) stock solution containing 56.8 mg/mL chlorine was purchased from Sangon Biotech Co., Ltd., Shanghai, China. Chlorination of chilling water was prepared by diluting NaClO stock solution using sterile Milli-Q water (Millipore, Billerica, MA, USA) and the chlorine concentration in chilling water was determined using a Palintest ChlorSense meter (CS 100, Gateshead, Tyne and Wear, UK).

To evaluate the effect of chlorine concentrations (0, 20, 50, and 100 mg/L) on the bacterial reduction, twelve inoculated chicken breast samples were submerged into the water bath. Three inoculated samples were removed for processing every 10 min within a 40 min treatment period. The samples were individually added to sterile stomacher bags (Seward, London, UK) containing 25 mL buffered peptone water (BPW, BD) and homogenized for 1 min in a Model 400 food stomacher (Seward, London, UK). The homogenates were diluted to appropriate concentrations for bacterial enumeration. The initial inoculation load of chicken breasts was  $5.6 \pm 0.1$  log CFU/g. The bacterial population on chicken samples at  $t = 0$  was determined using the inoculated samples without treatment. Each treatment was repeated three times on different days and duplicated plates were used in microbial tests for each sample.

To assess the effect of chlorine concentrations (0, 20, 50, and 100 mg/L) on the bacterial transfer rate from inoculated chicken breasts to non-inoculated chicken breasts via chilling water, twelve inoculated and twelve non-inoculated chicken breasts were submerged into the water bath together. Three inoculated and three non-inoculated samples were removed for processing every 10 min within a 40 min treatment period. The non-inoculated samples were individually added to sterile stomacher bags containing 25 mL of BPW and homogenized for 1 min in a Model 400 food stomacher. The homogenates were diluted to appropriate concentrations for bacterial enumeration. The initial inoculation load of chicken breast samples was  $5.6 \pm 0.1$  log CFU/g. The bacterial population on chicken samples at  $t = 0$  was determined using the inoculated samples without treatment. Each treatment was repeated three times on different days and duplicated plates were used in microbial tests for each sample.

To evaluate the combined effects of initial contamination levels (1, 2, 3, 4, and 5 log CFU/g), pre-chill incidences (3%, 10.2%, 21.5%, 32.8%, and 40%) and chlorine concentrations (0, 20, 50, 80, and 100 mg/L) on post-chill incidence of *Salmonella*, a response surface model based on the central composite design was developed with the JMP10 software (SAS Institute, Cary, NC, USA) to predict the post-chill incidence. In multivariable analyses, the traditional optimization technique of changing one variable at a time to study the variable response effect is impracticable and does not represent the interaction effect between different factors. Therefore, experimental statistical design considers one of the most useful methods in obtaining valuable and statistically significant models of a phenomenon by performing a minimum number of calculated experiments. It also considers interactions among the variables and can be used to optimize the operating parameters in multivariable analyses. Response surface methodology was used for the modeling and analysis of problems in which a response of

interest is influenced by several variables to optimize the same response. A favorite model in response surface methodology is the central composite design, which is efficient and flexible in providing adequate data on the effects of variables and overall experiment error even with a fewer number of experiments [23–25]. Tested variables (initial contamination level, pre-chill incidence, and chlorine concentration) were denoted as  $X_1$ ,  $X_2$ , and  $X_3$ , respectively, and each factor in the design was studied at five different levels ( $-\alpha$ ,  $-1$ ,  $0$ ,  $+1$ ,  $+\alpha$ ) (Table 1), with a total of 20 runs according to the experimental design (Table 2). Among the 30 chicken breasts, a total of one, three, seven, ten, and twelve chicken breasts were inoculated to obtain a pre-chill incidence of 3%, 10.2%, 21.5%, 32.8%, and 40%, respectively. Then, inoculated and non-inoculated samples were mixed together in the water bath for a 40 min chilling treatment. The samples were individually added to sterile stomacher bags containing 25 mL of BPW and homogenized for 1 min in a Model 400 food stomacher. The homogenates were diluted to appropriate concentrations for bacterial enumeration. The whole experiment was repeated twice on different days and duplicated plates were used in microbial tests for each chicken breast sample.

**Table 1.** Levels of variables in the experiment.

Variable	Range	Level				
		$-\alpha$	$-1$	$0$	$1$	$\alpha$
Initial contamination level (log CFU/g)	1–5	1	2	3	4	5
Pre-chill incidence (%)	3–40	3	10.2	21.5	32.8	40
Chlorine (mg/L)	0–100	0	20	50	80	100

**Table 2.** Experimental design and post-chill incidence of *Salmonella* in chilling process.

Run	Initial Contamination Level (log CFU/g)	Pre-Chill Incidence (%)	Chlorine (mg/L)	<sup>a</sup> Post-Chill Incidence (%)
1	2	32.8	80	43.3 ± 14.1 <sup>abcd</sup>
2	4	10.2	80	66.7 ± 4.7 <sup>abc</sup>
3	3	21.5	50	65 ± 2.4 <sup>abc</sup>
4	2	10.2	20	40 ± 12.4 <sup>bcd</sup>
5	3	21.5	50	68.4 ± 11.8 <sup>ab</sup>
6	4	32.8	20	91.7 ± 7.1 <sup>a</sup>
7	3	21.5	50	65 ± 11.8 <sup>abc</sup>
8	4	32.8	80	85 ± 2.4 <sup>a</sup>
9	2	10.2	80	35 ± 7.1 <sup>bcd</sup>
10	2	32.8	20	60 ± 28.3 <sup>bcd</sup>
11	3	21.5	50	70 ± 14.1 <sup>ab</sup>
12	4	10.2	20	90 ± 9.4 <sup>a</sup>
13	3	21.5	0	81.7 ± 21.2 <sup>a</sup>
14	3	40	50	78.4 ± 25.9 <sup>a</sup>
15	5	21.5	50	90 ± 4.7 <sup>a</sup>
16	3	21.5	50	65 ± 4.7 <sup>abc</sup>
17	3	3	50	40 ± 18.9 <sup>bcd</sup>
18	3	21.5	100	65 ± 2.4 <sup>bc</sup>
19	3	21.5	50	63.3 ± 4.7 <sup>abc</sup>
20	1	21.5	50	30 ± 4.7 <sup>cd</sup>

<sup>a</sup> Values are means ± standard deviations ( $n = 2$ ). Values followed by different superscript letters are significantly different ( $p < 0.05$ ).

#### 2.4. Bacterial Enumeration

In the preliminary test, the recovery of injured *Salmonella* was considered. XLT4 agar was compared with Tryptic Soy agar (TSA, BD) in plate counting and only less than 0.1 log CFU/g difference was observed, which was not significantly different ( $p > 0.05$ ) as determined by ANOVA. Therefore, populations of *Salmonella* were selectively enumerated on XLT4 agar. The homogenates were serially 10-fold diluted in BPW and a 50  $\mu$ L portion of appropriate dilutions was plated in duplicate onto the XLT4 agar using a spiral plater (WASP 2, Don Whitley Scientific, Shipley, UK). The plates were

incubated at 37 °C for 18 h. Colonies on XLT4 agar plates were enumerated by a ProtoCOL 3 automated colony counter (Synbiosis, Cambridge, UK). The limit of detection was one colony for 50 µL sample (1.3 log CFU/mL). The viable bacterial populations on samples were expressed as CFU/g.

When the bacterial population on samples were under the detection limit, enrichment test was conducted to determine whether there were *Salmonella* survivors in samples. One milliliter of homogenates and water sample were aseptically collected in 9 mL of Selenite Cystine Broth (SC, HopeBiol, Qing Dao, Shan Dong, China) and incubated at 37 °C for 24 h, respectively. The enriched samples were plated onto the XLT4 agar and incubated at 37 °C for 18 h. Black colonies on the XLT4 agar denoted that *Salmonella* survivors in samples.

### 2.5. Calculation of Bacterial Reduction, Bacterial Transfer Rate and Post-chill Incidence

Counts of surviving bacteria were log transformed, and the bacterial reduction ( $Y_{red}$ ) in bacterial population was calculated with the Equation (1):

$$Y_{red} = \log N_t - \log N_0 \quad (1)$$

where  $N_t$  (CFU/g) is the number of bacteria at time  $t$  (min) and  $N_0$  (CFU/g) is the initial number of bacteria.

The transfer rate (TR%) was defined as the percentage of bacterial transferred from the donor surface to recipient surface, which could be calculated by Equation (2).

$$TR(\%) = \frac{N_{recipient}}{N_{donor}} \times 100 \quad (2)$$

where,  $N_{recipient}$  and  $N_{donor}$  are bacterial population on the recipient surface and donor surface, respectively. In this study,  $N_{recipient}$  (CFU) is the number of *Salmonella* on non-inoculated chicken breast samples after chilling, and  $N_{donor}$  (CFU) is the initial microbial load on inoculated chicken breast samples.

Positive contamination of *Salmonella* on each chicken breast was recorded. Post-chill incidence ( $Y_{pc}$ ) could be calculated by Equation (3):

$$Y_{pc}(\%) = \frac{A}{B} \times 100 \quad (3)$$

where,  $A$  and  $B$  is number of positive chicken breasts and total number of chicken breasts, respectively.

### 2.6. Model Development

In the cases that no significant differences of bacterial reduction and transfer rate were observed, probability distribution model could provide all potential results [26]. Besides, bacterial reductions and transfer rates showed a large variation because of multiple uncertainties that were involved, as well as the inherent errors in microbial collection from surfaces and enumeration techniques. Hence, to describe the uncertainty, probability distributions were defined based on the Kolmogorov–Smirnov test to describe the uncertainty using @Risk 7.5 software (Palisade, Newfield, NY, USA).

The stepwise regression model was constructed recursively by adding or deleting one independent prediction at each time. A backward elimination was used which starts from a model with a full set of base functions and then gradually dropped out the predictor with the least effect in the model in a stepwise fashion [27]. For the prediction of post-chill incidence, a second order polynomial equation was fitted to the post-chill incidence by a backward stepwise regression. This resulted in an empirical model that related the response measured to the independent variables of the experiment. For a three-factor system the model is Equation (4):

$$Y_{pc} = K_0 + K_1X_1 + K_2X_2 + K_3X_3 + K_4X_1^2 + K_5X_2^2 + K_6X_3^2 + K_7X_1X_2 + K_8X_1X_3 + K_9X_2X_3 \quad (4)$$

where  $Y_{pc}$  is the post-chill incidence (%);  $X_1$ ,  $X_2$ , and  $X_3$  are initial contamination level (log CFU/g), pre-chill incidence (%) and chlorine concentration (mg/L), respectively.  $K_0$  is the intercept;  $K_1$ ,  $K_2$ , and  $K_3$ , are linear coefficients;  $K_4$ ,  $K_5$ , and  $K_6$  are squared coefficients;  $K_7$ ,  $K_8$ , and  $K_9$  are interaction coefficients.

### 2.7. Model Evaluation and Validation

ANOVA was used to evaluate significance and adequacy of the model. Fitting goodness of the model was characterized by correlation coefficient ( $R^2$ ) and the root mean square error (RMSE). Eight random independent trials were conducted with chicken carcasses to calculate the bias factors ( $B_f$ ) (Equation (5)) and accuracy factors ( $A_f$ ) (Equation (6)) for model validation (Table 3) [28]. The selected parameters were within the original range of the experimental design but not included in the development of the model [29].

$$B_f = 10^{\left[\frac{\sum_{i=1}^n \log\left(\frac{obs}{pred}/n\right)}{n}\right]} \quad (5)$$

$$A_f = 10^{\left[\frac{\sum_{i=1}^n |\log\left(\frac{obs}{pred}\right)|/n}{n}\right]} \quad (6)$$

where  $n$  is the number of trials,  $obs$  is the observed values of post-chill incidence (%), and  $pred$  is the predicted values of post-chill incidence (%).

**Table 3.** Eight independent trials to validate the model for post-chill incidence in chilling process.

Run	Initial Contamination Level (log CFU/g)	Pre-Chill Incidence (%)	Chlorine (mg/L)	Post-Chill Incidence (%)	
				Observed	Predicted
1	3	12.5	10	62.5 ± 8.8	69.1
2	3	37.5	10	100.0 ± 0.0	88.0
3	3	12.5	70	50.0 ± 0.0	52.0
4	3	37.5	70	87.5 ± 8.8	71.0
5	5	12.5	10	100.0 ± 0.0	100.6
6	5	37.5	10	100.0 ± 0.0	119.5
7	5	12.5	70	100.0 ± 0.0	83.5
8	5	37.5	70	100.0 ± 0.0	102.5

### 2.8. Color Measurements

Color changes of chicken breasts that occurred during chilling with 0, 20, 50, and 100 mg/L of chlorine were determined using a Chroma Meter CR 400 instrument (Minolta, Osaka, Japan). Samples were measured after a 40 min treatment. Values of  $L$ ,  $a$ , and  $b$  represented the lightness, redness and yellowness, respectively, were recorded. The total color difference ( $\Delta E$ ) was calculated according to Equation (7).  $\Delta E$  values of 1 to 2 mean that color change is perceived through close observation, and values of 3 to 10 mean changes is perceived at a glance [30]. All measurements were taken on five sites of each chicken breast sample.

$$\Delta E = \sqrt{\Delta L^2 + \Delta a^2 + \Delta b^2} \quad (7)$$

where  $\Delta L$ ,  $\Delta a$ , and  $\Delta b$  are the differences of lightness, redness, and yellowness between the treated and untreated samples, respectively.

### 2.9. Transmission Electron Microscopy (TEM)


In order to observe the morphology differences of the *Salmonella* cells with 0, 50, and 100 mg/L of chlorine, ultrastructural changes of bacterial cells were observed with a transmission electron microscope (TEM, Hitachi 7650, Ibaraki, Japan). The treated samples were removed from the water bath and placed into a sterile stomacher bag containing 25 mL BPW and then squeezed by hand to wash off attached bacteria. The wash solution for each treatment was collected into a 50 mL centrifuge tube and centrifuged at 6000×  $g$  for 10 min. The bacterial pellets were transferred into sterile 1.5

mL centrifuge tubes. Samples were fixed with 1 mL of 2.5% (*v/v*) glutaraldehyde (Sangon Biotech, Shanghai, China) and examined using a TEM as previously described [31].

### 3. Results and Discussion

#### 3.1. *Salmonella* Reduction at Different Chlorine Concentrations

In non-chlorinated water, bacterial reductions of *Salmonella* on chicken breasts after the 40 min treatments were  $0.49 \pm 0.1$  log CFU/g. The reductions of *Salmonella* on chicken breasts after the 40 min treatments were  $0.75 \pm 0.04$ ,  $0.74 \pm 0.08$ , and  $0.79 \pm 0.07$  log CFU/g with 20, 50, and 100 mg/L of chlorine, respectively (Figure 1). There were no significant differences of bacterial reductions among 20, 50, and 100 mg/L of chlorine and the treatment time was not a significant variable to inactivate the bacteria ( $p > 0.05$ ).



**Figure 1.** Bacterial reductions of *Salmonella* on chicken breasts with 20 (◆), 50 (■) and 100 (▲) mg/L of chlorine.

The effectiveness of chlorine on reduction of *Salmonella* on chicken was limited [8,32]. It is believed that the chlorine does not easily access the bacteria in ridges and crevices on poultry carcasses [15,33]. Besides, *Salmonella* on chicken surface cannot be removed effectively by chlorine due to the interference of oil between the sanitizers and the surface [32]. Generally, treatments with chlorine resulted in less than a 1 log reduction on carcasses [15,34,35]. For example, Yang et al. (2010) [21] found that the reductions of *Salmonella* with 10–50 mg/L of chlorine were  $< 0.5$  log CFU/cm<sup>2</sup> in chicken skin. Lee et al. (2013) [15] observed that the combination of 200 mg/L of chlorine and ultrasound significantly reduced *Salmonella* populations by 0.5–0.8 log CFU/g in chicken skin. Tamblyn and Conner (1997) [36] found that chlorine levels must be at least 400 mg/L to kill the attached *Salmonella* on broilers. In this study, bacterial reductions on chicken breasts were less than 1 log CFU/g with 20–100 mg/L of chlorine, showing the effectiveness of chlorine on reduction of *Salmonella* was limited, which was consistent with previous studies.

#### 3.2. Transfer of *Salmonella* at Different Chlorine Concentrations

The averages of TR% on chicken breasts at the end of 40 min treatments were 0.12, 0.08, 0.03, and 0.02% with 0, 20, 50, and 100 mg/L of chlorine, respectively. The average bacterial populations in chilling water at the end of 40 min treatments of 0, 20, 50, and 100 mg/L of chlorine were 4, 3.5,  $< 1.3$  (detection limit) and 0 log CFU/mL, respectively. Chlorine with a concentration of 0–20 mg/L failed to mitigate bacterial transfer due to a high contamination level in the washing water. Bacterial


populations in water significantly decreased after chilling with 50 mg/L of chlorine and no *Salmonella* survivor in water was found at 100 mg/L of chlorine. The results indicated that 50–100 mg/L of chlorine were effective in controlling *Salmonella* transfer in chilling process. Chilling water is an ideal medium for the potential spread of bacterial pathogens during processing. Therefore, the presence of a sanitizing agent such as sodium hypochlorite in the wash water is critical to preventing pathogen survival and transfer [37,38]. Similarly, Mead et al. (1994) [39] reported that less than 30 mg/L of chlorine did not prevent microbial cross-contamination on poultry. Chlorination of chilling water was effective to control cross-contamination, but was limited for reducing the bacteria attached on the chicken carcasses [21,39].

### 3.3. *Salmonella* Post-Chill Incidence under Different Initial Contamination Levels, Pre-Chill Incidences, and Chlorine Concentrations


The post-chill incidences of *Salmonella* on chicken breasts under different initial contamination levels, pre-chill incidences, and chlorine concentrations are presented in Table 2. The effect of initial contamination level, pre-chill incidence and chlorine concentration on the post-chill incidence was significant ( $p < 0.05$ ). In Table 2, with a 5 log CFU/g of *Salmonella* inoculated on chicken breasts, the post-chill incidence of *Salmonella* was 90%, but was reduced by 60% at initial contamination level of 1 log CFU/g. Besides, with the same initial contamination level and chlorine concentration, the post-chill incidence of *Salmonella* was 78.4% when washed at a 40% pre-chill incidence, but was reduced by 38.4% with a 3% pre-chill incidence. With the same initial contamination level and pre-chill incidence, the post-chill incidence of *Salmonella* was 81.7% in non-chlorinated chilling water, but was reduced to 23.3% after chilling with 50 mg/L of chlorine. Yang et al. (2002) [6] reported that with a 43.3% pre-chill contamination, the post-chill contamination of *Salmonella* was more than 90% without chlorination, but was reduced to 20% after chlorination of chill water with 50 mg/L of chlorine. Chlorination of chilling water could contribute significantly toward a reduction in *Salmonella* incidence on commercially processed carcasses [6,40]. However, when the pathogen has a resistance to chlorine, a result has been reported that no correlation between the incidence of *Salmonella* on post-chill poultry carcasses and the chlorine concentration in chilling process [16].

### 3.4. Model Development

There were not significant correlation between bacterial reductions with the 20–100 mg/L chlorine treatments and the chilling time ( $p > 0.05$ ). A Normal ( $-0.75, 0.1$ ) distribution could describe the bacterial reductions with 20–100 mg/L of chlorine (Figure 2). Normal distributions, fitted by the sampling data randomly collected before and after chilling in slaughterhouse, have been used to describe the *Campylobacter* reductions on chicken carcasses [41,42]. Besides, there were not significant correlation between bacterial transfer rates with the treatments of 50–100 mg/L of chlorine and the chilling time ( $p > 0.05$ ). Log transfer rates were essentially normally distributed and a Triangle ( $-2.5, -1.5, -1.1$ ) distribution could describe the TR with 50–100 mg/L of chlorine (Figure 3). Normal distributions have been used to describe the variation of TR in previous studies [43,44]. A paradigmatic study was reported by Chen et al. (2001) [43]. They investigated bacterial transfer rates between hands and other common surfaces involved in food preparation in kitchen, and found the distribution of the logarithmic transfer rates appears approximately normal.



**Figure 2.** Probability distribution of *Salmonella* reduction on chicken breasts in chilling water with 20–100 mg/L of chlorine.



**Figure 3.** Probability distribution of transfer rate (log%) among chicken breasts in chilling water with 50–100 mg/L of chlorine.

A backward stepwise regression was carried out using JMP software to develop a simplified response surface model only with the significant variables. By using ANOVA on the estimated parameters of all variables, we found statistical significance with three linear coefficients ( $X_1$ ,  $X_2$  and  $X_3$ ) ( $p < 0.001$ ), and the quadratic coefficient ( $X_3^2$ ) ( $p < 0.05$ ) for estimation of bacterial post-chill incidence. A summary of the estimated parameters for uncoded variables with significances is given in Table 4. The simplified model is shown as Equation (8).

$$Y_{pc} = 18.28 + 15.75X_1 + 0.757X_2 - 0.636X_3 + 0.0044X_3^2 \tag{8}$$


where  $Y_{pc}$  is post-chill incidence (%),  $X_1$  is the initial contamination level (log CFU/g),  $X_2$  is the pre-chill incidence (%), and  $X_3$  is the chlorine concentration (mg/L).

**Table 4.** The estimated parameters for variables with significance.

Factor	Value	Standard Error	Prob > F
Intercept	92.0	3.3	
$X_1$	31.0	2.8	< 0.0001 **
$X_2$	13.6	2.8	0.0002 *
$X_3$	-9.7	2.8	0.004 *
$X_3^2$	10.8	4.6	0.03 *

Asterisks indicate statistical significance by F test (\*\*  $p < 0.0001$ ; \*  $p < 0.05$ ).

The response surface plot in Figure 4 demonstrates that the effect of initial contamination level, pre-chill incidence and chlorine concentration on the post-chill incidence of *Salmonella* in chilling process. The response surface plot in Figure 4A describes the post-chill incidence of initial contamination level and pre-chill incidence at the fixed chlorine concentration of 50 mg/L (coded level of 0). The post-chill incidence of *Salmonella* was significantly reduced ( $p < 0.05$ ) as the initial contamination level and pre-chill incidence decreased, and the post-chill incidence was more sensitive to the initial contamination level (Figure 4A). Post-chill incidence was slightly sensitive to the change of chlorine concentration, as compared with initial contamination level and pre-chill incidence (Figure 4B,C). The goodness-of-fit of a predictive model has a satisfactory performance as evidenced by statistical indices (pseudo- $R^2 = 0.9$ ;  $p < 0.0001$ ; RMSE = 0.21).



**Figure 4.** Response surface plots describing the effect of initial contamination level and pre-chill incidence (A); initial contamination level and chlorine concentration (B); and pre-chill incidence and chlorine concentration (C) on post-chill incidence of *Salmonella* in chilling process.

### 3.5. Model Evaluation and Validation

The model has a good statistical performance as shown by pseudo- $R^2$  (0.9), probability value ( $p < 0.0001$ ) and the lack of fit test ( $p > 0.05$ ). A statistical validation is insufficient to evaluate accuracy of the model to predict post-chill incidence. Therefore, an external validation was carried out. The results of eight independent experiments (not included in the model development) shown in Table 3 were used to calculate the  $B_f$  and  $A_f$  based on Equations (5) and (6). The calculated  $B_f$  was 1.02, which lies in the acceptable range of 0.9–1.05 [28,45]. The  $A_f$  value was 1.11, revealing a merely 11% difference between the observations and predictions. These results indicated that the developed model could give a reliable prediction for post-chill incidence within the range of variables employed.

### 3.6. Color Changes during Chilling

For poultry products, color and visual appearance are important attributes to consumers [46]. The changes of color ( $\Delta L$ ,  $\Delta a$ ,  $\Delta b$ ) and  $\Delta E$  values of the treated chicken breasts are summarized in Table 5. The  $L$ ,  $a$  and  $b$  value of the chicken breasts color did not change with the treatments ( $p > 0.05$ ). There were no obvious color changes of chicken breasts after chilling with 20–100 mg/L of chlorine ( $\Delta E < 10$ ). In general, the color changes of chicken breasts were acceptable based on the visual inspection.


**Table 5.** Changes in the color of chicken breasts treated with different chlorine concentrations.

Chlorine (mg/L)	$\Delta L$	$\Delta a$	$\Delta b$	$\Delta E$
0	$5.4 \pm 1.0^a$	$1.4 \pm 0.4^a$	$3.3 \pm 0.9^a$	6.5
20	$7.0 \pm 1.9^a$	$2.2 \pm 0.9^a$	$1.6 \pm 0.7^a$	7.5
50	$5.6 \pm 1.6^a$	$0.8 \pm 0.9^a$	$2.8 \pm 1.6^a$	6.3
100	$5.5 \pm 2.2^a$	$1.7 \pm 1.0^a$	$2.8 \pm 1.4^a$	6.4

Mean values followed by different superscript letters are significantly different ( $p < 0.05$ ).

### 3.7. Morphological Changes Reveled by TEM

We closely examined the morphology changes of bacterial cells using TEM. *Salmonella* cells in the absence of chlorine treatment retained their round shapes (Figure 5A). However, after chlorine treatment, the cells were completely disrupted and leaked cytoplasmic material to the extracellular medium (Figure 5B,C). Chlorine reacts with water to form the active antimicrobial hypochlorous acid, which destroys microbial cells by hindering carbohydrate metabolism [46].



**Figure 5.** TEM images of *Salmonella* treated with 0 (A), 50 (B), and 100 mg/L (C) of chlorine.

## 4. Conclusions

Chlorine with concentrations of 50–100 mg/L was effective for reducing bacteria in chilling water to control cross-contamination, but was limited for reducing the bacteria attached on the chicken breasts. A Normal (−0.75, 0.1) distribution model could describe the bacterial reduction in chilling process and a Triangle (−2.5, −1.5, −1.1) distribution could describe the logarithm of transfer rate. For prediction

of post-chill incidence, the developed response surface model has a satisfactory performance, which could be used to provide the input for QMRA model of *Salmonella* in poultry supply chain in China.

**Author Contributions:** Conceptualization, X.X. and W.W.; methodology, X.X.; software, X.X.; validation, X.X., W.W. and Y.L.; formal analysis, X.X.; investigation, X.X.; resources, X.X.; data curation, X.X.; writing—original draft preparation, X.X.; writing—review and editing, X.X., W.W., J.Z., M.L., H.Y., W.F., and Y.L.; visualization, X.X.; supervision, W.W. and Y.L.; project administration, W.W. and Y.L.; funding acquisition, W.W. and Y.L.

**Funding:** This research was supported in part by National Natural Science Foundation of China (31601392), Walmart Foundation (SA1703164) and Walmart Food Safety Collaboration Center.

**Acknowledgments:** The authors thank Gaolan Ding and Zihan Xu for their help in the experiments. The authors also thank three poultry companies for providing the information on poultry processing.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Rajan, K.; Shi, Z.; Ricke, S.C. Current aspects of *Salmonella* contamination in the US poultry production chain and the potential application of risk strategies in understanding emerging hazards. *Crit. Rev. Microbiol.* **2017**, *43*, 370–392. [CrossRef] [PubMed]
2. Wang, H.; Duan, D.; Wu, Z.; Xue, S.; Xu, X.; Zhou, G. Primary concerns regarding the application of electrolyzed water in the meat industry. *Food Control.* **2019**, *95*, 50–56. [CrossRef]
3. Guo, C.; Hoekstra, R.M.; Schroeder, C.M.; Pires, S.M.; Ong, K.L.; Hartnett, E.; Scallan, E. Application of Bayesian techniques to model the burden of human salmonellosis attributable to US food commodities at the point of processing: Adaptation of a Danish model. *Foodborne Pathog. Dis.* **2011**, *8*, 509–516. [CrossRef] [PubMed]
4. Zhu, J.; Wang, Y.; Song, X.; Cui, S.; Xu, H.; Yang, B.; Huang, J.; Liu, G.; Chen, Q.; Zhou, G.; et al. Prevalence and quantification of *Salmonella* contamination in raw chicken carcasses at the retail in China. *Food Control.* **2014**, *44*, 198–202. [CrossRef]
5. Yang, B.; Xi, M.; Wang, X.; Cui, S.; Yue, T.; Hao, H.; Walls, I. Prevalence of *Salmonella* on raw poultry at retail markets in China. *J. Food Prot.* **2011**, *74*, 1724–1728. [CrossRef] [PubMed]
6. Yang, H.; Li, Y.; Griffis, C.L.; Waldroup, A.L. A probability model for cross-contamination by *Campylobacter jejuni* and *Salmonella* Typhimurium in poultry chilling process. *Appl. Eng. Agric.* **2002**, *18*, 717–724. [CrossRef]
7. James, C.; Vincent, C.; de Andrade Lima, T.I.; James, S.J. The primary chilling of poultry carcasses—a review. *Int. J. Refrig.* **2006**, *29*, 847–862. [CrossRef]
8. Northcutt, J.K.; Smith, D.P.; Musgrove, M.T.; Ingram, K.D.; Hinton, J.A. Microbiological impact of spray washing broiler carcasses using different chlorine concentrations and water temperatures. *Poultry Sci.* **2005**, *84*, 1648–1652. [CrossRef]
9. Jung, Y.; Jang, H.; Guo, M.; Gao, J.; Matthews, K.R. Sanitizer efficacy in preventing cross-contamination of heads of lettuce during retail crisping. *Food Microbiol.* **2017**, *64*, 179–185. [CrossRef]
10. Northcutt, J.; Smith, D.; Ingram, K.D.; Hinton, J.A.; Musgrove, M. Recovery of bacteria from broiler carcasses after spray washing with acidified electrolyzed water or sodium hypochlorite solutions. *Poultry Sci.* **2007**, *86*, 2239–2244. [CrossRef]
11. Nou, X.; Luo, Y. Whole-leaf wash improves chlorine efficacy for microbial reduction and prevents pathogen cross-contamination during fresh-cut lettuce processing. *J. Food Sci.* **2010**, *75*, 283–290. [CrossRef] [PubMed]
12. USDA-FSIS. Draft FSIS compliance guideline for Controlling Salmonella and Campylobacter in raw poultry. 2010. Available online: [https://www.fsis.usda.gov/wps/wcm/connect/1d562776-ebfb-4ea5-a19b-e994776a02c6/Compliance\\_Guideline\\_Controlling\\_Salmonella\\_Poultry.pdf?MOD=AJPERES](https://www.fsis.usda.gov/wps/wcm/connect/1d562776-ebfb-4ea5-a19b-e994776a02c6/Compliance_Guideline_Controlling_Salmonella_Poultry.pdf?MOD=AJPERES) (accessed on 1 August 2019).
13. Jun, W.; Guo, Y.C.; Ning, L.I. Prevalence and risk assessment of *Campylobacter jejuni* in chicken in China. *Biomed. Environ. Sci.* **2013**, *26*, 243–248.
14. Moore, A.; Nannapaneni, R.; Kiess, A.; Sharma, C.S. Evaluation of USDA approved antimicrobials on the reduction of *Salmonella* and *Campylobacter* in ground chicken frames and their effect on meat quality. *Poultry Sci.* **2017**, *96*, 2385–2392. [CrossRef] [PubMed]

15. Lee, N.Y.; Park, S.Y.; Kang, I.S.; Ha, S.D. The evaluation of combined chemical and physical treatments on the reduction of resident microorganisms and *Salmonella* Typhimurium attached to chicken skin. *Poultry Sci.* **2013**, *93*, 208–215. [[CrossRef](#)]
16. Parveen, S.; Taabodi, M.; Schwarz, J.G.; Oscar, T.P.; Harter-Dennis, J.; White, D.G. Prevalence and antimicrobial resistance of *Salmonella* recovered from processed poultry. *J. Food Prot.* **2007**, *70*, 2466–2472. [[CrossRef](#)]
17. Reiter, M.G.R.; Fiorese, M.L.; Moretto, G.; López, M.C.; Jordano, R. Prevalence of *Salmonella* in a poultry slaughterhouse. *J. Food Prot.* **2007**, *70*, 1723–1725. [[CrossRef](#)]
18. Thomson, J.E.; Bailey, J.S.; Cox, N.A.; Posey, D.A.; Carson, M.O. *Salmonella* on broiler carcasses as affected by fresh water input rate and chlorination of chiller water. *J. Food Prot.* **1979**, *42*, 954–955. [[CrossRef](#)]
19. Minami, A.W.; Chaicumpa, C.N.; Manas, S.; Samosornsuk, S.; Monden, K.; Takeshi, S.; Kawamoto, M. Prevalence of foodborne pathogens in open markets and supermarkets in Thailand. *Food Control.* **2010**, *21*, 221–226. [[CrossRef](#)]
20. Wang, W.; Li, M.; Li, Y. Modeling the thermoultrasound inactivation of *Vibrio parahaemolyticus* in raw peeled shrimps. *J. Food Prot.* **2013**, *76*, 1712–1718. [[CrossRef](#)]
21. Yang, H.; Wang, S.; Li, Y.; Johnson, M.G. Predictive models for the survival/death of *Campylobacter jejuni* and *Salmonella* Typhimurium in poultry scalding and chilling. *J. Food Sci.* **2010**, *67*, 1836–1843. [[CrossRef](#)]
22. NY/T1174-2006. Quality Management Practice for Broiler Slaughtering (ICS: 67.120.20 X18, issued by the Standardization Administration of the People's Republic of China). Available online: <http://down.foodmate.net/standard/sort/5/10836.html> (accessed on 10 May 2018).
23. Saeed, M.O.; Azizli, K.; Isa, M.H.; Bashir, M.J.K. Application of ccd in rsm to obtain optimize treatment of pome using fenton oxidation process. *J. Water Process Eng.* **2015**, *8*, e7–e16. [[CrossRef](#)]
24. Mohana, S.; Shrivastava, S.; Divecha, J.; Madamwar, D. Response surface methodology for optimization of medium for decolorization of textile dye direct black 22 by a novel bacterial consortium. *Bioresour. Technol.* **2008**, *99*, 562–569. [[CrossRef](#)] [[PubMed](#)]
25. Bezerra, M.A.; Santelli, R.E.; Oliveira, E.P.; Villar, L.S.; Escalera, L.A. Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta* **2008**, *76*, 965–977. [[CrossRef](#)] [[PubMed](#)]
26. Xiao, X.; Wang, W.; Zhang, X.; Zhang, J.; Liao, M.; Yang, H.; Li, Y. Modeling the reduction of *Salmonella* spp. on chicken breasts and wingettes during scalding for QMRA of the poultry supply chain in China. *Microorganisms* **2019**, *7*, 165. [[CrossRef](#)] [[PubMed](#)]
27. Liao, X.; Li, Q.; Yang, X.; Zhang, W.; Li, W. Multiobjective optimization for crash safety design of vehicles using stepwise regression model. *Struct. Multidiscip. Optim.* **2008**, *35*, 561–569. [[CrossRef](#)]
28. Ross, T. Indices for performance evaluation of predictive models in food microbiology. *J. Appl. Microbiol.* **1996**, *81*, 501–508. [[CrossRef](#)]
29. Wang, W.; Li, M.; Fang, W.; Pradhan, A.K.; Li, Y. A predictive model for assessment of decontamination effects of lactic acid and chitosan used in combination on *Vibrio parahaemolyticus* in shrimps. *Int. J. Food Microbiol.* **2013**, *167*, 124–130. [[CrossRef](#)]
30. Ramirez-Hernandez, A.; Brashears, M.M.; Sanchez-Plata, M.X. Efficacy of lactic acid, lactic acid-acetic acid blends, and peracetic acid to reduce *Salmonella* on chicken parts under simulated commercial processing conditions. *J. Food Prot.* **2018**, *81*, 17–24. [[CrossRef](#)]
31. Xie, L.; Shang, W.; Liu, C.; Zhang, Q.; Sunter, G.; Hong, J.; Zhou, X. Mutual association of broad bean wilt virus 2 VP37-derived tubules and plasmodesmata obtained from cytological observation. *Sci. Rep.* **2016**, *6*, 21552. [[CrossRef](#)]
32. Yang, H.; Li, Y.; Johnson, M.G. Survival and death of *Salmonella* Typhimurium and *Campylobacter jejuni* in processing water and on chicken skin during poultry scalding and chilling. *J. Food Prot.* **2001**, *64*, 770–776. [[CrossRef](#)]
33. Lillard, H.S. Factors affecting the persistence of *Salmonella* during the processing of poultry. *J. Food Prot.* **1989**, *52*, 829–832. [[CrossRef](#)] [[PubMed](#)]
34. Nagel, G.M.; Bauermeister, L.J.; Bratcher, C.L.; Singh, M.; McKee, S.R. *Salmonella* and *Campylobacter* reduction and quality characteristics of poultry carcasses treated with various antimicrobials in a post-chill immersion tank. *Int. J. Food Microbiol.* **2013**, *165*, 281–286. [[CrossRef](#)] [[PubMed](#)]
35. Russell, S.M.; Axtell, S. The effect of monochloramine versus chlorine on pathogenic, indicator, and spoilage bacteria associated with broiler chicken carcasses: A model, pilot scale, and industrial study. *J. Food Prot.* **2005**, *68*, 758–763. [[CrossRef](#)] [[PubMed](#)]

36. Tamblyn, K.C.; Conner, D.E. Bactericidal activity of organic acids in combination with transdermal compounds against *Salmonella* Typhimurium attached to broiler skin. *Food Microbiol.* **1997**, *14*, 477–484. [[CrossRef](#)]
37. Luo, Y.; Nou, X.; Millner, P.; Zhou, B.; Shen, C.; Yang, Y.; Shelton, D. A pilot plant scale evaluation of a new process aid for enhancing chlorine efficacy against pathogen survival and cross-contamination during produce wash. *Int. J. Food Microbiol.* **2012**, *158*, 133–139. [[CrossRef](#)] [[PubMed](#)]
38. Munther, D.; Wu, J. Enhanced surveillance on food-borne disease outbreaks: Dynamics of cross-contamination in biocidal wash procedure. *J. Theor. Biol.* **2013**, *321*, 28–35. [[CrossRef](#)]
39. Mead, G.C.; Hudson, W.R.; Hinton, M.H. Use of a marker organism in poultry processing to identify sites of cross-contamination and evaluate possible control measures. *Brit. Poultry Sci.* **1994**, *35*, 345–354. [[CrossRef](#)]
40. Lillard, H.S. Effect on broiler carcasses and water of treating chiller water with chlorine or chlorine dioxide. *Poultry Sci.* **1980**, *59*, 1761–1766. [[CrossRef](#)]
41. FAO/WHO. *Risk assessment of Campylobacter spp. in broiler chickens: Technical Report*; WHO Library: Geneva, Switzerland, 2009; Available online: [https://apps.who.int/iris/bitstream/handle/10665/43731/9789241547369\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/43731/9789241547369_eng.pdf) (accessed on 10 August 2019).
42. Hayama, Y.; Yamamoto, T.; Kasuga, F.; Tsutsui, T. Simulation model for *Campylobacter* cross-contamination during poultry processing at slaughterhouses. *Zoonoses Public Health* **2011**, *58*, 399–406. [[CrossRef](#)]
43. Chen, Y.; Jackson, K.M.; Chea, F.P.; Schaffner, D.W. Quantification and variability analysis of bacterial cross-contamination rates in common food service tasks. *J. Food Prot.* **2001**, *64*, 72–80. [[CrossRef](#)]
44. Jensen, D.A.; Friedrich, L.M.; Harris, L.J.; Danyluk, M.D.; Schaffner, D.W. Cross-contamination of *Escherichia coli* O157: H7 between lettuce and wash water during home-scale washing. *Food Microbiol.* **2015**, *46*, 428–433. [[CrossRef](#)] [[PubMed](#)]
45. Ross, T.; Dalgaard, P.; Tienungoon, S. Predictive modelling of the growth and survival of *Listeria* in fishery products. *Int. J. Food Microbiol.* **2000**, *62*, 231–245. [[CrossRef](#)]
46. Sharma, C.S.; Ates, A.; Joseph, P.; Nannapaneni, R.; Kiess, A. Reduction of *Salmonella* in skinless chicken breast fillets by lauric arginate surface application. *Poultry Sci.* **2013**, *92*, 1419–1424. [[CrossRef](#)] [[PubMed](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).