



Article

Gut Anaerobes Capable of Chicken Caecum Colonisation

Tereza Kubasova ¹, Miloslava Kollarcikova ¹, Magdalena Crhanova ¹, Daniela Karasova ¹, Darina Cejkova ¹, Alena Sebkova ¹, Jitka Matiasovicova ¹, Marcela Faldynova ¹, Frantisek Sisak ¹, Vladimir Babak ¹, Alexandra Pokorna ², Alois Cizek ^{2,3} and Ivan Rychlik ^{1,*}

¹ Veterinary Research Institute, 621 00 Brno, Czech Republic; kbasova@vri.cz (T.K.); kollarcikova@vri.cz (M.K.); crhanova@vri.cz (M.C.); karasova@vri.cz (D.K.); cejkova@vri.cz (D.C.); asebkova@vri.cz (A.S.); matiasovicova@vri.cz (J.M.); faldynova@vri.cz (M.F.); sisak@vri.cz (F.S.); babak@vri.cz (V.B.)

² Department of Infectious Diseases and Microbiology, Faculty of Veterinary Medicine, University of Veterinary and Pharmaceutical Sciences Brno, 612 42 Brno, Czech Republic; pokornaa@vfu.cz

³ Central European Institute of Technology (CEITEC), University of Veterinary and Pharmaceutical Sciences Brno, 612 42 Brno, Czech Republic; cizeka@vfu.cz

* Correspondence: rychlik@vri.cz; Tel.: +420 533331201

Received: 29 October 2019; Accepted: 18 November 2019; Published: 21 November 2019

Abstract: Chicks in commercial production are highly sensitive to enteric infections and their resistance can be increased by administration of complex adult microbiota. However, it is not known which adult microbiota members are capable of colonising the caecum of newly hatched chicks. In this study, we therefore orally inoculated chicks with pure cultures of 76 different bacterial isolates originating from chicken caecum on day 1 of life and determined their ability to colonise seven days later. The caecum of newly hatched chickens could be colonised by bacteria belonging to phyla Bacteroidetes, Proteobacteria, Synergistetes, or Verrucomicrobia, and isolates from class Negativicutes (phylum Firmicutes). On the other hand, we did not record colonisation with isolates from phyla Actinobacteria and Firmicutes (except for Negativicutes), including isolates from families Lachnospiraceae, Ruminococcaceae, Erysipelotrichaceae, and Lactobacillaceae. Representatives of genera commonly used in probiotics such as *Lactobacillus*, *Enterococcus*, or *Bacillus* therefore did not colonise the chicken intestinal tract after a single dose administration. Following challenge with *Salmonella enterica* serovar Enteritidis, the best protecting isolates increased the chicken's resistance to *S. Enteritidis* only tenfold, which, however, means that none of the tested individual bacterial isolates on their own efficiently protected chicks against *S. Enteritidis*.

Keywords: caecum; chicken; oral inoculation; colonisation; *Salmonella*; Bacteroidetes; Firmicutes

1. Introduction

In animal species where parents raise their offspring, parents alone act as an important source of gut microbiota. This is also the case for chickens, who evolved for millions of years to be hatched in a nest in contact with a hen. However, contact between hen and chicks has been interrupted in commercial production, and the intestinal tract of commercially hatched chicks is gradually colonised from environmental sources only [1–4]. However, if the chicks are provided microbiota from a hen experimentally, they can be colonised by adult-type microbiota from the very first days of life [5,6].

The specific development of chick gut microbiota in commercial production is a reason why commercially hatched chicks are highly sensitive to infections, such as pathogenic *Escherichia coli*, *Clostridium perfringens*, or *Salmonella*, while chicks inoculated with faecal microbiota of adult hens are resistant to these infections [5,7–9]. This phenomenon, known as competitive exclusion, is well

established in chickens [10]. However, administration of complex competitive exclusion microbiota from adult hens to newly hatched chickens, though effective in preventing infections, is not widely accepted. The reason is that such products may contribute to the spread of yet unknown pathogens. In agreement, we have shown that representatives of Campylobacteriales are present at a higher abundance in the chicks inoculated with the caecal content of donor hens than in the microbiota of the donor hen itself [6]. This issue can be avoided if products consisting of defined bacterial strains were used [11]. In poultry, such products contain mostly *Lactobacilli*, *Enterococci*, or *Bacilli* [12–14], although there are reports questioning their real protective effect against infections [15,16]. Whether this is due to the low microbial complexity of products consisting of one or a few strains or whether this is caused by inappropriate strain selection is not known.

Repeatedly confirmed efficacy of competitive exclusion products together with accumulating knowledge on chicken microbiota composition led us to systematic culture of individual chicken gut microbiota members [17,18]. Because previous studies indicated that not all bacterial species present in the faeces of adult hens efficiently colonise the chicken caecum [5,6], in this study, we tested pure cultures of 76 chicken gut anaerobes for their ability to colonise the caecum of newly hatched chicks. Using this approach, we addressed whether the tested bacterial isolates could (i) efficiently colonise the chicken caecum during the first week of life, and (ii) protect chicks against *Salmonella enterica* serovar Enteritidis infection. Conclusions were rather unexpected. The caecum of newly hatched chickens could be populated only with Gram-negative bacteria belonging to phyla Bacteroidetes, Proteobacteria, Synergistetes, or Verrucomicrobia, and Gram-positive isolates from class Negativicutes. On the other hand, chickens could not be colonised experimentally with Gram-positive isolates from phyla Actinobacteria or Firmicutes (except for those from class Negativicutes), including *Lactobacilli*, *Enterococci*, and *Bacilli*, which are commonly used as probiotics in chickens [12–14].

2. Materials and Methods

2.1. Ethics Statement

The handling of animals in the study was performed in accordance with current Czech legislation (Animal Protection and Welfare Act No. 246/1992 Coll. of the Government of the Czech Republic). The specific experiments were approved by the Ethics Committee of the Veterinary Research Institute followed by the Committee for Animal Welfare of the Ministry of Agriculture of the Czech Republic on 31 March 2016 (permit number 4/2016) and on 15 January 2018 (permit number MZe1922).

2.2. Bacterial Isolates

Seventy-six chicken gut anaerobes characterised previously [17] were used for oral inoculation of chicks on the day of hatching. Of these, 54 were used for oral inoculation of chickens as individual cultures. An additional 15 isolates were tested both as individual cultures and as a part of defined mixtures, and the remaining 7 isolates were tested only as a part of defined mixtures (Table S1).

2.3. Chicken Inoculation with Gut Anaerobes and *S. Enteritidis* Challenge

In all experiments, newly hatched male ISA Brown chicks were obtained from a local hatchery on the day of hatching. In total, 542 chicks were used in this study. Of these, 474 were inoculated with one of the tested isolates or their mixtures, and 68 chicks served as noninoculated controls across the whole study. The whole study was accomplished in 10 different experimental batches performed from 2016 to 2018 (Table S2). Chicks were reared in plastic boxes with free access to water and feed in air-conditioned rooms with a controlled light and temperature regime and with filtered air supply. Temperature was set to 30 °C during the first week of life and to 28 °C in the second week of life. The light regime was set to 24 h light in the first week of life and 22 h of light during the second week of

life. The same standard starter feed formula for raising chickens during the first days of life was provided to chicks in all experiments (Table S3).

Chicks in experimental groups were orally inoculated on day 1 of life with 100 μ L of a particular anaerobe (approx. 10^7 CFU) resuspended in prereduced anaerobically sterilised (PRAS) solution to $OD_{600nm} = 1$ (PRAS solution composition - 0.1 g magnesium sulfate heptahydrate, 0.2 g monobasic potassium phosphate, 0.2 g potassium chloride, 1.15 g dibasic sodium phosphate, 3.0 g sodium chloride, 1.0 g sodium thioglycolate, 0.5 g L-cysteine, 1000 mL distilled water; final pH 7.5 \pm 0.2 at 25 °C). The cultures were washed from Wilkins–Chalgren agar (WCHA) after 3 days of incubation under anaerobic conditions at 37 °C as described previously [17]. Chicks in control groups were kept under the same conditions, but in separate rooms and without any treatment on day 1. On day 8 of life, half of the chicks in each group were euthanized, and caecal contents were frozen at -20 °C to examine caecal microbiota composition. In addition, sections of caecal tissue were collected in RNALater (Qiagen, Hilden, Germany) and stored at -80 °C prior to RNA purification. The remaining chicks were orally infected with *S. Enteritidis*, and the experiment was terminated 4 days postinfection.

2.4. Oral Inoculation with Defined Mixtures of Gut Anaerobes

In addition, selected gut anaerobes were also tested in mixtures. The isolates were grown separately and were mixed in an anaerobic cabinet Bactron600 (Sheldon Manufacturing Inc., Cornelius, OR, USA) to form the inoculum just prior to administration. In the first experiment, the chicks were inoculated with a mixture of bacteria cultured for 10 days (instead of the routine 3-day culture), testing whether stationary phase cells or spore-enriched cultures might affect their ability to colonise (see Table S1 for the composition of L10 and C10 mixtures). Next, we tested whether incubation of spore forming bacteria under nutrient-limited conditions might increase their potential to colonise chicks. Therefore, seven isolates were grown for 3 days on WCHA agar and thereafter resuspended either in sterile distilled water or sterile PRAS solution for 2 days before mixing and using as inoculum (Clost mixture in Table S1). The rest of these experiments were performed exactly as described above for individual isolates, i.e., newly hatched chicks were inoculated on day 1 of life with the mixture, half of the inoculated chicks were euthanised on day 8 to check for colonisation, and the second half of chicks was challenged with *S. Enteritidis*. Four days later, i.e., when the chicks were 12 days old, the experiment was terminated, caecal contents were collected, and *Salmonella* counts were determined.

In the last experiment with defined mixtures, we tested whether inoculation of chicks older than one day and initially precolonised with mostly Gram-negative bacteria might enable Gram-positive isolates to successfully colonise. The chicks were therefore inoculated with a mixture designated as BVL (Table S1) on day 1, followed by Clost mixture on day 8 of life.

2.5. *Salmonella Enteritidis* Challenge

S. Enteritidis challenge was performed orally with 1×10^7 CFU *S. Enteritidis* 147 spontaneously resistant to nalidixic acid in 0.1 mL inoculum [19]. Four days after infection, the chicks were euthanised under chloroform anesthesia by cervical dislocation, and during necropsy, 0.5 g of caecum was collected to enumerate *S. Enteritidis*. In addition, sections of caecal tissue were collected in RNALater (Qiagen, Hilden, Germany) and stored at -80 °C prior to RNA purification.

2.6. Chicken Gene Expression

The chicken inflammatory response in the caecum was determined by quantitative reverse transcribed PCR (Qiagen, Hilden, Germany) quantifying the expression of extracellular fatty acid binding protein (ExFABP) as described previously [5]. This gene was selected due to its high basal expression in the caecum of chickens and also high induction following *S. Enteritidis* challenge [20].

2.7. Sequencing of V3/V4 Region of 16S rRNA Genes

Caecal contents were homogenised in a MagNALyzer (Roche, Prague, Czech Republic). The DNA was then extracted using a QIAamp DNA Stool Mini Kit according to the manufacturer's instructions (Qiagen, Hilden, Germany), and the DNA concentration was determined spectrophotometrically. DNA samples were diluted to 5 ng/mL and were used as a template in PCR with forward primer 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-MID-GT-cctacggnggcwgcag-3' and reverse primer 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-MID-GT gactachvgggtatctaacc-3'.

The sequences in italics served for index and adapter ligation whereas the sequences in low case letters allowed the amplification over the V3/V4 region of 16S rRNA genes. MID sequences represent different sequences of 5, 6, 7, or 9 base pairs in length, which were used to identify individual samples within the sequencing groups. PCR amplification was performed using a HotStarTaq Plus MasterMix kit (Qiagen, Hilden, Germany), and the resulting PCR products were purified using AMPure beads (Beckman Coulter, Prague, Czech Republic). In the next step, the concentration of PCR products was determined spectrophotometrically, the DNA was diluted to 100 ng/μL, and groups of 14 PCR products with different MID sequences were indexed with the same indices using Nextera XT Index Kit following the manufacturer's instructions (Illumina, Cambridge, UK). Prior to sequencing, the concentration of differently indexed samples was determined using a KAPA Library Quantification Complete kit (Kapa Biosystems, Boston, MA, USA). All indexed samples were diluted to 4 ng/μL, and 20 pM phiX DNA was added to the final concentration of 5% (v/v). Sequencing was performed using MiSeq Reagent Kit v3 and MiSeq apparatus according to the manufacturer's instructions (Illumina, Cambridge, UK).

Quality trimming of the raw reads was performed using TrimmomaticPE v0.32 [21] with the following parameters: Window size of 4 with 15 as average quality and minimal reads length 150 bp. The FASTQ files generated after quality trimming were uploaded into QIIME software [22]. Forward and reverse sequences were joined with minimum 8 bp overlap. In the next step, chimeric sequences were predicted by the slayer algorithm implemented in QIIME and excluded from subsequent analysis. The resulting sequences were then classified by RDP Seqmatch with an OTU (operational taxonomic units) discrimination level set to 97%. Principal coordinate analysis (PCoA) implemented in QIIME was used for data visualisation.

2.8. Statistics

To identify to which OTU an anaerobe used for inoculation belonged, a set of 76 "artificial" samples, each containing a single sequence of an anaerobe used for the inoculation, was included in the QIIME calculation. Partial 16S rRNA sequences of each anaerobe were thus assigned to particular OTUs, and the OTU abundance was compared in inoculated and control chickens by the nonparametric Mann–Whitney U test. In addition to statistical significance, differentially abundant OTUs must have been present in at least 0.1% average abundance in the microbiota of inoculated chicks, and the difference in their abundance in inoculated and control chicks must have been 10-fold or higher. The inflammatory response to colonisation with the tested anaerobes and to *S. Enteritidis* infection in inoculated and control chicks was evaluated by a Kruskal–Wallis test, followed by Dunn's post hoc test. The likelihood of the ability to colonise and protect against *S. Enteritidis* challenge was tested by Chi2-test. In all cases, comparisons with *p* values lower than 0.05 were considered significant.

3. Results

3.1. Inoculation of Newly Hatched Chickens with Individual Bacterial Isolates

Altogether, 76 isolates were used for oral inoculation of chicks on day 1 of life. Using PCoA for visualisation of gut microbiota composition in inoculated and control chicks on day 8 of life, the majority of inoculated chicks clustered together with noninoculated controls, i.e., isolates used for

their inoculation likely did not colonise (Figure 1A). When the abundance of particular OTU representing the isolates used for oral inoculation was compared in inoculated and control chickens by the Mann–Whitney U test, only 25 isolates were capable of colonisation of the chicken caecum during the first week of life. These included 18 isolates belonging to phylum Bacteroidetes, 3 isolates belonging to phylum Firmicutes class Negativicutes, two isolates of *Desulfovibrio* (phylum Proteobacteria), and *Akkermansia* and *Cloacibacillus* belonging to phyla Verrucomicrobia and Synergistetes, respectively (Table 1). Except for Negativicutes, we did not record colonisation of chicks seven days after inoculation with any isolate belonging to phylum Firmicutes and the families Lactobacillaceae, Lachnospiraceae, Ruminococcaceae, or Erysipelotrichaceae.

Although we tested each bacterial isolate only in 3–5 chicks, altogether, 59 out of 69 chicks inoculated with isolates belonging to phylum Bacteroidetes were efficiently colonised, while not a single chick out of 110 inoculated with any isolate belonging to phylum Firmicutes was colonised (except for those belonging to class Negativicutes). We also failed with successful colonisation of the caecum of newly hatched chicks with four tested isolates belonging to phylum Actinobacteria (Figure 1C). Unweighted PCoA analysis showed that successful colonisation was of a low effect on the composition of other microbiota members as colonised, noncolonised, and control chicks were randomly distributed in the plot (Figure 1B). The apparent separation of some noncolonised chicks from colonised chicks was caused by minor experiment-to-experiment variation, similar to a previous report [23] (Figure S1).




Figure 1. Gut microbiota composition in the caecum of newly hatched chicks. (a) Weighted principal coordinate analysis (PCoA) indicating gut anaerobes capable of colonisation. Chicks inoculated with isolates not capable of colonisation (dark blue dots) clustered together with control chicks (red dots). Dots outside this cluster represent chicks that were inoculated with isolates capable of caecum colonisation (light blue dots), and these are identified by the numbers of their An codes (see Table S1). Green dots—chicks inoculated with different defined mixtures. Not all isolates capable of colonisation can be seen in panel (a) because the projection of some successfully colonised chicks along PC3 resulted in an overlap with the cluster of control chicks. (b) According to unweighted PCoA, successful colonisation did not affect composition of the rest of the microbiota. The same colour coding is used in panels (a) and (b). (c). Cluster alignment based on the whole gene sequence of 16S rRNA genes and ability to colonise the chicken caecum during the first week of life. Isolates with a

in these mixtures. In the second experiment, Clost mixture was prepared by resuspension and incubation of individual isolates in nutrient-limited PRAS solution or in water for two days at room temperature in an anaerobic cabinet (see Table S1 for the composition of Clost mixture). However, even after such treatment, none of the seven isolates present in the Clost mixture colonised the chicken caecum.

To test the possibility that the caecum of newly hatched chicks was not permissive for Clostridiales due to the presence of residual oxygen, the chicks were first inoculated with BVL mixture containing bacteria mostly capable of colonisation (see Table S1 for the composition of BVL mixture), followed by inoculation with Clost mixture on day 8 of life. However, even in this experiment, none of the isolates present in the Clost mixture colonised the chicken caecum when checked on day 15 of life.

3.3. Chicken Response to Colonisation with Tested Anaerobes

If the tested isolates were commensals with probiotic potential, these should not have induced an inflammatory response characterised by increased expression of ExFABP [20]. Although ExFABP expression varied among the groups inoculated with different anaerobes, it was never induced to the levels induced by *S. Enteritidis* (compare γ -axis scaling in Figure 2A,B). Despite this, significantly higher ExFABP expression than in control chicks was recorded in the chicks inoculated with *Bacteroides clarus* An43, *Olsenella uli* An270, and *Enorma timonensis* An5 (Figure 2A). However, because this happened in a single experiment and two of these species did not even colonise the chicken caecum, we concluded that the ExFABP expression was induced by other factor(s) and not by the isolates used for inoculation.




Figure 2. Extracellular fatty acid binding protein (ExFABP) expression in the caecum of inoculated and challenged chickens. (a) ExFABP expression in the caecum of 8-day-old chicks inoculated with different anaerobes on day 1 of life. Groups of chicks were inoculated on the day of hatching, and seven days later, expression of ExFABP was determined in the caecum of inoculated chicks by quantitative RT-PCR. Light-blue columns: ExFABP expression in the chicks that were successfully colonised by indicated gut anaerobes. Dark blue columns: ExFABP expression in the chicks that were inoculated with anaerobes that did not colonise the chicken caecum. Red columns highlight ExFABP expression in control chicks included in each experiment batch (except for the experiment with L10

and C10 mixtures in which no noninoculated chicks were included). *—Significantly different expression from appropriate control (Cont Exp7) by Kruskal–Wallis test followed by Dunn’s post hoc test ($p < 0.05$). (b) ExFABP expression in the caecum of 12-day-old chicks inoculated with different anaerobes on day 1 of life and infected with *S. Enteritidis* on day 8. If the isolate used for inoculation on day 1 exhibited a protective effect, lower expression was expected in the colonised chicks compared to the controls. Light-blue columns: ExFABP expression in the chicks that were successfully colonised by the tested isolate. Dark blue columns: ExFABP expression in the chicks that were inoculated by isolates that did not colonise the chicken caecum. Red columns: ExFABP expression in noninoculated control chicks included in each experiment batch (except for the experiment with L10 and C10 mixtures in which no noninoculated control chicks were included). *—Significantly different expression from the appropriate control (Cont Exp2) by Kruskal–Wallis test followed by Dunn’s post hoc test ($p < 0.05$).

3.4. Chicken Response to *S. Enteritidis* Infection

Although *S. Enteritidis* infection increased ExFABP expression, there were only four significant differences in ExFABP expression between inoculated and control chickens (Figure 2B). Lower ExFABP expression in response to *S. Enteritidis* infection than in control chicks was recorded in chicks inoculated with [*Clostridium*] *saccharolyticum* An14, *Flavonifractor plautii* An52, or [*Eubacterium*] *cylindroides* An64. However, none of these anaerobes efficiently colonised chicken caecum. On the other hand, ExFABP expression in *Butyricimonas paravirosa* An62-colonised and *S. Enteritidis*-challenged chicks was significantly higher than in controls (Figure 2B). Because *Butyricimonas paravirosa* An62 was present in the caecum at the time of infection, this bacterium therefore did not protect chickens against *S. Enteritidis* infection and corresponding inflammatory response.

Evaluation of *S. Enteritidis* counts in inoculated and control chicks was complicated by the fact that 3–5 chicks per group were challenged, and in some cases, other bacterial species overgrew *S. Enteritidis* on Xylose-Lysine-Deoxycholate (XLD) plates (Figure S2). To characterise the interaction of *S. Enteritidis*, chicken host, and gut microbiota, we therefore combined *S. Enteritidis* counts in the caecum and expression of ExFABP. When these two parameters were plotted against each other, the groups of chicks were divided into those with *S. Enteritidis* counts higher or lower than 10^7 CFU/g of caecal content and exhibiting a high or low inflammatory response according to ExFABP expression. A threshold value for ExFABP expression was defined as 10 times higher than average ExFABP expression prior to *S. Enteritidis* infection. This resulted in the formation of four different clusters (Figure 3A).




Figure 3. *Salmonella* counts and chicken inflammatory response. (a) Each dot represents a group of chickens inoculated with a particular anaerobe whose position is defined by average *S. Enteritidis* counts (\log_{10} CFU/g) and inflammatory response determined by ExFABP expression. Cluster 1 (red dots) is formed by the chickens that were inoculated with gut anaerobes that did not protect them against *S. Enteritidis* challenge (high *S. Enteritidis* count and high ExFABP expression). Cluster 2

showing that Lachnospiraceae and Ruminococcaceae belong among the first colonisers [1,3,4]. Moreover, we also did not record Lachnospiraceae and Ruminococcaceae colonisation when these were administered to chicks already precolonised with *Bacteroides* species. We also considered that there could be a mutual dependence among bacterial species from families Lachnospiraceae, Ruminococcaceae, and Erysipelotrichaceae, but even when we inoculated these isolates in mixtures, we did not record efficient colonisation. Perhaps the time window allowing initial colonisation of Lachnospiraceae, Ruminococcaceae, or Erysipelotrichaceae is available earlier during the chicken life, shortly after hatching. This would be consistent with the fact that commercial, newly hatched chickens in fact represent a heterogeneous group of chicks that may differ in their age up to 48 h. However, we also consider the hypothesis that bacterial species from these families may not permanently colonise the chicken caecum and have to be continuously supplied from the environment.

No matter which of the explanations is/are correct, Gram-positive bacteria seem to be less suitable for probiotic products intended for a single dose administration, as these did not efficiently colonise chicken caecum under our experimental conditions. Instead, such products should consist mainly of Gram-negative bacteria, capable of efficient colonisation of chicken caecum, sources of which are limited in an environment. However, such products will have to consist of multiple carefully selected bacterial strains because individual strains only weakly protected chickens against pathogens like *Salmonella*.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Figure S1: Unweighted PCoA visualisation of gut microbiota composition in the caecum of newly hatched chicks. Figure S2: *Salmonella* counts in the caecum of 12-day-old chicks inoculated with different anaerobes on day 1 of life and challenged with *S. Enteritidis* on day 8. Table S1: OTU table of OTUs forming at least 0.05% of all microbiota in at least one of the tested samples. Table S2: List of strains used in this study. Table S3: Bacterial strains and numbers of chicks used in different experiments. Table S4: Feed composition, light, and temperature regime used during rearing.

Author Contributions: J.M., A.P., and A.C. were responsible for anaerobic culture. T.K., M.K., M.C., and D.K. purified DNA from caecal samples and nutrient broths and performed 16S rRNA sequencing. D.C., T.K., and V.B. were responsible for data analysis and their graphical presentation. A.S., M.F., and F.S. were responsible for animal experiments and infections. I.R. designed the study and wrote the manuscript. All authors read and approved the final manuscript.

Funding: This work has been supported by RVO0518 of the Czech Ministry of Agriculture and project CZ.02.1.01/0.0/0.0/16_025/0007404 of the Czech Ministry of Education. A.C. and A.P. were supported by CEITEC 2020 project LQ1601 of the Czech Ministry of Education under the National Sustainability Programme II and QK1810462 project of the Czech Ministry of Agriculture. The funders had no role in the study design, data collection, and analysis, the decision to publish, or preparation of the manuscript.

Acknowledgments: Authors would like to thank Peter Eggenhuizen for English language corrections.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Awad, W.A.; Mann, E.; Dzieciol, M.; Hess, C.; Schmitz-Esser, S.; Wagner, M.; Hess, M. Age-related differences in the luminal and mucosa-associated gut microbiome of broiler chickens and shifts associated with *Campylobacter jejuni* infection. *Front. Cell. Infect. Microbiol.* **2016**, *6*, 154. doi:10.3389/fcimb.2016.00154.
2. Ranjitkar, S.; Lawley, B.; Tannock, G.; Engberg, R.M. Bacterial succession in the broiler gastrointestinal tract. *Appl. Environ. Microbiol.* **2016**, *82*, 2399–2410. doi:10.1128/AEM.02549-15.
3. Videnska, P.; Sedlar, K.; Lukac, M.; Faldynova, M.; Gerzova, L.; Cejkova, D.; Sisak, F.; Rychlik, I. Succession and replacement of bacterial populations in the caecum of egg laying hens over their whole life. *PLoS ONE* **2014**, *9*, e115142, doi:10.1371/journal.pone.0115142.
4. Videnska, P.; Rahman, M.M.; Faldynova, M.; Babak, V.; Matulova, M.E.; Prukner-Radovic, E.; Krizek, I.; Smole-Mozina, S.; Kovac, J.; Szmolka, A.; et al. Characterization of egg laying hen and broiler fecal microbiota in poultry farms in Croatia, Czech Republic, Hungary and Slovenia. *PLoS ONE* **2014**, *9*, e110076, doi:10.1371/journal.pone.0110076.

5. Varmuzova, K.; Kubasova, T.; Davidova-Gerzova, L.; Sisak, F.; Havlickova, H.; Sebkova, A.; Faldynova, M.; Rychlik, I. Composition of gut microbiota influences resistance of newly hatched chickens to *Salmonella* enteritidis infection. *Front. Microbiol.* **2016**, *7*, 957, doi:10.3389/fmicb.2016.00957.
6. Polansky, O.; Sekelova, Z.; Faldynova, M.; Sebkova, A.; Sisak, F.; Rychlik, I. Important metabolic pathways and biological processes expressed by chicken cecal microbiota. *Appl. Environ. Microbiol.* **2015**, *82*, 1569–1576, doi:10.1128/AEM.03473-15.
7. Kaldhusdal, M.; Schneitz, C.; Hofshagen, M.; Skjerve, E. Reduced incidence of *Clostridium perfringens*-associated lesions and improved performance in broiler chickens treated with normal intestinal bacteria from adult fowl. *Avian Dis.* **2001**, *45*, 149–156.
8. Salvat, G.; Lalande, F.; Humbert, F.; Lahellec, C. Use of a competitive exclusion product (Broilact) to prevent *Salmonella* colonization of newly hatched chicks. *Int. J. Food Microbiol.* **1992**, *15*, 307–311.
9. Nisbet, D. Defined competitive exclusion cultures in the prevention of enteropathogen colonisation in poultry and swine. *Antonie Van Leeuwenhoek* **2002**, *81*, 481–486.
10. Rantala, M.; Nurmi, E. Prevention of the growth of *Salmonella infantis* in chicks by the flora of the alimentary tract of chickens. *Br. Poult. Sci.* **1973**, *14*, 627–630, doi:10.1080/00071667308416073.
11. Stavric, S. Defined cultures and prospects. *Int. J. Food Microbiol.* **1992**, *15*, 245–263.
12. Baldwin, S.; Hughes, R.J.; Hao Van, T.T.; Moore, R.J.; Stanley, D. At-hatch administration of probiotic to chickens can introduce beneficial changes in gut microbiota. *PLoS ONE* **2018**, *13*, e0194825, doi:10.1371/journal.pone.0194825.
13. Beirao, B.C.B.; Ingberman, M.; Favaro, C., Jr.; Mesa, D.; Bittencourt, L.C.; Fascina, V.B.; Caron, L.F. Effect of an *Enterococcus faecium* probiotic on specific IgA following live *Salmonella* Enteritidis vaccination of layer chickens. *Avian Pathol.* **2018**, *47*, 325–333, doi:10.1080/03079457.2018.1450487.
14. Mazanko, M.S.; Gorlov, I.F.; Prazdnova, E.V.; Makarenko, M.S.; Usatov, A.V.; Bren, A.B.; Chistyakov, V.A.; Tutelyan, A.V.; Komarova, Z.B.; Mosolova, N.I.; et al. *Bacillus* probiotic supplementations improve laying performance, egg quality, hatching of laying hens, and sperm quality of roosters. *Probiotics Antimicrob. Proteins* **2018**, *10*, 367–373, doi:10.1007/s12602-017-9369-4.
15. Weinack, O.M.; Snoeyenbos, G.H.; Soerjadi-Liem, A.S. Further studies on competitive exclusion of *Salmonella typhimurium* by lactobacilli in chickens. *Avian Dis.* **1985**, *29*, 1273–1276.
16. Adler, H.E.; DaMassa, A.J. Effect of ingested *Lactobacilli* on *Salmonella infantis* and *Escherichia coli* and on intestinal flora, pasted vents, and chick growth. *Avian Dis.* **1980**, *24*, 868–878.
17. Medvecky, M.; Cejkova, D.; Polansky, O.; Karasova, D.; Kubasova, T.; Cizek, A.; Rychlik, I. Whole genome sequencing and function prediction of 133 gut anaerobes isolated from chicken caecum in pure cultures. *BMC Genom.* **2018**, *19*, 561, doi:10.1186/s12864-018-4959-4.
18. Crhanova, M.; Karasova, D.; Juricova, H.; Matiasovicova, J.; Jahodarova, E.; Kubasova, T.; Seidlerova, Z.; Cizek, A.; Rychlik, I. Systematic culturomics shows that half of chicken caecal microbiota members can be grown in vitro except for two lineages of *Clostridiales* and a single lineage of *Bacteroidetes*. *Microorganisms* **2019**, *7*, 496, doi:10.3390/microorganisms7110496.
19. Methner, U.; Barrow, P.A.; Gregorova, D.; Rychlik, I. Intestinal colonisation-inhibition and virulence of *Salmonella phoP*, *rpoS* and *ompC* deletion mutants in chickens. *Vet. Microbiol.* **2004**, *98*, 37–43.
20. Matulova, M.; Rajova, J.; Vlasatikova, L.; Volf, J.; Stepanova, H.; Havlickova, H.; Sisak, F.; Rychlik, I. Characterization of chicken spleen transcriptome after infection with *Salmonella enterica* serovar Enteritidis. *PLoS ONE* **2012**, *7*, e48101, doi:10.1371/journal.pone.0048101.
21. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **2014**, *30*, 2114–2120. doi:10.1093/bioinformatics/btu170.
22. Caporaso, J.G.; Kuczynski, J.; Stombaugh, J.; Bittinger, K.; Bushman, F.D.; Costello, E.K.; Fierer, N.; Pena, A.G.; Goodrich, J.K.; Gordon, J.I.; et al. QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **2010**, *7*, 335–336. doi:10.1038/nmeth.f.303.
23. Stanley, D.; Hughes, R.J.; Geier, M.S.; Moore, R.J. Bacteria within the gastrointestinal tract microbiota correlated with improved growth and feed conversion: Challenges presented for the identification of performance enhancing probiotic bacteria. *Front. Microbiol.* **2016**, *7*, 187, doi:10.3389/fmicb.2016.00187.
24. Impey, C.S.; Mead, G.C.; George, S.M. Competitive exclusion of salmonellas from the chick caecum using a defined mixture of bacterial isolates from the caecal microflora of an adult bird. *J. Hyg. Lond.* **1982**, *89*, 479–490.

25. Kubasova, T.; Kollarcikova, M.; Crhanova, M.; Karasova, D.; Cejkova, D.; Sebkova, A.; Matiasovicova, J.; Faldynova, M.; Pokorna, A.; Cizek, A.; et al. Contact with adult hen affects development of caecal microbiota in newly hatched chicks. *PLoS ONE* **2019**, *14*, e0212446, doi:10.1371/journal.pone.0212446.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).