

Table S1. Pollutant concentration: List of the compounds detected in the contaminated soil, in which *Trichoderma lixii* MUT3171 was isolated. The concentration is expressed in mg/kg of dry soil. TPH: total petroleum hydrocarbons.

Pollutant	Concentration
2-metilnaphthalene	50.00
1-metilnaphthalene	28.26
naphthalene	62.81
acenaphtilene	0.44
acenaphthene	10.96
fluorene	11.83
phenanthrene	96.28
anthracene	6.67
fluoranthene	51.01
pyrene	32.91
benzo(a)anthracene	10.11
chrysene	5.97
benzo(b+I)fluoranthene	3.96
benzo(k)fluoranthene	1.50
benzo(a)pyrene	1.93
indeno[1,2,3-cd]pyrene	1.24
dibenzo(ac+ah)anthracene	0.32
benzo(ghi)perylene	1.02
dibenzo(al)pyrene	0.50
dibenzo(ae)pyrene	0.14
dibenzo(ai)pyrene	0.05
dibenzo(ah)pyrene	<0.01
TPHs	377.91

Table S2. Strains used for comparison: List of *Trichoderma* species used in the comparative analysis, with relative GenBank accessions and references.

Species	Accession	References
<i>Trichoderma guizhouense</i> NJAU 4742	GCA_002022785.1	[1]
<i>Trichoderma harzianum</i> CBS 226 95	GCA_003025095.1	[2]
<i>Trichoderma harzianum</i> TR274	GCA_002838845.1	[3]
<i>Trichoderma harzianum</i> T6776	GCA_000988865.1	[4]
<i>Trichoderma arundinaceum</i> IBT40837	GCA_003012105.1	[5]
<i>Trichoderma asperellum</i> CBS 433 97	GCA_003025105.1	[2]
<i>Trichoderma citrinoviride</i> TUCIM 6016	GCA_003025115.1	[2]
<i>Trichoderma gamsii</i> T6085	GCA_001481775.2	[2]
<i>Trichoderma longibrachiatum</i> ATCC 18648	GCA_003025155.1	[2]
<i>Trichoderma parareesei</i> CBS 125925	GCA_001050175.1	[6]
<i>Trichoderma reesei</i> QM6a	GCA_000167675.2	[7]
<i>Trichoderma virens</i> Gv29 8	GCA_000170995.2	[8]
<i>Trichoderma harzianum</i> TR1	GCA_002894145.1	[9]

References

- Meng, X.; Miao, Y.; Liu, Q.; Ma, L.; Guo, K.; Liu, D.; Ran, W.; Shen, Q. TgSWO from *Trichoderma guizhouense* NJAU4742 promotes growth in cucumber plants by modifying the root morphology and the cell wall architecture. *Microb. Cell Factories* **2019**, *18*, 148–15, doi:10.1186/s12934-019-1196-8.
- Druzhinina, I.S.; Chenthama, K.; Zhang, J.; Atanasova, L.; Yang, D.; Miao, Y.; Rahimi, M.J.; Grujic, M.; Cai, F.; Pourmehdi, S.; et al. Massive lateral transfer of genes encoding plant cell wall-degrading enzymes to the mycoparasitic fungus *Trichoderma* from its plant-associated hosts. *PLoS Genet.* **2018**, *14*, e1007322, doi:10.1371/journal.pgen.1007322.
- Steindorff, A.S.; Ramada, M.H.S.; Coelho, A.S.; Miller, R.N.G.; Pappasjr, G.J.; Ulhoa, C.J.; Noronha, E.F. Identification of mycoparasitism-related genes against the phytopathogen *Sclerotinia sclerotiorum* through transcriptome and expression profile analysis in *Trichoderma harzianum*. *BMC Genom.* **2014**, *15*, 204, doi:10.1186/1471-2164-15-204.
- Baroncelli, R.; Zapparata, A.; Piaggeschi, G.; Sarrocco, S.; Vannacci, G. Draft whole-genome sequence of *Trichoderma gamsii* T6085, a promising biocontrol agent of *Fusarium* head blight on wheat. *Genome Announc.* **2016**, *4*, 4, doi:10.1128/genomea.01747-15.
- Proctor, R.H.; McCormick, S.P.; Kim, H.-S.; Cardoza, R.E.; Stanley, A.M.; Lindo, L.; Kelly, A.; Brown, D.W.; Lee, T.; Vaughan, M.M.; et al. Evolution of structural diversity of trichothecenes, a family of toxins produced by plant pathogenic and entomopathogenic fungi. *PLOS Pathog.* **2018**, *14*, e1006946, doi:10.1371/journal.ppat.1006946.
- Baroncelli, R.; Piaggeschi, G.; Fiorini, L.; Bertolini, E.; Zapparata, A.; Pè, M.E.; Sarrocco, S.; Vannacci, G. Draft whole-genome sequence of the biocontrol agent *Trichoderma harzianum* T6776. *Genome Announc.* **2015**, *3*, 3, doi:10.1128/genomea.00647-15.
- Yang, D.; Pomraning, K.R.; Kopchinskiy, A.; Aghcheh, R.K.; Atanasova, L.; Chenthama, K.; Baker, S.E.; Zhang, R.; Shen, Q.; Freitag, M.; et al. Genome sequence and annotation of *Trichoderma parareesei*, the ancestor of the cellulase producer *Trichoderma reesei*. *Genome Announc.* **2015**, *3*, doi:10.1128/genomea.00885-15.
- Martinez, D.; Berka, R.M.; Henrissat, B.; Saloheimo, M.; Arvas, M.; Baker, S.E.; Chapman, J.; Chertkov, O.; Coutinho, P.M.; Cullen, D.; et al. Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocrea jecorina*). *Nat. Biotechnol.* **2008**, *26*, 553–560, doi:10.1038/nbt1403.
- Kubicek, C.P.; Herrera-Estrella, A.; Seidl, V.; A Martinez, D.; Druzhinina, I.S.; Thon, M.R.; Zeilinger, S.; Casas-Flores, S.; Horwitz, B.A.; Mukherjee, P.K.; et al. Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*. *Genome Biol.* **2011**, *12*, R40, doi:10.1186/gb-2011-12-4-r40.
- Gardiner, D.; Vos, C.; Kazan, K.; Harvey, P. Commonwealth Scientific and Industrial Research Organisation, Queensland, Australia. Unpublished Work. **2018**.