




Article

Isoleucine Enhanced the Function of the Small Intestinal Mucosal Barrier in Weaned Piglets to Alleviate Rotavirus Infection

Rongkun Zhao ^{1,†}, Changsheng Jiang ^{1,†}, Yuchen Yuan ¹, Shen Zhang ¹, Ahmed H. Ghonaim ^{2,3} , Chuanyan Che ¹, Xiaojin Li ¹, Mengmeng Jin ¹, Erhui Jin ¹ , Xiangfang Zeng ⁴, Shenghe Li ^{1,*}  and Man Ren ^{1,*}

- ¹ Anhui Provincial Key Laboratory of Animal Nutritional Regulation and Health, College of Animal Science, Anhui Science and Technology University, Fengyang 233100, China; rongkunzhao@126.com (R.Z.); jiangcs@ahstu.edu.cn (C.J.); 18855065915@163.com (Y.Y.); 15755797521@163.com (S.Z.); checy@ahstu.edu.cn (C.C.); lixj@ahstu.edu.cn (X.L.); jinmm@ahstu.edu.cn (M.J.); jineh@ahstu.edu.cn (E.J.)
- ² National Key Laboratory of Agricultural Microbiology, College of Animal Sciences and Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China; a.ghonaim@webmail.hzau.edu.cn
- ³ Desert Research Center, Cairo 11435, Egypt
- ⁴ State Key Laboratory of Animal Nutrition, College of Animal Science and Technology, China Agricultural University, Beijing 100193, China; zengxf@cau.edu.cn
- * Correspondence: lish@ahstu.edu.cn (S.L.); renman@yeah.net (M.R.)
- † These authors contributed equally to this work.

Simple Summary: Rotavirus is a major cause of diarrhea in young children and animals, especially piglets, leading to substantial economic losses in the global pig industry. Isoleucine, a branched-chain amino acid, plays an important role in regulating nutrient metabolism and has been shown to improve diarrhea. This study aimed to evaluate the effects of isoleucine supplementation on the mucosal immune barrier of the small intestine in rotavirus-infected weaned piglets. The results showed that the addition of isoleucine to the diet mitigated the effects of rotavirus infection on the intestinal morphology and the mucosal barrier, as well as the physiological function of weaned piglets, and improved the antioxidant and immune functions of these piglets to a certain extent. These findings offer valuable insights contributing to a deeper understanding of the role of isoleucine.



Citation: Zhao, R.; Jiang, C.; Yuan, Y.; Zhang, S.; Ghonaim, A.H.; Che, C.; Li, X.; Jin, M.; Jin, E.; Zeng, X.; et al.

Isoleucine Enhanced the Function of the Small Intestinal Mucosal Barrier in Weaned Piglets to Alleviate Rotavirus Infection. *Animals* **2024**, *14*, 3146.

<https://doi.org/10.3390/ani14213146>

Academic Editor: Christophe R. Casteleyn

Received: 12 September 2024

Revised: 20 October 2024

Accepted: 21 October 2024

Published: 2 November 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Background: Rotavirus (RV) is a major cause of diarrhea in young children and animals, especially piglets, leading to substantial economic losses in the global pig industry. Isoleucine (Ile), a branched-chain amino acid, plays an important role in regulating nutrient metabolism and has been shown to improve diarrhea. This study aimed to evaluate the effects of Ile supplementation on the mucosal immune barrier of the small intestine in RV-infected weaned piglets. Methods: Forty-eight 21-day-old weaned piglets were randomly divided into three dietary treatments (each treatment was subdivided into two groups, eight replicates per group), with 0%, 0.5%, or 1% Ile added for 15 days, and then, one group from each treatment was challenged with RV. Results: The results showed that 1% Ile added to the diet promoted the healthy development of the intestinal mucosa. Ile could restore the reduced villus height in the ileum and the goblet cell number in the duodenum and ileum to normal levels, improving the intestinal epithelial tight junctions in RV-infected piglets. Additionally, Ile increased the activity of lipase, amylase, and sucrase, as well as superoxide dismutase (SOD) and glutathione (GSH), along with the expression of SIgA, DEFβ1, and DEFβ2 in parts of the small intestine. Conclusions: The addition of Ile to the diet mitigated the effects of RV infection on intestinal morphology and mucosal barrier function, as well as the physiological functions of weaned piglets, and improved the antioxidant and immune functions of the piglets to some extent. These findings offer valuable insights, contributing to a deeper understanding of the role of Ile in supporting intestinal health.

Keywords: isoleucine; rotavirus; small intestine; mucosal barrier; weaned piglet

1. Introduction

Rotavirus (RV) is a major pathogen that causes acute enteritis in young children and animals, including nursing and weaned piglets [1–3]. It is globally prevalent, infecting many species, including humans [4], pigs, cattle [5], dogs [6], and sheep [7]. Some RV genotypes have also shown zoonotic potential [8,9]. Piglets are highly susceptible to rotavirus infection, which is associated with high mortality rates [10]. RV adversely impacts the health and growth of pigs in the global pig industry, especially in developing countries, resulting in significant economic losses [1,11]. Currently, rehydration (oral and intravenous) remains the main treatment strategy for rotavirus gastroenteritis (RVGE), and no specific drugs are available [12]. The virus is transmitted via the fecal–oral route and results in the destruction of mature small intestinal enterocytes, leading to villus atrophy [8,13], which subsequently causes malabsorption and diarrhea [14]. However, the effects of RV infection on the function and structure of the porcine intestinal mucosal barrier are not fully understood.

Isoleucine (Ile) is a branched-chain amino acid that is essential for humans and pigs, and it can regulate protein metabolism, fat deposition, and glucose utilization in the body [15,16]. A previous study has shown that Ile supplementation can alleviate acute diarrhea caused by malnutrition in children [17]. Additionally, Ile serves as an important energy substrate and is essential for immune protein synthesis, thereby improving immune function [18]. Previous studies have reported that Ile can improve growth performance, feed intake, feed utilization, relative gut length, and intestinal morphology in hybrid catfish [19]. Furthermore, it can regulate the intestinal immune function by increasing the excretion of β -defensins via the activation of the Sirt1/ERK/90RSK signaling pathway [20,21]. Thus, isoleucine has the potential to prevent pathogen invasion by boosting immunity.

In recent years, the potential of nutrients to prevent and treat viral infections has attracted the attention of many scientists, given their multiple beneficial effects on the host, such as maintaining normal intestinal function, protecting the intestinal tract from pathogenic bacteria, and modulating immunity [22–24]. However, studies on the effects of dietary Ile on the intestinal tract of enteropathogenic virus-infected piglets are limited. Given the multiple beneficial effects of Ile, the present study aimed to preliminarily investigate its effects on the intestinal tissue structure, digestive capacity, and immune function in RV-infected weaned piglets.

2. Materials and Methods

2.1. Animal Ethics Statement

All animal experiments were approved by the Animal Ethics Committee of Anhui Science and Technology University, under protocol number AK2023013. All experimental procedures were carried out in strict accordance with the “Guidelines for the Care and Use of Test Animals” of Anhui Province.

2.2. Animal Experiment

The rotavirus used in this study was a tissue-culture-adapted Ohio State University strain (ATCC#VR-893). RV preparation and virus titer determination [tissue culture infective dose 50 (TCID₅₀)] were conducted as described previously [25].

Forty-eight healthy Duroc \times Landrace \times Yorkshire weaned barrows (6.88 ± 0.54 kg) at 21 days old were randomly divided into 6 groups (eight replicates in each group). Each piglet was individually housed in a metabolic cage with a self-feeder and a nipple watering device, in rooms with moderate temperature control.

Piglets had free access to feed and water during the trial period. The control (CON) and RV groups received the basal diet, while the 0.5% Ile and RV + 0.5% Ile groups were fed the basal diet containing 0.5% Ile. The 1% Ile and RV + 1% Ile groups were fed a basal diet containing 1% Ile. On the 15th day of feeding, all piglets were orogastrically inoculated with 5 mL of 100 mmol/L sodium bicarbonate (NaHCO₃) solution. Twenty minutes later, the infected group was orogastrically inoculated with 5 mL of RV (1×10^6 TCID₅₀/mL),

and the uninfected group was orogastrically inoculated with 5 mL of 0.9% sodium chloride (NaCl) solution. At 5 days post infection (dpi), all piglets were euthanized, and samples were collected. The animal experimental design and treatments are shown in Table 1.

Table 1. The animal experimental design.

Ile supplementation (%) Groups	RV (–)			RV (5 mL)		
	0 CON	0.5 0.5% Ile	1 1% Ile	0 RV	0.5 RV + 0.5% Ile	1 RV + 1% Ile

The basal diets met the NRC (National Research Council) standard nutritional requirements for 7–11 kg pigs [26]. In this study, 0, 0.5%, and 1% L-isoleucine was added to the basal diet. Alanine was included for isonitrogen treatment. The L-Ile and alanine in the diets replaced the corn starch in the basal diet in equal amounts. The specific ingredient composition and nutritional levels are shown in Table 2.

Table 2. Basal ration composition and nutrient levels.

Items	Groups		
	CON/ RV	0.5% Ile/ RV + 0.5% Ile	1% Ile/ RV + 0.5% Ile
Ingredients (%)			
Corn	61.27	61.11	60.95
Peeled soybean meal, 49% CP	6	6	6
Soybean protein concentrate, 65%	10	10	10
Fish meal, 64.5% CP	5	5	5
Deproteinized whey powder, 5.5% CP	5	5	5
Soybean oil	3	3	3
Corn starch	3.3	3.3	3.3
Glucose	2	2	2
Limestone	0.7	0.7	0.7
CaHPO ₄	1	1	1
NaCl	0.3	0.3	0.3
Chloride choline	0.15	0.15	0.15
Vitamin premix ¹	0.03	0.03	0.03
Mineral premix ²	0.3	0.3	0.3
L-lysine-HCL	0.43	0.43	0.43
L-threonine	0.08	0.08	0.08
L-tryptophan	0.4	0.4	0.4
DL-methionine	0.06	0.06	0.06
L-valine	0.3	0.3	0.3
L-isoleucine	0	0.5	1
L-alanine	0.68	0.34	0
Total	100	100	100
Calculated nutrient levels (%)			
Crude protein	19.79	19.79	19.79
Calcium	0.8	0.8	0.8
Total phosphorus	0.65	0.65	0.65
Available phosphorus	0.48	0.48	0.48
Na	0.21	0.21	0.21
Cl	0.33	0.33	0.33
Lysine	1.35	1.35	1.35
Threonine	0.79	0.79	0.79
Tryptophan	0.22	0.22	0.22
Methionine	0.39	0.39	0.39

Table 2. Cont.

Items	Groups		
	CON/ RV	0.5% Ile/ RV + 0.5% Ile	1% Ile/ RV + 0.5% Ile
Valine	0.86	0.86	0.86
Isoleucine	0.72	1.22	1.72
Leucine	1.56	1.56	1.56
Leucine/isoleucine	217	128	91
Metabolizable energy (MJ/kg)	14.05	14.05	14.05

¹ The vitamin premix provides the following per kg of diet: VA 9000 IU, VD3 3000 IU, VE 20.0 IU, VK3 3.0 mg, VB1 1.5 mg, VB2 4.0 mg, VB6 3.0 mg, VB12 0.2 mg, niacin 30.0 mg, pantothenic 15.0 mg, folic acid 0.75 mg, and biotin 0.1 mg. ² The mineral premix provides the following per kg of diet: Fe 100 mg, Cu 6 mg, Zn 100 mg, Mn 4 mg, I 0.14 mg, and Se 0.3 mg.

2.3. Sample Collection

At 5 dpi, all piglets were euthanized via intracardiac injection of Na pentobarbital (50 mg/kg of body weight) and exsanguinated. The abdomen was immediately opened and flushed with 0.9% NaCl solution. Approximately 2 cm from the midpoint of the duodenum, jejunum and ileum were collected, and the fatty tissue on the surface of the intestine was peeled off to avoid extrusion during the sampling process. The samples were then put into 4% paraformaldehyde fixative for preservation. Simultaneously, the intestinal tubes were cut longitudinally, and the contents were flushed out with 0.9% NaCl solution, and then, the mucosa of the duodenum, jejunum, and ileum were scraped and put into liquid nitrogen for preservation.

2.4. Intestinal Morphology

Paraffin-embedded fixed intestinal segments were sectioned into 5 µm sections. Sections were deparaffinized in xylene and then hydrated in a concentration gradient of 100%, 95%, 85%, and 70% alcohol. The sections were then stained with hematoxylin and eosin using an H&E stain kit (Solarbio, Beijing, China) according to the manufacturer's protocol to assess tissue morphology. Examination of sections was performed at least in 10 well-oriented intact villi and their associated crypts using light microscopy. Goblet cells were stained using a periodic acid–Schiff (PAS) stain kit (Solarbio, Beijing, China) according to the manufacturer's instructions. Villus height and crypt depth were measured, and the number goblet cells was counted by Image J 1.53a software.

2.5. Western Blot

The intestinal mucosal protein was extracted by RIPA lysis buffer with 1% PMSF (Beyotime, Haimen, China) at 4 °C. The protein concentration was measured using a BCA Protein Assay Kit (Thermo Fisher, Waltham, MA, USA). A total of 20 µg of protein from each sample was loaded onto a 10% SDS-PAGE gel and then transferred to a polyvinylidene difluoride (PVDF) membrane. To prevent nonspecific binding, the PVDF membranes were blocked using 5% bovine serum albumin (BSA) at room temperature for 2 h. Next, the membranes were incubated overnight at 4 °C with the appropriate primary antibody. Following incubation, the membranes were washed three times with Tris-buffered saline with Tween 20 (TBST) and then incubated with either HRP-linked goat anti-rabbit or HRP-linked goat anti-mouse antibodies at room temperature for 2 h. Visualization was performed using an ECL detection kit (Solarbio, Beijing, China). The antibodies used for the Western blot are listed in Table 3.

Table 3. Antibodies used in this study.

Antibodies	Supplier	Catalog No.
β -Actin Monoclonal antibody	Proteintech, Wuhan, China	66009-1-Ig
Claudin1 Polyclonal antibody		13050-1-AP
Occludin Polyclonal antibody		27260-1-AP
MUC-1 Rabbit pAb	Abclonal, Wuhan, China	A0333
HRP-conjugated Goat anti-Rabbit IgG (H + L)		AS014
HRP-conjugated Goat anti-Mouse IgG (H + L)		AS003

2.6. Digestive Enzyme and Antioxidant Activity Assessment

The activities of the intestinal mucosal lipase, amylase, sucrase, total antioxidant capacity (T-AOC), superoxide dismutase (SOD), and glutathione (GSH) were assessed using kits according to the manufacturer's protocol. All kits were purchased from Nanjing Jiancheng Bioengineering Institute (Table 4).

Table 4. Kits for antioxidant and digestive enzyme activity assays.

Items	Supplier	Catalog No.	Function
Total antioxidant capacity (T-AOC) assay kit (ABTS method)	Nanjing Jiancheng Bioengineer Institute, Nanjing, China	A015-2-1	Measures total antioxidant capacity
Superoxide dismutase (SOD) assay kit (WST-1 method)		A001-3-1	Measures the activity of superoxide dismutase
Glutathione peroxidase (GSH-PX) assay kit (colorimetric method)		A015-2-1	Measures the activity of glutathione peroxidase
Lipase (LPS) assay kit		A054-2-1	Measures the activity of lipase
α -Amylase (AMS) assay kit		C016-1-1	Measures the activity of amylase
Sucrase assay kit		A082-2-1	Measures the activity of sucrase

2.7. Cytokine Index Assessment

The expressions of interferon- γ (IFN- γ), secretory immunoglobulin A (SIgA), and defensins (DEF β) 1 and 2 were measured in the intestinal mucosa using kits following the manufacturer's protocol. All kits were purchased from Beijing Dakome Co., Beijing, China (Table 5).

Table 5. Kits for cytokine expression assays.

Items	Supplier	Catalog No.	Function
Porcine IFN- γ ELISA kit	Dakome, Beijing, China	DKM2552A	Measures the expression of interferon gamma
Porcine SIgA ELISA kit		DKM9851A	Measures the expression of secretory immunoglobulin A
Porcine DEF β 1 ELISA kit		DKM0031PA	Measures the expression of defensin beta 1
Porcine DEF β 2 ELISA kit		DKM0035PA	Measures the expression of defensin beta 2

2.8. Statistical Analysis

Data were analyzed by two-way analysis of variance (ANOVA), with model factors including the effects of L-isoleucine administration, RV infection, and their interaction. When significant effects were present, the Tukey test was used to compare means amongst treatments with significant effects. Data are presented as means \pm standard deviations (SDs). p values of <0.05 were considered statistically significant. All statistical analysis was performed using GraphPad Prism 8 software.

3. Results

3.1. Effects of Ile Supplementation on Small Intestinal Morphology in RV-Infected Piglets

To evaluate the effect of Ile on small intestinal morphology in RV-infected piglets, the histomorphology of the small intestine was assessed. As shown in Figure 1 and Table 6, compared with the CON group, 1% Ile increased the villus height ($p < 0.05$) and villus height/crypt depth ratio ($p < 0.05$) in the duodenum of RV-uninfected piglets. In contrast, compared with the CON group, RV infection significantly decreased the villus height/crypt depth ratio in the duodenum ($p < 0.05$), jejunum ($p < 0.05$), and ileum ($p < 0.05$), increased the crypt depth of the jejunum ($p < 0.05$), and obviously decreased the villus height of the

ileum ($p < 0.05$). However, compared with the RV group, 1% Ile significantly increased the villus height ($p < 0.05$) and villus height/crypt depth ratio ($p < 0.05$) in the ileum of RV-infected piglets. These data suggest that Ile could ameliorate the damage caused by RV infection to the histological structure of the small intestine.




Figure 1. Effects of Ile supplementation on small intestinal morphology in RV-infected piglets. Representative histologic HE-stained micrographs. Scale bars are 200 μ m.

Table 6. Effects of Ile supplementation on small intestinal morphology in RV-infected piglets (n = 8).

Items	RV−			RV+			p-Value		
	CON	0.5% Ile	1% Ile	CON	0.5% Ile	1% Ile	Ile	RV	Ile × RV
Duodenum									
VH, μm	363.08 ± 30.61 ^{bc}	424.15 ± 58.60 ^{ab}	465.15 ± 54.35 ^a	313.46 ± 28.87 ^c	367.85 ± 21.24 ^{bc}	392.71 ± 38.65 ^{abc}	<0.01	<0.01	0.81
CD, μm	268.16 ± 55.60	272.68 ± 42.56	250.91 ± 44.67	305.45 ± 31.8	274.62 ± 9.48	299.67 ± 22.47	0.74	0.07	0.49
VH-to-CD ratio	1.40 ± 0.27 ^b	1.56 ± 0.14 ^{ab}	1.87 ± 0.14 ^a	1.03 ± 0.10 ^c	1.34 ± 0.10 ^{bc}	1.32 ± 0.23 ^{bc}	<0.01	<0.01	0.21
Jejunum									
VH, μm	318.91 ± 67.11 ^{ab}	375.96 ± 48.74 ^a	320.94 ± 62.54 ^{ab}	239.90 ± 36.69 ^b	237.02 ± 29.47 ^b	308.89 ± 53.03 ^{ab}	0.23	<0.01	0.02
CD, μm	219.65 ± 51.15 ^c	235.69 ± 35.93 ^c	251.49 ± 38.37 ^{bc}	316.02 ± 66.53 ^{ab}	290.41 ± 43.67 ^{abc}	345.50 ± 26.61 ^a	0.14	<0.01	0.46
VH-to-CD ratio	1.46 ± 0.07 ^{ab}	1.60 ± 0.08 ^a	1.29 ± 0.25 ^b	0.78 ± 0.14 ^c	0.83 ± 0.14 ^c	0.91 ± 0.21 ^c	0.18	<0.01	0.02
Ileum									
VH, μm	330.68 ± 41.11 ^{ab}	346.27 ± 38.43 ^a	348.04 ± 54.77 ^a	223.58 ± 30.72 ^c	256.86 ± 62.47 ^{bc}	311.49 ± 13.57 ^{ab}	0.06	<0.01	0.21
CD, μm	224.92 ± 24.55	207.09 ± 12.53	211.05 ± 26.45	204.24 ± 45.95	223.58 ± 31.05	215.27 ± 14.97	0.98	>0.99	0.36
VH-to-CD ratio	1.47 ± 0.02 ^a	1.67 ± 0.14 ^a	1.64 ± 0.09 ^a	1.12 ± 0.14 ^b	1.14 ± 0.18 ^b	1.45 ± 0.12 ^a	<0.01	<0.01	0.02

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. VH, villus height; CD, crypt depth. ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

3.2. Effects of Ile Supplementation on Mucosal Barrier Function of the Small Intestine in RV-Infected Piglets

The primary function of goblet cells is to synthesize and secrete mucus, which plays an important role in inhibiting viral infection. Figure 2 and Table 7 show the periodic acid–Schiff (PAS) staining performed to observe the distribution and number of goblet cells in small intestinal tissue. The results showed that, compared with the CON group, RV infection decreased the number of goblet cells in the duodenum and ileum ($p < 0.05$). However, 0.5% and 1% Ile supplementation did not significantly affect the number of goblet cells in the three segments of the small intestine in RV-uninfected piglets ($p > 0.05$). Additionally, compared with the RV group, 1% Ile supplementation increased the number of goblet cells in the duodenum and ileum of RV-infected piglets ($p < 0.05$). Based on these results, it can be concluded that 1% Ile supplementation significantly improves the histological structure and increases the number of goblet cells in the small intestine of RV-infected piglets, while 0.5% Ile has minimal effect on RV-induced histological changes and goblet cell number. Therefore, the following studies focused solely on 1% Ile supplementation.

To further investigate the effect of isoleucine on the small intestinal mucosal barrier of RV-infected weaned piglets, Figure 3 and Table 8 illustrate the expression of the tight junction markers Claudin 1 and Occludin, as well as the mucin Muc 1, which is a component of intestinal mucus, determined by Western blot analysis. The results showed that compared with the CON group, RV infection significantly decreased the expression of Claudin 1 in the duodenum ($p < 0.05$) and jejunum ($p < 0.05$), decreased the expression of Occludin in the jejunum ($p < 0.05$), and decreased the expression of Muc 1 in both the duodenum ($p < 0.05$) and jejunum ($p < 0.05$). In addition, compared with the CON group, 1% Ile significantly increased the expression of Claudin 1 and Occludin in the duodenum ($p < 0.05$) and ileum ($p < 0.05$), as well as the expression of Muc 1 in the ileum ($p < 0.05$). Furthermore, compared with the RV group, 1% Ile increased the expression of Claudin 1 and Muc 1 in the duodenum ($p < 0.05$) and jejunum ($p < 0.05$) and enhanced the expression of Occludin in the jejunum ($p < 0.05$) of RV-infected piglets. These data indicate that Ile promotes the differentiation and mucus secretion of small intestinal goblet cells, improves intestinal epithelial tight junctions, and alleviates the impairment of small intestinal barrier function by RV infection to a certain extent.





Figure 2. Effects of Ile supplementation on mucosal barrier function of the small intestine in RV-infected piglets. Representative histologic PAS-stained micrographs. Scale bars are 200 μ m.

Table 7. Effects of Ile supplementation on goblet cell number of the small intestine in RV-infected piglets (n = 8).

Items	RV−			RV+			<i>p</i> -Value		
	CON	0.5% Ile	1% Ile	CON	0.5% Ile	1% Ile	Ile	RV	Ile × RV
Duodenum GCs number/villus	13.83 ± 2.79 ^a	15.00 ± 1.50 ^a	13.33 ± 1.75 ^a	6.83 ± 1.94 ^b	9.83 ± 1.47 ^b	10.00 ± 1.41 ^b	0.03	<0.01	0.06
Jejunum GCs number/villus	15.83 ± 3.37	17.40 ± 3.05	16.50 ± 3.94	13.00 ± 1.63	13 ± 2.37	13.83 ± 2.99	0.81	0.05	0.77
Ileum GCs number/villus	15.33 ± 0.82 ^a	14.83 ± 2.14 ^a	14.33 ± 3.14 ^a	6.17 ± 2.32 ^c	10.83 ± 1.17 ^b	9.33 ± 1.63 ^{bc}	0.06	<0.01	0.01

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. GC, goblet cell. ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

**Figure 3.** Effects of Ile supplementation on the mechanical barrier of the small intestinal mucosa in RV-infected weaned piglets. The expressions of Claudin 1, Occludin, and Muc 1 were measured in the duodenum (A), jejunum (B), and ileum (C) using Western blot analysis.**Table 8.** Effects of Ile supplementation on the mechanical barrier of the small intestinal mucosa in RV-infected weaned piglets (n = 8).

Items	RV−		RV+		<i>p</i> -Value		
	CON	1% Ile	CON	1% Ile	Ile	RV	Ile × RV
Duodenum							
Claudin 1	1.00 ± 0.06 ^b	1.37 ± 0.10 ^a	0.73 ± 0.03 ^c	1.13 ± 0.06 ^b	<0.01	<0.01	0.78
Occludin	1.00 ± 0.25 ^b	1.74 ± 0.10 ^a	1.01 ± 0.18 ^b	1.38 ± 0.20 ^{ab}	<0.01	0.14	0.12
Muc 1	1.00 ± 0.05 ^a	1.08 ± 0.04 ^a	0.39 ± 0.02 ^c	0.51 ± 0.03 ^b	<0.01	<0.01	0.30
Jejunum							
Claudin 1	1.00 ± 0.08 ^a	1.06 ± 0.03 ^a	0.33 ± 0.06 ^c	0.69 ± 0.03 ^b	<0.01	<0.01	<0.01
Occludin	1.00 ± 0.04 ^a	1.14 ± 0.10 ^a	0.55 ± 0.14 ^b	0.95 ± 0.08 ^a	<0.01	<0.01	0.05
Muc 1	1 ± 0.01 ^a	1.09 ± 0.05 ^a	0.47 ± 0.09 ^b	0.98 ± 0.10 ^a	<0.01	<0.01	<0.01
Ileum							
Claudin 1	1.00 ± 0.05 ^b	1.57 ± 0.16 ^a	0.81 ± 0.05 ^b	0.77 ± 0.03 ^b	<0.01	<0.01	<0.01
Occludin	1.00 ± 0.03 ^b	1.31 ± 0.09 ^a	1.07 ± 0.05 ^b	0.91 ± 0.10 ^b	0.12	0.01	<0.01
Muc 1	1.00 ± 0.12 ^b	1.42 ± 0.05 ^a	1.08 ± 0.10 ^b	1.08 ± 0.08 ^b	<0.01	0.04	<0.01

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. Data show the results of gray-scale value quantification, normalized by β-actin (relative). ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

3.3. Effects of Ile Supplementation on Digestive Enzymes of the Small Intestinal Mucosa in RV-Infected Piglets

Table 9 demonstrates the effects of dietary supplementation with Ile on the activity of digestive enzymes in the small intestinal mucosa in RV-infected piglets. Compared with the CON group, RV infection significantly decreased the activity of lipase ($p < 0.05$) in the duodenum, reduced the activity of amylase in the jejunum, and decreased the activity of sucrase in the duodenum, jejunum, and ileum ($p < 0.05$). Compared with the CON group, 1% Ile increased the activity of amylase and sucrase ($p < 0.05$) in the ileum of the RV-uninfected piglets. Additionally, compared with the RV group, 1% Ile could increase

the activity of lipase ($p < 0.05$) in the duodenum and the activity of sucrase ($p < 0.05$) in the ileum of the RV-infected piglets. These findings demonstrate that Ile was able to partially reverse the RV-induced reductions in the activity of digestive enzymes in the small intestinal mucosa.

Table 9. Effects of Ile supplementation on digestive enzymes of the small intestinal mucosa in the RV-infected weaned piglets ($n = 8$).

Items	RV−		RV+		<i>p</i> -Value		
	CON	1% Ile	CON	1% Ile	Ile	RV	Ile × RV
Duodenum							
Lipase, U/g prot	20.45 ± 3.14 ^a	20.70 ± 2.68 ^a	14.69 ± 2.33 ^b	19.63 ± 3.28 ^a	0.04	0.01	0.06
Amylase, U/mg prot	0.10 ± 0.03	0.11 ± 0.03	0.08 ± 0.04	0.08 ± 0.03	0.71	0.08	0.71
Sucrase, U/mg prot	67.97 ± 17.58 ^{ab}	78.80 ± 13.88 ^a	32.00 ± 18.59 ^c	49.27 ± 16.00 ^{bc}	0.05	<0.01	0.64
Jejunum							
Lipase, U/g prot	19.58 ± 1.68 ^{ab}	21.64 ± 2.03 ^a	17.07 ± 2.14 ^{bc}	16.10 ± 2.05 ^c	0.51	<0.01	0.08
Amylase, U/mg prot	0.12 ± 0.02 ^a	0.15 ± 0.02 ^a	0.07 ± 0.02 ^b	0.11 ± 0.04 ^{ab}	<0.01	<0.01	0.65
Sucrase, U/mg prot	145.82 ± 20.92 ^a	156.18 ± 28.54 ^a	109.97 ± 19.69 ^b	127.50 ± 37.65 ^{ab}	0.23	0.01	0.75
Ileum							
Lipase, U/g prot	14.23 ± 2.78 ^{ab}	16.61 ± 2.47 ^a	11.34 ± 1.11 ^b	14.49 ± 1.34 ^{ab}	<0.01	0.01	0.65
Amylase, U/mg prot	0.10 ± 0.01 ^b	0.14 ± 0.01 ^a	0.07 ± 0.01 ^c	0.10 ± 0.01 ^b	<0.01	<0.01	0.24
Sucrase, U/mg prot	101.24 ± 22.50 ^b	139.25 ± 26.20 ^a	54.52 ± 13.29 ^c	91.13 ± 17.06 ^b	<0.01	<0.01	0.93

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

3.4. Effects of Ile Supplementation on Antioxidant Capacity of the Small Intestinal Mucosa in RV-Infected Piglets

Rotavirus infection can induce oxidative stress [27]; meanwhile, isoleucine possesses antioxidative properties [28]. Therefore, we speculated that Ile may mitigate the oxidative stress induced by RV. Table 10 depicts the effects of Ile on the antioxidant expression of the small intestinal mucosa in RV-infected piglets. Compared with the CON group, the expression of GSH ($p < 0.05$) was significantly decreased in the ileum of the RV group. Compared with the CON group, 1% Ile supplementation increased the expression of T-AOC ($p < 0.05$) in the jejunum and GSH ($p < 0.05$) in the ileum of RV-uninfected piglets. During viral infection, 1% Ile supplementation significantly increased the expression of SOD ($p < 0.05$) in the duodenum and GSH ($p < 0.05$) in the jejunum of RV-infected piglets compared with the RV group. These data suggest that Ile could ameliorate the reduced antioxidant expression of the small intestine in piglets caused by RV infection.

Table 10. Effects of Ile supplementation on antioxidant expression of the small intestinal mucosa in RV-infected weaned piglets ($n = 8$).

Items	RV−		RV+		<i>p</i> -Value		
	CON	1% Ile	CON	1% Ile	Ile	RV	Ile × RV
Duodenum							
T-AOC, U/mg prot	0.87 ± 0.10	0.86 ± 0.11	0.85 ± 0.09	0.92 ± 0.06	0.36	0.54	0.23
SOD, U/mg prot	5.15 ± 1.04 ^b	7.35 ± 1.59 ^a	4.56 ± 0.49 ^b	7.11 ± 1.29 ^a	<0.01	0.33	0.68
GSH, U/mg prot	0.87 ± 0.54 ^{ab}	1.24 ± 0.36 ^a	0.72 ± 0.31 ^b	1.08 ± 0.16 ^{ab}	0.01	0.24	0.97
Jejunum							
T-AOC, U/mg prot	0.73 ± 0.12 ^b	0.97 ± 0.10 ^a	0.69 ± 0.13 ^b	0.81 ± 0.16 ^{ab}	0.01	0.28	0.83
SOD, U/mg prot	6.87 ± 1.31	6.41 ± 1.98	6.22 ± 2.44	8.00 ± 3.14	0.43	0.57	0.18
GSH, U/mg prot	1.92 ± 0.38 ^{ab}	2.14 ± 0.74 ^{ab}	1.37 ± 0.64 ^b	2.2 ± 0.52 ^a	0.03	0.31	0.21
Ileum							
T-AOC, U/mg prot	0.81 ± 0.10 ^b	0.92 ± 0.04 ^a	0.80 ± 0.05 ^b	0.82 ± 0.08 ^b	0.02	0.04	0.09
SOD, U/mg prot	3.19 ± 2.36 ^{ab}	4.86 ± 1.61 ^a	2.57 ± 0.47 ^b	2.46 ± 1.21 ^b	0.17	0.01	0.12
GSH, U/mg prot	1.46 ± 0.54 ^b	3.07 ± 0.74 ^a	0.46 ± 0.23 ^c	0.95 ± 0.56 ^{bc}	<0.01	<0.01	0.05

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. T-AOC, total antioxidant capacity; GSH, glutathione peroxidase; SOD, superoxide dismutase. ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

3.5. Effects of Ile Supplementation on Cytokines of Small Intestinal Mucosa in RV-Infected Piglets

During viral infection, the body enters an antiviral state by producing immune cytokines like interferons and immunoglobulins [29]. Notably, interferon gamma (IFN- γ) and secretory immunoglobulin A (SIgA) play important roles in intestinal mucosal immunity [30,31]. Table 11 shows the effects of Ile on the expression of IFN- γ and SIgA in the small intestinal mucosa of RV-infected piglets. Compared with the CON group, RV infection significantly increased the expression of IFN- γ ($p < 0.05$) in the jejunum and ileum. Compared with the RV group, 1% Ile supplementation increased the expression of IFN- γ ($p < 0.05$) in the duodenum of RV-infected piglets. Moreover, we found that the expression of SIgA in the RV+1% Ile group was higher than that of the other three groups, particularly in the duodenum ($p < 0.05$) and jejunum ($p < 0.05$) compared with the RV group. These findings reveal that Ile could modulate the small intestinal mucosal immune function following RV infection in weaned piglets.

Table 11. Effects of Ile supplementation on cytokine expression of the small intestinal mucosa in RV-infected weaned piglets (n = 8).

Items	RV−		RV+		Ile	p-Value	
	CON	1% Ile	CON	1% Ile		RV	Ile × RV
Duodenum							
IFN- γ, pg/mg prot	45.64 ± 12.56 ^b	46.72 ± 13.67 ^b	48.47 ± 5.51 ^b	69.94 ± 14.90 ^a	0.04	0.02	0.07
SIgA, μg/mg prot	2.31 ± 0.37 ^b	2.50 ± 0.58 ^b	2.08 ± 0.24 ^b	3.42 ± 0.73 ^a	<0.01	0.12	0.01
Jejunum							
IFN- γ, pg/mg prot	40.50 ± 11.7 ^b	48.60 ± 12.35 ^b	86.95 ± 11.70 ^a	82.00 ± 17.55 ^a	0.78	<0.01	0.25
SIgA, μg/mg prot	2.23 ± 0.73 ^b	2.36 ± 0.21 ^b	2.64 ± 0.83 ^b	3.70 ± 0.65 ^a	0.08	<0.01	0.20
Ileum							
IFN- γ, pg/mg prot	58.50 ± 13.95 ^b	59.85 ± 13 ^b	83.80 ± 16.65 ^a	86.35 ± 6.30 ^a	0.72	<0.01	0.91
SIgA, μg/mg prot	2.04 ± 0.29	2.50 ± 0.54	2.22 ± 0.63	2.79 ± 0.48	0.06	0.36	0.83

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. IFN- γ , interferon gamma; SIgA, secretory immunoglobulin A. ^{a,b} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

3.6. Effects of Ile Supplementation on Defensins of the Small Intestinal Mucosa in RV-Infected Piglets

Previous studies have reported that beta defensins play an important role in resisting multiple viral infections [32–34]. In this study, ELISA kits were used to evaluate the expression of porcine beta defensins 1 and 2 in the mucosa of the duodenum, jejunum, and ileum. As shown in Table 12, compared with the CON group, RV infection significantly decreased the expression of DEF β 1 ($p < 0.05$) in the jejunum and decreased the expression of DEF β 2 in the duodenum ($p < 0.05$). Additionally, compared with the CON group, 1% Ile supplementation increased the expression of DEF β 1 ($p < 0.05$) in the ileum of RV-uninfected piglets. Furthermore, compared with the RV group, 1% Ile significantly increased the expression of DEF β 1 ($p < 0.05$) in the jejunum and increased the expression of DEF β 2 ($p < 0.05$) in duodenum of RV-infected piglets. These data suggest that Ile could modulate the defensin expression in the small intestinal mucosa of weaned piglets following RV infection.

Table 12. Effects of Ile supplementation on defensin expression of the small intestinal mucosa in RV-infected weaned piglets (n = 8).

Items	RV−		RV+		Ile	p-Value	
	CON	1% Ile	CON	1% Ile		RV	Ile × RV
Duodenum							
DEFβ1, pg/mg prot	59.36 ± 8.80 ^{ab}	67.53 ± 8.72 ^a	49.88 ± 6.10 ^b	50.94 ± 7.07 ^b	0.16	<0.01	0.28
DEFβ2, pg/mg prot	68.09 ± 6.65 ^b	77.57 ± 6.43 ^a	54.64 ± 8.09 ^c	72.83 ± 7.31 ^{ab}	<0.01	0.01	0.15

Table 12. Cont.

Items	RV−		RV+		p-Value		
	CON	1% Ile	CON	1% Ile	Ile	RV	Ile × RV
Jejunum							
DEFβ1, pg/mg prot	32.29 ± 5.33 ^a	33.54 ± 4.19 ^a	23.36 ± 4.69 ^b	31.96 ± 4.84 ^a	0.02	0.01	0.08
DEFβ2, pg/mg prot	68.06 ± 9.57	70.39 ± 7.94	61.16 ± 4.77	69.06 ± 8.50	0.13	0.22	0.40
Ileum							
DEFβ1, pg/mg prot	18.29 ± 5.06 ^b	26.45 ± 6.01 ^a	18.80 ± 3.58 ^{ab}	20.01 ± 3.93 ^{ab}	0.03	0.14	0.09
DEFβ2, pg/mg prot	55.69 ± 7.73	60.28 ± 5.99	57.75 ± 9.10	61.15 ± 4.18	0.18	0.61	0.84

RV−, rotavirus-uninfected; RV+, rotavirus-infected; CON, L-alanine-supplemented diet; Ile, L-isoleucine-supplemented diet. DEFβ1, defensin beta 1; DEFβ2, defensin beta 2. ^{a,b,c} In the same row, values with different letter superscripts mean a significant difference ($p < 0.05$).

4. Discussion

Isoleucine, an essential functional branched-chain amino acid, plays a vital role in the regulation of energy homeostasis, nutrient metabolism (protein, carbohydrates, and lipid), intestinal health, and immunity in humans and animals [15,18]. In our preliminary experiments, the addition of 2.5 mg/mL isoleucine to drinking water improved the morphology and structure of rat intestinal mucosa [35]. In this study, we found that the addition of 1% Ile to the diet also significantly increased the duodenal villus height of weaned piglets and significantly elevated the jejunal villus height reduced by RV infection. The overall effect of adding 1% Ile was better than that of 0.5% Ile. The results reveal that the addition of Ile to the diet promoted intestinal villi growth and alleviated the negative effects of rotaviruses on the morphology and structure of the intestinal tract of weaned piglets to a certain extent.

In terms of small intestinal immunity, the intestinal mucosal barrier is the first line of defense against endogenous and exogenous stimuli and pathogenic bacteria. The mucus of the intestinal mucosal barrier is mainly formed and secreted by goblet cells [36]. There is increasing evidence that certain enteropathogenic viruses, such as porcine epidemic diarrhea virus [37], murine astrovirus [38], and human enterovirus 71 [39], can inhibit the formation of goblet cells and affect their function. For example, porcine delta coronavirus infection severely damages the intestinal mucosal barrier by activating the Notch signaling pathway and inhibiting goblet cell differentiation and mucus secretion [40]. Goblet cells and mucins play an important role in actively defending against rotavirus infection. The number of goblet cells was significantly reduced in the duodenum and jejunum of rotavirus-infected mice [41]. In addition, porcine epidemic diarrhea virus (PEDV) infection reduced the number of goblet cells in the small intestine of piglets and resulted in a small number of epithelial goblet cells being virus-positive, suggesting that PEDV infection possibly leads to an impaired mucus layer and increased susceptibility to secondary enteric bacterial infection [42]. In this study, we found that RV infection significantly reduced the number of goblet cells throughout the small intestinal segments of weaned piglets. Notably, Ile restored the goblet cell counts reduced by RV infection. These findings suggest that RV infection inhibited the mucus production by goblet cells in the small intestine, while Ile alleviated this negative effect.

Tight junctions consist of multiple transmembrane proteins that connect epithelial cells. Intestinal tight junction proteins such as Claudin and Occludin play important roles in the formation and function of the intestinal mechanical barrier [43,44]. In the intestine, the transmembrane mucin Muc1 is an important member of the physical barrier, influencing epithelial cell morphology and receptor function [45]. In this study, 1% Ile increased the expression of Claudin 1 in the duodenum and jejunum, Occludin in the jejunum, and Muc 1 in the duodenum and jejunum of RV-infected piglets. This may be due to the fact that Ile enhances the immunity of weaned piglets, inhibiting the invasion of RV into small intestine epithelial cells and alleviating the damage to the tight junction mechanical barrier caused by RV infection, thus ensuring normal barrier function.

Enzyme activity in the digestive tract is an important factor influencing intestinal health and digestibility [46]. Different breeds of pigs exhibit significant differences in

growth performance and small intestinal development, mainly due to differences in digestive enzymes and nutrient transporter proteins [47]. Various digestive enzymes are synthesized in the intestines and secreted into the intestinal lumen primarily by the pancreas, with their levels regulated by nutrient intake. A previous study mentioned that the duodenal instillation of Ile increases pancreatic secretion, especially α -amylase [48]. In addition, the activities of lipase and sucrase are key markers for evaluating intestinal epithelial development or maturation [49,50]. In this study, we found that isoleucine significantly increased the activity of amylase and sucrase in the ileum of the small intestines of the RV-uninfected weaned piglets. However, RV infection resulted in varying reductions in lipase, amylase, and sucrase activity. Furthermore, the 1% Ile-supplemented group with RV infection exhibited significantly increased duodenal lipase and ileal sucrase activity compared with the RV group. These results reveal that Ile has a modulating effect on small intestinal digestive enzymes and can attenuate the reduction in digestive enzyme activity caused by RV infection to a certain extent, although the specific mechanisms need further study.

Reactive oxygen species (ROS) play an important role in cellular signaling and homeostasis in vivo [51]. However, the excessive accumulation of ROS in organisms leads to the inactivation of proteins, enzymes, and phospholipids in cellular membranes and the activation of pro-inflammatory signaling pathways. The intestine is a primary target organ affected by ROS [52]. RV interactions with mitochondria lead to the increased production of ROS [27,53]. Thus, maintaining antioxidant capacity during RV infection is important for the inhibition of replication within viral vectors. In our previous study, we concluded that drinking water supplemented with an appropriate amount of Ile could improve serum antioxidant capacity in rats [54]. In this study, we found that Ile significantly increased the expression of T-AOC in the jejunum and the expression of GSH in the ileum of the uninfected group. Moreover, the addition of 1% Ile in the RV-infected group significantly increased the expression of SOD in the duodenum as well as the expression of GSH in the jejunum.

IFN- γ is a pro-inflammatory cytokine produced during intestinal inflammation and has an important role in the intestinal inflammatory response [55]. In rotavirus-infected piglets, the concentration of IFN- γ significantly increased compared with that in healthy piglets [29]. Additionally, the addition of L-Ile to the diet increased the levels of IFN- γ in the serum of RV-infected weaned piglets [56]. SIgA is a polymeric immunoglobulin composed of dimeric IgA, a J chain, and a secretory component, most of which are generated by the gut and exert effects in situ [31]. In this study, we found that RV infection significantly increased the IFN- γ expression in both the jejunum and ileum, confirming that viral infection activates the antiviral state of the weaned piglet. The addition of 1% Ile increased the expression of interferon in the duodenum of the RV-infected group, indicating that isoleucine enhances small intestinal mucosal immunity after RV infection by increasing duodenal IFN- γ expression. Furthermore, we also found that with the addition of 1% Ile, the expression of SIgA in the duodenum and jejunum was significantly higher in the infected group compared with the uninfected group. These results demonstrate that Ile could promote the secretion of pro-inflammatory factors and regulate the immune ability of the organism to a certain extent. However, the specific mechanisms by which Ile influences intestinal immunity in piglets require further investigation.

Defensins are antimicrobial peptides produced by leukocytes and epithelial cells, acting directly on virus particles and host cells while playing an important role as antiviral agents [57]. Our previous study showed that Ile treatment enhanced the expression of porcine β -defensin-1 (pBD-1), pBD-2, pBD-3, pBD-114, and pBD-129 in the jejunum and ileum [58]. In this study, we found that 1% Ile increased the expression of DEF β 1 in the ileum. RV infection decreased the expression of DEF β 1 in the jejunum and the expression of DEF β 2 in the duodenum. Furthermore, the addition of 1% Ile in the RV-infected group significantly increased the expression of DEF β 1 in the jejunum and increased the expression

of DEF β 2 in the duodenum. These findings suggest that Ile may inhibit RV infection by increasing jejunal DEF β 1 expression and duodenal DEF β 2 expression.

5. Conclusions

In summary, the addition of 1% Ile to the diet promoted the healthy development of the intestinal mucosa. Following RV infection, Ile was able to restore the reduced villus height in the ileum and the number of goblet cells in both the duodenum and ileum to normal levels, while also improving intestinal epithelial tight junctions. Additionally, Ile increased the activity of lipase, amylase, sucrase, SOD, and GSH, as well as the expression of SIgA, DEF β 1, and DEF β 2 in various segments of the small intestine. Therefore, the inclusion of Ile in the diet can alleviate the effects of RV on intestinal morphology and barrier function, as well as the physiological functions of weaned piglets. Moreover, Ile has the ability to enhance the antioxidant capacity and immune function of weaned piglets to a certain extent.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ani14213146/s1>.

Author Contributions: R.Z. and C.J.: conceived and designed the experiments, performed the experiments, and wrote this manuscript. C.C. and X.L.: contributed reagents, materials, and analysis tools or data. Y.Y. and S.Z.: performed the experiments. M.J. and A.H.G.: analyzed and interpreted the data and edited this manuscript. E.J. and X.Z.: analyzed and interpreted the data. M.R. and S.L.: conceived and designed the experiments and provided financial support. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Veterinary Science Peak Discipline Project of Anhui Science and Technology University (XK-XJGF002); The University Synergy Innovation Program of Anhui Province (GXXT-2022-076); Major Special Projects of Science and Technology in Anhui Province (201903a06020002); Excellent Research Innovation Team in Universities in Anhui Province (2022AH010088); and Outstanding Youth Research Project of Natural Science in Colleges and Universities in Anhui Province (2022AH030146).

Institutional Review Board Statement: All animal experiments were approved by the Animal Ethics Committee of Anhui Science and Technology University, with the protocol number AK2023013. All experimental procedures were carried out in strict accordance with the “Guidelines for the Care and Use of Test Animals” of Anhui Province.

Informed Consent Statement: Informed consent was obtained from all subjects involved in this study.

Data Availability Statement: The original contributions presented in this study are included in this article/Supplementary Material; further inquiries can be directed to the corresponding authors.

Conflicts of Interest: We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, and there is no professional or other personal interest of any nature or kind in any product, service, and/or company that could be construed as influencing the content of this paper.

References

1. Vlasova, A.N.; Amimo, J.O.; Saif, L.J. Porcine Rotaviruses: Epidemiology, Immune Responses and Control Strategies. *Viruses* **2017**, *9*, 48. [[CrossRef](#)] [[PubMed](#)]
2. Ren, X.; Saleem, W.; Haes, R.; Xie, J.; Theuns, S.; Nauwynck, H.J. Milk lactose protects against porcine group A rotavirus infection. *Front. Microbiol.* **2022**, *13*, 989242. [[CrossRef](#)] [[PubMed](#)]
3. Ghonaim, A.H.; Hopo, M.G.; Ghonaim, N.H.; Jiang, Y.; He, Q.; Li, W. The Epidemiology of Circulating Rotavirus Associated with Diarrhea in Egyptian Kids and Calves: A Review. *Zoonoses* **2023**, *3*, 985. [[CrossRef](#)]
4. Esona, M.D.; Gautam, R. Rotavirus. *Clin. Lab. Med.* **2015**, *35*, 363–391. [[CrossRef](#)]
5. Karayel-Hacioglu, I.; Timurkan, M.O.; Pellegrini, F.; Marton, S.; Gul, B.; Banyai, K.; Martella, V.; Alkan, F. Whole-genome analysis of a rare G15P 21 group A rotavirus detected at a dairy cattle farm. *J. Gen. Virol.* **2022**, *103*, 001808. [[CrossRef](#)]
6. Chen, Y.; Wu, R.; Mi, W.; Ghonaim, A.H.; Ren, W.; Yang, L.; Ruan, S.; He, Q.; Chen, H.; Jiang, Y. Molecular evolution and genetic characteristics of G3P 3 group A canine rotavirus isolated in Wuhan, China. *J. Gen. Virol.* **2022**, *103*, 001784. [[CrossRef](#)]

7. Timurkan, M.O.; Alkan, F. Identification of rotavirus A strains in small ruminants: First detection of G8P 1 genotypes in sheep in Turkey. *Arch. Virol.* **2020**, *165*, 425–431. [[CrossRef](#)]
8. Estes, M.K.; Kang, G.; Zeng, C.Q.; Crawford, S.E.; Ciarlet, M. Pathogenesis of rotavirus gastroenteritis. In *Gastroenteritis Viruses: Novartis Foundation Symposium*; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2001; Volume 238, pp. 82–96; discussion pp. 96–100. [[CrossRef](#)]
9. Ghonaim, A.H.; Yi, G.; Lei, M.; Xie, D.; Ma, H.; Yang, Z.; Usama, U.; Wu, H.; Jiang, Y.; Li, W.; et al. Isolation, characterization and whole-genome analysis of G9 group A rotaviruses in China: Evidence for possible Porcine-Human interspecies transmission. *Virology* **2024**, *597*, 110129. [[CrossRef](#)]
10. Zhang, H.; Zhang, Z.; Wang, Y.; Wang, X.; Xia, M.; Wu, H. Isolation, molecular characterization and evaluation of the pathogenicity of a porcine rotavirus isolated from Jiangsu Province, China. *Arch. Virol.* **2015**, *160*, 1333–1338. [[CrossRef](#)]
11. Kumar, D.; Shepherd, F.K.; Springer, N.L.; Mwangi, W.; Marthaler, D.G. Rotavirus Infection in Swine: Genotypic Diversity, Immune Responses, and Role of Gut Microbiome in Rotavirus Immunity. *Pathogens* **2022**, *11*, 1078. [[CrossRef](#)]
12. Jiang, L.; Tang, A.; Song, L.; Tong, Y.; Fan, H. Advances in the development of antivirals for rotavirus infection. *Front. Immunol.* **2023**, *14*, 1041149. [[CrossRef](#)] [[PubMed](#)]
13. Amimo, J.O.; Raev, S.A.; Chepngeno, J.; Mainga, A.O.; Guo, Y.; Saif, L.; Vlasova, A.N. Rotavirus Interactions With Host Intestinal Epithelial Cells. *Front. Immunol.* **2021**, *12*, 793841. [[CrossRef](#)]
14. Yang, H.; Fan, X.; Mao, X.; Yu, B.; He, J.; Yan, H.; Wang, J. The protective role of prebiotics and probiotics on diarrhea and gut damage in the rotavirus-infected piglets. *J. Anim. Sci. Biotechnol.* **2024**, *15*, 61. [[CrossRef](#)]
15. Nie, C.; He, T.; Zhang, W.; Zhang, G.; Ma, X. Branched Chain Amino Acids: Beyond Nutrition Metabolism. *Int. J. Mol. Sci.* **2018**, *19*, 954. [[CrossRef](#)]
16. Le Couteur, D.G.; Solon-Biet, S.M.; Cogger, V.C.; Ribeiro, R.; de Cabo, R.; Raubenheimer, D.; Cooney, G.J.; Simpson, S.J. Branched chain amino acids, aging and age-related health. *Ageing Res. Rev.* **2020**, *64*, 101198. [[CrossRef](#)]
17. Alam, N.H.; Ashraf, H.; Gyr, N.E.; Meier, R.F. Mo1061 Evaluation of the Efficacy of L-Isoleucine and L-Arginine Supplemented Food in the Treatment of Acute Diarrhea in Malnourished Children. *Gastroenterology* **2013**, *144*, S-566. [[CrossRef](#)]
18. Gu, C.; Mao, X.; Chen, D.; Yu, B.; Yang, Q. Isoleucine Plays an Important Role for Maintaining Immune Function. *Curr. Protein Pept. Sci.* **2019**, *20*, 644–651. [[CrossRef](#)]
19. Zhao, Y.; Yan, M.Y.; Jiang, Q.; Yin, L.; Zhou, X.Q.; Feng, L.; Liu, Y.; Jiang, W.D.; Wu, P.; Zhao, J.; et al. Isoleucine improved growth performance, and intestinal immunological and physical barrier function of hybrid catfish *Pelteobagrus vachelli* x *Leiostichus longirostris*. *Fish Shellfish Immunol.* **2021**, *109*, 20–33. [[CrossRef](#)]
20. Zhang, S.; Zeng, X.; Ren, M.; Mao, X.; Qiao, S. Novel metabolic and physiological functions of branched chain amino acids: A review. *J. Anim. Sci. Biotechnol.* **2017**, *8*, 10. [[CrossRef](#)]
21. Ren, M.; Zhang, S.; Liu, X.; Li, S.; Mao, X.; Zeng, X.; Qiao, S. Different Lipopolysaccharide Branched-Chain Amino Acids Modulate Porcine Intestinal Endogenous beta-Defensin Expression through the Sirt1/ERK/90RSK Pathway. *J. Agric. Food Chem.* **2016**, *64*, 3371–3379. [[CrossRef](#)]
22. Zhang, Y.; Zhan, L.; Zhang, L.; Shi, Q.; Li, L. Branched-Chain Amino Acids in Liver Diseases: Complexity and Controversy. *Nutrients* **2024**, *16*, 1875. [[CrossRef](#)] [[PubMed](#)]
23. Alkhatib, A. Antiviral Functional Foods and Exercise Lifestyle Prevention of Coronavirus. *Nutrients* **2020**, *12*, 2633. [[CrossRef](#)] [[PubMed](#)]
24. Santos, F.R.D.; Dos, S.C.; Maranhao, M.L.A.B.; Espinola, C.C.M.; Franco, O.L. Immunonutrition effects on coping with COVID-19. *Food Funct.* **2021**, *12*, 7637–7650. [[CrossRef](#)]
25. Mao, X.; Gu, C.; Hu, H.; Tang, J.; Chen, D.; Yu, B.; He, J.; Yu, J.; Luo, J.; Tian, G. Dietary *Lactobacillus rhamnosus* GG Supplementation Improves the Mucosal Barrier Function in the Intestine of Weaned Piglets Challenged by Porcine Rotavirus. *PLoS ONE* **2016**, *11*, e0146312. [[CrossRef](#)]
26. NRC (National Research Council). *Nutrient Requirements of Swine*, 11th ed.; National Academic Press: Washington, DC, USA, 2012.
27. Guerrero, C.A.; Acosta, O. Inflammatory and oxidative stress in rotavirus infection. *World J. Virol.* **2016**, *5*, 38–62. [[CrossRef](#)]
28. Wu, S.; Liu, X.; Cheng, L.; Wang, D.; Qin, G.; Zhang, X.; Zhen, Y.; Wang, T.; Sun, Z. Protective Mechanism of Leucine and Isoleucine against H(2)O(2)-Induced Oxidative Damage in Bovine Mammary Epithelial Cells. *Oxid. Med. Cell. Longev.* **2022**, *2022*, 4013575. [[CrossRef](#)]
29. Chethan, G.E.; Garkhal, J.; Sircar, S.; Malik, Y.P.S.; Mukherjee, R.; Sahoo, N.R.; Agarwal, R.K.; De, U.K. Immunomodulatory potential of beta-glucan as supportive treatment in porcine rotavirus enteritis. *Vet. Immunol. Immunopathol.* **2017**, *191*, 36–43. [[CrossRef](#)]
30. Beaupaire, C.; Smyth, D.; McKay, D.M. Interferon-gamma regulation of intestinal epithelial permeability. *J. Interferon Cytokine Res.* **2009**, *29*, 133–144. [[CrossRef](#)]
31. Li, Y.; Jin, L.; Chen, T. The Effects of Secretory IgA in the Mucosal Immune System. *Biomed. Res. Int.* **2020**, *2020*, 2032057. [[CrossRef](#)]
32. Huang, J.; Liu, X.; Sun, Y.; Huang, C.; Wang, A.; Xu, J.; Zhou, H.; Li, L.; Zhou, R. Porcine beta-defensin 2 confers enhanced resistance to swine flu infection in transgenic pigs and alleviates swine influenza virus-induced apoptosis possibly through interacting with host SLC25A4. *Antivir. Res.* **2022**, *201*, 105292. [[CrossRef](#)]

33. Huang, J.; Qi, Y.; Wang, A.; Huang, C.; Liu, X.; Yang, X.; Li, L.; Zhou, R. Porcine beta-defensin 2 inhibits proliferation of pseudorabies virus in vitro and in transgenic mice. *Viol. J.* **2020**, *17*, 18. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Vemula, S.V.; Amen, O.; Katz, J.M.; Donis, R.; Sambhara, S.; Mittal, S.K. Beta-defensin 2 enhances immunogenicity and protection of an adenovirus-based H5N1 influenza vaccine at an early time. *Virus Res.* **2013**, *178*, 398–403. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Ren, M.; Yang, Y.N.; Zhao, C.F.; Che, C.Y.; Li, S.H. The regulatory mechanism of isoleucine on the mucosal morphology and structure of rat intestine in Chinese. *Chin. Sci. Bull.* **2022**, *67*, 520–527. [\[CrossRef\]](#)
36. Paone, P.; Cani, P.D. Mucus barrier, mucins and gut microbiota: The expected slimy partners? *Gut* **2020**, *69*, 2232–2243. [\[CrossRef\]](#)
37. Li, L.; Fu, F.; Guo, S.; Wang, H.; He, X.; Xue, M.; Yin, L.; Feng, L.; Liu, P. Porcine Intestinal Enteroids: A New Model for Studying Enteric Coronavirus Porcine Epidemic Diarrhea Virus Infection and the Host Innate Response. *J. Virol.* **2019**, *93*, e01682-18. [\[CrossRef\]](#)
38. Cortez, V.; Boyd, D.F.; Crawford, J.C.; Sharp, B.; Livingston, B.; Rowe, H.M.; Davis, A.; Alsallaq, R.; Robinson, C.G.; Vogel, P.; et al. Astrovirus infects actively secreting goblet cells and alters the gut mucus barrier. *Nat. Commun.* **2020**, *11*, 2097. [\[CrossRef\]](#)
39. Holly, M.K.; Smith, J.G. Adenovirus Infection of Human Enteroids Reveals Interferon Sensitivity and Preferential Infection of Goblet Cells. *J. Virol.* **2018**, *92*, e00250-18. [\[CrossRef\]](#)
40. Zhang, S.; Zhang, S.; Hou, Y.; Huang, Y.; Cai, J.; Wang, G.; Cao, Y.; Chen, Z.; Fang, X.; Bao, W. Porcine Deltacoronavirus Infection Disrupts the Intestinal Mucosal Barrier and Inhibits Intestinal Stem Cell Differentiation to Goblet Cells via the Notch Signaling Pathway. *J. Virol.* **2023**, *97*, e0068923. [\[CrossRef\]](#)
41. Boshuizen, J.A.; Reimerink, J.H.; Korteland-van, M.A.M.; van Ham, V.J.; Bouma, J.; Gerwig, G.J.; Koopmans, M.P.; Buller, H.A.; Dekker, J.; Einerhand, A.W. Homeostasis and function of goblet cells during rotavirus infection in mice. *Virology* **2005**, *337*, 210–221. [\[CrossRef\]](#)
42. Jung, K.; Saif, L.J. Goblet cell depletion in small intestinal villous and crypt epithelium of conventional nursing and weaned pigs infected with porcine epidemic diarrhea virus. *Res. Vet. Sci.* **2017**, *110*, 12–15. [\[CrossRef\]](#)
43. Groschwitz, K.R.; Hogan, S.P. Intestinal barrier function: Molecular regulation and disease pathogenesis. *J. Allergy Clin. Immunol.* **2009**, *124*, 3–20; quiz 21–22. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Assimakopoulos, S.F.; Scopa, C.D.; Charonis, A.; Spiliopoulou, I.; Georgiou, C.; Nikolopoulou, V.; Vagianos, C.E. Experimental obstructive jaundice disrupts intestinal mucosal barrier by altering occludin expression: Beneficial effect of bombesin and neurotensin. *J. Am. Coll. Surg.* **2004**, *198*, 748–757. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Jonathan, M.; Bruce, K.R.; Judith, A.V. Mucins, Mucus, and Goblet Cells. *Chest* **2018**, *154*, 169–176. [\[CrossRef\]](#)
46. Zhang, G.G.; Yang, Z.B.; Wang, Y.; Yang, W.R.; Zhou, H.J. Effects of dietary supplementation of multi-enzyme on growth performance, nutrient digestibility, small intestinal digestive enzyme activities, and large intestinal selected microbiota in weanling pigs. *J. Anim. Sci.* **2014**, *92*, 2063–2069. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Cheng, Y.; Ding, S.; Azad, M.A.K.; Song, B.; Kong, X. Small Intestinal Digestive Functions and Feed Efficiency Differ in Different Pig Breeds. *Animals* **2023**, *13*, 1172. [\[CrossRef\]](#)
48. Liu, K.; Shen, J.; Cao, Y.; Cai, C.; Yao, J. Duodenal infusions of isoleucine influence pancreatic exocrine function in dairy heifers. *Arch. Anim. Nutr.* **2018**, *72*, 31–41. [\[CrossRef\]](#)
49. Wong, H.; Schotz, M.C. The lipase gene family. *J. Lipid Res.* **2002**, *43*, 993–999. [\[CrossRef\]](#)
50. Senftleber, N.K.; Ramne, S.; Moltke, I.; Jorgensen, M.E.; Albrechtsen, A.; Hansen, T.; Andersen, M.K. Genetic Loss of Sucrase-Isomaltase Function: Mechanisms, Implications, and Future Perspectives. *Appl. Clin. Genet.* **2023**, *16*, 31–39. [\[CrossRef\]](#)
51. Yang, B.; Chen, Y.; Shi, J. Reactive Oxygen Species (ROS)-Based Nanomedicine. *Chem. Rev.* **2019**, *119*, 4881–4985. [\[CrossRef\]](#)
52. Brieger, K.; Schiavone, S.; Miller, F.J., Jr.; Krause, K.H. Reactive oxygen species: From health to disease. *Swiss Med. Wkly.* **2012**, *142*, w13659. [\[CrossRef\]](#)
53. Foo, J.; Bellot, G.; Pervaiz, S.; Alonso, S. Mitochondria-mediated oxidative stress during viral infection. *Trends Microbiol.* **2022**, *30*, 679–692. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Gong, B.S.; Ren, M.; Jin, E.H.; Li, Q.Q.; Li, X.R.; Li, S.H. Effect of Isoleucine on the Serum Antioxidant Levels and the Structure of Spleen in Rats. *J. Anhui Sci. Technol. Univ.* **2015**, *29*, 21–25. (In Chinese)
55. Kak, G.; Raza, M.; Tiwari, B.K. Interferon-gamma (IFN-gamma): Exploring its implications in infectious diseases. *Biomol. Concepts* **2018**, *9*, 64–79. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Mao, X.; Gu, C.; Ren, M.; Chen, D.; Yu, B.; He, J.; Yu, J.; Zheng, P.; Luo, J.; Luo, Y.; et al. l-Isoleucine Administration Alleviates Rotavirus Infection and Immune Response in the Weaned Piglet Model. *Front. Immunol.* **2018**, *9*, 1654. [\[CrossRef\]](#)
57. Holly, M.K.; Diaz, K.; Smith, J.G. Defensins in Viral Infection and Pathogenesis. *Annu. Rev. Virol.* **2017**, *4*, 369–391. [\[CrossRef\]](#)
58. Ren, M.; Cai, S.; Zhou, T.; Zhang, S.; Li, S.; Jin, E.; Che, C.; Zeng, X.; Zhang, T.; Qiao, S. Isoleucine attenuates infection induced by E. coli challenge through the modulation of intestinal endogenous antimicrobial peptide expression and the inhibition of the increase in plasma endotoxin and IL-6 in weaned pigs. *Food Funct.* **2019**, *10*, 3535–3542. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.