

Review

Methanogens Through Time and Space: Impact on Earth's Planetary Evolution and Biogeochemistry [†]

Paxton Tomko ¹, Cesar Ivan Ovando-Ovando ², Pierre Boussagol ^{1,3}, Michel Geovanni Santiago-Martínez ² and Pieter T. Visscher ^{1,2,4,*} 

¹ Department of Marine Sciences, The University of Connecticut, Groton, CT 06340, USA; paxton.tomko@uconn.edu (P.T.); pierre.boussagol@u-bourgogne.fr (P.B.)

² Department of Molecular and Cell Biology, The University of Connecticut, Storrs, CT 06269, USA; cesar.ovando@uconn.edu (C.I.O.-O.); geo_santiagom@uconn.edu (M.G.S.-M.)

³ Department Biogéosciences, Université Bourgogne Franche-Comté, 21000 Dijon, France

⁴ Department of Earth Sciences, The University of Connecticut, Groton, CT 06340, USA

* Correspondence: pieter.visscher@uconn.edu

[†] Dedicated to the memory of Ronald S. Oremland, whose career advanced our knowledge of the biogeochemistry of methanogens.

Abstract

Methanogens, or methanogenic archaea (MA), are among the most ancient and widely distributed microorganisms, characterized by a unique metabolism that generates methane (CH₄) as the terminal product of anaerobic respiration. Their ability to grow and/or survive across a wide range of environmental conditions has made methanogens key contributors to biogeochemical cycles throughout most of Earth's history. Most importantly, these oxygen-sensitive microorganisms have regulated the climate since the early Archean and impacted biogeochemical cycles throughout Earth's history by producing the potent greenhouse gas, CH₄, while consuming H₂, CO₂, and small organic molecules. Hence, methanogens are attributed a key role in the start and end of several Proterozoic glaciations and mass extinction events. Their specific roles in the long-term carbon cycle that focus on CH₄ production are well-established, but, in contrast, only very few studies report on interactions with CaCO₃ and long-term carbon storage. Methanogens evolved early during Earth's history, likely during the Archaean Eon, in layered benthic microbial communities called microbial mats. When lithified, these mats form microbialites that represent some of the earliest evidence of life in the fossil record, dating back >3.5 Gy. Methanogens are an integral part of contemporary microbial mats and have been identified both in the anoxic and oxic zones of these sedimentary ecosystems; however, their adaptations to apparently unfavorable oxic conditions and their role in the precipitation of carbonate in mats are unclear. In addition to an important role in the evolution of our planet by producing CH₄, methanogens may also produce a biosignature that could be relevant for astrobiology research. This review will discuss the diversity, physiology, and ecology of methanogens in detail to clarify their role in some of the major biogeochemical processes and ecological climatic events through the fluctuating environmental conditions on Earth through geologic time.

Keywords: methanogens; archaea; microbial mats; early life; exobiology; climate regulation; mass extinctions



Academic Editor: Alberto G. Fairén

Received: 23 December 2025

Revised: 21 March 2026

Accepted: 25 March 2026

Published: 1 April 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article

distributed under the terms and

conditions of the [Creative Commons](#)

[Attribution \(CC BY\)](#) license.

1. The Biogeochemistry of Methanogens

1.1. Introduction

Methanogens are strict anaerobes with highly specialized biochemistry. All methanogens belong to the domain Archaea and produce methane as an end-product of their metabolism. Methanogenesis serves as their primary energy-conserving pathway, an oxygen-independent process at the limits of what is thermodynamically feasible for sustaining life [1–4]. In addition to CH₄, some heterotrophic methanogens also produce carbon dioxide as an end-product (see Section 1.4.1), contributing significantly to greenhouse gas emissions and thereby regulating the Earth's temperature [2,5], and potentially that of other planets [6].

For most of geological history, prokaryotes have dominated our planet [7], filling a diverse array of ecological niches [8,9]. Methanogens are a prime example of an ancient and versatile group of microorganisms that thrive in diverse environments, sometimes under polyextreme conditions [3,10]. Their metabolic versatility, unique biogeochemistry, and interactions with other microorganisms, such as fermenters, syntrophs, methane oxidizers, and sulfate reducers, may have facilitated microbial life to develop throughout Earth's history [11,12]. Furthermore, their exemplary ecological success through geologic time and space (i.e., distribution across diverse and extreme conditions) suggests that methanogens could survive conditions elsewhere in the solar system and possibly beyond, making them relevant model organisms for astrobiology [10,13].

1.2. Methanogenic Habitats

Methanogens are characterized by the production of methane (CH₄) as the obligatory product of their oxygen-independent metabolism [3]. Methanogens are found in nearly all anoxic environments on Earth, and many anaerobic niches, including deep sea, freshwater, and terrestrial sediments, marshes, swamps, permafrost soils, geothermal springs, hydrothermal vents, the deep biosphere, the rumen of cows and intestines of other mammals (including humans), arthropods, and human-made environments such as landfills, (anaerobic) digesters, and rice paddies [3,10,14–23]. Some methanogens have also been found in oxic environments as well [24], such as the surface of biofilms and microbial mats [4,24,25], the upper oxic layer of water bodies [26–28], mangroves [29], and oxygenated soils [30]. The survival of methanogens has also been assessed under a variety of exoplanetary, artificial, and space conditions, including Martian soil analogues [31,32], under simulated conditions of ocean worlds (such as Enceladus) [13], on the outside of spacecraft during interplanetary travel [10], and in clean rooms [33].

1.3. Physicochemical Boundaries of Methanogenesis

Methanogens have a unique biochemistry, which makes them well-suited for life in extreme niches [10]. As such, methanogens have been isolated from environments with a wide range of physicochemical and thermochemical conditions (Table 1), including temperature [34,35], pH [3,36], salinity [16], and pressure gradients [32,35,37–39]. Some methanogens can also withstand desiccation [32], freeze/thaw cycles [34], and survive high doses of ionizing and UV radiation [40]. Methanogens are found in a wide range of extreme environments and tolerate multiple extreme conditions, making them versatile polyextremophiles [4,10,14,16]. For example, barophilic and thermophilic methanogens, members of the *Methanopyrus* genus, *Methanocaldococcus jannaschii*, and *Methanothermococcus thermolithotrophicus*, have been isolated from hydrothermal vent sites, with *Methanopyrus kandleri* strain 116 being the most heat-tolerant microorganism cultured, growing at temperatures up to 122 °C and pressures up to 20 MPa [35]. On the opposite extreme, psychrophilic methanogens are abundant in wetlands and arctic permafrost [3,10] and can thrive in sub-atmospheric pressure [41]. Soda and saline lakes host alkaliphilic and

halophilic methanogens belonging to the genera *Methanohalobium*, *Methanohalophilus* and *Methanosalsum* [16].

Methanogens are widely distributed and well adapted to thrive in microbial mat ecosystems [4,42–44], where they could be exposed to high and fluctuating concentrations of oxygen (O₂) despite the traditional belief that they are obligate anaerobes or oxygen-sensitive microorganisms [25]. Their unique biology allows them to adapt and thrive in a range of challenging environments. For example, hydrogenotrophic methanogens, under pH and thermal stress, low H₂, iron, phosphate, and trace metals can split their metabolic investments to maintain homeostasis or add biomass, a process referred to as ‘uncoupling’ [10]. Some methanogens can reverse reactions of some steps in methanogenesis to maximize their energy yield when substrates for respiration are sparse [15,45,46] or enter a dormancy-like low-activity state from which they can rapidly revive [47]. Such mechanisms may have played a role in their evolution through conditions in Earth’s past, such as through glaciations and the oxygenation of Earth’s atmosphere (see Section 2).

Table 1. Growth limits for methanogens and examples of species that are found at these limits. Definitions of extremes, in terms of the known limits for life [8,48] and the limits that have been identified for methanogens. The geologic relevance highlights some of the environments on early Earth, in the present, and on other planets where these conditions could be/have been found.

Environmental Condition	Extremophile Type (and Growth Definition(s))	Growth Limits for Methanogens	Examples of Species Found Within Extreme	Geologic Relevance
Temperature	(1) Hyperthermophile (>80 °C) (2) Thermophile (60–80 °C) (3) Psychrophile (<15 °C)	(1 & 2) 122 °C [2,35]. (3) −2.5 °C [49].	(1 & 2) <i>Methanopyrus kandleri</i> strain 116, <i>Methanopyrus</i> genus, <i>Methanocaldococcus jannaschii</i> , and <i>Methanothermococcus thermolithotrophicus</i> [35]. (3) <i>Candidatus Methanoflorens stordalenmirensis</i> (freeze/thawing environments) [34], <i>Methanococcoides burtonii</i> [49].	(1) Hydrothermal vents (modern day [35], potential vents of Europa and Enceladus [13], hot springs/geothermal springs [20]). (2) Archean oceans [50]. (3) Permafrost and ice; modern day and during glaciation events, like at the last glacial maxima; deep sediments [34,51], terrestrial surface of Mars [37], oceans of Europa and Enceladus [10,48].
Radiation	(1) Exposure to UV (2) Ionizing radiation	(1) 24 h of 254 nm UV radiation [52].	(1) <i>Methanococcus maripaludis</i> and <i>Methanobacterium formicicum</i> [40,53]. (2) <i>Methanosarcina</i> spp. [52].	(1) High UV radiation during the Archean [48]; surface of Mars [40]. (2) Ionizing radiation from Ur decay in nuclear geysers, possible location for the origin of life [54].
Light	(1) Exposure to visible light (380–750 nm)	(1) Unknown limit; growth inhibition, particularly in the blue region of the spectrum (≈370–430 nm) [55]. Reduced sensitivity in association with photosynthetic bacteria [56].	(1) <i>Methanosarcina</i> and <i>Methanocella</i> spp. (observed in soil) [56]; <i>Methanosarcina barkeri</i> in a cocultured biofilm with <i>Synechocystis</i> spp. strain PCC6803 [24].	(1) Surface environments on Earth (from ancient to modern day).
Salinity	(1) Halophile (2–5 M NaCl)	(1) Moderate/extreme halophiles growth 2–3 M [16].	(1) <i>Methanosarcinales</i> spp., <i>Methanohalobium</i> , <i>Methanohalophilus</i> and <i>Methanosalsum</i> spp., <i>Methanonatronarchaeum thermophilum</i> and <i>Candidatus Methanohalarchaeum thermophilum</i> [16,39].	(1) Archean oceans; 1.5–2 times modern salinity [57]; saline lakes [16].

Table 1. Cont.

Environmental Condition	Extremophile Type (and Growth Definition(s))	Growth Limits for Methanogens	Examples of Species Found Within Extreme	Geologic Relevance
Pressure	(1) Survivability to low pressure (2) Barophile (obligate high pressure)/Piezophile (adaptation to high pressure)	(1) Low pressure 6–143 mbar [37]. (2 & 3) 200 bar [35], 500 bar [39].	(1) <i>Methanothermobacter wolfeii</i> , <i>Methanosarcina barkeri</i> , <i>Methanobacterium formicicum</i> , <i>Methanococcus maripaludis</i> [32,37,38]. (2) <i>Methanopyrus kandleri</i> strain 116, <i>Methanopyrus</i> genus, <i>Methanocaldococcus jannaschii</i> , and <i>Methanothermococcus thermolithotrophicus</i> [35,38].	(1) Terrestrial surface of Mars [38]. (2) Hydrothermal vents (modern day [35] and possible location for the origin of life [58]; deep marine and continental subsurface sediments [59,60]).
pH	(1) Alkaliphile (2) Acidophile	(1) pH > 10.2 [3,36]. (2) pH < 3 [3].	(1) <i>Methanohalobium</i> , <i>Methanohalophilus</i> and <i>Methanosalsum</i> [36].	(1) Soda lakes [36].
Hydrogen concentration	(1) High pH_2 (2) Low pH_2 (3) Fluctuations in H_2 concentration	(1) $>6 \times 10^{-2}$ bar [61] *. (2 & 3) As low as 0.1 Pa pH_2 [62]; fluctuating H_2 (from low pH_2 to overpressure) [14].	(1) Species name unknown. Plausibly, early hydrogenotrophs. (2 & 3) <i>Methanobacterium bryantii</i> , <i>Methanoculleus bourgensis</i> MAB1, and <i>Methanosarcina barkeri</i> [62], <i>Methanobrevibacter</i> spp., <i>Methanomicrobium</i> spp., and <i>Methanosarcinales</i> [14].	(1) Earth's secondary atmosphere [61]. (2) Titan [63]. (3) Cow rumen [14]; deep biosphere [62].
Oxygen	(1) Aerotolerant (tolerates some (high to low) amount of O_2)	(1) Unknown limit; exposure up to 300 μM O_2 in lithifying microbial mat [64].	(1) <i>Methanosarcina barkeri</i> [24]; methanogens in microbial mats [25]. Class II methanogens [65].	(1) Euxinic oceans of Proterozoic [50]; modern day (and ancient) biofilms and microbial mats [24,25].
Metals	(1) Arsenic (2) Cadmium	(1) Unknown limit; methane production by methanogens observed under high arsenic concentrations (0.8–1.5 mM) in microbial mats [66,67]. (2) Unknown limit; 10 μM of $CdCl_2$ activates the rate of methanogenesis; ≥ 100 μM of $CdCl_2$ inhibits cell growth [68].	(1) <i>Methanomassiliicoccus</i> spp. [69] (2) <i>Methanosarcina acetivorans</i> [68]. Other divalent metals (Co, Zn, Cu, Fe) also activate the rate of methanogenesis at 10 μM .	(1 & 2) Early oceans [70], rice paddies [71], modern day anoxic microbial mats in the Atacama, Chile [66,67].

* pH_2 of Earth's atmosphere at the time of early methanogen evolution.

1.3.1. Temperature Extremes

Temperature on Earth through time and space shows huge variations ranging from extreme cold or psychrophilic conditions for life (growth at <15 °C) to thermophilic (60–80 °C) and hyperthermophilic (>80 °C) conditions [8]. These temperature extremes not only exist in specific environments (e.g., modern-day hydrothermal vents [35,58]). Arctic and Antarctic permafrost and water (ice) [34,72] and deep and high-altitude sediments [51,73], but also as widespread environmental conditions of the past and present, such as during glaciations (e.g., “Snowball Earth”) [74,75], through mass extinctions (rapid warming/cooling) [76], and in the Archean oceans [57,77]. Such extremes in temperature are also relevant for potential life in other locations in our solar system, such as on the Martian surface and in the oceans of Europa, Enceladus, and Titan [10,48], and beyond (exoplanets).

Methanogenesis, like all metabolisms, is a temperature-dependent process and therefore both substrate use and community compositions of methanogens are dependent on temperature [2]. For example, at temperatures higher than 30 °C, methanogens tend to utilize H_2 to produce CH_4 rather than acetate as a growth substrate at lower temperatures [2]. Adaptations of methanogens to become psychrophiles, thermophiles, and hyperthermophiles are associated with modifications at the tRNA and protein level [10]. Psychrophilic *Methanococcoides burtonii* displays increased flexibility of polynucleotides at low temperatures associated with tRNA due to modifications with dihydrouridine [78]. The same study found tRNA modifications (e.g., methylation of nucleosides) in a thermophilic species of archaeon, *Stetteria hydrogenophila*, which can stabilize tRNA to accommodate growth under high temperatures [78]. At the protein level, *Methanococcoides burtonii* dis-

plays cold shock proteins such as Elongation Factor 2, which is unstable at higher temperatures [49]. Similarly, a so-called “Dead box” RNA helicase gene is expressed in this species at 4 °C [79], a process which is also used by the thermophile, *Thermococcus kodakarensis*, when experiencing “cold” stress at 60 °C [80]. Modifications are made to another cold shock protein (Csp), where the CspA homolog domain helps cells cope with cold stress [81]. Similar to adaptations in bacteria, all archaea must maintain membrane fluidity in cold environments by regulating the unsaturation of membrane lipids [82]. Low permeability and liquid crystalline phases of isoprenoid ether lipid membranes at temperatures ranging from 0 to 100 °C are characteristic of archaeal membranes. They allow for stability at a wider range of temperatures than that found in bacteria [77]. Still, there is less membrane lipid variation observed in thermophilic methanogens than in bacteria [83].

1.3.2. UV and Ionizing Radiation

The surface of Mars experiences high doses of UV radiation from the sun, which may restrict extant life to the subsurface and could erase surface biosignatures of ancient life over geologic time [84]. Life on the Archean Earth also experienced higher levels of UV radiation due to the lack of an ozone layer in the upper atmosphere [85]. This condition favors life to develop and evolve in microbial mats [4]. Exposure to UV radiation leads to the limited survival of *Methanococcus maripaludis* and *Methanobacterium formicicum* under simulated Mars conditions [40]. Moreover, *Methanosarcina solegelidi*, isolated from the Siberian permafrost, has a similar response to UV-C and X-rays as *Deinococcus radiodurans* [53], presumably the organism with the highest resistance to these conditions.

While the exact mechanisms by which methanogens adapt to UV stress are unknown, DNA repair mechanisms could play a role [40]. Physical barriers can protect cells from UV radiation, including FeS, which was abundant on early Earth [50], in layers of soil and sediment surrounding the cells [40], through exopolymeric substances (EPS), and in photopigments in microbial mats containing methanogens [4,74]. *Methanosarcina* spp. survives radiation and desiccation [52] by deploying enzymes and DNA repair mechanisms to protect against oxidative stress [52].

1.3.3. Visible Light

The sun’s luminosity has increased through geological time, while short-wavelength radiation has decreased [86]. Light that reaches the Earth’s surface is a critical environmental factor that regulates a wide range of biological processes. At the molecular level, visible light (wavelengths from 380 to 750 nm) is perceived by sensory photoreceptor proteins that detect specific wavelengths and convert light signals into biochemical responses [87,88]. These photoreceptors span a broad region of the electromagnetic spectrum, ranging from near-ultraviolet (near-UV) to near-infrared (NIR) wavelengths [88].

In methanogens, visible light from both solar and artificial sources negatively affects growth, morphology, and physiology, and in some species, it can even trigger cell lysis and death [55]. Conversely, the absence of visible light has long been considered a critical parameter for the optimal cultivation of multiple strains of methanogens isolated from diverse environments [89–91].

Recent mesocosm experiments, conducted in artificial ponds containing environmental sediments, showed that methane emissions by methanogens are two- to three-fold higher at night compared to daytime levels [92–94]. A decade-long monitoring of microbial diversity and functional activity demonstrated that sunlight modulates the production rates of methane and carbon dioxide, indicating sunlight-dependent metabolic changes in methanogens [94]; however, it is still unclear whether this effect is due to temperature fluctuations or direct exposure to sunlight.

Despite their sensitivity to visible light, methanogens coexist with oxygenic photosynthetic bacteria in various natural habitats, including microbial mats, soil crusts, and the aerobic epilimnion of oligotrophic lakes [24,25,56,95–97]. The coexistence of methanogens and phototrophs may be more common and widespread in nature than previously recognized [24,98,99]. Anoxygenic photosynthetic bacteria likely act as photosensitizers, promoting the CO_2 -to- CH_4 conversion in association with anaerobic methanogens when cocultured in anoxic biofilms [23]. Taken together, these observations suggest that visible light exerts a far more complex influence on methanogens than previously understood, not only constraining their physiology but also enabling novel ecological interactions.

1.3.4. Salinity

Methanogens in the Archean would have evolved in an ocean 1.5–2 times more saline [57] than the average of 35 g.kg^{-1} today [100]. Higher salinity would have affected not only the ecophysiology and evolution of methanogens but also could have aided in the warming of the early oceans due to a lower gas solubility [100].

There are two primary mechanisms archaea utilize to maintain osmotic balance in halophilic environments. The first is a so-called ‘salt out’ approach in which an organic osmotic solute, such as glycine betaine, dimethylsulfoniopropionate, ectoine, or hydroxyectoine, is utilized internally by the cell to maintain lower cellular concentrations of NaCl than in its surroundings [16,100,101]. These compatible solutes provide protection for the cell by balancing osmotic pressure without interfering with biochemical processes [16,102,103]. The second is a ‘salt in’ method, through high intracellular concentrations of K^+ , which is employed by species of the genus *Methanonatronarchaeum* [39,101]. This genus contains extremely halophilic methanogen species *Methanonatronarchaeum thermophilum* and *Candidatus Methanohalarchaeum thermophilum*, which are novel in terms of their metabolism, utilizing C_1 compounds as an electron acceptor and H_2 as an electron donor for methanogenesis [16]. These species have abundant UspA family stress response proteins, their membrane lipids contain pleckstrin homology domains, and archaeal histones may reinforce the stability of RNA, DNA, and proteins—all of which may aid in their adaptations to high salt concentrations [16].

1.3.5. Pressure

Studies simulating low-pressure conditions on Mars reveal the survival of methanogens in 6–400 mbar [37,38]. The physiological and physiochemical adaptations of methanogens in low-pressure environments are unknown, due to the lack of relevance in Earth environments. Survival in low-pressure environments is in part attributed to the rigidity and structure of archaeal lipid membranes [37]. Some species, like *Methanoscarchina* spp. are flexible in their electron donor choice and grow in multicellular clusters, which makes them more resistant to desiccation and possibly low gas pressures [37].

High pressure, or barophilic conditions, are better studied in the context of the Earth’s biosphere, for example, in the deep-sea or in oceanic and terrestrial subsurface sediments [59,60]. Barophilic pressure conditions often coincide with other extremes, such as hyperthermophilic conditions at hydrothermal vents [35,39,59]. Morphological adaptations to high temperatures are thus present in methanogens growing at high pressures [39,59]. Some species, like *Methanococcus thermolithotrophicus*, typically coccoidal cells, have been observed to grow in an elongated shape under high hydrostatic pressure [39].

1.3.6. Hydrogen Concentration

Low and high partial pressure of hydrogen ($p\text{H}_2$) is found in a variety of environments on Earth and is highly relevant in astrobiology, such as the low $p\text{H}_2$ atmosphere of Saturn’s moon Titan [63]. A high $p\text{H}_2$ was characteristic of Earth’s early secondary atmosphere,

with a $p\text{H}_2$ of over 6×10^{-2} bar [61]; conditions under which early methanogens evolved. The evolution of nitrogen fixation over 3.2 billion years ago [104] may have contributed to an abundance of H_2 [105], which could have supported early methanogenesis.

At the community level, methanogens adapt well to fluctuations in H_2 found in the rumens of cows and sheep, which change from low concentration H_2 to supersaturated conditions [14,106]. Some of the species found in these communities, like *Methanobacteriales*, lack cytochromes and are associated with short doubling times and mesophilic temperatures [14]. Lithotrophic methanogenesis (i.e., use of H_2/CO_2 ; see Section 1.4.1) is most common in animal gut communities [14]. Trace metals, such as nickel used in hydrogenases, may aid in adaptation to low H_2 conditions [62].

1.3.7. Oxygen Exposure

Methanogens are considered oxygen-sensitive microorganisms and obligate anaerobes but are exposed to high levels of oxygen in modern-day microbial mats [4,25] and potentially in oxic environments such as surface waters, including those of the Proterozoic eon. Presence of O_2 in the environment expanded the redox states of elements and allowed for greater variation in electron donor/acceptor availability (i.e., aerobic metabolism) [107]. Exposure to O_2 in biofilms [24], microbial mats [25], and other terrestrial and aquatic systems [26–28,30] may challenge the doctrine that methanogens are strict obligate anaerobes: some species are clearly more aerotolerant than previously thought [108]. Ye and colleagues (2024) observed enhanced methanogenesis in lab-grown cocultured biofilms of a cyanobacterium, *Synechocystis* PCC6803, and a methanogenic archaeon, *Methanosarcina barkeri*. During dark cycles, syntrophic methanogenesis was driven by inorganic products (H_2/CO_2) produced through respiration by cyanobacteria, while *M. barkeri* reduced the $p\text{H}_2$, enhancing *Synechocystis* PCC6803 growth [24]. During the day, active photosynthesis oxygenated the biofilm, triggering the generation of reactive oxygen species (ROS) by *M. barkeri*, which oxidized methyl radicals, eventually resulting in the abiotic production of CH_4 [24,109]. The ROS produced in the presence of O_2 can also be used as an organic substrate for CH_4 production by *M. barkeri* [24]. Additionally, diel fluctuations in metabolic activity by *M. barkeri* established Fe-redox cycling, which enhanced growth [24].

1.3.8. Metals

The abundance and redox state of elements have changed radically through Earth's evolution, predominantly due to the oxygenation of the oceans and atmosphere [70]. Methanogens can transform the redox state of metals in processes that lead to their solubilization or biomineralization [110]. This dynamic interaction is an adaptation to their need for certain elements to carry out their physiological functions [111]. A considerable number of metals are indispensable for cellular function, given their well-established biological roles. These include Fe, Na, K, Ca, Cu, Zn, Mn, Mo, Co, and Mg [112]. Collectively, these elements constitute the basis of cellular metabolism and are fundamental to processes such as enzymatic catalysis, biological polymer folding, osmoregulation, membrane transport, and signal transduction [113]. Consequently, intracellular biochemistry can produce a “chemical fossil record” that reflects the geochemical conditions of the early Earth [114]. The use of metals can be viewed as a primitive metabolic strategy that has influenced the evolution of protocells since the very origin of life [115]. Indeed, all classes of biological molecules, from nucleic acids to proteins and antibiotics, interact with metal ions, which explains why living cells invest considerable effort in tightly regulating their intracellular concentrations [113].

There is strong scientific evidence that metals (both essential and non-essential) play a crucial role in the global carbon cycle, particularly in the production of methane and the

consumption of CO₂ and H₂ [68,116–121]. All methanogens require Fe, Ca, Co, and Ni for optimal growth, while some also need Zn, Se, Mo, and/or W [117,122,123]. Iron can serve as a carrier or reactant in electron transfer, especially in the form of iron–sulfur ([Fe-S]) clusters [111]. The greater reliance on [Fe-S] clusters in methanogens has been suggested to result from increased bioavailability of Fe and S in anoxic early Earth environments (see Section 2.1) [124]. Nickel is an important cofactor present in methyl-coenzyme M reductase (Mcr); the final enzyme in the methanogenesis pathway and is a key component of the cofactor F₄₃₀ [117]. One hypothesis suggests that the decline in nickel availability around 2.7 billion years ago contributed to the reduction in atmospheric methane levels, thereby weakening the methane greenhouse and paving the way for the Great Oxidation Event (see Section 2.2) [125,126]. Cobalt is found in corrinoid cofactors (cobalamides), which are present within methyltransferases required for the transfer of the methyl group from methyl-tetrahydromethanopterin (methyl-H₄MPT) to coenzyme M (CoM) [127]. Calcium and magnesium are required at low concentrations for methanogen growth [128,129], but at high concentrations disrupt cellular processes, such as quorum sensing (i.e., cell-to-cell communication) [130] and electron transport [131]. Zinc is present in the heterodisulfide reductase (Hdr), where it plays a structural and catalytic role in the reduction in CoM-S-S-CoB, thereby recycling the cofactors required for the activity of the methyl-coenzyme M reductase enzyme [132]. Molybdenum is involved in catalyzing the first step of hydrogenotrophic methanogenesis through formylmethanofuran dehydrogenase, as well as the Wood–Ljungdahl pathway of CO₂ fixation. Similarly, the oxidation of formate to CO₂ and hydrogen (H₂) is catalyzed by formate dehydrogenase, which is Mo-dependent [133,134]. Under conditions of Mo limitation, tungsten can substitute for molybdenum due to its similar chemical properties [135].

Methanogens exhibit distinct responses to heavy metals. Although some of these are required for growth, others are well-known for their inhibitory effects on these microorganisms. Heavy metals are generally defined as metallic elements with a high atomic weight and a density at least five times greater than that of water [136]. The term “heavy metals” refers to both essential and nonessential trace elements, which may exert toxic effects depending on their intrinsic properties, chemical speciation, and concentration [137]. This definition classifies cadmium, copper, chromium, mercury, nickel, lead, and zinc as heavy metals. The abundance of these heavy elements in the dissolved state may have supported the evolution of methanogens early in Earth’s history [70]. It should be noted that the term “heavy metals” includes certain metalloids, such as arsenic and selenium, based on the historical assumption that heaviness correlates with toxicity [136]. As a consequence of such long-standing interactions, microorganisms are equipped with at least one As-detoxifying mechanism to prevent its toxicity [138]. This phenomenon may reflect the central role of arsenic toxicity in the evolution and survival of life throughout Earth’s history [139–141]. Methylation of arsenic, a common detoxification pathway, has been documented in methanogens [71]. Although arsenic can inhibit acetoclastic and hydrogenotrophic methanogenesis [142,143], recent reports indicate that arsenic can also promote acetoclastic methane production at low concentrations (10 µM) [144]. The effect of As on methane production seems site-specific. For example, in permanently anoxic microbial mats in the Atacama, Chile, methane was produced in the presence of 0.8–1.5 mM total arsenic [66,67]. A similar stimulating effect has been observed with Cd, which can increase methanogenesis rates up to nine-fold in pure cultures at concentrations of 10 µM [68,145,146]. In contrast, metals such as copper (Cu) and chromium (Cr) inhibit methanogenesis, although they do not necessarily lead to a significant reduction in biomass [117,147].

1.4. Evolution and Diversity of Methanogenic Pathways

As mentioned, methanogenesis is one of the oldest metabolisms on our planet, with estimates placing the onset no later than 3.8 to 3.5 Gya [6,50,148,149] (Figure 1) in Earth's secondary, $p\text{H}_2$ -rich atmosphere [50,100]. Phylogenetic analyses suggest that methanogenesis may have evolved earlier, possibly around 4.1 to 3.8 Gya [7], during the late heavy bombardment [150]. They diverged within the phylum Euryarchaeota ~3.51 Gya [149] based on estimates of the date of horizontal gene transfer (HGT) events between methanogens and cyanobacteria, which were calibrated to chemo- and microfossil evidence [149]. There is limited geochemical evidence of methanogenesis before their divergence within Euryarchaeota in the Archean. To date, the only geochemical evidence that places the evolution of methanogenesis during the Archean is CH_4 -rich fluid inclusions in metamorphosed rocks from the Pilbara craton, dated back to ~3.5 Gy [148] and confirmed by Mißbach and colleagues [151], as well as from isotopic signatures of ^{13}C from ~2.7 Ga-old kerogens [152]. Lipid biomarkers of archaeal membranes have been identified in the rock record dating back to 2.7 Gy [153]. However, certain abiotic processes could produce similar isotopic signatures, and the biogenicity of methane in Archean samples remains unresolved [66].




Figure 1. Timeline of early methanogen evolution and key geologic events [74,75,126,148,151,154–157]. The hydrogenotrophic and methylothermophilic metabolisms are ancient pathways of methanogenesis [65], with the aceticlastic pathway evolving ~474 Mya after an HGT event with clostridia, transferring two essential genes for the pathway (acetate kinase (*actA*) and phosphoacetyl transferase (*pta*)) to methanogens [149,158]. There are two evolution scenarios for the origin of Class I and Class II methanogen lineages. The first is a monophyly scenario in which Class II methanogens (those with a higher tolerance to oxygen stress due to the use of cytochrome/methanopheasine-mediated electron transfer mechanisms in the methanogenesis pathway, rather than flavin/ferredoxin-mediated electron bifurcation used by Class I methanogens. Class II methanogens also possess a wider variety of antioxidant enzymes and repair mechanisms that help them cope with reactive oxygen species [65]),

diverging from Class I around the onset of the GOE [65]. The second, alternative timeline depicts a dual lineage scenario in which Class I and Class II methanogens evolved separately [159]. The geologic timeline shows major events discussed in this review that have impacted methanogen evolution, as well as early geologic signatures and phylogenetic evidence (*) of methanogen activity and evolution.

1.4.1. Methanogenic Pathways

Methanogens are a diverse group of microorganisms that belong to several different taxa within the domain Archaea [10,149,160]. Methanogens produce CH_4 anaerobically [3]; however, they are not the only microorganisms to produce this gas [109].

There are three primary pathways associated with methanogenesis (Figure 2), also called the Wolfe cycle (see Sections 1.4.2–1.4.4), but methanogens are often metabolically flexible (i.e., mixotrophic), with some capable of utilizing more than one energy-generating pathway. For example, *Methanoscarchina acetivorans* is a model organism for aceticlastic methanogenesis, but can also grow on certain C_1 compounds, such as methanol, methylamines, and methyl sulfides [15]. While methanogenesis lies on the thermodynamic threshold of life ($\Delta G^0 = -5$ to -20 kJ mol^{-1}) [3,161–163], the energetic yield can be modified by combining methane production with other metabolic reactions. Coupling of one of the primary methanogenic pathways with other microbial metabolisms increases the overall energy yield, such as in syntrophic methanogenesis (e.g., with cable bacteria) ([11,164,165], interspecies hydrogen transfer [166,167], and reverse methanogenesis coupled to sulfate reduction (anaerobic methane oxidation) [167,168]. Energetic yield of the various reactions varies, depending on temperature and actual concentration of the reactants and products [2,161,168].




Figure 2. Comparison of the three most common methanogenesis pathways and their key intermediates. The hydrogenotrophic, methylo-trophic, and aceticlastic pathways differ in carbon substrates, electron sources, and metabolic intermediates but converge at methyl-coenzyme M (methyl-CoM), the immediate precursor of methane (CH_4). The hydrogenotrophic pathway involves the stepwise reduction of CO_2 through formyl-, methenyl-, methylene-, and methyl-bound intermediates on tetrahydrosarcinapterin

(H₄Spt). In the methylotrophic pathway, methylated compounds, including methanol, methylamines, and methyl sulfides, are converted directly to Ni-containing methyl-CoM, with internal electron transfer supporting methane formation. The acetoclastic pathway proceeds via activation of acetate to acetyl-CoA and cleavage into a methyl group and CO₂, with electrons derived from oxidation of the carbonyl group. Solid arrows indicate carbon flow, while annotated arrows denote electron transfer steps associated with reductive reactions. CoM, coenzyme M; H₄Spt, tetrahydrosarcinapterin; MF, methanofuran. Modified from [169]; red arrows CO₂ reduction, blue arrows indicate methylotrophy, orange arrows show the acetoclastic pathway.


1.4.2. Acetoclastic Methanogenesis

Acetoclastic (also referred to as acetoclastic) methanogenesis is perhaps the most common form of methane production in the environment [170,171], as acetate is likely abundant in many habitats [2,15,170]. The acetoclastic pathway is a chemoorganoheterotrophic process of methane production that utilizes acetate with a yield of $\Delta G^{0'} = -11.0 \text{ kJ mol}^{-1}$ [161], or acetate-like compounds, such as pyruvate or propionate (Equation (1)) [3,161,172,173]. For detailed information on the thermodynamics for various growth substrates, see [15,161].



1.4.3. Methylotrophic Methanogenesis

C₁ compounds (e.g., methanol, methylamines, methyl sulfides and thiols, methyl halogens, and methylated aromatics) get reduced through the methylotrophic pathway (Equation (2)) (another mode of chemoorganoheterotrophic growth) [3,161,172,173]. The energy yield of these compounds depends largely on the number of methyl groups available; for example, the $\Delta G^{0'}$ of methanol is $-290.0 \text{ kJ mol}^{-1}$, dimethylamine is $-167.0 \text{ kJ mol}^{-1}$, and trimethylamine is $-510.9 \text{ kJ mol}^{-1}$ [2,161].



1.4.4. Lithotrophic Methanogenesis

Lithotrophic methanogenesis (i.e., hydrogenotrophic methanogenesis) is a chemolithoautotrophic pathway where CO₂ (from CO₂(g) or HCO₃[−]/CO₂ (aq), CaCO₃ (S), carbon monoxide, formate, or some alcohols paired with CO₂) is reduced using H₂ as an electron source (Equation (3)) [161,174–176]. In some cases, calcium carbonate is used as a source of electron acceptor (i.e., as a source of CO₂) [174]. The $\Delta G^{0'}$ of Equation (3), using H₂(g) as an electron donor, is $-193.0 \text{ kJ mol}^{-1}$ [2,161].



In terms of their evolution, the hydrogenotrophic pathway is likely more ancient than acetoclastic and methylotrophic methanogenesis [50]. While methylotrophic methanogenesis was likely next to evolve [50], hydrogenotrophic methanogenesis may have been supported by the abundance of H₂ in an early anoxic atmosphere. Estimates for Archean atmospheric hydrogen concentrations range from a conservative 1000 ppm [50] to 10,000 ppm [5] or even 300,000 ppm [177], which are all well above modern levels of 530 ppb [178]. Lithotrophic methanogens also share hydrogenases with the Last Universal Common Ancestor (LUCA), which suggests LUCA likely utilized H₂ as part of its metabolism [50,179,180].

1.5. Anaerobic Oxidation of Methane (ANME)

ANAerobic MEthanotrophic (ANME) archaea and methanogens both encode the methyl-coenzyme M reductase (Mcr) system, including the *mcrA* gene. In methanogens, Mcr catalyzes the terminal step of methanogenesis (Figure 2), whereas in ANME archaea the same core enzymatic machinery operates in reverse to drive anaerobic methane oxidation [181–183]. Therefore, although ANME archaea are phylogenetically related to methanogens and share much of the core methanogenic metabolic machinery, they are functionally distinct.

ANME archaea typically form obligate syntrophic consortia with sulfate-reducing bacteria (SRBs) and, in some systems, with nitrate- or metal-reducing microbial partners, rather than with methanogens (see Section 2.2.2) [184–186]. These consortia allow electron transfer from methane oxidation to external electron acceptors, such as sulfate, mediated by diffusible intermediates or direct electron transfer mechanisms (DIETs) between species [45,187–189]. Although methanogens and ANMEs frequently coexist in the same anoxic environments (Figure 3) and may interact through the methane cycle, stable and obligate syntrophic associations between ANMEs and methanogens remain unresolved [184,190].

Phylogenetic analyses of *mcrA* and other methyl-coenzyme M reductase subunits indicate that ANME archaea and methanogens inherited the *mcr* operon vertically from a common methanogenic ancestor, rather than via recent horizontal gene transfer. The congruence between *mcr* phylogenies and archaeal core genes, together with the high conservation of operon structure and cofactor biosynthesis, supports the view that ANME evolved anaerobic methane oxidation through an ancestral methanogenesis [181,184,191]. Although divergent Mcr-like enzymes involved in alkane oxidation reveal ancient diversification within the Mcr family, there is no evidence of an exchange of Mcr between ANME and methanogens [192,193].




Figure 3. Spatial distribution of methane cycling by methanogens (MA), aerobic (MO), and anaerobic methane oxidation (AMO). MA exist and remain active in a wide range of oxic and anoxic environments, ranging from the deep biosphere [23,194] to their presence as bioaerosols in the atmosphere ([195]). The methane produced by MA supports methane oxidation (MO) in the water column and sediment–water interface. AMO is carried out in anaerobic niches in these habitats, as



well as the deep biosphere, indirectly by AMO bacteria (such as sulfate reduction, arsenate reduction, etc.) and in consortia with ANME. Bubbles show the degassing of methane production in sediments and from clathrates. The arrow in the left panel shows methane fluxes to the atmosphere and the lithosphere; the circular arrow indicates methane production and consumption (i.e., cycling).

2. Signatures of Methanogens Through Geologic Time

The ecophysiological background of methanogens reviewed above is instrumental in understanding the impact that these microbes have had on the evolution of our planet. From early Earth to modern times, methanogens have altered Earth's biosphere, atmosphere, and lithosphere. While there are challenges associated with biogeochemical evidence of methanogenesis in deep time (i.e., through the Archean and Proterozoic eons), indicators of methanogen activity in the rock record include depleted $\delta^{13}\text{C}$ isotopes -27‰ to -38‰ [196], of which the activity of biological methane oxidation (both anaerobic and aerobic) can alter these signatures [50], further fractionating ^{13}C isotopes [152]. In some cases, the isotopic fractionation effect deviates, e.g., hyperthermophilic methanogenesis in environmental extremes of high heat ($\sim 122^\circ\text{C}$) and hydrostatic pressure ($>40\text{ MPa}$), yields a fractionation of $<-12\text{‰}$ [35]. Methanogen-specific lipids have been used as an indicator of methanogens in sedimentary structures, both ancient and modern [148,197,198]. It should be noted that abiotic fractionation can also produce a negative $\delta^{13}\text{C}$ -methane value. In Fischer–Tropsch synthesis reactions, the methane that forms can have isotopic C values as low as -45‰ , depending on the iron mineral used in the reaction [199].

2.1. The Archean Eon (4–2.5 Gy)

The atmosphere of the Archean was defined by pervasive anoxia. Methane was delivered to early Earth's atmosphere by volcanism (which released H_2 , CH_4 , CO_2 , and NH_3), metamorphic reactions such as serpentinization (Equation (4)) [200] and Fischer–Tropsch synthesis reactions, which often concur in hydrothermal vents (Equation (5)) [199], and through impacts with comets and asteroids [6,201].



However, biogenic CH_4 production was likely the single most important source [6,50,148,196]. Large amounts of microbially produced methane in the atmosphere (100–15,000 times present concentrations) (Figure 3) [50] made up a major component of the carbon cycle and contributed to a greenhouse effect along with high atmospheric $p\text{CO}_2$ (0.6 bar) [61]. This would have supported early methanogenesis [202]. The high concentration of greenhouse gases kept global temperatures elevated [50,203], despite 20–25% lower solar radiation than today [50]. Secondary ion mass spectrometric analysis of kerogens recovered from 3.4 Gy Apex chert of Northwestern Western Australia revealed isotopic signatures consistent with methanogenesis, suggesting that this metabolism was likely an important process in the Archean biosphere [196]. This is consistent with the earliest evidence of methanogenesis from hydrothermal deposits dating back to 3.5 Gy [148].

2.1.1. Sulfur Cycling

The Archaeal oceans contained high levels of iron sulfides (FeS and FeS_2), hydrogen sulfide (H_2S), zero-valent sulfur (S^0), and possibly thiosulfate ($\text{S}_2\text{O}_3^{2-}$) [50,204]. All these reduced sulfur compounds can fulfill the nutritional requirements for methanogens [50,205,206]. Sulfur fractionation occurs during enzymatic S^0 reduction with H_2 [207]. Low sulfate and thus the absence of sulfate-reducing microbes in the early Archean [208,209], as well as a

lesser degree of $\delta^{34}\text{S}$ fractionation observed in the fossil record from that eon [207], has been used to corroborate an early (i.e., Paleoarchean) presence of assimilatory sulfur reduction by methanogens [50]. Furthermore, proteins of lithotrophic methanogenic species that contain Fe-S clusters are thought to be among the earliest inorganic catalysts, as they can self-aggregate when present in high concentrations, like those found in the iron and sulfide-rich Archean oceans [50].

In addition to inorganic sulfur, organic sulfur compounds made up an important part of the cycling of this element during the Archean as well [210]. Both biotic and abiotic reactions produce low-molecular-weight organic sulfur compounds [44]. Of these organosulfur compounds, methanethiol (CH_3SH) and dimethylsulfide ($(\text{CH}_3)_2\text{S}$) are important sources of electrons during methane generation in microbial mats [43,44]. Considerable amounts of methylated sulfides were produced by volcanoes and hydrothermal vents during the Hadean and Archean [210,211], and could thus have supported the evolution of methylotrophic methanogens early during Earth's history. A reconstruction of microbial metabolisms in microbialites of the Dresser Formation, Pilbara, Australia, suggests that sulfate- and arsenate-reducing microbes may have competed with methanogens for metabolic substrates [154,210,211], a notion that is based on the presence of sulfate and arsenic in these 3.5 Gy old microbial sediments. A role for methanogens in the arsenic cycle is further supported by the use of methyl arsenicals, which support methane production in some cases, thus generating energy in methylotrophic methanogens [212].

2.1.2. The Global Nickel Famine

Hydrogenotrophic and methylotrophic methanogens rely on the presence of several metals as cofactors (see Section 1.3.8), notably nickel used in F_{430} , iron in ferredoxins, and cobalt for methylcobalamine (Figure 2). The cooling of Earth's mantle during the Neoarchean (~2.7 Gya) caused a decrease in komatiitic volcanism, and consequently, an increase in iron but a significant reduction in the nickel and cobalt supply in magma [213], resulting in a sharp decline in the Ni:Fe ratio in banded iron formations [126,214]. By 2.5 Gya, the concentration of these metals dropped to <200 nM, and the rate of methanogenesis declined (see Section 2.1.3). This resulted in less greenhouse warming and a reduction in the formation of organic hazes (Figure 4). Reduced levels of these hazes increased the light regime at Earth's surface, allowing cyanobacteria and other phototrophs to thrive [215]. This global-scale nickel famine [215] was one of the critical drivers that set the stage for the Great Oxidation Event.




Figure 4. Impact of Ni-depletion during the Global Nickel Famine [126,214,215], 2.7–2.5 Gya. (A) Pre-GNF scenario, where methane production by hydrogenotrophic and methylotrophic MA produced

CH₄, possibly fixed N₂. The resulting flux to the atmosphere sparked photochemical reactions in the Paleoarchean atmosphere that led to the formation of organic hazes (reducing surface light and temperature). **(B)** The change in volcanism reduced Ni inputs during the start of the GNF, causing nickel-starved conditions. Reduced MA activity led to a decrease in organic haze formation, thus more light for photosynthesis, and a reduction in photochemical organic N production. Assuming cyanobacteria engaged in oxygenic photosynthesis, this set the stage for the Great Oxidation Event (~2.4 Gya) to occur. “+” indicates a positive impact, “-” a negative impact, and “?” an unknown effect.

2.1.3. Environmental Conditions and Photochemistry

Methanogens in the Archean would have experienced several ‘extreme’ conditions, like increased temperatures [57,77], higher amounts of UV surface radiation [85], and higher salinity in the oceans [57,61]. This may have been ‘normal’ for methanogens that had evolved at the time. Some localized environments may have hosted methanogens and could have been potential locations for the origin of life, such as hyperthermophilic conditions at hydrothermal vents [58] or exposure to ionizing radiation in nuclear geysers [54]. The average global ocean temperatures were likely >50 °C and could have supported thermophiles as potentially the most ancient extremophilic mode of life [57]. This is further supported by some studies that suggest that LUCA could have been a thermophilic microbe [180,216]. Alternatively, studies focusing on hydrogen and inorganic phosphate ($\delta^{18}\text{O}_\text{P}$) isotopes report a much lower ocean temperature of 26 to 40 °C [217,218]. Under this scenario, or if it evolved in a non-marine system, LUCA may have been a mesophile. A mesophilic origin of LUCA is also supported by the thermoreduction hypothesis [219]. Increased ocean temperatures could have a greater potential for methane production from hydrogenotrophic and mixotrophic methanogens [193]; however, an ocean temperature of 40 °C could have potentially been (near) optimal for methane production from both acetoclastic and hydrogenotrophic species [2,220].

Exposure to UV radiation at the surface of the planet was greater during the Archean due to the lack of ozone in the upper atmosphere [85]. Physical barriers to this radiation, perhaps protected microbes in the ocean, such as the high abundance of iron [40]. A variety of photopigments in microbial mats and microbialites could have quenched this damaging high-energy radiation [4]. Salinity reconstructions of the Archean oceans suggest that salinity was 1.5 to 2 times higher than it is at present [57], which would have supported halophilic species.

The impact methanogens had on the Archean carbon cycle not only affected the increased global temperature at the time but had a lasting effect on the evolution of Earth’s secondary and tertiary atmospheres, as methane likely supported the irreversible oxidation of the atmosphere, oceans, and terrestrial systems in the Proterozoic [50]. Methane was removed in the Archean upper atmosphere by photochemical breakdown of CH₄ into H· and CH₃· by UV radiation, providing a source of OH radicals in the upper atmosphere [50,221] (Equation (6)).



While it is still debated, anaerobic methane oxidation (possibly by methanogens and heterotrophic acetogens [15,222]) coupled to sulfate, arsenate, nitrite or iron reduction [154,223] could have played a role in the removal of methane in the late Archean [152] (Figure 5B). Methanogens may also have been critical in the formation of an organic haze. When the atmospheric ratio of CH₄:CO₂ exceeds 0.1, a photochemical reaction of these gases using energy from UV radiation produces a variety of n-alkanes [224]. Atmospheric photolysis of CH₄ and N₂ formed HCN and NH₄OH [225], adding an essential nutrient for life to the global oceans. This Archean organic atmospheric haze was not as thick as, e.g., the current haze on

Saturn's moon Titan (which has a $\text{CH}_4:\text{CO}_2$ ratio > 1) and may have had a climate-stabilizing effect on our planet [226].




Figure 5. Methane transformations in the carbon cycle during the Archean and Proterozoic eons. (A) Atmospheric CH_4 and O_2 concentrations from ~4.0 Gy to 0.5 Gy, adapted from [61]. (B) The role of CH_4 in the iron-rich oceans of the Archean, when there was a large flux of CH_4 from methanogenesis (MA) in the oceans; sinks of methane included photochemical breakdown in the upper atmosphere and possible anaerobic oxidation of methane (AOM). The CH_4 production exceeded removal, which

resulted in a strong greenhouse effect and warming characteristic of the Archean atmosphere. Arrows show the cycling of CH_4 , including the flux of CH_4 to the atmosphere by MA and the removal and decomposition of CH_4 . (C) CH_4 and the carbon cycle in the Paleo-Proterozoic. Oxygen produced by cyanobacterial photosynthesis emerges as a novel oxidant for microbial removal (i.e., methane oxidation; MO) (sink) of CH_4 from the atmosphere. Reduced hydrothermal activity in the Paleo-Proterozoic could have contributed, in part, to decreased CH_4 , though methanogen activity in sediments (bubbles) was still an important sedimentary microbial process. (D) Atmospheric CH_4 is further reduced during the Meso-Neo-Proterozoic due to increased removal by oxic and anoxic methane oxidation, i.e., reverse methanogenesis coupled to sulfate and/or arsenate reduction, as well as through photochemical reaction of CH_4 with O_3 (through $\text{OH}^\cdot$). Clathrates (bubbles) could have been abundant and an important source and sink for CH_4 . The Proterozoic saw a rise in thrombolites due to increased competition and bioturbation of microbial mats [227]. (E) CH_4 production by syntrophic microbial processes through carbon degradation in sediments during the Archean and Meso-Neo-Proterozoic Eons. H_2 and organic products (such as CH_3COOH) from fermentation (Ferm) could have supported MA in sediments. Anaerobic oxidation of methane (AOM) by sulfate reduction (SR; shown here, but possibly also respiration using other electron donors such as nitrate, selenate, or arsenate (see Section 1.3.8) could have contributed to the removal of CH_4 in sediments and promoted mineral formation (such as (arseno)pyrite). Active MA combined with other microbial processes [228] can drive carbonate mineral precipitation (such as CaCO_3) in sediments and microbial mats.

2.1.4. Methanogens in Early Hydrothermal Vents

Hydrothermal vents were important sites for the evolution and diversification of early life [58,154,229–231]. The oldest fossilized hydrothermal vents date back >1.43 Gy [232], but these sites must have existed much longer, possibly 4.3 Gya [233]. Vents (fluid) of the Archean were likely highly alkaline (pH 8.5 to >12) [234] and high temperature (~290 to 350 °C) [154,230], vastly different than the black smokers (350 °C to >450 °C; pH 2–3) and white carbonate chimneys (50–90 °C; pH 9–11) found in today's oceans [58]. Similarly, the physicochemical gradient of vent fluid, in their interaction with the surrounding neutral to acidic seawater, was the reverse of what is observed today, where vent fluids interact with a more alkaline ocean [230]. Still, modern-day vent systems may provide relics of ancient biogeochemical interactions (for further reading, see [58]).

Archean vent fluids were rich in iron and sulfide [154,235], CO_2 and CaCO_3 [230], and SiO_2 [230]. While the Fe content of ancient vent fluid and, likewise, a hydrothermal contribution to Archean banded iron formation is debated [230,236], serpentinization in vent fluids supplied abundant H_2 , C_{1-4} intermediates, and metals, which could have supported hydrogenotrophic methanogens, acetogens, and sulfate reducers as early metabolisms [155,180,231,237]. For example, transition metals Co, Ni, and Fe from vents could have been a source of early cofactors in the acetyl-CoA (Wood–Ljungdahl) pathway found in modern-day methanogens, acetogens [237,238], and possibly in LUCA [180,231]. Additionally, some methanogens can fix N_2 using dinitrogenase reductase [239]. Such nitrogenases are ancient and were likely important for early biochemistry at hydrothermal vents [231]. Mo may have also been abundant in Peridotite-hosted alkaline systems [240]. Hydrothermal activity in the Archean could have also been a source of oxidized compounds, such as SO_4^{2-} , which could have expanded element cycles and the potential for microbial metabolisms (i.e., sulfate reduction) [154,234].

2.1.5. Microbial Mats of the Archean

Microbial mats are semi-closed sedimentary biofilms comprising bacteria and archaea, in a matrix made up of extracellular polymeric substances [4]. Microbial mats in the Archean were likely dominated by H_2S - and H_2 -utilizing microbes, including lithotrophic methanogens and perhaps methane oxidizers, as well as other early het-

erotrophs [4,67,154,241–243]. However, it is thought that the earliest microbial mats may have been chemolithotrophic rather than autotrophic [4,67,154]. Microbial mats could have been among the earliest stable habitats [243], and the combination of a complex geochemical composition, a high cell density, and a close proximity of organisms would have made them conducive sites for the evolution of early life [4,67,243] and the diversification of metabolisms [154]. At the onset of the Neoproterozoic, cyanobacteria in microbial mats combined two photosystems [244] and the oxygenation of Earth's oceans, atmosphere, and protocontinents took off [245]. Although the cyanobacteria date back ~2.8 Gy, their photosynthesis can also be supported by reduced sulfur species, and thus their first appearance cannot be used as evidence for oxygen production. In fact, anoxygenic photosynthesis (e.g., using reduced sulfur as an electron donor) likely prevailed for tens of millions of years in these phototrophs [246]. Microbially induced sedimentary systems [247] and stromatolites [248] record the preservation of microbial mats in the fossil record. The earliest evidence for this is found in the Pilbara dating back ~3.5 Gy [241,249,250]. Even though the true nature of the 3.7 Gy old stromatolitic structures remains under discussion to date [251–253], these fossilized lithified microbial mats, which include stromatolites, thrombolites, dendrolites, and leiolites, are the main systems that provide insight into the biogeochemistry of the Archean world [4,248,254].

2.2. The Proterozoic

2.2.1. The Great Oxygenation Event and the Paleoproterozoic (2.5 to 1.6 Gy)


The evolution of oxygenic photosynthesis ~2.5 Gya spurred the permanent oxidation of the planet (Figure 5A) and led to greater oxidation states of the sulfur cycle [50,204] and expansion to the carbon cycle through the evolution of new metabolic pathways [50]. The abundance of methane left over from the Archean likely contributed to the loss of H₂ from photolysis and the irreversible, but slow, oxidation of the atmosphere, terrestrial environments, and Earth's oceans [209]. Anoxia in the atmosphere was prolonged due to icehouse conditions [255]. Reduced atmospheric CO₂ could have contributed to quicker removal (through oxidation) of methane by O₂ [255] (Figure 5C). This relatively rapid removal of CH₄ likely contributed greatly to/was an important factor in the reduction in global temperatures [50,203]. The oxidation of sulfide increased the sulfate concentration in the oceans and set the stage for the evolution of sulfur metabolisms 2.3–2.4 Gya [204].

2.2.2. The “Boring Billion” (1.8 to 0.8 Gy)


The “Boring Billion” refers to a period characterized by extreme geologic (minimal tectonic activity), climatic (warm, lacking global glaciations), and biological stability [256,257]. A main reason for stasis in eukaryotic biodiversity was that the increase in oxygen transformed abundant pyrite into sulfate, which supported a robust sulfur cycle. The onset of oxidation of reducing pre-GOE conditions (including vast pyrite reservoirs) resulted in euxinic, stratified oceans with oxygenated surface waters and sulfidic, anoxic bottom waters [50]. Anoxic oceans contained an abundance of hydrogen sulfide, which some methanogens used as a sulfur source [50]. Reactions between the oceanic ferrous iron (Fe(II)) and atmospheric O₂ lead to the precipitation of iron sulfide minerals, like pyrite. This limited the availability of Fe(II) and other trace metals, such as nickel [258], which are essential for methanogen growth [62]. Some species of methanogens can reductively dissolve nickel and iron sulfide minerals, making them more bioavailable in euxinic conditions [118]. This capacity could have been used to survive in the anoxic zone of the oceans during the Boring Billion, which would have supported a continuing methane flux to the atmosphere [203]. However, methane concentrations were presumably reduced to less than 10 ppm in the mid Proterozoic [203] from Archean levels in the magnitude of 10² to

10^3 ppm [50,221], either through removal of large amounts of CH_4 from the atmospheric reservoir and/or a reduction in CH_4 flux to the atmosphere [203].

Sulfate reduction-supported anaerobic oxidation of methane (see ANME; Section 1.5) could have limited the impact of methanogenesis on the global climate [203], in which CH_4 is removed via anaerobic methane oxidation (Figure 5E) (Equation (7)):



Sulfate-reducing bacteria (SRB) consume H_2 (Equation (8)) produced during methanogenesis, lowering the availability of this for methanogens [46]:



This syntrophic relationship is energetically feasible with an overall $\Delta G^0 = -16.6 \text{ kJ.mol}^{-1}$ [259]. A similar role in anaerobic methane oxidation has been attributed to arsenate, iron, manganese, and nitrate reducers (see Section 1.3.8) [260]. Anaerobic methane oxidation thus created a sink for methane in the Proterozoic oceans (Figure 5D), removing most of this gas before it could escape into the atmosphere [203]. Simultaneously, methane removal by UV (Equation (6)) slowed down when oxygenation (and consequently O_3 production) of the atmosphere commenced [220]. These two events combined may help explain the steady but severely reduced temperature during the Boring Billion [256].

2.2.3. Methanogens Through Glaciations

The Ending of Marinoan “Snowball” Earth (650–635 Mya)

The continual decrease in the greenhouse gas concentration, including CO_2 and CH_4 , through the mid-Proterozoic and Neoproterozoic eras triggered extensive low latitude glaciation, and perhaps the most extensive ice age of the planet [74,75]. Deglaciation and recovery of primary production were part of a feedback loop, which resulted in the reestablishment of strong euxinic oceanic conditions [74,75]. This was the result of extensive continental weathering following the accumulation of atmospheric CO_2 during the glacial period [74]. A large flux of CH_4 into the atmosphere played an important role in deglaciation [74] from active methanogenesis that may have utilized methyl sulfide compounds from seawater [74,75], volcanic activity [261], and the release of methane from methane clathrates [74,262]. Though still debated, methanogenesis likely played a key role in the formation of these methane clathrates [262]. Methanogens that produce CH_4 under low temperature and/or high pressure are known to contribute to clathrate formation [263]. The production and consecutive release of methane from clathrates to the atmosphere has also been invoked later in the more recent events like the melting of the Laurentide–Cordilleran ice sheet [264].

2.2.4. Methanogens Through Mass Extinctions

Widespread and regional anoxic and euxinic ocean conditions have been characteristic of multiple extinction events, such as those found in Western Tethys and Panthalassa at the end Triassic mass extinction (~201.6 Mya) [155,156] and during the Permian mass extinction (252 Mya) [157]. A rise in global temperature following rapid release of greenhouse gases (CO_2 and CH_4) [76,155,156], via volcanism (e.g., the Siberian Traps) [265], destabilization of methane reservoirs (e.g., methane clathrates/hydrates and permafrost) [155,157], and/or microbial sources (i.e., methanogenesis) [155]. Reservoirs could have fueled methanogen activity, creating a positive feedback loop that further contributed to a global rise in temperature, which in turn led to the melting and release of additional methane. The End Triassic mass extinction experienced increases in acetate and H_2 availability and a decrease

in ocean sulfate concentrations that may have contributed to increased methane flux from methanogen activity in the oceans [155]. Likewise, following the Permian mass extinction, an euxinic photic zone and low ocean oxygen concentrations led to a buildup of organic material in sediments [266]. The degradation of this organic matter was likely both a substrate for methanogenesis and a source of acetate, which could have supported the evolution of the modern-day high efficiency acetoclastic pathway [266] in *Methanoscarchina* spp. [158]. Increased CH₄ and H₂S fluxes from the oceans during the Permian reduced atmospheric O₂ and O₃ by ~14% and consequently decreased CH₄ oxidation in the atmosphere [155]. Under lower atmospheric O₂ conditions like this, aerobic and anaerobic methane oxidation may play a critical role in the removal of CH₄ in environments with and without oxygen, respectively [267]. Similar conditions may have been present during other mass extinction events [268], even though methanogenic activity during these times is not discussed extensively in the literature.

2.2.5. Resurgence of Methanogens and Microbialites Following Mass Extinctions

Methanogens often thrive following mass extinction events, significantly delaying ecosystem recovery, notably in marine environments [266]. The Siberian Traps replenished oceanic nickel [265], thereby stimulating massive CH₄ production in a “Global Nickel Feast”. This in turn increased temperatures, lowering gas (including O₂) in the water column. Aerobic oxidation of CH₄ further depletes O₂, while anaerobic oxidation produces toxic sulfide. This led to a scenario with a lethally hot photic zone and euxinic bottom waters [269]. Thus, methanogens contributed to persistent toxic conditions, resulting in “dead zones”. Consequently, re-establishment of complex life was difficult, slowing the return of diverse ecosystems [270].

There is often an increase in the abundance of microbialites following mass extinction events [271] due to a carbonate-saturated, alkaline ocean, increased availability of nutrients (e.g., nitrogen-fixing cyanobacteria, weathering) [242,243], and a lack of competition with decimated eukaryote populations (i.e., decreased bioturbation and grazing pressure) [271–275]. Following the Permian–Triassic extinction event, the abundance of organic carbon combined with dysoxic conditions likely supported sulfate reduction [276] and methanogenesis [266], and sulfidic conditions possibly caused a shift from oxygenic to anoxygenic photosynthesis [277] by *Gakhumella*-like cyanobacteria [278]. However, early Triassic oceans may have had a low oceanic sulfate concentration [279], which would have profited methanogens [266]. Importantly, all three anaerobic metabolisms favor lithification of mats, potentially forming stromatolites [4,228]. A similar resurgence of microbialites under anoxic or dysoxic conditions, possibly supported in part by methanogens, occurred, e.g., during the Early Silurian, following the Late Ordovician Extinction Event [280] and the end-Devonian mass extinction [281].

2.3. Past, Present, and Future—Methanogens in Climate Change

Methanogens still play an essential role in the modern-day climate, producing over 70% of methane emissions annually [282,283]. Although methane has a lower abundance in the atmosphere (~1.4 to 1.9 ppm; [50,284]) than CO₂, at ~360 to 390 ppm [285,286], it is about 30 times more potent as a greenhouse gas [3]. Methanogens process in excess of 500 Tg C y^{−1} through their metabolism in the production of CH₄ and biomass [287]. However, methanogens may play an additional role in the global carbon cycle by mediating precipitation and dissolution of carbonate minerals [288], thus impacting long-term storage of atmospheric CO₂ (Figure 6) [176]. This makes their specific role in the carbon cycle more complex than previously thought.






Figure 6. The multiple roles of methanogens in the modern-day carbon cycle. Processes shown in red contribute to greenhouse gas production. Part of the methane is removed through microbial methane oxidation (A)MO and photochemical reactions, where e is a generic electron donor (e.g., sulfate, nitrate, arsenate, iron, or manganese) that is reduced to eH . Processes depicted in green sequester CO_2 , storing it long term as carbonates. Dashed arrows show the potential contribution of methanogen metabolism in the precipitation of carbonate minerals, which contributes to the storage of carbon in the lithosphere. The metabolic activity of methanogens may locally alter pH (a part of energy generation) and produce copious amounts of EPS (seen as a veil surrounding the cells in the microphotograph (yellow arrow)). These EPS bind cations (e.g., Ca, Mg) and act as a nucleation site, while elevated pH enhances the carbonate precipitation in microbial mats [228]. This contribution by methanogens may ultimately impact the production of microbialites [154]. Solid arrows show known effects; dashed lines indicate hypothetical effects.

Lithotrophic metabolisms are believed to promote the precipitation of carbonate minerals [288] (Equation (3)) where removal of CO_2 favors the production of CO_3^{2-} ions, increasing pH (Equation (9)) in the surrounding environment. This could promote the precipitation of carbonates, such as CaCO_3 [4,228] (Equation (10); sum of Equations (9) and (10); Equation (11)).



However, other types of methanogenesis have different effects on the CaCO_3 budget. Methanogens produce exopolymeric substances (EPS) [52,108,289], which could also aid in the process of mineralization [228,248]. On the other hand, acetoclastic (Equation (1)) and methylotrophic (Equation (2)) species may promote the dissolution of CaCO_3 minerals through a decrease in pH, where CO_2 is produced as a product of the metabolisms (Equations (1) and (2)). The increase in CO_2 would favor HCO_3^- ions (Equation (12)) and promote the dissolution of CaCO_3 minerals (Equation (13); net for acetoclastic methanogen-

esis) Equation (14)), in which case, carbonate minerals could serve as a potential carbon source for methanogen growth [176].



This dual role that methanogens play in the global carbon cycle and their metabolic and biogeochemical role in CaCO_3 production remains unstudied, leaving gaps in our understanding of the capacity of microbes to regulate planetary climates. Currently, studies of methanogen-driven carbonate mineralization are limited to wastewater [289–291], carbonates in authigenic marine sediments [260,292], and Mn carbonates formed as a product of the methanogenic pathway [293]. Methanogens are abundant in many sediments, including microbial mats [25,44], which makes mat ecosystems prime locations to study methanogenic CaCO_3 precipitation [43,294]. Methanogens occupy a unique niche within mats, relying mostly on non-competitive methylated compounds for growth, such as methylamines and methylsulfides are not used by SRB [44,295,296].

3. Methanogens in Microbial Mats as Biosignatures for Life

Organisms in microbial mats must cope with a range of not only extreme but also fluctuating conditions [4]. Sulfide and oxygen gradients in these systems fluctuate over diel cycles [4,228,248], with O_2 peaking during the early afternoon, along with peak photosynthesis by cyanobacteria (Figure 7), reaching daily peaks of >600% O_2 saturation in lithifying mats [4,64]. In contrast, peak sulfide concentrations are found near the mat surface, peaking at the end of the night, associated with sulfate reduction activity [4,228].




Figure 7. Relative concentration of O_2 , CH_4 , and sulfide with depth in microbial mats and expected distribution of methane oxidizers (MO; gray), cyanobacteria (Cyano; green), sulfate reducers (SR; red), methanogens (MA; dark blue), and anaerobic methane oxidizers (AMO, including ANME; light blue). Profiles and abundance are estimates of relative daytime concentration. Modified from [25,53].

While the modern-day diversity of archaea in microbial mats is limited [294,297], methanogens have been identified (via their metabolic activity) in the anoxic but also the oxic zone of microbial mats [25,43,44,64,298]. Microbial mats are often found in extreme environments; in hypersaline conditions, when the salinity exceeds 180 g.kg^{-1} , methanogens can outcompete other anaerobic heterotrophs, notably SRB [64]. Desiccation impacts microbial mat communities through wetting and drying cycles [64] that enhance compatible solute production. As noted above (Sections 1.4 and 2.3), several of these compounds are precursors to methylamines and methylsulfides used by methanogens [43,298]. Desiccation has been proposed as a survival strategy in methanogens [52]. These authors reported that the viability of *Methanosarcinales* was significantly enhanced by EPS, and that desiccated organisms had a higher resistance to high temperature and, to a lesser extent, also to oxidative conditions. These properties may prove useful in extraterrestrial environments, such as Mars (see Section 4) [38].

3.1. The Evolution of Microbial Mats and Stromatolites

Microbial mats are organosedimentary systems made up of complex microbial communities embedded in an exopolymeric matrix [248]. Some mat communities facilitate microbially induced CaCO_3 precipitation and produce lithifying mats [4,228,288]. The precipitation potential is determined by the combined metabolic activities of the entire microbial community and by the environmental (physicochemical) conditions [4,299]. Complete element cycling, notably of C, O, and S, is very efficient in these semi-closed, self-sustaining ecosystems [288]. Microbial mats could have been among the first complex microbial communities to evolve on Earth [243], in shallow ponds near hot springs [154,300], appearing in the fossil record as stromatolites as early as $\sim 3.5\text{--}3.7$ Gya [251,253]. A variety of available chemical elements supporting a range of metabolisms, combined with a close proximity of an assortment of microorganisms [301] characteristic of mats, could have been instrumental in the diversification of microbial life.

Both lithified and non-lithified (soft) mats' biofilm systems were more abundant during the Archean and Proterozoic eons, but in modern times have been forced to environments with extreme conditions likely due to competition with eukaryotes [4,227]. The presence of methanogenic activity in the oldest microbial mats and stromatolites has been suggested, mainly based on carbon isotopic evidence and metal availability [66,154,302,303]. Methanogens are still involved in the formation of modern microbialites [288], although their actual role in the lithification process remains largely unknown.

3.2. The Conundrum of Methanogenesis and Oxygen in Mats

It is yet unclear how methanogens cope with the fluctuating conditions in microbial mats. Notably, the presence of supersaturated oxygen concentrations during the early afternoon creates a challenge for these anaerobic organisms [25] due to the oxygen-sensitivity of methyl-coenzyme M reductase and other metalloenzymes in methanogenesis [304]. Although methanogens possess biochemical properties, such as robust ether-linked membrane lipids [305] or pseudomurein cell walls or proteinaceous sheaths [306] that make these microbes more resistant to extreme conditions (e.g., pH, temperature, mechanical stress, salinity), other survival mechanisms can be employed. Methanogens do not form endospores but survive in a dormant state followed by rapid recovery [307]. Such a state of reduced activity is accomplished during desiccation. Additionally, there are specific solutions to oxygen-related stress: although glutathione has not been found in methanogens, enzymes

like superoxide dismutase, catalase, and peroxidase cope directly with reactive oxygen species, and repair mechanisms of oxygen damage (e.g., disulfide reductases) also exist [108,308]. Other physiological adaptations include modified electron transport chains, production of protein-rich EPS, formation of cell clusters, polyphosphate accumulation, or a metabolic shift to methylated substrate use [52,108,306]. Quorum sensing has been reported in some methanogens [108,130], which hypothetically could allow for cyanobacteria, methanotrophs, and methanogens to live in close proximity while metabolically active, e.g., in the surface layer of a microbial mat (Figure 8). Such a model was proposed previously for the coexistence of sulfate reducers, sulfide oxidizers, and cyanobacteria [309]. Last but not least, other ecological adaptations may enable methanogens to thrive in oxic conditions. These include formation of consortia [310], syntrophism [311], and interspecies transfer of electrons [312].




Figure 8. Hypothetical interactions between cyanobacteria, methanogens, and methane oxidizers in the oxic layer of a microbial mat. Quorum sensing (red arrows) between cyanobacteria, methanogens (MA), and aerobic methane oxidizers (MOs) would allow high activities of MA by coordinating the metabolisms of all three functional groups of microbes despite highly unfavorable (i.e., toxic, O₂-containing) conditions for MA. A similar scenario was proposed to explain sulfate reduction near the surface of microbial mats [309] as observed by several researchers (see [4]). Modified from [308]. “+” indicates a positive effect, “-” a negative effect, and “+/-” either a positive or a negative effect, and “?” an unknown effect.

4. Methanogens in the Search for Extraterrestrial Life

The fundamental biochemical principles observed on Earth are hypothesized to be universal characteristics of life throughout the cosmos [313]. Methanogens are of prime interest for possible life on exoplanets. The primary product of their metabolism, methane, is an ideal gas for detection in remote atmospheres due to the unique absorption spectrum and short photochemical lifespan [6]. To date, methane has been detected on several planetary bodies in our solar system that may be fit for supporting methanogenesis-like life, including Mars [314], Saturn's moon Enceladus [315], and Titan [316]. The detection of methane on other planets and its short residency time in planetary atmospheres suggests sources for an active methane flux on these planetary bodies, which on Earth, primarily is of biological origin [6]. The potential of methane, through the greenhouse effect, to warm planetary atmospheres may expand the possibility for life beyond the "Goldilocks" zone [10].

Many studies have assessed methanogen survival under Mars-like conditions [31,37,38,40,317] and their potential habitability on ice worlds like Enceladus [13]. The extreme environments in which many methanogens are found are used as analogs for extraterrestrial bodies [10,36,197]. For example, soda lakes are studied for their potential as analogs to Mars's ancient ocean [36,318]. Similarly, hydrothermal vents are expected to exist on other worlds, like Enceladus, where products of serpentinization reactions and silica-rich particles have been detected [13]. Thermophilic methanogens may be able to survive at hydrothermal vent sites on Enceladus, and the predicted temperature and pressure associated with this ecological extreme [13]. Survival of methanogens to conditions on Mars (e.g., low pressure at the surface (6 mbar), desiccated, acidic soil, high UV radiation at the surface, and freezing surface conditions and permafrost due to lack of an atmosphere) [31,37,38,40,84,317,319,320] has supported the idea of potential subsurface life on the planet [37], or more broadly, a subsurface source for methane [321]. Microbial mats themselves are useful biosignatures due to their self-sustainability, resistance to extremes, and abundance on early Earth [197].

5. Conclusions

Methanogens were among the earliest microorganisms to evolve on Earth and continue to thrive in a wide range of both seemingly "non-extreme" and "extreme" environments today, several of which are similar to those throughout Earth's history (Figure 9). Microbial mats are a prime example of ecosystems that evolved with our planet (or, arguably, changed our planet's properties) and were present in abundance during the Archean and Proterozoic eons. Methanogens were likely part of the early stromatolite communities and continued to be ubiquitous in microbial mat ecosystems through time [243]. The transition from strictly oxic conditions during the day and anoxic ones during the night may have supported methanogens to be well-adapted to a wide range of sometimes fluctuating conditions (e.g., diel fluctuations of O₂, sulfide, pH, and light gradients) [4]. The lack of a proper understanding of the ecophysiological capabilities and flexibilities (e.g., EPS properties, stress responses, viral–host relationships, cell-to-cell communication) is a major gap in the understanding of the role and evolution of methanogens and their impact on our planet through geologic time. Their elementary metabolism (e.g., inorganic anaerobic respiration), efficient use of a minimal energy yield, and the simplicity of their main metabolic product (i.e., CH₄)—which is observable remotely and supports a diversity of photochemical reactions—make methanogens the ultimate candidates for astrobiological studies and for understanding microbial life on early Earth.




Figure 9. Summary of the diverse roles of methanogens (MA) in biogeochemical processes, including positive and negative feedback, through geologic time that impacted climate. The major processes methanogens participate in include (i) consumption of organic carbon (e.g., XCH_3 , or methylated compounds), producing biomass and CH_4 and fractionating stable isotopes of carbon; (ii) consumption of H_2/CO_2 (e.g., from Earth's early atmosphere), potentially changing the redox balance; (iii) precipitation of minerals (e.g., carbonates); (iv) use and fractionation of trace elements (e.g., Ni) that stimulated their growth and metabolic activity; (v) the release of CH_4 to the environment, stimulating methane-consuming microbes; (vi) the release of CH_4 to the atmosphere, acting as a greenhouse gas, contributing to organic haze formation that yield organic nitrogen, thereby regulating climate and, consequently, play a role in beginning and/or ending mass extinctions and the ecological recovery after these icehouse events. Note that the strong isotopic fractionation of carbon and nickel by methanogens can be used to reconstruct early ecosystems on Earth. The production of methane and fractionation of carbon is a potential biosignature that can be used in the search for life beyond our planet. " XCH_3 " represents a generic methyl group donor for CH_4 production, "+" indicates a positive effect, "++" a strong positive effect, "-" a negative effect, and "?" an unknown role.

Funding: This research received no external funding.

Data Availability Statement: No original data were used for the preparation of this manuscript.

Acknowledgments: PV acknowledges support from the NASA Exobiology NNN22ZDA001N-WXO grant. PT received support from the National Science Foundation Graduate Research Fellowship Grant No. 2136520. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science

Foundation. PB acknowledges support of AAPG-1-HARMI-ANR. This work was also supported by Start-Up funds provided to MGS-M by the Department of Molecular and Cell Biology (MCB) and the College of Liberal Arts and Sciences (CLAS) at the University of Connecticut (UConn). We thank two anonymous reviewers for their constructive comments that helped improved this manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Megonigal, J.P.; Mines, M.E.; Visscher, P.T. Anaerobic Metabolism: Linkages to Trace Gases and Aerobic Processes. *Biogeochemistry* **2004**, *8*, 317–424. [\[CrossRef\]](#)
2. Conrad, R. Complexity of Temperature Dependence in Methanogenic Microbial Environments. *Front. Microbiol.* **2023**, *14*, 1232946. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Buan, N.R. Methanogens: Pushing the Boundaries of Biology. *Emerg. Top. Life Sci.* **2018**, *2*, 629–646. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Visscher, P.T.; Gallagher, K.L.; Bouton, A.; Vennin, E.; Thomazo, C.; White, R.A., III; Burns, B.P. *Treatise on Invertebrate Paleontology, Part B, Volume 2, Chapter 3: Microbial Mats*; Kansas University Press: Lawrence, KS, USA, 2022; Volume 163, pp. 24–56.
5. Kadoya, S.; Catling, D.C. Constraints on Hydrogen Levels in the Archean Atmosphere Based on Detrital Magnetite. *Geochim. Cosmochim. Acta* **2019**, *262*, 207–219. [\[CrossRef\]](#)
6. Thompson, M.A.; Krissansen-Totton, J.; Wogan, N.; Telus, M.; Fortney, J.J. The Case and Context for Atmospheric Methane as an Exoplanet Biosignature. *Proc. Natl. Acad. Sci. USA* **2022**, *119*, e2117933119. [\[CrossRef\]](#)
7. Battistuzzi, F.U.; Feijao, A.; Hedges, S.B. A Genomic Timescale of Prokaryote Evolution: Insights into the Origin of Methanogenesis, Phototrophy, and the Colonization of Land. *BMC Evol. Biol.* **2004**, *4*, 44. [\[CrossRef\]](#)
8. Rothschild, L.; Mancinelli, R. Life in Extreme Environments. *Nature* **2001**, *6823*, 1092–1101. [\[CrossRef\]](#)
9. Lyons, T.W.; Tino, C.J.; Fournier, G.P.; Anderson, R.E.; Leavitt, W.D.; Konhauser, K.O.; Stüeken, E.E. Co-evolution of Early Earth Environments and Microbial Life. *Nat. Rev. Microbiol.* **2024**, *22*, 572–586. [\[CrossRef\]](#)
10. Taubner, R.S.; Schleper, C.; Firneis, M.G.; Rittmann, S.K.-M. Assessing the Ecophysiology of Methanogens in the Context of Recent Astrobiological and Planetological Studies. *Life* **2015**, *5*, 1652–1686. [\[CrossRef\]](#)
11. Jung, H.; Yu, H.; Lee, C. Direct Interspecies Electron Transfer Enables Anaerobic Oxidation of Sulfide to Elemental Sulfur Coupled with CO₂-Reducing Methanogenesis. *iScience* **2023**, *26*, 107504. [\[CrossRef\]](#)
12. Yin, Q.; Gu, M.; Hermanowicz, S.W.; Hu, H.; Wu, G. Potential Interactions between Syntrophic Bacteria and Methanogens via Type IV Pili and Quorum-Sensing Systems. *Environ. Int.* **2020**, *138*, 105650. [\[CrossRef\]](#)
13. Taubner, R.S.; Pappenreiter, P.; Zwicker, J.; Smrzka, D.; Pruckner, C.; Kolar, P.; Bernacchi, S.; Seifert, A.H.; Krajete, A.; Bach, W.; et al. Biological Methane Production under Putative Enceladus-like Conditions. *Nat. Commun.* **2018**, *9*, 748. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Morgavi, D.P.; Forano, E.; Martin, C.; Newbold, C.J. Microbial Ecosystem and Methanogenesis in Ruminants. *Animal* **2010**, *4*, 1024–1036. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Ferry, J.G. *Methanosarcina acetivorans*: A Model for Mechanistic Understanding of Aceticlastic and Reverse Methanogenesis. *Front. Microbiol.* **2020**, *11*, 1806. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Sorokin, D.Y.; Makarova, K.S.; Abbas, B.; Ferrer, M.; Golyshin, P.N.; Galinski, E.A.; Ciordia, S.; Mena, M.C.; Merkel, A.Y.; Wolf, Y.I.; et al. Discovery of Extremely Halophilic, Methyl-Reducing Euryarchaea Provides Insights into the Evolutionary Origin of Methanogenesis. *Nat. Microbiol.* **2017**, *2*, 17081. [\[CrossRef\]](#)
17. Protasov, E.; Nonoh, J.O.; Kastle Silva, J.M.; Mies, U.S.; Hervé, V.; Dietrich, C.; Lang, K.; Mikulski, L.; Platt, K.; Poehlein, A.; et al. Diversity and Taxonomic Revision of Methanogens and Other Archaea in the Intestinal Tract of Terrestrial Arthropods. *Front. Microbiol.* **2023**, *14*, 1281628. [\[CrossRef\]](#)
18. Wen, X.; Yang, S.; Horn, F.; Winkel, M.; Wagner, D.; Liebner, S. Global Biogeographic Analysis of Methanogenic Archaea Identifies Community-Shaping Environmental Factors of Natural Environments. *Front. Microbiol.* **2017**, *8*, 1339. [\[CrossRef\]](#)
19. Merkel, A.Y.; Podosokorskaya, O.A.; Sokolova, T.G.; Bonch-Osmolovskaya, E.A. Diversity of Methanogenic Archaea from the 2012 Terrestrial Hot Spring (Valley of Geysers, Kamchatka). *Microbiology* **2016**, *85*, 342–349. [\[CrossRef\]](#)
20. Wang, J.; Qu, Y.N.; Evans, P.N.; Guo, Q.; Zhou, F.; Nie, M.; Jin, Q.; Zhang, Y.; Zhai, X.; Zhou, M.; et al. Evidence for Nontraditional Mcr-Containing Archaea Contributing to Biological Methanogenesis in Geothermal Springs. *Sci. Adv.* **2023**, *9*, eadg6004. [\[CrossRef\]](#)
21. Schulz, S.; Conrad, R. Influence of Temperature on Pathways to Methane Production in the Permanently Cold Profundal Sediment of Lake Constance. *FEMS Microbiol. Ecol.* **1996**, *20*, 1–14. [\[CrossRef\]](#)
22. Simankova, M.V.; Kotsyurbenko, O.R.; Lueders, T.; Nozhevnikova, A.N.; Wagner, B.; Conrad, R.; Friedrich, M.W. Isolation and Characterization of New Strains of Methanogens from Cold Terrestrial Habitats. *Syst. Appl. Microbiol.* **2003**, *26*, 312–318. [\[CrossRef\]](#) [\[PubMed\]](#)

23. Inagaki, F.; Hinrichs, K.U.; Kubo, Y.; Bowles, M.W.; Heuer, V.B.; Hong, W.L.; Hoshino, T.; Ijiri, A.; Imachi, H.; Ito, M.; et al. DEEP BIOSPHERE: Exploring Deep Microbial Life in Coal-Bearing Sediment down to ~2.5 Km below the Ocean Floor. *Science* **2015**, *349*, 420–424. [[CrossRef](#)] [[PubMed](#)]
24. Ye, J.; Zhuang, M.; Hong, M.; Zhang, D.; Ren, G.; Hu, A.; Yang, C.; He, Z.; Zhou, S. Methanogenesis in the Presence of Oxygenic Photosynthetic Bacteria May Contribute to Global Methane Cycle. *Nat. Commun.* **2024**, *15*, 5682. [[CrossRef](#)]
25. Buckley, D.H.; Baumgartner, L.K.; Visscher, P.T. Vertical Distribution of Methane Metabolism in Microbial Mats of the Great Sippewissett Salt Marsh. *Environ. Microbiol.* **2008**, *10*, 967–977. [[CrossRef](#)] [[PubMed](#)]
26. Bogard, M.J.; Del Giorgio, P.A.; Boutet, L.; Chaves, M.C.G.; Prairie, Y.T.; Merante, A.; Derry, A.M. Oxic Water Column Methanogenesis as a Major Component of Aquatic CH₄ Fluxes. *Nat. Commun.* **2014**, *5*, 5350. [[CrossRef](#)]
27. Patel, L.; Singh, R.; Thottathil, S.D. Contribution of Photosynthesis-Driven Oxic Methane Production to the Methane Cycling of a Tropical River Network. *ACS EST Water* **2024**, *4*, 2836–2847. [[CrossRef](#)]
28. Cynar, F.J.; Yayanos, A.A. Enrichment and Characterization of a Methanogenic Bacterium from the Oxic Upper Layer of the Ocean. *Curr. Microbiol.* **1991**, *23*, 89–96. [[CrossRef](#)]
29. Zhang, C.J.; Pan, J.; Liu, Y.; Duan, C.H.; Li, M. Genomic and Transcriptomic Insights into Methanogenesis Potential of Novel Methanogens from Mangrove Sediments. *Microbiome* **2020**, *8*, 94. [[CrossRef](#)]
30. Angle, J.C.; Morin, T.H.; Solden, L.M.; Narrow, A.B.; Sith, G.J.; Borton, M.A.; Rey-Sanchez, C.; Daly, R.A.; Mirfenderesgi, G.; Hoyt, D.W.; et al. Methanogenesis in Oxygenated Soils Is a Substantial Fraction of Wetland Methane Emissions. *Nat. Commun.* **2017**, *8*, 1567. [[CrossRef](#)]
31. Kendrick, M.G.; Kral, T.A. Survival of Methanogens during Desiccation: Implications for Life on Mars. *Astrobiology* **2006**, *6*, 546–551. [[CrossRef](#)]
32. Kral, T.A.; Travis Aitheide, S. Methanogen Survival Following Exposure to Desiccation, Low Pressure and Martian Regolith Analogs. *Planet. Space Sci.* **2013**, *89*, 167–171. [[CrossRef](#)]
33. Moissl-Eichinger, C. Archaea in Artificial Environments: Their Presence in Global Spacecraft Clean Rooms and Impact on Planetary Protection. *ISME J.* **2011**, *5*, 209–219. [[CrossRef](#)] [[PubMed](#)]
34. Mondav, R.; Woodcroft, B.J.; Kim, E.H.; McCalley, C.K.; Hodgkins, S.B.; Crill, P.M.; Chanton, J.; Hurst, G.B.; Verberkmoes, N.C.; Saleska, S.R.; et al. Discovery of a Novel Methanogen Prevalent in Thawing Permafrost. *Nat. Commun.* **2014**, *5*, 3212. [[CrossRef](#)] [[PubMed](#)]
35. Takai, K.; Nakamura, K.; Toki, T.; Tsunogai, U.; Miyazaki, M.; Miyazaki, J.; Hirayama, H.; Nakagawa, S.; Nunoura, T.; Horikoshi, K. Cell Proliferation at 122 °C and Isotopically Heavy CH₄ Production by a Hyperthermophilic Methanogen under High-Pressure Cultivation. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 10949–10954. [[CrossRef](#)]
36. Oremland, R.S.; Marsh, L.; Des Marais, D.J. Methanogenesis in Big Soda Lake, Nevada: An Alkaline, Moderately Hypersaline Desert Lake. *Appl. Environ. Microbiol.* **1982**, *43*, 462. [[CrossRef](#)]
37. Mickol, R.L.; Kral, T.A. Low Pressure Tolerance by Methanogens in an Aqueous Environment: Implications for Subsurface Life on Mars. *Orig. Life Evol. Biosph.* **2017**, *47*, 511–532. [[CrossRef](#)]
38. Kral, T.A.; Altheide, T.S.; Lueders, A.E.; Schuerger, A.C. Low Pressure and Desiccation Effects on Methanogens: Implications for Life on Mars. *Planet. Space Sci.* **2011**, *59*, 264–270. [[CrossRef](#)]
39. Bernhardt, G.; Jaenicke, R.; Ludemann, H.; Konig, H.; Stetter, K.O. High Pressure Enhances the Growth Rate of the Thermophilic Archaeobacterium *Methanococcus thermolithotrophicus* without Extending Its Temperature Range. *Appl. Environ. Microbiol.* **1988**, *54*, 1258–1261. [[CrossRef](#)]
40. Sinha, N.; Kral, T.A. Effect of UVC Radiation on Hydrated and Desiccated Cultures of Slightly Halophilic and Non-Halophilic Methanogenic Archaea: Implications for Life on Mars. *Microorganisms* **2018**, *6*, 43. [[CrossRef](#)]
41. Harris, R.L.; Schuerger, A.C. Hydrogenotrophic Methanogenesis at 7–12 Mbar by *Methanosarcina barkeri* under Simulated Martian Atmospheric Conditions. *Sci. Rep.* **2025**, *15*, 2880. [[CrossRef](#)]
42. Wong, H.L.; Visscher, P.T.; White, R.A., III; Smith, D.L.; Patterson, M.M.; Burns, B.P. Dynamics of Archaea at Fine Spatial Scales in Shark Bay Mat Microbiomes. *Sci. Rep.* **2017**, *7*, 46160. [[CrossRef](#)]
43. Visscher, P.T.; Baumgartner, L.K.; Buckley, D.H.; Rogers, D.R.; Hogan, M.E.; Raleigh, C.D.; Turk, K.A.; Des Marais, D.J. Dimethyl Sulphide and Methanethiol Formation in Microbial Mats: Potential Pathways for Biogenic Signatures. *Environ. Microbiol.* **2003**, *5*, 296–308. [[CrossRef](#)]
44. Visscher, P.T.; Quist, P.; Van Gernerden, H. Methylated Sulfur Compounds in Microbial Mats: In Situ Concentrations and Metabolism by a Colorless Sulfur Bacterium. *Appl. Environ. Microbiol.* **1991**, *57*, 1758–1763. [[CrossRef](#)] [[PubMed](#)]
45. Timmers, P.H.A.; Welte, C.U.; Koehorst, J.J.; Plugge, C.M.; Jetten, M.S.M.; Stams, A.J.M. Reverse Methanogenesis and Respiration in Methanotrophic Archaea. *Archaea* **2017**, *2017*, 1654237. [[CrossRef](#)] [[PubMed](#)]
46. Boetius, A.; Ravensschlag, K.; Schubert, C.J.; Rickert, D.; Widdel, F.; Gieseke, A.; Amann, R.; Jürgensen, B.B.; Witte, U.; Pfannkuche, O. A Marine Microbial Consortium Apparently Mediating Anaerobic Oxidation of Methane. *Nature* **2000**, *407*, 623–626. [[CrossRef](#)] [[PubMed](#)]

47. Yan, M.; Fotidis, I.A.; Jégliot, A.; Treu, L.; Tian, H.; Palomo, A.; Zhu, X.; Angelidaki, I. Long-Term Preserved and Rapidly Revived Methanogenic Cultures: Microbial Dynamics and Preservation Mechanisms. *J. Clean. Prod.* **2020**, *263*, 121577. [\[CrossRef\]](#)
48. Merino, N.; Aronson, H.S.; Bojanova, D.P.; Feyhl-Buska, J.; Wong, M.L.; Zhang, S.; Giovannelli, D. Living at the Extremes: Extremophiles and the Limits of Life in a Planetary Context. *Front. Microbiol.* **2019**, *10*, 780. [\[CrossRef\]](#)
49. Thomas, T.; Kumar, N.; Cavicchioli, R. Effects of Ribosomes and Intracellular Solutes on Activities and Stabilities of Elongation Factor 2 Proteins from Psychrotolerant and Thermophilic Methanogens. *J. Bacteriol.* **2001**, *183*, 1974–1982. [\[CrossRef\]](#)
50. Liu, Y.; Beer, L.L.; Whitman, W.B. Methanogens: A Window into Ancient Sulfur Metabolism. *Trends Microbiol.* **2012**, *20*, 251–258. [\[CrossRef\]](#)
51. Nozhevnikova, A.N.; Zepp, K.; Vazquez, F.; Zehnder, A.J.B.; Holliger, C. Evidence for the Existence of Psychrophilic Methanogenic Communities in Anoxic Sediments of Deep Lakes. *Appl. Environ. Microbiol.* **2003**, *69*, 1832–1835. [\[CrossRef\]](#)
52. Anderson, K.L.; Apolinario, E.E.; Sowers, K.R. Desiccation as a Long-Term Survival Mechanism for the Archaeon *Methanosarcina barkeri*. *Appl. Environ. Microbiol.* **2012**, *78*, 1473. [\[CrossRef\]](#)
53. Morozova, D.; Moeller, R.; Rettberg, P.; Wagner, D. Enhanced Radiation Resistance of *Methanosarcina soligelidi* SMA-21, a New Methanogenic Archaeon Isolated from a Siberian Permafrost-Affected Soil in Direct Comparison to *Methanosarcina barkeri*. *Astrobiology* **2015**, *15*, 951–960. [\[CrossRef\]](#)
54. Ebisuzaki, T.; Maruyama, S. Nuclear Geyser Model of the Origin of Life: Driving Force to Promote the Synthesis of Building Blocks of Life. *Geosci. Front.* **2016**, *8*, 275–298. [\[CrossRef\]](#)
55. Olson, K.D.; McMahon, C.W.; Wolfe, R.S. Light Sensitivity of Methanogenic Archaeobacteria. *Appl. Environ. Microbiol.* **1991**, *57*, 2683–2686. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Angel, R.; Matthies, D.; Conrad, R. Activation of Methanogenesis in Arid Biological Soil Crusts Despite the Presence of Oxygen. *PLoS ONE* **2011**, *6*, e20453. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Knauth, L.P. Temperature and Salinity History of the Precambrian Ocean: Implications for the Course of Microbial Evolution. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **2005**, *219*, 53–69. [\[CrossRef\]](#)
58. Martin, W.; Baross, J.; Kelley, D.; Russell, M.J. Hydrothermal Vents and the Origin of Life. *Nat. Rev. Microbiol.* **2008**, *6*, 805–814. [\[CrossRef\]](#)
59. Horikochi, K. Barophiles: Deep-Sea Microorganisms Adapted to an Extreme Environment. *Curr. Opin. Microbiol.* **1998**, *1*, 291–295. [\[CrossRef\]](#)
60. Van Dam, F.; Westmeijer, G.; Rezaei Somee, M.; Ketzer, M.; Kietäväinen, R.; Ono, S.; Bertilsson, S.; McIntosh, J.C.; Dopson, M.; Drake, H. Active methylotrophic methanogenesis by a microbial consortium enriched from a terrestrial meteorite impact crater. *Mbio* **2026**, *17*, e03017-25. [\[CrossRef\]](#)
61. Catling, D.C.; Zahnle, K.J. The Archean Atmosphere. *Sci. Adv.* **2020**, *6*, eaax1420. [\[CrossRef\]](#)
62. Neubeck, A.; Sjöberg, S.; Price, A.; Callac, N.; Schnürer, A. Effect of Nickel Levels on Hydrogen Partial Pressure and Methane Production in Methanogens. *PLoS ONE* **2016**, *11*, e0168357. [\[CrossRef\]](#)
63. Strobel, D.F. Photochemistry of the Reducing Atmospheres of Jupiter, Saturn and Titan. *Int. Rev. Phys. Chem.* **1983**, *3*, 145–176. [\[CrossRef\]](#)
64. Visscher, P.T.; Dupraz, C.; Braissant, O.; Gallagher, K.L.; Glunk, C.; Casillas, L.; Reed, R.E.S. *Biogeochemistry of Carbon Cycling in Hypersaline Mats: Linking the Present to the Past Through Biosignatures*; Springer: Dordrecht, The Netherlands, 2010; pp. 443–468. [\[CrossRef\]](#)
65. Lyu, Z.; Lu, Y. Metabolic Shift at the Class Level Sheds Light on Adaptation of Methanogens to Oxidative Environments. *ISME J.* **2018**, *12*, 411–423. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Lepot, K. Signatures of Early Microbial Life from the Archean (4 to 2.5 Ga) Eon. *Earth-Sci. Rev.* **2020**, *209*, 103296. [\[CrossRef\]](#)
67. Visscher, P.T.; Gallagher, K.L.; Bouton, A.; Farias, M.E.; Kurth, D.; Sancho-Tomás, M.; Philippot, P.; Somogyi, A.; Medjoubi, K.; Vennin, E.; et al. Modern Arsenotrophic Microbial Mats Provide an Analogue for Life in the Anoxic Archean. *Commun. Earth Environ.* **2020**, *1*, 24. [\[CrossRef\]](#)
68. Lira-Silva, E.; Santiago-Martínez, M.G.; Hernández-Juárez, V.; García-Contreras, R.; Moreno-Sánchez, R.; Jasso-Chávez, R. Activation of Methanogenesis by Cadmium in the Marine Archaeon *Methanosarcina acetivorans*. *PLoS ONE* **2012**, *7*, e48779. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Chen, C.; Li, L.; Wang, Y.; Dong, X.; Zhao, F.J. Methylotrophic Methanogens and Bacteria Synergistically Demethylate Dimethylarsenate in Paddy Soil and Alleviate Rice Straighthead Disease. *ISME J.* **2023**, *17*, 1851–1861. [\[CrossRef\]](#)
70. Anbar, A.D.; Knoll, A.H. Proterozoic Ocean Chemistry and Evolution: A Bioinorganic Bridge? *Science* **2002**, *297*, 1137–1142. [\[CrossRef\]](#)
71. Chen, C.; Li, L.; Huang, K.; Zhang, J.; Xie, W.Y.; Lu, Y.; Dong, X.; Zhao, F.J. Sulfate-Reducing Bacteria and Methanogens Are Involved in Arsenic Methylation and Demethylation in Paddy Soils. *ISME J.* **2019**, *13*, 2523–2535. [\[CrossRef\]](#)

72. Karr, E.A.; Sattley, W.M.; Rice, M.R.; Jung, D.O.; Madigan, M.T.; Achenbach, L.A. Diversity and Distribution of Sulfate-Reducing Bacteria in Permanently Frozen Lake Fryxell, McMurdo Dry Valleys, Antarctica. *Appl. Environ. Microbiol.* **2005**, *71*, 6353–6359. [\[CrossRef\]](#)
73. Zhang, G.; Jiang, N.; Liu, X.; Dong, X. Methanogenesis from Methanol at Low Temperatures by a Novel Psychrophilic Methanogen, “*Methanolobus sychrophilus*” sp. nov., Prevalent in Zoige Wetland of the Tibetan Plateau. *Appl. Environ. Microbiol.* **2008**, *74*, 6114–6120. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Zhao, Z.; Shen, B.; Zhu, J.M.; Lang, X.; Wu, G.; Tan, D.; Pei, H.; Huang, T.; Ning, M.; Ma, H. Active Methanogenesis during the Melting of Marinoan Snowball Earth. *Nat. Commun.* **2021**, *12*, 955. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Peng, Y.; Bao, H.; Jiang, G.; Crockford, P.; Feng, D.; Xiao, S.; Kaufman, A.J.; Wang, J. A Transient Peak in Marine Sulfate after the 635-Ma Snowball Earth. *Proc. Natl. Acad. Sci. USA* **2022**, *119*, e2117341119. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Song, H.; Kemp, D.B.; Tian, L.; Chu, D.; Song, H.; Dai, X. Thresholds of Temperature Change for Mass Extinctions. *Nat. Commun.* **2021**, *12*, 4694. [\[CrossRef\]](#) [\[PubMed\]](#)
77. McGunnigle, J.P.; Cano, E.J.; Sharp, Z.D.; Muehlenbachs, K.; Cole, D.; Hardman, M.F.; Stachel, T.; Pearson, D.G. Triple Oxygen Isotope Evidence for a Hot Archean Ocean. *Geology* **2022**, *50*, 991–9915. [\[CrossRef\]](#)
78. Noon, K.R.; Guymon, R.; Crain, P.F.; McCloskey, J.A.; Thomm, M.; Lim, J.; Cavicchioli, R. Influence of Temperature on tRNA Modification in Archaea: *Methanococcoides burtonii* (Optimum Growth Temperature [T_{opt}], 23 °C) and *Stetteria hydrogenophila* (T_{opt}, 95 °C). *J. Bacteriol.* **2003**, *185*, 5483–5490. [\[CrossRef\]](#)
79. Lim, J.; Thomas, T.; Cavicchioli, R. Low Temperature Regulated DEAD-Box RNA Helicase from the Antarctic Archaeon, *Methanococcoides burtonii*. *J. Mol. Biol.* **2000**, *297*, 553–567. [\[CrossRef\]](#)
80. Shimada, Y.; Fukuda, W.; Akada, Y.; Ishida, M.; Nakayama, J.; Imanaka, T.; Fujiwara, S. Property of Cold Inducible DEAD-Box RNA Helicase in Hyperthermophilic Archaea. *Biochem. Biophys. Res. Commun.* **2009**, *389*, 622–627. [\[CrossRef\]](#)
81. Saunders, N.F.W.; Thomas, T.; Curmi, P.M.G.; Mattick, J.S.; Kuczek, E.; Slade, R.; Davis, J.; Franzmann, P.D.; Boone, D.; Rusterholtz, K.; et al. Mechanisms of Thermal Adaptation Revealed from the Genomes of the Antarctic. *Genome Res.* **2003**, *13*, 1580–1588. [\[CrossRef\]](#)
82. Nichols, D.S.; Miller, M.R.; Davies, N.W.; Goodchild, A.; Raftery, M.; Cavicchioli, R. Cold Adaptation in the Antarctic Archaeon *Methanococcoides burtonii* Involves Membrane Lipid Unsaturation. *J. Bacteriol.* **2004**, *186*, 8508–8515. [\[CrossRef\]](#)
83. Koga, Y. Thermal Adaptation of the Archaeal and Bacterial Lipid Membranes. *Archaea* **2012**, *2012*, 789652. [\[CrossRef\]](#)
84. Cheptsov, V.S.; Vorobyova, E.A.; Osipov, G.A.; Manucharova, N.A.; Polyanskaya, L.M.; Gorlenko, M.V.; Pavlov, A.K.; Rosanova, M.S.; Lomasov, V.N. Microbial Activity in Martian Analog Soils after Ionizing Radiation: Implications for the Preservation of Subsurface Life on Mars. *AIMS Microbiol.* **2018**, *4*, 541–562. [\[CrossRef\]](#)
85. Cockell, C.S. Biological Effects of High Ultraviolet Radiation on Early Earth—A Theoretical Evaluation. *J. Theor. Biol.* **1998**, *193*, 717–729. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Zahnle, K.J.; Walker, J.C.G. The Evolution of Solar Ultraviolet Luminosity. *Rev. Geophys.* **1982**, *20*, 280–292. [\[CrossRef\]](#)
87. Ohlendorf, R.; Möglich, A. Light-Regulated Gene Expression in Bacteria: Fundamentals, Advances, and Perspectives. *Front. Bioeng. Biotechnol.* **2022**, *10*, 1029403. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Elahi, Y.; Baker, M.A.B. Light Control in Microbial Systems. *Int. J. Mol. Sci.* **2024**, *25*, 4001. [\[CrossRef\]](#)
89. Katayama, T.; Kamagata, Y. Cultivation of Methanogens. In *Hydrocarbon and Lipid Microbiology Protocols*; Springer: Berlin/Heidelberg, Germany, 2015; pp. 177–195. [\[CrossRef\]](#)
90. Paula, F.S.; Chin, J.P.; Schnürer, A.; Müller, B.; Manesiotis, P.; Waters, N.; Macintosh, K.A.; Quinn, J.P.; Connolly, J.; Abram, F.; et al. The Potential for Polyphosphate Metabolism in Archaea and Anaerobic Polyphosphate Formation in *Methanosarcina mazei*. *Sci. Rep.* **2019**, *9*, 17101. [\[CrossRef\]](#)
91. Mullaymeri, A.; Payr, M.; Wunderer, M.; Eva Maria, E.M.; Wagner, A.O. Shaken Not Stirred—Effect of Different Mixing Modes during the Cultivation of Methanogenic Pure Cultures. *Curr. Res. Microb. Sci.* **2025**, *8*, 100386. [\[CrossRef\]](#)
92. Jeffrey, L.C.; Maher, D.T.; Johnston, S.G.; Kelaher, B.P.; Steven, A.; Tait, D.R. Wetland Methane Emissions Dominated by Plant-Mediated Fluxes: Contrasting Emissions Pathways and Seasons within a Shallow Freshwater Subtropical Wetland. *Limnol. Oceanogr.* **2019**, *64*, 1895–1912. [\[CrossRef\]](#)
93. Zhang, M.; Xiao, Q.; Zhang, Z.; Gao, Y.; Zhao, J.; Pu, Y.; Wang, W.; Xiao, W.; Liu, S.; Lee, X. Methane Flux Dynamics in a Submerged Aquatic Vegetation Zone in a Subtropical Lake. *Sci. Total Environ.* **2019**, *672*, 400–409. [\[CrossRef\]](#)
94. Zhu, Y.; Purdy, K.J.; Eyice, Ö.; Shen, L.; Harpenslager, S.F.; Yvon-Durocher, G.; Dumbrell, A.J.; Trimmer, M. Disproportionate Increase in Freshwater Methane Emissions Induced by Experimental Warming. *Nat. Clim. Change* **2020**, *10*, 685–690. [\[CrossRef\]](#)
95. Bolhuis, H.; Cretoiu, M.S.; Stal, L.J. Molecular Ecology of Microbial Mats. *FEMS Microbiol. Ecol.* **2014**, *90*, 335–350. [\[CrossRef\]](#) [\[PubMed\]](#)
96. Grossart, H.P.; Frindte, K.; Dziallas, C.; Eckert, W.; Tang, K.W. Microbial Methane Production in Oxygenated Water Column of an Oligotrophic Lake. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 19657–19661. [\[CrossRef\]](#) [\[PubMed\]](#)

97. Wang, S.; Sun, P.; Liu, J.; Xu, Y.; Dolfing, J.; Wu, Y. Distribution of Methanogenic and Methanotrophic Consortia at Soil-Water Interfaces in Rice Paddies across Climate Zones. *iScience* **2023**, *26*, 105851. [\[CrossRef\]](#)
98. Huang, L.; Liu, X.; Rensing, C.; Yuan, Y.; Zhou, S.; Nealson, K.H. Light-Independent Anaerobic Microbial Oxidation of Manganese Driven by an Electrosynthetic Coculture. *ISME J.* **2023**, *17*, 163–171. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Kalathil, S.; Rahaman, M.; Lam, E.; Augustin, T.L.; Greer, H.F.; Reisner, E. Solar-Driven Methanogenesis through Microbial Ecosystem Engineering on Carbon Nitride. *Angew. Chem. Int. Ed.* **2024**, *63*, e202409192. [\[CrossRef\]](#)
100. Olson, S.; Jansen, M.F.; Abbot, D.S.; Halevy, I.; Goldblatt, C. The Effect of Ocean Salinity on Climate and Its Implications for Earth's Habitability. *Geophys. Res. Lett.* **2022**, *49*, e2021GL095748. [\[CrossRef\]](#)
101. Bardavid, R.E.; Oren, A. The Amino Acid Composition of Proteins from Anaerobic Halophilic Bacteria of the Order Halanaerobiales. *Extremophiles* **2012**, *16*, 567–572. [\[CrossRef\]](#)
102. Welsh, D.T. Ecological Significance of Compatible Solute Accumulation by Micro-Organisms: From Single Cells to Global Climate. *FEMS Microbiol. Rev.* **2000**, *24*, 263–290. [\[CrossRef\]](#)
103. Empadinhas, N.; da Costa, M.S. Osmoadaptation Mechanisms in Prokaryotes: Distribution of Compatible Solutes. *Int. Microbiol.* **2008**, *11*, 151–161. [\[CrossRef\]](#)
104. Stüeken, E.E.; Buick, R.; Guy, B.M.; Koehler, M.C. Isotopic Evidence for Biological Nitrogen Fixation by Molybdenum-Nitrogenase from 3.2 Gyr. *Nature* **2015**, *520*, 666–669. [\[CrossRef\]](#) [\[PubMed\]](#)
105. Hoehler, T.M.; Bebout, B.M.; Des Marais, D.J. The Role of Microbial Mats in the Production of Reduced Gases on the Early Earth. *Nature* **2001**, *412*, 324–327. [\[CrossRef\]](#) [\[PubMed\]](#)
106. Wang, M.; Ungerfeld, E.M.; Wang, R.; Zhou, C.S.; Basang, Z.Z.; Ao, S.M.; Tan, Z.L. Supersaturation of Dissolved Hydrogen and Methane in Rumen of Tibetan Sheep. *Front. Microbiol.* **2016**, *7*, 850. [\[CrossRef\]](#) [\[PubMed\]](#)
107. Catling, D.C. The Great Oxidation Event Transition. In *Treatise on Geochemistry: Second Edition*; Elsevier: Amsterdam, The Netherlands, 2014; Volume 6, pp. 177–195. [\[CrossRef\]](#)
108. Jasso-Chávez, R.; Santiago-Martínez, M.G.; Lira-Silva, E.; Pineda, E.; Zepeda-Rodríguez, A.; Belmont-Díaz, J.; Encalada, R.; Saavedra, E.; Moreno-Sánchez, R. Air-Adapted *Methanosarcina acetivorans* Shows High Methane Production and Develops Resistance against Oxygen Stress. *PLoS ONE* **2015**, *10*, e0117331. [\[CrossRef\]](#)
109. Ernst, L.; Steinfeld, B.; Barayeu, U.; Klintzsch, T.; Kurth, M.; Grimm, D.; Dick, T.P.; Rebele, J.G.; Bischofs, I.B.; Keppler, F. Methane Formation Driven by Reactive Oxygen Species across All Living Organisms. *Nature* **2022**, *603*, 482–487. [\[CrossRef\]](#)
110. Eliani-Russak, E.; Tik, Z.; Uzi-Gavrilov, S.; Meijler, M.M.; Sivan, O. The Reduction of Environmentally Abundant Iron Oxides by the Methanogen *Methanosarcina barkeri*. *Front. Microbiol.* **2023**, *14*, 1197299. [\[CrossRef\]](#)
111. Glass, J.B.; Orphan, V.J. Trace Metal Requirements for Microbial Enzymes Involved in the Production and Consumption of Methane and Nitrous Oxide. *Front. Microbiol.* **2012**, *3*, 61. [\[CrossRef\]](#)
112. Rossetto, D.; Mansy, S.S. Metals Are Integral to Life as We Know It. *Front. Cell Dev. Biol.* **2022**, *10*, 864830. [\[CrossRef\]](#)
113. Capdevila, D.A.; Edmonds, K.A.; Giedroc, D.P. Metallochaperones and Metalloregulation in Bacteria. *Essays Biochem.* **2017**, *61*, 177–200. [\[CrossRef\]](#)
114. Summons, R.E.; Welander, P.V.; Gold, D.A. Lipid Biomarkers: Molecular Tools for Illuminating the History of Microbial Life. *Nat. Rev. Microbiol.* **2022**, *20*, 174–185. [\[CrossRef\]](#)
115. Mulkidjanian, A.Y.; Bychkov, A.Y.; Dibrova, D.V.; Galperin, M.Y.; Koonin, E.V. Origin of First Cells at Terrestrial, Anoxic Geothermal Fields. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, E821–E830. [\[CrossRef\]](#) [\[PubMed\]](#)
116. Ünal, B.; Perry, V.R.; Sheth, M.; Gomez-Alvarez, V.; Chin, K.J.; Nüsslein, K. Trace Elements Affect Methanogenic Activity and Diversity in Enrichments from Subsurface Coal Bed Produced Water. *Front. Microbiol.* **2012**, *3*, 175. [\[CrossRef\]](#) [\[PubMed\]](#)
117. Abdel Azim, A.; Rittmann, S.K.M.R.; Fino, D.; Bochmann, G. The Physiological Effect of Heavy Metals and Volatile Fatty Acids on *Methanococcus maripaludis* S2. *Biotechnol. Biofuels* **2018**, *11*, 301. [\[CrossRef\]](#) [\[PubMed\]](#)
118. Spietz, R.L.; Payne, D.; Boyd, E.S. Methanogens Acquire and Bioaccumulate Nickel during Reductive Dissolution of Nickelian Pyrite. *Appl. Environ. Microbiol.* **2023**, *89*, e00991-23. [\[CrossRef\]](#)
119. Glass, J.B.; Chen, S.; Dawson, K.S.; Horton, D.R.; Vogt, S.; Ingall, E.D.; Twining, B.S.; Orphan, V.J. Trace Metal Imaging of Sulfate-Reducing Bacteria and Methanogenic Archaea at Single-Cell Resolution by Synchrotron X-Ray Fluorescence Imaging. *Geomicrobiol. J.* **2018**, *35*, 81–89. [\[CrossRef\]](#)
120. Holmes, D.E.; Tang, H.; Woodard, T.; Liang, D.; Zhou, J.; Liu, X.; Lovley, D.R. Cytochrome-Mediated Direct Electron Uptake from Metallic Iron by *Methanosarcina acetivorans*. *mLife* **2022**, *1*, 443–447. [\[CrossRef\]](#)
121. Wagner, T.; Ermler, U.; Shima, S. The Methanogenic CO₂ Reducing-and-Fixing Enzyme Is Bifunctional and Contains 46 [4Fe-4S] Clusters. *Science* **2016**, *354*, 114–117. [\[CrossRef\]](#)
122. Demirel, B.; Scherer, P. Trace Element Requirements of Agricultural Biogas Digesters during Biological Conversion of Renewable Biomass to Methane. *Biomass Bioenergy* **2011**, *35*, 992–998. [\[CrossRef\]](#)
123. Rother, M.; Krzycki, J.A. Selenocysteine, Pyrrolysine, and the Unique Energy Metabolism of Methanogenic Archaea. *Archaea* **2010**, *2010*, 453642. [\[CrossRef\]](#)

124. Johnson, C.; England, A.; Munro-Ehrlich, M.; Colman, D.R.; DuBois, J.L.; Boyd, E.S. Pathways of Iron and Sulfur Acquisition, Cofactor Assembly, Destination, and Storage in Diverse Archaeal Methanogens and Alkanotrophs. *J. Bacteriol.* **2021**, *203*, 117–138. [\[CrossRef\]](#)
125. Bini, E. Archaeal Transformation of Metals in the Environment. *FEMS Microbiol. Ecol.* **2010**, *73*, 1–16. [\[CrossRef\]](#) [\[PubMed\]](#)
126. Konhauser, K.O.; Pecoits, E.; Lalonde, S.V.; Papineau, D.; Nisbet, E.G.; Barley, M.E.; Arndt, N.T.; Zahnle, K.; Kamber, B.S. Oceanic Nickel Depletion and a Methanogen Famine before the Great Oxidation Event. *Nature* **2009**, *458*, 750–753. [\[CrossRef\]](#) [\[PubMed\]](#)
127. Gottschalk, G.; Thauer, R.K. The Na⁺-Translocating Methyltransferase Complex from Methanogenic Archaea. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2001**, *1505*, 28–36. [\[CrossRef\]](#)
128. Boone, D.R.; Mah, R.A. Effects of Calcium, Magnesium, PH, and Extent of Growth on the Morphology of Methanosarcina Mazei S-6. *Appl. Environ. Microbiol.* **1987**, *53*, 1699–1700. [\[CrossRef\]](#)
129. Vanček, M.; Vidová, M.; Majerník, A.I.; Šmigář, P. Methanogenesis Is Ca²⁺ Dependent in *Methanothermobacter thermautotrophicus* Strain DeltaH. *FEMS Microbiol. Lett.* **2006**, *258*, 269–273. [\[CrossRef\]](#)
130. Zhang, G.; Zhang, F.; Ding, G.; Li, J.; Guo, X.; Zhu, J.; Zhou, L.; Cai, S.; Liu, X.; Luo, Y.; et al. Acyl Homoserine Lactone-Based Quorum Sensing in a Methanogenic Archaeon. *ISME J.* **2012**, *6*, 1336–1344. [\[CrossRef\]](#)
131. Yepez-Ceron, O.D.; McCarthy, P.; Patterson, S.; Casey, E.; Dereli, R.K. The Role of Calcium in Anaerobic Treatment: The Good, the Bad and the Ugly. *Water Res. X* **2025**, *28*, 100381. [\[CrossRef\]](#)
132. Thauer, R.K.; Kaster, A.K.; Goenrich, M.; Schick, M.; Hiromoto, T.; Shima, S. Hydrogenases from Methanogenic Archaea, Nickel, a Novel Cofactor, and H₂ Storage. *Annu. Rev. Biochem.* **2010**, *79*, 507–536. [\[CrossRef\]](#)
133. Leimkühler, S. Metal-Containing Formate Dehydrogenases, a Personal View. *Molecules* **2023**, *28*, 5338. [\[CrossRef\]](#)
134. Mota, C.S.; Rivas, M.G.; Brondino, C.D.; Moura, I.; Moura, J.J.G.; González, P.J.; Cerqueira, N.M.F.S.A. The Mechanism of Formate Oxidation by Metal-Dependent Formate Dehydrogenases. *J. Biol. Inorg. Chem.* **2011**, *16*, 1255–1268. [\[CrossRef\]](#)
135. Payne, D.; Keller, L.M.; Larson, J.; Bothner, B.; Colman, D.R.; Boyd, E.S. Alternative Sources of Molybdenum for Methanococcus Maripaludis and Their Implication for the Evolution of Molybdoenzymes. *Commun. Biol.* **2024**, *7*, 1337. [\[CrossRef\]](#) [\[PubMed\]](#)
136. Tchounwou, P.B.; Yedjou, C.G.; Patlolla, A.K.; Sutton, D.J. Heavy Metals Toxicity and the Environment. In *Molecular, Clinical and Environmental Toxicology: Volume 3: Environmental Toxicology*; Springer: Berlin/Heidelberg, Germany, 2012; Volume 101, p. 133. [\[CrossRef\]](#)
137. Marcovecchio, J.E.; Botté, S.E. Heavy Metals, Major Metals, Trace Elements. In *Handbook of Water Analysis*, 2nd ed.; Nollet, L.M.L., Ed.; CRC Press: Boca Ratón, FL, USA; Taylor & Francis Group LCC: Boca Ratón, FL, USA, 2007; pp. 289–326.
138. Slyemi, D.; Bonnefoy, V. How Prokaryotes Deal with Arsenic. *Environ. Microbiol. Rep.* **2011**, *4*, 571–586. [\[CrossRef\]](#) [\[PubMed\]](#)
139. Van Lis, R.; Nitschke, W.; Duval, S.; Schoepp-Cothenet, B. Arsénics as Bioenergetic Substrates. *Biochim. Biophys. Acta Bioenerg.* **2013**, *1827*, 176–188. [\[CrossRef\]](#) [\[PubMed\]](#)
140. Ben Fekih, I.; Zhang, C.; Li, Y.P.; Zhao, Y.; Alwathnani, H.A.; Saquib, Q.; Rensing, C.; Cervantes, C. Distribution of Arsenic Resistance Genes in Prokaryotes. *Front. Microbiol.* **2018**, *9*, 2473. [\[CrossRef\]](#)
141. Fru, E.C.; Somogyi, A.; El Albani, A.; Medjoubi, K.; Aubineau, J.; Robbins, L.J.; Lalonde, S.V.; Konhauser, K.O. The Rise of Oxygen-Driven Arsenic Cycling at ca. 2.48 Ga. *Geology* **2019**, *47*, 243–246. [\[CrossRef\]](#)
142. Sierra-Alvarez, R.; Cortinas, I.; Yenal, U.; Field, J.A. Methanogenic Inhibition by Arsenic Compounds. *Appl. Environ. Microbiol.* **2004**, *70*, 5688. [\[CrossRef\]](#)
143. Rodriguez-Freire, L.; Moore, S.E.; Sierra-Alvarez, R.; Field, J.A. Adaptation of a Methanogenic Consortium to Arsenite Inhibition. *Water Air Soil. Pollut.* **2015**, *226*, 414. [\[CrossRef\]](#)
144. Liang, Y.; Yan, Y.; Shi, L.; Wang, M.; Yuan, X.; Wang, S.; Ye, L.; Yan, Z. Molecular Basis of Thioredoxin-Dependent Arsenic Transformation in Methanogenic Archaea. *Environ. Sci. Technol.* **2025**, *59*, 443–453. [\[CrossRef\]](#)
145. Jasso-Chavez, R.; Lira-Silva, E.; González-Sánchez, K.; Larios-Serrato, V.; Mendoza-Monzoy, D.L.; Pérez-Villatoro, F.; Morett, E.; Vega-Segura, A.; Torres-Márquez, M.E.; Zepeda-Rodríguez, A.; et al. Marine Archaeon *Methanosarcina acetivorans* Enhances Polyphosphate Metabolism Under Persistent Cadmium Stress. *Front. Microbiol.* **2019**, *10*, 2432. [\[CrossRef\]](#)
146. Lira-Silva, E.; Santiago-Martínez, M.G.; García-Contreras, R.; Zepeda-Rodríguez, A.; Marín-Hernández, A.; Moreno-Sánchez, R.; Jasso-Chávez, R. Cd²⁺ Resistance Mechanisms in *Methanosarcina acetivorans* Involve the Increase in the Coenzyme M Content and Induction of Biofilm Synthesis. *Environ. Microbiol. Rep.* **2013**, *5*, 799–808. [\[CrossRef\]](#)
147. Singh, R.; Dong, H.; Liu, D.; Zhao, L.; Marts, A.R.; Farquhar, E.; Tierney, D.L.; Almquist, C.B.; Briggs, B.R. Reduction of Hexavalent Chromium by the Thermophilic Methanogen *Methanothermobacter thermautotrophicus*. *Geochim. Cosmochim. Acta* **2014**, *148*, 442. [\[CrossRef\]](#) [\[PubMed\]](#)
148. Ueno, Y.; Yamada, K.; Yoshida, N.; Maruyama, S.; Isozaki, Y. Evidence from Fluid Inclusions for Microbial Methanogenesis in the Early Archaeal Era. *Nature* **2006**, *440*, 516–519. [\[CrossRef\]](#) [\[PubMed\]](#)
149. Wolfe, J.M.; Fournier, G.P. Horizontal Gene Transfer Constrains the Timing of Methanogen Evolution. *Nat. Ecol. Evol.* **2018**, *2*, 897–903. [\[CrossRef\]](#) [\[PubMed\]](#)
150. Lowe, D.R.; Byerly, G.R. The Terrestrial Record of Late Heavy Bombardment. *New Astron. Rev.* **2018**, *81*, 39–61. [\[CrossRef\]](#)

151. Mißbach, H.; Duda, J.P.; van den Kerkhof, A.M.; Lüders, V.; Pack, A.; Reitner, J.; Thiel, V. Ingredients for Microbial Life Preserved in 3.5 Billion-Year-Old Fluid Inclusions. *Nat. Commun.* **2021**, *12*, 1101. [\[CrossRef\]](#)
152. Hinrichs, K.U. Microbial Fixation of Methane Carbon at 2.7 Ga: Was an Anaerobic Mechanism Possible? *Geochim. Geophys. Geosyst.* **2002**, *3*, 1–10. [\[CrossRef\]](#)
153. Brocks, J.J.; Logan, G.A.; Buick, R.; Summons, R.E. Archean Molecular Fossils and the Early Rise of Eukaryotes. *Science* **1999**, *285*, 1033–1036. [\[CrossRef\]](#)
154. Baumgartner, R.J.; Van Kranendonk, M.J.; Caruso, S.; Campbell, K.A.; Dobson, M.J.; Teece, B.L.; Verrall, M.; Homann, M.; Lalonde, S.; Visscher, P.T. Pyritic Stromatolites from the Paleoarchean Dresser Formation, Pilbara Craton: Resolving Biogenicity and Hydrothermally Influenced Ecosystem Dynamics. *Geobiology* **2024**, *22*, e12610. [\[CrossRef\]](#)
155. Wang, Y.; Fu, X.; Wang, J.; Wei, H.; Zeng, Y.; Ahmed, M.S.; Mansour, A. Methane Emissions during Marine Environmental Perturbations before the End-Triassic Mass Extinction. *Facies* **2025**, *71*, 11. [\[CrossRef\]](#)
156. Petryshyn, V.A.; Greene, S.E.; Farnsworth, A.; Lunt, D.J.; Kelley, A.; Gammariello, R.; Ibarra, Y.; Bottjer, D.J.; Tripathi, A.; Corsetti, F.A. The Role of Temperature in the Initiation of the End-Triassic Mass Extinction. *Earth. Sci. Rev.* **2020**, *208*, 103266. [\[CrossRef\]](#)
157. Brand, U.; Blamey, N.; Garbelli, C.; Griesshaber, E.; Posenato, R.; Angiolini, L.; Azmy, K.; Farabegoli, E.; Came, R. Methane Hydrate: Killer Cause of Earth's Greatest Mass Extinction. *Palaeoworld* **2016**, *25*, 496–507. [\[CrossRef\]](#)
158. Fournier, G.P.; Gogarten, J.P. Evolution of Acetoclastic Methanogenesis in Methanosarcina via Horizontal Gene Transfer from Cellulolytic Clostridia. *J. Bacteriol.* **2008**, *190*, 1124–1127. [\[CrossRef\]](#) [\[PubMed\]](#)
159. Luo, H.; Sun, Z.; Arndt, W.; Shi, J.; Friedman, R.; Tang, J. Gene Order Phylogeny and the Evolution of Methanogens. *PLoS ONE* **2009**, *4*, e6069. [\[CrossRef\]](#) [\[PubMed\]](#)
160. Kohtz, A.J.; Petrosian, N.; Krukenberg, V.; Jay, Z.J.; Pilhofer, M.; Hatzenpichler, R. Cultivation and Visualization of a Methanogen of the Phylum Thermoproteota. *Nature* **2024**, *632*, 1118–1123. [\[CrossRef\]](#) [\[PubMed\]](#)
161. Cozannet, M.; Le Guellec, S.; Alain, K. A Variety of Substrates for Methanogenesis. *Case Stud. Chem. Environ. Eng.* **2023**, *8*, 100533. [\[CrossRef\]](#)
162. Hoehler, T.M.; Alperin, M.J.; Albert, D.B.; Martens, C.S. Apparent Minimum Free Energy Requirements for Methanogenic Archaea and Sulfate-Reducing Bacteria in an Anoxic Marine Sediment. *FEMS Microbiol. Ecol.* **2001**, *38*, 33–41. [\[CrossRef\]](#)
163. Vital, M.; Howe, A.C.; Tiedje, J.M. Revealing the Bacterial Butyrate Synthesis Pathways by Analyzing (Meta)Genomic Data. *mBio* **2014**, *5*, e00889-14. [\[CrossRef\]](#)
164. Hamilton, J.J.; Calixto Contreras, M.; Reed, J.L. Thermodynamics and H₂ Transfer in a Methanogenic, Syntrophic Community. *PLoS Comput. Biol.* **2015**, *11*, e1004364. [\[CrossRef\]](#)
165. Kato, S.; Hashimoto, K.; Watanabe, K. Methanogenesis Facilitated by Electric Syntrophy via (Semi)Conductive Iron-Oxide Minerals. *Environ. Microbiol.* **2012**, *14*, 1646–1654. [\[CrossRef\]](#)
166. Ishii, S.; Kosaka, T.; Hori, K.; Hotta, Y.; Watanabe, K. Coaggregation Facilitates Interspecies Hydrogen Transfer between *Pelotomaculum thermopropionicum* and *Methanothermobacter thermautotrophicus*. *Appl. Environ. Microbiol.* **2005**, *71*, 7838–7845. [\[CrossRef\]](#)
167. Meulepas, R.J.W.; Jagersma, C.G.; Khadem, A.F.; Stams, A.J.M.; Lens, P.N.L. Effect of Methanogenic Substrates on Anaerobic Oxidation of Methane and Sulfate Reduction by an Anaerobic Methanotrophic Enrichment. *Appl. Microbiol. Cell Physiol.* **2010**, *87*, 1499–1506. [\[CrossRef\]](#)
168. Stams, A.J.M.; Plugge, C.M. Electron Transfer in Syntrophic Communities of Anaerobic Bacteria and Archaea. *Nat. Rev. Microbiol.* **2009**, *7*, 568–577. [\[CrossRef\]](#) [\[PubMed\]](#)
169. Galagan, J.E.; Nusbaum, C.; Roy, A.; Endrizzi, M.G.; Macdonald, P.; Fitzhugh, W.; Calvo, S.; Engels, R.; Smirnov, S.; Atnoor, D.; et al. The Genome of *M. Acetivorans* Reveals Extensive Metabolic and Physiological Diversity. *Genome Res.* **2002**, *12*, 532–542. [\[CrossRef\]](#) [\[PubMed\]](#)
170. Oremland, R.S.; Marsh, L.M.; Polcin, S. Methane Production and Simultaneous Sulphate Reduction in Anoxic, Salt Marsh Sediments. *Nature* **1982**, *296*, 143–145. [\[CrossRef\]](#)
171. Cappenberg, T.E.; Prins, R.A. Interrelations between Sulfate-Reducing and Methane-Producing Bacteria in Bottom Deposits of a Fresh-Water Lake. III. Experiments with ¹⁴C-Labeled Substrates. *Antonie Van Leeuwenhoek* **1974**, *40*, 457–469. [\[CrossRef\]](#)
172. Ferry, J.G. Fundamentals of Methanogenic Pathways That Are Key to the Biomethanation of Complex Biomass. *Curr. Opin. Biotechnol.* **2011**, *22*, 351. [\[CrossRef\]](#)
173. Sorokin, D.Y.; Banciu, H.L.; Muyzer, G. Functional Microbiology of Soda Lakes. *Curr. Opin. Microbiol.* **2015**, *25*, 88–96. [\[CrossRef\]](#)
174. Wormald, R.M.; Hopwood, J.; Humphreys, P.N.; Mayes, W.; Gomes, H.I.; Rout, S.P. Methanogenesis from Mineral Carbonates, a Potential Indicator for Life on Mars. *Geosciences* **2022**, *12*, 138. [\[CrossRef\]](#)
175. Kral, T.A.; Birch, W.; Lavender, L.E.; Virden, B.T. Potential Use of Highly Insoluble Carbonates as Carbon Sources by Methanogens in the Subsurface of Mars. *Planet. Space Sci.* **2014**, *101*, 181–185. [\[CrossRef\]](#)
176. Fiore, N.A.; Kohtz, A.J.; Miller, D.N.; Antony-Babu, S.; Pan, D.; Lahey, C.; Huang, X.; Lu, Y.; Buan, N.R.; Weber, K.A. Microbial Methane Production from Calcium Carbonate at Moderately Alkaline pH. *Commun. Earth Environ.* **2025**, *6*, 85. [\[CrossRef\]](#)

177. Tian, F.; Toon, O.B.; Pavlov, A.A.; De Sterck, H. A Hydrogen-Rich Early Earth Atmosphere. *Science* **2005**, *308*, 1014–1017. [[CrossRef](#)] [[PubMed](#)]
178. Novelli, P.C.; Lang, P.M.; Masarie, K.A.; Hurst, D.F.; Myers, R.; Elkins, J.W. Molecular Hydrogen in the Troposphere: Global Distribution and Budget. *J. Geophys. Res. Atmos.* **1999**, *104*, 30427–30444. [[CrossRef](#)]
179. Mei, R.; Kaneko, M.; Imachi, H.; Nobu, M.K. The Origin and Evolution of Methanogenesis and Archaea Are Intertwined. *PNAS Nexus* **2023**, *2*, pgad023. [[CrossRef](#)]
180. Weiss, M.C.; Soussa, F.L.; Mrnjavac, N.; Neukirchen, S.; Roettger, M.; Nelson-Sathi, S.; Martin, W.F. The Physiology and Habitat of the Last Universal Common Ancestor. *Nat. Microbiol.* **2016**, *1*, 16116. [[CrossRef](#)] [[PubMed](#)]
181. Hallam, S.J.; Girguis, P.R.; Preston, C.M.; Richardson, P.M.; DeLong, E.F. Identification of Methyl Coenzyme M Reductase A (McrA) Genes Associated with Methane-Oxidizing Archaea. *Appl. Environ. Microbiol.* **2003**, *69*, 5483–5491. [[CrossRef](#)]
182. Garcia, P.S.; Gribaldo, S.; Borrel, G. Diversity and Evolution of Methane-Related Pathways in Archaea. *Annu. Rev. Microbiol.* **2022**, *76*, 727–755. [[CrossRef](#)]
183. Chadwick, G.L.; Dury, G.A.; Nayak, D.D. Physiological and Transcriptomic Response to Methyl-Coenzyme M Reductase Limitation in *Methanosarcina acetivorans*. *Appl. Environ. Microbiol.* **2024**, *90*, e02220-23. [[CrossRef](#)]
184. Knittel, K.; Boetius, A. Anaerobic Oxidation of Methane: Progress with an Unknown Process. *Annu. Rev. Microbiol.* **2009**, *63*, 311–334. [[CrossRef](#)]
185. Weber, H.S.; Habicht, K.S.; Thamdrup, B. Anaerobic Methanotrophic Archaea of the ANME-2d Cluster Are Active in a Low-Sulfate, Iron-Rich Freshwater Sediment. *Front. Microbiol.* **2017**, *8*, 619. [[CrossRef](#)]
186. Wegener, G.; Laso-Pérez, R.; Orphan, V.J.; Boetius, A. Anaerobic Degradation of Alkanes by Marine Archaea. *Annu. Rev. Microbiol.* **2022**, *76*, 553–577. [[CrossRef](#)]
187. McGlynn, S.E. Minireview of Energy Metabolism during Anaerobic Methane Oxidation in ANME Archaea. *Microbes Environ.* **2017**, *32*, 5–13. [[CrossRef](#)]
188. Coon, C.R.; Williams, L.C.; Matthews, A.; Diaz, R.; Kevorkian, R.T.; LaRowe, D.E.; Steen, A.D.; Lapham, L.L.; Lloyd, K.G. Control of Hydrogen Concentrations by Microbial Sulfate Reduction in Two Contrasting Anoxic Coastal Sediments. *Front. Microbiol.* **2024**, *15*, 1455857. [[CrossRef](#)] [[PubMed](#)]
189. Chadwick, G.L.; Skennerton, C.T.; Laso-Pérez, R.; Leu, A.O.; Speth, D.R.; Yu, H.; Morgan-Lang, C.; Hatzenpichler, R.; Goudeau, D.; Malmstrom, R.; et al. Comparative Genomics Reveals Electron Transfer and Syntrophic Mechanisms Differentiating Methanotrophic and Methanogenic Archaea. *PLoS Biol.* **2022**, *20*, e3001508. [[CrossRef](#)] [[PubMed](#)]
190. Ruff, S.E.; Biddle, J.F.; Teske, A.P.; Knittel, K.; Boetius, A.; Ramette, A. Global Dispersion and Local Diversification of the Methane Seep Microbiome. *Proc. Nat. Acad. Sci. USA* **2015**, *112*, 4015–4020. [[CrossRef](#)] [[PubMed](#)]
191. Borrel, G.; Adam, P.S.; McKay, L.J.; Chen, L.X.; Sierra-García, I.N.; Sieber, C.M.K.; Letourneur, Q.; Ghoulane, A.; Andersen, G.L.; Li, W.J.; et al. Wide Diversity of Methane and Short-Chain Alkane Metabolisms in Uncultured Archaea. *Nat. Microbiol.* **2019**, *4*, 603–613. [[CrossRef](#)]
192. Evans, P.N.; Boyd, J.A.; Leu, A.O.; Woodcroft, B.J.; Parks, D.H.; Hugenholtz, P.; Tyson, G.W. An Evolving View of Methane Metabolism in the Archaea. *Nat. Rev. Microbiol.* **2019**, *17*, 219–232. [[CrossRef](#)]
193. Wang, Y.; Xie, R.; Hou, J.; Lv, Z.; Li, L.; Hu, Y.; Huang, H.; Wang, F. The Late Archaeal to Early Proterozoic Origin and Evolution of Anaerobic Methane-Oxidizing Archaea. *mLife* **2022**, *1*, 96–100. [[CrossRef](#)]
194. Escudero, C.; Oggerin, M.; Amils, R. The Deep Continental Subsurface: The Dark Biosphere. *Int. Microbiol.* **2018**, *21*, 3–14. [[CrossRef](#)]
195. Smith, D.J.; Ravichandar, J.D.; Jain, S.; Griffin, D.W.; Yu, H.; Tan, Q.; Thissen, J.; Lusby, T.; Nicoll, P.; Shedler, S.; et al. Airborne Bacteria in Earth's Lower Stratosphere Resemble Taxa Detected in the Troposphere: Results from a New NASA Aircraft Bioaerosol Collector (ABC). *Front. Microbiol.* **2018**, *9*, 390777. [[CrossRef](#)]
196. Schopf, J.W.; Kitajima, K.; Spicuzza, M.J.; Kudryavtsev, A.B.; Valley, J.W. SIMS Analyses of the Oldest Known Assemblage of Microfossils Document Their Taxon-Related Carbon Isotope Compositions. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 53–58. [[CrossRef](#)]
197. Foster, J.S.; Mobberley, J.M. *Past, Present, and Future: Microbial Mats as Models for Astrobiological Research*; Springer: Dordrecht, The Netherlands, 2010; pp. 563–582. [[CrossRef](#)]
198. Xia, L.; Cao, J.; Hu, W.; Zhi, D.; Tang, Y.; Li, E.; He, W. Coupling of Paleoenvironment and Biogeochemistry of Deep-Time Alkaline Lakes: A Lipid Biomarker Perspective. *Earth. Sci. Rev.* **2021**, *213*, 103499. [[CrossRef](#)]
199. Zhao, Z.; Feng, Q.; Lu, H.; Peng, P.; Zhang, T.; Hsu, C.S. Stable Carbon Isotopic Fractionation and Hydrocarbon Generation Mechanism of CO₂ Fischer–Tropsch-Type Synthesis under Hydrothermal Conditions. *Energy Fuels* **2021**, *35*, 11909–11919. [[CrossRef](#)]
200. Rezaee, R. Quantifying Natural Hydrogen Generation Rates and Volumetric Potential in Onshore Serpentinization. *Geosciences* **2025**, *3*, 112. [[CrossRef](#)]
201. Kasting, J.F.; Siefert, J.L. Life and the Evolution of Earth's Atmosphere. *Science* **2002**, *296*, 1066–1068. [[CrossRef](#)]

202. Ding, Y.; Wang, M.; Du, X.; He, X.; Xu, T.; Liu, X.; Song, F. The Impact of Elevated CO₂ on Methanogen Abundance and Methane Emissions in Terrestrial Ecosystems: A Meta-Analysis. *iScience* **2024**, *27*, 111504. [\[CrossRef\]](#)
203. Olson, S.L.; Reinhard, C.T.; Lyons, T.W. Limited Role for Methane in the Mid-Proterozoic Greenhouse. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 11447–11452. [\[CrossRef\]](#)
204. Canfield, D.E.; Raiswell, R. The Evolution of the Sulfur Cycle. *Am. J. Sci.* **1999**, *299*, 697–723. [\[CrossRef\]](#)
205. Steward, K.F.; Payne, D.; Kincannon, W.; Johnson, C.; Lensing, M.; Fausset, H.; Németh, B.; Shepard, E.M.; Broderick, W.E.; Broderick, J.B.; et al. Proteomic Analysis of *Methanococcus voltae* Grown in the Presence of Mineral and Nonmineral Sources of Iron and Sulfur. *Microbiol. Spectr.* **2022**, *10*, e01893-22. [\[CrossRef\]](#)
206. Payne, D.; Spietz, R.L.; Boyd, E.S. Reductive Dissolution of Pyrite by Methanogenic Archaea. *ISME J.* **2021**, *15*, 3498–3507. [\[CrossRef\]](#)
207. Canfield, D.E. Biogeochemistry of Sulfur Isotopes. *Rev. Mineral. Geochem.* **2001**, *43*, 607–636. [\[CrossRef\]](#)
208. Fakhraee, M.; Crockford, P.W.; Bauer, K.W.; Pasquier, V.; Sugiyama, I.; Katsev, S.; Raven, M.R.; Gomes, M.; Philippot, P.; Crowe, S.A.; et al. The History of Earth's Sulfur Cycle. *Nat. Rev. Earth Environ.* **2025**, *6*, 106–125. [\[CrossRef\]](#)
209. Ebadiad, S.; Lyons, T.W.; Fournier, G.P. Archean Methane Cycling and Life's Co-Evolution: Intertwining Early Biogeochemical Processes and Ancient Microbial Metabolism. In *The Archean Earth*, 2nd ed.; Homann, M., Lyons, T.W., Ernst, R.E., Heubeck, C., Stueken, E.E., Webb, A.A.G., Papineau, D., Mason, P.R.D., Mazumder, R., Altermann, W., Eds.; Elsevier: Amsterdam, The Netherlands, 2026; pp. 445–469.
210. Fakhraee, M.; Katsev, S. Organic Sulfur Was Integral to the Archean Sulfur Cycle. *Nat. Commun.* **2019**, *10*, 4556–4558. [\[CrossRef\]](#) [\[PubMed\]](#)
211. Reeves, E.P.; McDermott, J.M.; Seewald, J.S. The Origin of Methanethiol in Midocean Ridge Hydrothermal Fluids. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 5474–5479. [\[CrossRef\]](#) [\[PubMed\]](#)
212. Xiao, X.; Wang, Y.; Huang, F.; Yang, Y.; Long, Y.; Fang, C.; Hu, L. Uncover Co-Evolution of Arsenic Transformation and Methanogens in Landfills. *J. Hazard. Mater.* **2025**, *497*, 139623. [\[CrossRef\]](#) [\[PubMed\]](#)
213. Fontecilla-Camps, J.C. Nickel and the Origin and Early Evolution of Life. *Metallomics* **2022**, *14*, 16. [\[CrossRef\]](#)
214. Dreher, C.L.; Schad, M.; Robbins, L.J.; Konhauser, K.O.; Kappler, A.; Joshi, P. Microbial processes during deposition and diagenesis of Banded Iron Formations. *PalZ* **2021**, *95*, 593–610. [\[CrossRef\]](#)
215. Konhauser, K.O.; Robbins, L.J.; Pecoits, E.; Peacock, C.; Kappler, A.; Lalonde, S.V. The Archean Nickel Famine Revisited. *Astrobiology* **2015**, *15*, 804–815. [\[CrossRef\]](#)
216. Moody, E.R.R.; Álvarez-Carretero, S.; Mahendrarajah, T.A.; Clark, J.W.; Betts, H.C.; Dombrowski, N.; Szánthó, L.L.; Boyle, R.A.; Daines, S.; Chen, X.; et al. The Nature of the Last Universal Common Ancestor and Its Impact on the Early Earth System. *Nat. Ecol. Evol.* **2024**, *8*, 1654–1666. [\[CrossRef\]](#)
217. Hren, M.T.; Tice, M.M.; Chamberlain, C.P. Oxygen and Hydrogen Isotope Evidence for a Temperate Climate 3.42 Billion Years Ago. *Nature* **2009**, *462*, 205–208. [\[CrossRef\]](#)
218. Blake, R.E.; Chang, S.J.; Lepland, A. Phosphate Oxygen Isotopic Evidence for a Temperate and Biologically Active Archaeal Ocean. *Nature* **2010**, *464*, 1029–1032. [\[CrossRef\]](#)
219. Penny, D.; Poole, A. The Nature of the Last Universal Common Ancestor. *Curr. Opin. Genet. Dev.* **1999**, *9*, 672–677. [\[CrossRef\]](#) [\[PubMed\]](#)
220. Liu, R.; Gong, H.; Xu, Y.; Cai, C.; Hua, Y.; Li, L.; Dai, L.; Dai, X. The Transition Temperature (42 °C) from Mesophilic to Thermophilic Micro-Organisms Enhances Biomethane Potential of Corn Stover. *Sci. Total Environ.* **2021**, *759*, 143549. [\[CrossRef\]](#) [\[PubMed\]](#)
221. Catling, D.C.; Zahnle, K.J.; McKay, C.P. Biogenic Methane, Hydrogen Escape, and the Irreversible Oxidation of Early Earth. *Science* **2001**, *293*, 839–843. [\[CrossRef\]](#) [\[PubMed\]](#)
222. Yang, S.; Lv, Y.; Liu, X.; Wang, Y.; Fan, Q.; Yanf, Z.; Boon, N.; Wang, F.; Xiao, X.; Zhang, Y. Genomic and Enzymatic Evidence of Acetogenesis by Anaerobic Methanotrophic Archaea. *Nat. Commun.* **2020**, *11*, 3941. [\[CrossRef\]](#)
223. Ettwig, K.F.; Zhu, B.; Speth, D.; Keltjens, J.T.; Jetten, M.S.M.; Kartal, B. Archaea Catalyze Iron-Dependent Anaerobic Oxidation of Methane. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 12792–12796. [\[CrossRef\]](#)
224. Pavlov, A.A.; Kasting, J.F.; Eigenbrode, J.L.; Freeman, K.H. Organic Haze in Earth's Early Atmosphere: Source of Low-¹³C Late Archean Kerogens? *Geology* **2001**, *29*, 1003–1006. [\[CrossRef\]](#)
225. Tian, F.; Kasting, J.F.; Zahnle, K. Revisiting HCN Formation in Earth's Early Atmosphere. *Earth Planet. Sci. Lett.* **2011**, *308*, 417–423. [\[CrossRef\]](#)
226. Wolf, E.T.; Toon, O.B. Fractal Organic Hazes Provided an Ultraviolet Shield for Early Earth. *Science* **2010**, *328*, 1266–1268. [\[CrossRef\]](#)
227. Walter, M.R.; Heys, G.R. Links between the Rise of the Metazoa and the Decline of Stromatolites. *Precambrian Res.* **1985**, *29*, 149–174. [\[CrossRef\]](#)

228. Dupraz, C.; Visscher, P.T. Microbial Lithification in Marine Stromatolites and Hypersaline Mats. *Trends Microbiol.* **2005**, *13*, 429–438. [[CrossRef](#)]
229. Colman, D.R.; Templeton, A.S.; Spear, J.R.; Boyd, E.S. Microbial Ecology of Serpentinite-Hosted Ecosystems. *ISME J.* **2025**, *19*, 29. [[CrossRef](#)] [[PubMed](#)]
230. Shibuya, T.; Komiya, T.; Nakamura, K.; Takai, K.; Maruyama, S. Highly Alkaline, High-Temperature Hydrothermal Fluids in the Early Archean Ocean. *Precambrian Res.* **2010**, *182*, 230–238. [[CrossRef](#)]
231. Preiner, M.; Xavier, J.C.; Sousa, F.L.; Zimorski, V.; Neubeck, A.; Lang, S.Q.; Chris Greenwell, H.; Kleinermaans, K.; Tüysüz, H.; McCollom, T.M.; et al. Serpentinization: Connecting Geochemistry, Ancient Metabolism and Industrial Hydrogenation. *Life* **2018**, *8*, 41. [[CrossRef](#)] [[PubMed](#)]
232. Li, J.; Kusky, T. World's largest known Precambrian fossil black smoker chimneys and associated microbial vent communities, North China: Implications for early life. *Gondwana Res.* **2001**, *12*, 84–100. [[CrossRef](#)]
233. Dodd, M.S.; Papineau, D.; Grenne, T.; Slack, J.F.; Rittner, M.; Pirajno, F.; O'Neil, J.; Little, C.T.S. Evidence for Early Life in Earth's Oldest Hydrothermal Vent Precipitates. *Nature* **2017**, *543*, 60–64. [[CrossRef](#)]
234. Aoyama, S.; Nishizawa, M.; Miyazaki, J.; Shibuya, T.; Ueno, Y.; Takai, K. Recycled Archean Sulfur in the Mantle Wedge of the Mariana Forearc and Microbial Sulfate Reduction within an Extremely Alkaline Serpentine Seamount. *Earth Planet. Sci. Lett.* **2018**, *491*, 109–120. [[CrossRef](#)]
235. Tosca, N.J.; Tutolo, B.M. Hydrothermal Vent Fluid-Seawater Mixing and the Origins of Archean Iron Formation. *Geochim. Cosmochim. Acta* **2023**, *352*, 51–68. [[CrossRef](#)]
236. Bjerrum, C.J.; Canfield, D.E. Ocean Productivity before about 1.9 Gyr Ago Limited by Phosphorus Adsorption onto Iron Oxides. *Nature* **2002**, *417*, 159–162. [[CrossRef](#)]
237. Martin, W.; Russell, M.J. On the Origin of Biochemistry at an Alkaline Hydrothermal Vent. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2007**, *362*, 1887–1926. [[CrossRef](#)]
238. Mrnjavac, N.; Schwander, L.; Brabender, M.; Martin, W.F. Chemical Antiquity in Metabolism. *Acc. Chem. Res.* **2024**, *57*, 2267–2278. [[CrossRef](#)]
239. Bae, H.S.; Morrison, E.; Chanton, J.P.; Ogram, A. Methanogens Are Major Contributors to Nitrogen Fixation in Soils of the Florida Everglades. *Appl. Environ. Microbiol.* **2018**, *84*, e02222-17. [[CrossRef](#)] [[PubMed](#)]
240. Evans, G.N.; Ji, S.; Kaçar, B.; Anbar, A.D.; Seyfried, W.E. Transition Metals in Alkaline Lost City Vent Fluids Are Sufficient for Early-Life Metabolisms. *Geochim. Cosmochim. Acta* **2024**, *385*, 61–73. [[CrossRef](#)]
241. Noffke, N.; Christian, D.; Wacey, D.; Hazen, R.M. Microbially Induced Sedimentary Structures Recording an Ancient Ecosystem in the ca. 3.48 Billion-Year-Old Dresser Formation, Pilbara, Western Australia. *Astrobiology* **2013**, *13*, 1103–1124. [[CrossRef](#)] [[PubMed](#)]
242. Gutierrez-Preciado, A.; Saghai, A.; Moreira, D.; Zivanovic, Y.; Dschamps, P.; Lopez-Garcia, P. Functional Shifts in Microbial Mats Recapitulate Early Earth Metabolic Transitions. *Nat. Ecol. Evol.* **2018**, *2*, 1700–1708. [[CrossRef](#)] [[PubMed](#)]
243. Nisbet, E.G.; Fowler, C.M.R. Archaeal Metabolic Evolution of Microbial Mats. *Proc. R. Soc. Lond. B Biol. Sci.* **1999**, *266*, 2375–2382. [[CrossRef](#)]
244. Blank, C.E.; Sánchez-Baracaldo, P. Timing of Morphological and Ecological Innovations in the Cyanobacteria—A Key to Understanding the Rise in Atmospheric Oxygen. *Geobiology* **2010**, *8*, 1–23. [[CrossRef](#)]
245. Knoll, A.H.; Nowak, M.A. The Timetable of Evolution. *Sci. Adv.* **2017**, *3*, e1603076. [[CrossRef](#)]
246. Hamilton, T.L.; Bryant, D.A.; Macalady, J.L. The Role of Biology in Planetary Evolution: Cyanobacterial Primary Production in Low-Oxygen Proterozoic Oceans. *Environ. Microbiol.* **2016**, *18*, 325–340. [[CrossRef](#)]
247. Noffke, N.; Gerdes, G.; Klenke, T.; Krumbein, W.E. Microbially Induced Sedimentary Structures: A New Category within the Classification of Primary Sedimentary Structures. *J. Sediment. Res.* **2001**, *71*, 649–656. [[CrossRef](#)]
248. Dupraz, C.; Reid, P.R.; Braissant, O.; Decho, A.W.; Normal, S.R.; Visscher, P.T. Processes of Carbonate Precipitation in Modern Microbial Mats. *Earth. Sci. Rev.* **2009**, *96*, 141–162. [[CrossRef](#)]
249. Allwood, A.C.; Walter, M.R.; Kamber, B.S.; Marshall, C.P.; Burch, I.W. Stromatolite Reef from the Early Archean Era of Australia. *Nature* **2006**, *441*, 714–718. [[CrossRef](#)]
250. Allwood, A.C.; Walter, M.R.; Burch, I.W.; Kamber, B.S. 3.43 Billion-Year-Old Stromatolite Reef from the Pilbara Craton of Western Australia: Ecosystem-Scale Insights to Early Life on Earth. *Precambrian Res.* **2007**, *158*, 198–227. [[CrossRef](#)]
251. Nutman, A.P.; Bennett, V.C.; Friend, C.R.L.; Van Kranendonk, M.J.; Chivas, A.R. Rapid Emergence of Life Shown by Discovery of 3,700-Million-Year-Old Microbial Structures. *Nature* **2016**, *537*, 535–538. [[CrossRef](#)]
252. Van Kranendonk, M.J.; Nutman, A.P.; Friend, C.R.L.; Bennett, V.C. A Review of 3.7 Ga Stromatolites from the Isua Supracrustal Belt, West Greenland. *Earth. Sci. Rev.* **2025**, *262*, 105034. [[CrossRef](#)]
253. Allwood, A.C.; Rosing, M.T.; Flannery, D.T.; Hurowitz, J.A.; Heirwegh, C.M. Reassessing Evidence of Life in 3,700-Million-Year-Old Rocks of Greenland. *Nature* **2018**, *563*, 241–244. [[CrossRef](#)] [[PubMed](#)]
254. Allwood, A.C.; Grotzinger, J.P.; Knoll, A.H.; Burch, I.W.; Anderson, M.S.; Coleman, M.L.; Kanik, I. Controls on Development and Diversity of Early Archean Stromatolites. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 9548–9555. [[CrossRef](#)] [[PubMed](#)]

255. Watanabe, Y.; Tajika, E.; Ozaki, K. Biogeochemical Transformations after the Emergence of Oxygenic Photosynthesis and Conditions for the First Rise of Atmospheric Oxygen. *Geobiology* **2023**, *21*, 537–555. [\[CrossRef\]](#)
256. Mukherjee, I.; Large, R.R.; Corkrey, R.; Danyushevsky, L.V. The Boring Billion, a Slingshot for Complex Life on Earth. *Sci. Rep.* **2018**, *8*, 4432. [\[CrossRef\]](#)
257. Buick, R.; Des Marais, D.J.; Knoll, A.H. Stable Isotopic Compositions of Carbonates from the Mesoproterozoic Bangemall Group, Northwestern Australia. *Chem. Geol.* **1995**, *123*, 153–171. [\[CrossRef\]](#)
258. Anbar, A.D. Oceans. Elements and Evolution. *Science* **2008**, *322*, 1481–1483. [\[CrossRef\]](#)
259. Caldwell, S.L.; Laidler, J.R.; Brewer, E.A.; Eberly, J.O.; Sandborgh, S.C.; Colwell, F.S. Anaerobic Oxidation of Methane: Mechanisms, Bioenergetics, and the Ecology of Associated Microorganisms. *Environ. Sci. Technol.* **2008**, *42*, 6791–6799. [\[CrossRef\]](#) [\[PubMed\]](#)
260. Beckmann, S.; Farag, I.F.; Zhao, R.; Christman, G.D.; Prouty, N.G.; Biddle, J.F. Expanding the Repertoire of Electron Acceptors for the Anaerobic Oxidation of Methane in Carbonates in the Atlantic and Pacific Ocean. *ISME J.* **2021**, *15*, 2523–2536. [\[CrossRef\]](#) [\[PubMed\]](#)
261. Stott, L.D.; Harazin, K.M.; Quintana Krupinski, N.B. Hydrothermal Carbon Release to the Ocean and Atmosphere from the Eastern Equatorial Pacific during the Last Glacial Termination. *Environ. Res. Lett.* **2019**, *14*, 025007. [\[CrossRef\]](#)
262. Buffett, B.A. Clathrate Hydrates. *Annu. Rev. Earth Planet. Sci.* **2000**, *28*, 477–507. [\[CrossRef\]](#)
263. Huard, D.J.E.; Johnson, A.M.; Fan, Z.; Kenney, L.G.; Xu, M.; Drori, R.; Gumbart, J.C.; Dai, S.; Lieberman, R.L.; Glass, J.B. Molecular Basis for Inhibition of Methane Clathrate Growth by a Deep Subsurface Bacterial Protein. *PNAS Nexus* **2023**, *2*, pgad268. [\[CrossRef\]](#)
264. Weitemeyer, K.A.; Buffett, B.A. Accumulation and Release of Methane from Clathrates below the Laurentide and Cordilleran Ice Sheets. *Glob. Planet. Change* **2006**, *53*, 176–187. [\[CrossRef\]](#)
265. Burgess, S.D.; Black, B.A. The Anatomy and Lethality of the Siberian Traps Large Igneous Province. *Annu. Rev. Earth Planet. Sci.* **2025**, *53*, 567–594. [\[CrossRef\]](#)
266. Rothman, D.H.; Fournier, G.P.; French, K.L.; Alm, E.J.; Boyle, E.A.; Cao, C.; Summons, R.E. Methanogenic Burst in the End-Permian Carbon Cycle. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 5462–5467. [\[CrossRef\]](#)
267. Strock, K.E.; Krewson, R.B.; Hayes, N.M.; Deemer, B.R. Oxidation Is a Potentially Significant Methane Sink in Land-Terminating Glacial Runoff. *Sci. Rep.* **2024**, *14*, 23389. [\[CrossRef\]](#)
268. Luo, G.; Liu, D.; Yang, H. Microbes in Mass Extinction: An Accomplice or a Savior? *Natl. Sci. Rev.* **2024**, *11*, nwad291. [\[CrossRef\]](#)
269. Song, H.; Wignall, P.B.; Chu, D.; Tong, J.; Sun, Y.; Song, H.; He, W.; Tian, L. Anoxia/High Temperature Double Whammy during the Permian-Triassic Marine Crisis and Its Aftermath. *Sci. Rep.* **2014**, *4*, 4132. [\[CrossRef\]](#)
270. Lau, K.V.; Maher, K.; Altiner, D.; Kelley, B.M.; Kump, L.R.; Lehrmann, D.J.; Silva-Tamayo, J.C.; Weaver, K.L.; Yu, M.; Payne, J.L. Marine Anoxia and Delayed Earth System Recovery after the End-Permian Extinction. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 2360–2365. [\[CrossRef\]](#) [\[PubMed\]](#)
271. Mata, S.A.; Bottjer, D.J. Microbes and Mass Extinctions: Paleoenvironmental Distribution of Microbialites during Times of Biotic Crisis. *Geobiology* **2012**, *10*, 3–24. [\[CrossRef\]](#) [\[PubMed\]](#)
272. Shubert, J.; Bottjer, D.J. Early Triassic Stromatolites as Post-Mass Extinction Disaster Forms. *Geology* **1992**, *10*, 883–886. [\[CrossRef\]](#)
273. Kirton, J.M.C.; Woods, A.D. Stromatolites from the Lower Triassic Virgin Limestone at Blue Diamond, NV USA: The Role of Dysoxia, Enhanced Calcification and Nutrient Availability in the Growth of Post-Extinction Microbialites. *Glob. Planet. Change* **2021**, *198*, 103429. [\[CrossRef\]](#)
274. Baud, A.; Cirilli, S.; Marcoux, J. Biotic Response to Mass Extinction: The Lowermost Triassic Microbialites. *Facies* **1997**, *36*, 238–242. [\[CrossRef\]](#)
275. Vennin, E.; Olivier, N.; Brayard, A.; Bour, I.; Thomazo, C.; Escarguel, G.; Fara, E.; Bylund, K.G.; Jenks, J.F.; Stephen, D.A.; et al. Microbial Deposits in the Aftermath of the End-Permian Mass Extinction: A Diverging Case from the Mineral Mountains (Utah, USA). *Sedimentology* **2015**, *62*, 753–792. [\[CrossRef\]](#)
276. Schobben, M.; Stebbins, A.; Ghaderi, A.; Strauss, H.; Korn, D.; Korte, C. Flourishing Ocean Drives the End-Permian Marine Mass Extinction. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 10298–10303. [\[CrossRef\]](#)
277. Klatt, J.M.; Haas, S.; Yilmaz, P.; de Beer, D.; Polerecky, L. Hydrogen Sulfide Can Inhibit and Enhance Oxygenic Photosynthesis in a Cyanobacterium from Sulfidic Springs. *Environ. Microbiol.* **2015**, *17*, 3301–3313. [\[CrossRef\]](#)
278. Wu, S.; Chen, Z.Q.; Fang, Y.; Pei, Y.; Yang, H.; Ogg, J. A Permian-Triassic Boundary Microbialite Deposit from the Eastern Yangtze Platform (Jiangxi Province, South China): Geobiologic Features, Ecosystem Composition and Redox Conditions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **2017**, *486*, 58–73. [\[CrossRef\]](#)
279. Luo, G.; Kump, L.R.; Wang, Y.; Tong, J.; Arthur, M.A.; Yang, H.; Huang, J.; Yin, H.; Xie, S. Isotopic Evidence for an Anomalously Low Oceanic Sulfate Concentration Following End-Permian Mass Extinction. *Earth Planet. Sci. Lett.* **2010**, *300*, 101–111. [\[CrossRef\]](#)
280. Sheehan, P.M.; Harris, M.T. Microbialite Resurgence after the Late Ordovician Extinction. *Nature* **2004**, *430*, 75–78. [\[CrossRef\]](#) [\[PubMed\]](#)

281. Yao, L.; Aretz, M.; Chen, J.; Webb, G.E.; Wang, X. Global Microbial Carbonate Proliferation after the End-Devonian Mass Extinction: Mainly Controlled by Demise of Skeletal Bioconstructors. *Sci. Rep.* **2016**, *6*, 39694. [[CrossRef](#)] [[PubMed](#)]
282. Liu, Y.; Whitman, W.B. Metabolic, Phylogenetic, and Ecological Diversity of the Methanogenic Archaea. *Ann. N. Y. Acad. Sci.* **2008**, *1125*, 171–189. [[CrossRef](#)]
283. Ehhalt, D.H. The Atmospheric Cycle of Methane. *Tellus A Dyn. Meteorol. Oceanogr.* **1974**, *26*, 58. [[CrossRef](#)]
284. Skeie, R.B.; Hodnebrog, Ø.; Myhre, G. Trends in Atmospheric Methane Concentrations since 1990 Were Driven and Modified by Anthropogenic Emissions. *Commun. Earth Environ.* **2023**, *4*, 317. [[CrossRef](#)]
285. Veron, J.E.N.; Hoegh-Guldberg, O.; Lenton, T.M.; Lough, J.M.; Obura, D.O.; Pearce-Kelly, P.; Sheppard, C.R.C.; Spalding, M.; Stafford-Smith, M.G.; Rogers, A.D. The Coral Reef Crisis: The Critical Importance of . *Mar. Pollut. Bull.* **2009**, *58*, 1428–1436. [[CrossRef](#)]
286. Ehleringer, J.R.; Cerling, T.E. Atmospheric CO₂ and the Ratio of Intercellular to Ambient CO₂ Concentrations in Plants. *Tree Physiol.* **1993**, *15*, 105–111. [[CrossRef](#)]
287. Dlugokencky, E.J.; Nisbet, E.G.; Fisher, R.; Lowry, D. Global Atmospheric Methane: Budget, Changes and Dangers. *Philos. Trans. R. Soc. A* **2011**, *369*, 2058–2072. [[CrossRef](#)]
288. Visscher, P.T.; Stolz, J.F. Microbial Mats as Bioreactors: Populations, Processes, and Products. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **2005**, *1*, 87–100. [[CrossRef](#)]
289. Veiga, M.C.; Jain, M.K.; Wu, W.M.; Hollingsworth, R.I.; Zeikus, J.G. Composition and Role of Extracellular Polymers in Methanogenic Granules. *Appl. Environ. Microbiol.* **1997**, *63*, 403–407. [[CrossRef](#)] [[PubMed](#)]
290. Su, F.; Yang, Y.Y. Microbially Induced Carbonate Precipitation via Methanogenesis Pathway by a Microbial Consortium Enriched from Activated Anaerobic Sludge. *J. Appl. Microbiol.* **2021**, *131*, 236–256. [[CrossRef](#)] [[PubMed](#)]
291. Sowers, K.R.; Baron, S.F.; Ferry, J.G. *Methanosarcina acetivorans* Sp. Nov., an Acetotrophic Methane-Producing Bacterium Isolated from Marine Sediments. *Appl. Environ. Microbiol.* **1984**, *47*, 971–978. [[CrossRef](#)] [[PubMed](#)]
292. Loyd, S.J.; Marissa, N. Smirnoff Progressive Formation of Authigenic Carbonate with Depth in Siliciclastic Marine Sediments Including Substantial Formation in Sediments Experiencing. *Chem. Geol.* **2022**, *594*, 120775. [[CrossRef](#)]
293. Cai, S.; Wei, G.; Xian, H.; Zhu, J.; Xu, Y.G. Methanogen-Mediated Precipitation of Mn Carbonates at the Expense of Mn Oxides. *Geophys. Res. Lett.* **2024**, *51*, e2023GL106921. [[CrossRef](#)]
294. Ramírez-Arenas, P.J.; Latisnere-Barragán, H.; García-Maldonado, J.Q.; López-Cortés, A. Highly Diverse–Low Abundance Methanogenic Communities in Hypersaline Microbial Mats of Guerrero Negro BCS, Assessed through Microcosm Experiments. *PLoS ONE* **2024**, *19*, e0303004. [[CrossRef](#)]
295. Oremland, R.S.; Polcin, S. Methanogenesis and Sulfate Reduction: Competitive and Noncompetitive Substrates in Estuarine Sediments. *Appl. Environ. Microbiol.* **1982**, *44*, 1270. [[CrossRef](#)]
296. Mitterer, R.M. Methanogenesis and Sulfate Reduction in Marine Sediments: A New Model. *Earth Planet. Sci. Lett.* **2010**, *295*, 358–366. [[CrossRef](#)]
297. Vogt, J.C.; Abed, R.M.M.; Albach, D.C.; Palinska, K.A. Bacterial and Archaeal Diversity in Hypersaline Cyanobacterial Mats Along a Transect in the Intertidal Flats of the Sultanate of Oman. *Microb. Ecol.* **2018**, *75*, 331–347. [[CrossRef](#)]
298. Visscher, P.T.; Taylor, B.F.; Kiene, R.P. Microbial Consumption of Dimethyl Sulfide and Methanethiol in Coastal Marine Sediments. *FEMS Microbiol. Ecol.* **1995**, *18*, 145–154. [[CrossRef](#)]
299. Visscher, P.T.; Reid, R.P.; Bebout, B.M.; Hoefft, S.E.; Macintyre, I.G.; Thompson, J.A. Formation of Lithified Micritic Laminae in Modern Marine Stromatolites (Bahamas); the Role of Sulfur Cycling. *Am. Mineral.* **1998**, *83*, 1482–1493. [[CrossRef](#)]
300. Djokic, T.; VanKranendonk, M.J.; Campbel, K.A.; Walter, M.R.; Ward, C.R. Earliest Signs of Life on Land Preserved in ca. 3.5 Ga Hot Spring Deposits. *Nat. Commun.* **2017**, *8*, 15263. [[CrossRef](#)] [[PubMed](#)]
301. Baumgartner, L.K.; Dupraz, C.; Buckley, D.H.; Spear, J.R.; Pace, N.R.; Visscher, P.T. Microbial Species Richness and Metabolic Activities in Hypersaline Microbial Mats: Insight into Biosignature Formation Through Lithification. *Astrobiology* **2009**, *9*, 861–874. [[CrossRef](#)] [[PubMed](#)]
302. Hohl, S.V.; Bian, X.; Viehmann, S.; Yang, S.C.; Raad, R.J.; Meister, P.; John, S.G. A Novel Biomarker for Deep-Time Methanogenesis—Perspectives from Nickel Isotope Fractionation in Modern Microbialites. *Earth Planet. Sci. Lett.* **2025**, *666*, 119492. [[CrossRef](#)]
303. Coffey, J.M.; Flannery, D.T.; Walter, M.R.; George, S.C. Sedimentology, Stratigraphy and Geochemistry of a Stromatolite Biofacies in the 2.72 Ga Tumbiana Formation, Fortescue Group, Western Australia. *Precambrian Res.* **2013**, *236*, 282–296. [[CrossRef](#)]
304. Scheller, S.; Goenrich, M.; Thauer, R.K.; Jaun, B. Methyl-Coenzyme M Reductase from Methanogenic Archaea: Isotope Effects on the Formation and Anaerobic Oxidation of Methane. *J. Am. Chem. Soc.* **2013**, *135*, 14975–14984. [[CrossRef](#)]
305. Van de Vossenberg, J.L.; Driessen, A.J.; Konings, W.N. The Essence of Being Extremophilic: The Role of the Unique Archaeal Membrane Lipids. *Extremophiles* **1998**, *2*, 163–170. [[CrossRef](#)]
306. Myers, T.; Dykstra, C.M. Teaching Old Dogs New Tricks: Genetic Engineering Methanogens. *Appl. Environ. Microbiol.* **2024**, *90*, e02247-23. [[CrossRef](#)]

307. Conrad, R. Methane Production in Soil Environments—Anaerobic Biogeochemistry and Microbial Life between Flooding and Desiccation. *Microorganisms* **2020**, *8*, 881. [\[CrossRef\]](#)
308. Horne, A.J.; Lessner, D.J. Assessment of the Oxidant Tolerance of *Methanosarcina acetivorans*. *FEMS Microbiol. Lett.* **2013**, *343*, 13–19. [\[CrossRef\]](#)
309. Decho, A.W.; Norman, R.S.; Visscher, P.T. Quorum Sensing in Natural Environments: Emerging Views from Microbial Mats. *Trends Microbiol.* **2010**, *18*, 73–80. [\[CrossRef\]](#)
310. Guerrero-Cruz, S.; Cremers, G.; van Alen, T.A.; Op den Camp, H.J.M.; Jetten, M.S.M.; Rasigraf, O.; Vaksmaa, A. Response of the Anaerobic Methanotroph “*Candidatus Methanoperedens nitroreducens*” to Oxygen Stress. *Appl. Environ. Microbiol.* **2018**, *84*, e01832–18. [\[CrossRef\]](#) [\[PubMed\]](#)
311. Magalhães, C.P.; Duarte, M.S.; Pereira, M.A.; Stams, A.J.M.; Cavaleiro, A.J. Facultative Anaerobic Bacteria Enable Syntrophic Fatty Acids Degradation under Micro-Aerobic Conditions. *Bioresour. Technol.* **2025**, *417*, 131829. [\[CrossRef\]](#) [\[PubMed\]](#)
312. Zhao, Y.-C.; Chong, S.+; Zhao, X.-M.; Du, J.-X.; Zou, L.; Yong, Y.-C. Unnatural Direct Interspecies Electron Transfer Enabled by Living Cell-Cell Click Chemistry. *Angew. Chem.* **2024**, *136*, e202402318. [\[CrossRef\]](#)
313. Cockell, C.S. The Similarity of Life across the Universe. *Mol. Biol. Cell* **2016**, *27*, 1553. [\[CrossRef\]](#)
314. Formisano, V.; Atreya, S.; Encrenaz, T.; Ignatiev, N.; Giuranna, M. Detection of Methane in the Atmosphere of Mars. *Science* **2004**, *306*, 1758–1761. [\[CrossRef\]](#)
315. Waite, J.H.; Combi, M.R.; Ip, W.; Thomas, E.C.; McNutt, R.L.; Kasprzak, W.; Yelle, R.; Luhmann, J.; Niemann, H.; Gell, D.; et al. Cassini Ion and Neutral Mass Spectrometer: Enceladus Plume Composition and Structure. *Science* **2006**, *311*, 1419–1422. [\[CrossRef\]](#)
316. Niemann, H.B.; Atreya, S.K.; Bauer, S.J.; Carignan, G.R.; Demick, J.E.; Frost, R.L.; Gautier, D.; Haberman, J.A.; Harpold, D.N.; Hunten, D.M.; et al. The Abundances of Constituents of Titan’s Atmosphere from the GCMS Instrument on the Huygens Probe. *Nature* **2005**, *438*, 779–784. [\[CrossRef\]](#)
317. Mickol, R.L.; Takagi, Y.A.; Kral, T.A. Survival of Non-Psychrophilic Methanogens Exposed to Martian Diurnal and 48-h Temperature Cycles. *Planet. Space Sci.* **2018**, *157*, 63–71. [\[CrossRef\]](#)
318. Hurowitz, J.A.; Catling, D.C.; Fischer, W.W. High Carbonate Alkalinity Lakes on Mars and Their Potential Role in an Origin of Life Beyond Earth. *Elements* **2023**, *19*, 37–44. [\[CrossRef\]](#)
319. Butturini, A.; Benaiges-Fernandez, R.; Fors, O.; García-Castellanos, D. Potential Habitability of Present-Day Martian Subsurface for Earth-Like Methanogens. *Astrobiology* **2025**, *25*, 253–268. [\[CrossRef\]](#)
320. Amils, R.; Fernández-Remolar, D.; Ipbsl Team. Río Tinto: A geochemical and mineralogical terrestrial analogue of Mars. *Life* **2014**, *4*, 511–534. [\[CrossRef\]](#)
321. Yung, Y.L.; Chen, P.; Nealson, K.; Atreya, S.; Beckett, P.; Blank, J.G.; Ehlmann, B.; Eiler, J.; Etiope, G.; Ferry, J.G.; et al. Methane on Mars and Habitability: Challenges and Responses. *Astrobiology* **2018**, *18*, 1221–1242. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.