

Article

Comparison of Freshwater Mussels *Unio tumidus* and *Unio crassus* as Biomonitors of Microplastic Contamination of Tisza River (Hungary)

Wael Almeshal ¹, Anita Takács ², László Aradi ³, Sirat Sandil ⁴, Péter Dobosy ^{4,5}, and Gyula Záray ^{4,5,*}

¹ Doctoral School of Environmental Science, Eötvös Loránd University, 1053 Budapest, Hungary

² Institute of Aquatic Ecology, Centre for Ecological Research, 1113 Budapest, Hungary

³ Lithosphere Fluid Research Laboratory, Department of Petrology and Geochemistry, Institute of Geography and Earth Sciences, Faculty of Science, Eötvös Loránd University, 1053 Budapest, Hungary

⁴ Institute of Aquatic Ecology, Centre for Ecological Research, H-1113 Budapest, Hungary

⁵ National Laboratory for Water Science and Water Security, Institute of Aquatic Ecology, Centre for Ecological Research, H-1113 Budapest, Hungary

* Correspondence: zaray.gyula@ecolres.hu




Figure S1. Stereomicroscopic picture and Raman spectrum of blue fiber found in *Unio tumidus* mussel collected at sampling site Szeged and Raman spectrum of indigo dye reference material.





Figure S2. Stereomicroscopic picture and Raman spectrum of a fiber found in *Unio tumidus* mussel collected at sampling site Szeged and Raman spectrum of PET reference material.






Figure S3. Stereomicroscopic picture and Raman spectrum of a thick fiber found in *Unio tumidus* mussel collected at sampling site Szeged and Raman spectrum of cellulose reference material




Figure S4. Stereomicroscopic picture and Raman spectrum of a fragment found in *Unio tumidus* mussel collected at sampling site Szeged and Raman spectrum of polyamide reference material.