

Article

Finding Targets in Non-Radial FDH Models: A Hybrid Technique Based on STEM and Extended Ratio Based Approach

Mehdi Abbasi ^{1,*}, Mohammad Reza Mozaffari ², Peter F. Wanke ³ and Mohamad Amin Kaviani ⁴

¹ Department of Industrial Engineering, Shiraz Branch, Islamic Azad University, Shiraz 73, Iran

² Department of Mathematics, Shiraz Branch, Islamic Azad University, Shiraz 73, Iran; mozaffari854@yahoo.com

³ Business Analytics and Economics Research Unit, COPPEAD Graduate Business School, Federal University of Rio de Janeiro, Rua Paschoal Lemme, 355, Rio de Janeiro 21941-901, Brazil; peter@coppead.ufrj.br

⁴ Department of Management and Technology, Université du Québec à Montréal, Montreal, QC H0H-H9X, Canada; Kaviani.mohammadamin@courrier.uqam.ca or aminkaviani1366@yahoo.com

* Correspondence: abbasi.iau@gmail.com or abbasi_m@iaushiraz.ac.ir

Abstract: Performance evaluation using interactive methods and extended ratio-based approaches can be very important for some organizations. Free disposal hull models can be created if there is no concern for convexity, and using non-radial DEA models can simultaneously create more logical and practical situations for finding DMU targets. In this paper, a new hybrid technique based on the additive slack based method and enhanced Russel measure in variable return to scale technology has been proposed. The proposed technique can find decision making unit targets in non-radial free disposal hull models using the step method. Furthermore, the extended ratio-based approach in the proposed technique has been applied to find DMU targets of related non-radial free disposal hull models without solving any mathematical programming models. Finally, targets of Fars province pharmaceutical distributing companies were found by applying the proposed hybrid technique.

Keywords: data envelopment analysis (DEA); additive slack based method; non-radial free disposal hull (FDH); step method (STEM); ratio-based approach (RBA)

Citation: Abbasi, M.; Mozaffari, M.R.; Wanke, P.F.; Kaviani, M.A. Finding Targets in Non-Radial FDH Models: A Hybrid Technique Based on STEM and Extended Ratio Based Approach. *Appl. Sci.* **2021**, *11*, 10626. <https://doi.org/10.3390/app112210626>

Academic Editors: João Carlos de Oliveira Matias and Radu Godina

Received: 18 September 2021

Accepted: 8 November 2021

Published: 11 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The accurate estimation of production possibility set (PPS) boundaries is crucial for performance analysis and efficient estimation. Different papers propose alternative approaches to handle the issue of estimating PPSs and their respective boundaries. Non-parametric data envelopment analysis (DEA) is possibly one of the most used linear programming (LP) approaches to build up piecewise PPS boundaries. DEA is a useful tool to evaluate decision making units (DMUs). Efficiency criterion can be considered as a number between 0 and 1 for evaluating a DMU in DEA. If the efficiency criterion for a DMU is 1, the mentioned DMU is efficient, else, it is inefficient. Evaluating DMUs in input-oriented, output-oriented, and combined-oriented radial and non-radial models were developed from a proposal by Farrell (1957) [1], and were then followed by the development of the CCR model by Charnes et al. (1978) [2]. The CCR model was then developed into the BCC model by Banker et al. (1984) [3]. Additive models were then suggested to separate efficient and non-efficient DMUs [4]. Tone (2001) proposed a slack based model which evaluates DMUs considering the relationship between CCR models [5]. The Russel Graph Model (RGM) and the relationship between SBM and additive DEA models are

very important subjects that have been studied [6]. These models determine the benchmark for efficient DMUs, in addition to calculating efficiency and non-efficiency criterion of DMUs. Over three decades, extensive and useful studies on DEA have been undertaken to calculate DMU efficiencies [7,8] and to find DMU benchmarks [9,10]. Chen and Zhu (2020) completed efficient and non-efficient definitions on the basis of slack variables using the slack based method, and showed that additive slacks-based models (ASBM) and enhanced RGM are equal. Moreover, the authors showed that the simultaneous use of ASBM and network DEA models can create a comparable DEA score. Finding DMU targets in ASBM by eliminating convexity conditions can be investigated in practical studies [11]. The use of non-radial FDH models based on ASBM can also be beneficial in practical studies.

Specifically, Free Disposal Hull (FDH) is a subclass of DEA models where DMUs are not projected on the piecewise convex envelope, but are projected on the actual maximal attainable boundary, which results in a staircase shape for the single input–output case. In other words, FDH, which was first introduced by Deprins et al. (1984), evaluates DMU efficiency by considering the closest inner approximation of the true non-convex (disposable) boundary [12]. Many studies have been investigated FDH models. Soleimani-Damaneh et al. estimated returns-to-scale in FDH models [13]. Soleimani-Damaneh and Rashidi proposed a polynomial-time algorithm to estimate returns to scale in FDH models [14]. Mostafaei and Soleimani-Damaneh proposed the definition, characterization and calculation of global sub-increasing and global sub-decreasing returns to scale in FDH technologies [15]. Fukuyama et al. measured efficiency with non-convex FDH technology [16]. Manzari Tavakoli and Mostafaei studied FDH efficiency scores of units with network structures [17]. Arfa et al. measured the efficiency of hospital cardiology wards using the FDH approach [18]. Kerstens and Van De Woestyne reviewed solution methods for non-convex FDH models and give some critical comments [19]. Soleimani-Damaneh and Mostafaei identified the anchor points in FDH models [20]. Mirmozaffari et al. proposed an improved DEA model based on SBM and FDH models [21]. One issue that is frequently neglected in FDH models is the identification of DMU targets, which is a cumbersome task due to boundary non-convexity, especially when the number of inputs and outputs increase. A possible approach is to define a multiple objective function for measuring the closeness among the DMU under analysis and its eventual targets [22].

Multiple objective linear programming (MOLP) is a form of multiple objective decision making (MODM). In MODM problems, more than one objective is considered in regard to the opinions of the decision maker (DM). Interactive methods (IMs) are a kind of MODM and MOLP methods. IMs explore the criterion space on the progressive definition of the DM's preferences at each iteration [23]. IMs have been used in some reported works [24,25]. Traditional DEA models tend to ignore the DM's preferences and value judgment in the computation of the DMU targets, completely. The use of IMs allows the obtainment of DMU targets which have perfect adaptation for the DM's preferences [24]. An IM was applied for the extension of DEA to effectiveness analysis [26]. The step method (STEM) that was introduced in 1971 [27] is an IMs in MOLP, and has been reported in several studies [28–31]. To the best of our knowledge, STEM has not yet been used by researchers to find targets in non-radial FDH models which consider ASBM and enhanced Russell measures in variable return to scale technology (the first research gap).

Multiple criteria decision making (MCDM) is divided into multiple attribute decision making (MADM) and MODM. In situations where the data are fuzzy, a combination of DEA and fuzzy MCDM [32–34] can be used for the development of the proposed technique. In this study, as the data of the second case study were deterministic, it was not necessary to use fuzzy methods.

FDH is a well-known subclass of DEA models and is based on two distinctive features that are reflected in the PPS boundary. First, FDH ensures that efficiency evaluations are affected only by actually observed performances. Secondly, FDH relies on the non-

convexity assumption which satisfies free disposability in PPS. There is inherent computational complexity to solve FDH models. As a matter of fact, FDH models are mixed 0–1 LP, and solving them is difficult. In this regard, a ratio-based approach (RBA) is proposed to solve radial FDH models without solving any mathematical programming models [13]. This approach has been employed by some other studies [15,20]. To the best of our knowledge, RBA is one of the most suitable suggested methods to find DMU targets of radial FDH models without solving any mathematical programming models. DMU target finding of non-radial FDH models without solving any mathematical programming models can be considered as another research gap. It can be achieved by extending RBA. As ASBM relates to enhanced Russell measures, finding non-radial DMU targets has been possible using extended RBA.

In Fars province pharmaceutical distributing companies (the second case study), there was a variable return to scale assumption. Moreover, as the combination of pharmaceutical distributing companies was impossible, using FDH models was beneficial. Non-radial FDH models based on ASBM in variable return to scale technology can therefore be considered in the proposal for a technique for finding DMU targets. Therefore, considering mentioned research gaps, two research questions have considered as follows:

- (1) Is it possible to propose a technique to find all DMU targets in non-radial FDH models based on ASBM using IM?
- (2) Is it possible to find DMU targets of non-radial FDH models in the proposed technique without solving any mathematical models?

A hybrid technique to answer the above two research questions with the following properties has been proposed as the innovation of this research:

- (a) DMU target finding in non-radial FDH models based on ASBM are more realistic because they are based on a non-convexity assumption,
- (b) Proposing a new LP formulation of ASBM,
- (c) Applying IMs instead of regular DEA methods to find FDH models targets which have more adaptation to the DM's preferences,
- (d) Finding the required DMU targets in FDH models using an algorithm that works by checking some conditions for DMUs without solving of any mathematical models.

To the best of our knowledge, proposing a hybrid technique with the mentioned properties has not been reported until now. It is notable that, according to the practical view, the proposed technique will be beneficial if finding DMU targets in a studied organization is useful. According to the theoretical view, as the proposed technique works based on mathematical modelling, considering assumptions and determining suitable parameters to compose related models are important subjects, too. Therefore, to implement and generalize the results, practical and theoretical views should be simultaneously considered.

There are several outlier detection methods, such as parametric robust regression in statistics [35] and non-parametric k-means in data mining [36]. Moreover, a predictive DEA model for outlier detection was proposed, and a comprehensive set of simulation experiments were conducted to examine the relative performance of the suggested method with two popular mentioned methods under the influence of five factors. The results provide users with practical guidelines on how to choose appropriate methods to detect outliers [37]. Outlier detection and investigation of the sensitivity of the modeling approach to outliers can be applied for the development of the proposed technique.

The paper is structured in following sections: first, the background on ASBM, MOLP, STEM, and RBA is provided. After that, a new hybrid technique is introduced to find DMU targets in non-radial FHD models based on ASBM. Finally, the presented technique is applied in two real case studies.

2. Background

In this section, ASBM, STEM to solve MOLP, and RBA for finding targets of radial FDH models are briefly described. The purpose of this section is to introduce the theoretical basis for finding targets of non-radial FDH models based on ASBM using STEM and extended RBA.

2.1. Additive Slack Based Model

Suppose DMU_j , $j = 1, \dots, n$ by consuming m inputs $x_{1j}, \dots, x_{mj}$, $j = 1, \dots, n$ can produce s outputs $y_{1j}, \dots, y_{sj}$, $j = 1, \dots, n$. The background of ASBM can be related to the additive model in Charnes et al. (1985) [4] and Green et al. (1997) [6]. Therefore, efficiency for output r of DMU_o is defined as $\frac{y_{ro}}{y_{ro} + t_r} \leq 1$ and non-efficiency for output r of DMU_o is defined as $1 - \frac{y_{ro}}{y_{ro} + t_r} = \frac{t_r}{y_{ro} + t_r} \leq 1$. Furthermore, efficiency for input i of DMU_o is defined as $\frac{x_{io} - s_i}{x_{io}} \leq 1$ and non-efficiency for output r of DMU_o is defined as $1 - \frac{x_{io} - s_i}{x_{io}} = \frac{s_i}{x_{io}} \leq 1$. Therefore, by suggesting model (1), the relationship between non-efficiency calculated by Greek et al. (1997) [6] and ASBM based on efficiency and non-efficiency definitions have presented [11].

$$\begin{aligned} \text{Min } & \frac{1}{s+m} \left(\sum_{r=1}^s \frac{y_{ro}}{y_{ro} + t_r} + \sum_{i=1}^m \frac{x_{io} - s_i}{x_{io}} \right), \\ \text{s. t. } & \\ & \sum_{j=1}^n \mu_j x_{ij} + s_i = x_{io}, i = 1 \dots m, \\ & \sum_{j=1}^n \mu_j y_{rj} - t_r = y_{ro}, r = 1 \dots s, \\ & \sum_{j=1}^n \mu_j = 1, \\ & \mu_j, s_i, t_r \geq 0, i = 1 \dots m, r = 1 \dots s. \end{aligned} \quad (1)$$

Model (1) is a nonlinear mathematical programming model containing linear constraints and linear fractional objective function. In this regard, model (1) by Chen and Zhu (2020) [11] is equivalent to RGM developed by Fare et al. (1985). Moreover, finding DMU targets using slack variables are important because focus is given to the summation of slack variables. In this regard, the projection of DMU with respect to Model (1) is calculated by $\sum_{j=1}^n \mu_j^* x_{ij} = x_{io} - s_i^*$, $i = 1 \dots m$, and $\sum_{j=1}^n \mu_j^* y_{rj} = y_{ro} + t_r^*$, $r = 1 \dots s$, which μ_j^* , s_i^* , and t_r^* are optimal solutions of Model (1).

2.2. MOLP and STEM

A general formulation of the MOLP problem is given in model 2.

$$\begin{aligned} \text{Max } G(u) &= (g_1(u), \dots, g_k(u)) = (C_1^T u, \dots, C_k^T u) \\ \text{s. t. } u &\in W = \{u \in R_+^v | Au \leq b\} \end{aligned} \quad (2)$$

where $G(u) = (g_1(u), \dots, g_k(u))$ represents the objective function vector. Linear objective functions are denoted by $C_l^T u$ where $u = (u_1, \dots, u_v)$ is the decision-making vector. The symbol T is a transposed vector. $g_l(u)$, C_l , and c_l are the l^{th} objective function, the vector of decision-making variable coefficients in the l^{th} objective function, and the coefficients of l^{th} objective function namely C_l , $l = 1, \dots, k$, per n existing variables, respectively. W is the feasible region of the MOLP problem and k is the number of the objective function. The decision-making variable multiples matrix is denoted by A , while b represents the right-hand side vector of the constraints. $Au \leq b$ is the constraints of the feasible region and R_+^v , representing the Euclidean space comprising all nonnegative vectors in a v -dimensional space. In this MOLP problem, the l^{th} objective function is formulated as $C_l^T u =$

$c_{1l}u_1 + c_{2l}u_2 + \dots + c_{vl}u_v$, $l = 1, \dots, k$. The vector $u^* \in W$ is considered as an efficient (non-dominated) solution, if there does not exist another $u \in W$, such that $g_l(u) \geq g_l(u^*)$ for all l and $g_l(u) > g_l(u^*)$ for at least one l .

STEM is an IM that can be used to solve MOLP problems. It works based on the obtained information from DM preferences and reduces the feasible region, step by step. STEM relies on DMs information to identify feasible and efficient solutions during the procedure. STEM includes following steps [27,30]:

Step 0: building-up the pay-off table

Objective functions should be optimized separately as follows (cf. model 3 and Table 1):

$$\begin{aligned} \text{Max } g_l(u) &= C_l^T u \\ \text{s. t. } u &\in W = \{u \in R_+^v | Au \leq b\} \end{aligned} \quad (3)$$

Table 1. The pay-off table.

	1	2	...	l	...	k
1	g_1^*		...		...	z_{1k}
2	z_{21}	g_2^*	...	z_{2l}	...	z_{2k}
.	.	.	.	.	.	.
.	.	.	.	.	.	.
.	.	.	.	.	.	.
l	z_{l1}	z_{l2}	...	g_l^*	...	z_{lk}
.	.	.	.	.	.	.
.	.	.	.	.	.	.
.	.	.	.	.	.	.
k	z_{k1}	z_{k2}	...	z_{kl}	...	g_k^*

The diagonal elements, represented by g_l^* , are the optimal solutions for the single $g_l(u) = C_l^T(u)$, $l = 1, \dots, k$ problem obtained through the solving of model 3. z_{dl} values are the results for d^{th} objective function, computed upon the optimal solution obtained for l^{th} objective function, $l = 1, \dots, k$, $d = 1, \dots, k$, $d \neq l$.

Step 1: Calculation Phase

The computation of coefficients β_l , $l = 1 \dots k$, is the cornerstone to compute the relative importance (Equation (4)) of each distance from the optimal objective function value. Suppose that π_l denotes the relative importance of the distance between objective functions and their optimal values. Although these coefficients are locally meaningful, they cannot capture the overall importance, unlike other utility models. It is therefore necessary to solve model 5, where the solution obtained in the p^{th} iteration is denoted by $G^p = (g_1(u^p), g_2(u^p), \dots, g_k(u^p))$.

$$\pi_l = \begin{cases} \left(\frac{g_l^* - g_l^{\min}}{g_l^*} \right) \left[\frac{1}{\sqrt{\sum_{d=1}^v (C_{dl})^2}} \right] & \text{if } g_l^* > 0 \\ \left(\frac{g_l^{\min} - g_l^*}{g_l^*} \right) \left[\frac{1}{\sqrt{\sum_{d=1}^v (C_{dl})^2}} \right] & \text{if } g_l^* \leq 0 \end{cases} \quad (4)$$

and $\beta_l = \frac{\pi_l}{\sum_{d=1}^k \pi_d}$, $l = 1, \dots, k$. Model 5 is given as follows:

$$\begin{aligned} \text{Min } h \\ \text{s. t. } h &\geq (g_l^* - g_l(u)) \cdot \beta_l \quad l = 1, \dots, k, \end{aligned}$$

$$u \in W^p, h \geq 0. \text{ Where } W^p = \{u | Au \leq b, u \geq 0\} \cup W^{p-1} \quad (5)$$

W^p represents the feasible region in the p^{th} iteration. In order to find the vector $u \in W^p$, which provides the minimum of maximum distance between the objective function vector of $G(u) = (g_1(u), g_2(u), \dots, g_k(u))$ and its optimal vector, $G^* = (g_1^*, \dots, g_k^*)$, h should be minimized. h indicates the maximum distance of the functions from their optimal values based on their relative importance for each individual feasible solution in the feasible region. In other words, h indicates the closest possible distance to the optimal value of the l^{th} objective function, that is g_l^* . Before proceeding to Step 2, the minimum value in column l of the pay-off table should be picked up. It is denoted as $g_l^{\min}$.

Step 2: Decision Phase

In this phase, the DM provides relative importance information with respect to the solution collected during the first step of the p^{th} iteration, that is $G^p = (g_1(u^p), g_2(u^p), \dots, g_k(u^p))$, where u^p denotes the feasible solution in the p^{th} iteration. If all objective function values are satisfied, in light of DM preferences, the best compromise solution is obtained, and the STEM algorithm finished. Otherwise, the DM should modify some of the $g_l(u^p) = g_l^p$ to confirm that the values of the l^{th} objective function in the p^{th} iteration is satisfied. In other words, this modification amount, Δg_q , is necessary to collectively improve other remaining objective functions. Thus, the feasible region should be also adjusted for the next iteration.

$$W^{p+1} = W^p \cup \left\{ u \mid \begin{array}{l} g_q(u) \geq g_q(u^p) - \Delta g_q \\ g_l(u) \geq g_l(u^p), \quad l = 1, \dots, k, l \neq q \end{array} \right\} \quad (6)$$

Δg_l denotes the number of modifications made to the l^{th} objective function in order to improve the other objective functions, and W^{p+1} denotes the feasible region in iteration $p + 1$. When the coefficients in the subsequent iterations are computed, the coefficients within π_l should be zeroed. Therefore, other values for π_l and $\beta_l, l = 1, \dots, k, l \neq k$ should be re-determined using Equation (4), before re-solving model 5 in the $p = p + 1$ iteration.

2.3. Ratio Based Approach

The ratio based approach is an approach to find DMU targets of radial FDH models without solving any mathematical programming models. To the best of our knowledge, Soleimani-Damaneh et al. (2006) [13] were the first researchers to propose RBA to find targets of radial FDH models. They considered a set of n peer DMUs ($DMU_j, j = 1, \dots, n$), such that each DMU_j produces multiple outputs $y_{rj} > 0$ ($r = 1, \dots, s$) by utilizing multiple inputs $x_{ij} > 0$ ($i = 1, \dots, m$). Considering $DMU_o(x_o, y_o)$ ($o = 1, \dots, n$) as the unit under assessment, the basic input-oriented and output-oriented linear mixed-integer radial FDH model under variable returns-to-scale technology are shown by models 7 and 8, respectively [13,20].

$$\theta_o = \text{Min } \theta, \quad \text{s.t.} \quad (7)$$

$$\begin{aligned} \sum_{j=1}^n \mu_j x_{ij} &\leq \theta x_{io}, \quad i = 1 \dots m, \\ \sum_{j=1}^n \mu_j y_{rj} &\geq y_{ro}, \quad r = 1 \dots s, \\ \mu_j &= \delta \omega_j, \quad \omega_j \in \{0,1\}, \quad j = 1, \dots, n, \\ \delta &= 1, \quad \sum_{j=1}^n \omega_j = 1. \\ \varphi_o &= \text{Max } \varphi, \end{aligned}$$

$$\begin{aligned}
 & \text{s. t} \\
 & \sum_{j=1}^n \mu_j x_{ij} \leq x_{io}, i = 1 \dots m, \\
 & \sum_{j=1}^n \mu_j y_{rj} \geq \varphi y_{ro}, r = 1 \dots s, \\
 & \mu_j = \delta \omega_j, \quad \omega_j \in \{0,1\}, j = 1, \dots, n, \\
 & \delta = 1, \sum_{j=1}^n \omega_j = 1.
 \end{aligned} \tag{8}$$

In model 7, DMU_o is called radial input-oriented FDH-efficient if $\theta_o = 1$. Moreover, in model 8, DMU_o is called radial output-oriented FDH-efficient if $\varphi_o = 1$. The targets of radial models 7 and 8 can be found by computing some simple ratios using RBA. After considering DMU_o ($o = 1, \dots, n$) as an under assessment DMU, for $j = 1, \dots, n$, $\lambda_{jo} = \max_r \{y_{ro}/y_{rj}\}$ and $\lambda^{jo} = \min_i \{x_{io}/x_{ij}\}$ are defined. The optimized objective function in radial models 7 and 8 is then calculated as $\theta_o = \min_{j=1, \dots, n, y_j \geq y_o} \left\{ \max_i \{x_{ij}/x_{io}\} \right\}$ and $\varphi_o = \max_{j=1, \dots, n, x_j \leq x_o} \left\{ \min_r \{y_{rj}/y_{ro}\} \right\}$, respectively [13]. The details on RBA validity have been reported previously [13,14,19]. The targets of radial FDH models 7 and 8 can be found by RBA.

3. A New Hybrid Technique for Finding DMU Targets in Non-Radial FHD Models

In this section, a hybrid technique including two main parts is proposed. At first, an algorithm containing two interactive stages is introduced to find DMU targets in non-radial FDH models. Applied models are obtained based on ASBM. The first interactive stage is proposed to determine efficient DMUs and their targets. STEM is used in the second interactive stage to find targets of other DMUs. Finding DMU targets of three kinds of non-radial FDH models was required in the first part of hybrid technique. RBA is one of the suitable suggested methods to find DMU targets of radial FDH models. In the second part of the hybrid technique, extended RBA is proposed to find DMU targets of non-radial FDH models without solving any mathematical models. Extended RBA, included two steps, found alternative DMU targets just by checking some conditions for DMUs in non-radial FDH models without solving any mathematical models. In the first step, DMUs of feasible region were found. In the second step, the optimum objective function value was calculated and DMU targets were found.

3.1. The Interactive Algorithm

In this section, an algorithm is proposed to find DMU targets in non-radial FDH models. The interactive algorithm (the first part of proposed hybrid technique) contains two interactive stages. The purpose of the first interactive stage is to determine efficient DMUs and their targets. The second interactive stage determines the target of other DMUs using STEM. DMU targets are found through solving three non-radial FDH models in interactive algorithm. Applied models are obtained based on ASBM.

Classical DEA models have been created based on decreasing inputs and increasing outputs. Moreover, only one variable has been defined for decreasing all inputs and only one another variable has been defined for increasing all outputs in classical radial models. However, in non-radial classical models, different variables have been defined separately for decreasing each input, and different variables have been defined separately for increasing different outputs. In addition, all DMUs in the research area should be considered, and the relationship $n \geq 3(m + s)$ exists between the number of DMUs (n), number of inputs (m), and number of outputs (s) parameters. In this regard, the main improvements that are obtained from solving DEA models relates to non-efficient DMUs. Therefore, decreasing the number of DMUs may increase the number of efficient DMUs [38]. In

DEA models, choosing appropriate inputs and outputs is an important step which may significantly affect the efficient frontier. In this regard, inputs and outputs should be relevant to the research area, gathering related data for DMUs should be possible, and experts should confirm them. It is considerable that using real data is preferred rather than gathering from experts' opinions. However, if gathering real data is not impossible, experts' judgment is used. In this situation, experts should have enough information about the problem. As classical DEA models are created based on decreasing inputs and increasing outputs, this subject should be considered for selecting inputs and outputs as well. Moreover, all DMUs in the research area should be considered, and the establishment of the relationship $n \geq 3(m + s)$ between inputs, outputs, and DMU numbers is recommended. As a matter of fact, in situations where $n < 3(m + s)$, the number of efficient DMUs may be increased. Therefore, a set of non-duplicated DMU_j, $j = 1, \dots, n$, that utilizes m positive inputs, x_{ij} , $i = 1, \dots, m$, to produce s positive outputs, y_{rj} , $r = 1, \dots, s$, are considered. Moreover, objective functions e_l , $l = 1, \dots, m + s$, and decision making variables, u_v , $v = 1, \dots, n + m + s$ are defined as

$$(e_1(u), \dots, e_m(u), e_{m+1}(u), \dots, e_{m+s}(u)) = \left(\frac{-w_1}{m} \left(\frac{x_{10}-s_1}{x_{10}} \right), \dots, \frac{-w_m}{m} \left(\frac{x_{m0}-s_m}{x_{m0}} \right), -\frac{w_{m+1}}{s} \left(\frac{y_{(m+1)0}}{y_{(m+1)0}+t_{m+1}} \right), \dots, -\frac{w_{m+s}}{s} \left(\frac{y_{(m+s)0}}{y_{(m+s)0}+t_{m+s}} \right) \right)$$

and $(u_1, \dots, u_n, u_{n+1}, \dots, u_{n+m}, u_{n+m+1}, \dots, u_{n+m+s}) = (\lambda_1, \dots, \lambda_n, s_1, \dots, s_m, t_1, \dots, t_s)$, respectively.

By modifying objective functions coefficients and turning min into max objective functions, a multi objective non-linear FDH model based on ASBM is formulated as follows (model 9):

$$\text{Max } E(e_1(u), \dots, e_m(u), e_{m+1}(u), \dots, e_{m+s}(u)) = \left(\frac{-w_1}{m} \left(\frac{x_{10}-s_1}{x_{10}} \right), \dots, \frac{-w_m}{m} \left(\frac{x_{m0}-s_m}{x_{m0}} \right), -\frac{w_{m+1}}{s} \left(\frac{y_{(m+1)0}}{y_{(m+1)0}+t_{m+1}} \right), \dots, -\frac{w_{m+s}}{s} \left(\frac{y_{(m+s)0}}{y_{(m+s)0}+t_{m+s}} \right) \right) \quad (9)$$

$$\text{s.t. } (\lambda_1, \dots, \lambda_n, s_1, \dots, s_m, t_1, \dots, t_s) \in S =$$

$$\left\{ (\lambda_1, \dots, \lambda_n) \in R_+^n \left| \begin{array}{l} \sum_{j=1}^n \lambda_j x_{ij} + s_i = x_{i0}, \quad i = 1 \dots m, \\ \sum_{j=1}^n \lambda_j y_{rj} - t_r = y_{r0}, \quad r = 1 \dots s, \\ \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, \quad j = 1 \dots n, \\ s_i \geq 0, \quad i = 1 \dots m, \quad t_r \geq 0, \quad r = 1 \dots s. \end{array} \right. \right\}$$

It is considerable that model 9 is a non-linear mathematical model because $e_{m+1}(u), \dots, e_{m+s}(u)$ are non-linear. By considering $\text{Max } E(-1/(e_1(u)), \dots, -1/(e_m(u)), -1/(e_{m+1}(u)), \dots, -1/(e_{m+s}(u)))$ instead of $\text{Max } E(e_{m+1}(u), \dots, e_{m+s}(u))$, the equivalent multi objective linear FDH model is composed as model 10.

$$\text{Max } E(e_1(u), \dots, e_m(u), e_{m+1}(u), \dots, e_{m+s}(u)) = \left(\frac{-w_1}{m} \left(\frac{x_{10}-s_1}{x_{10}} \right), \dots, \frac{-w_m}{m} \left(\frac{x_{m0}-s_m}{x_{m0}} \right), \frac{w_{m+1}}{s} \left(\frac{y_{(m+1)0}+t_{m+1}}{y_{(m+1)0}} \right), \dots, \frac{w_{m+s}}{s} \left(\frac{y_{(m+s)0}+t_{m+s}}{y_{(m+s)0}} \right) \right) \quad (10)$$

$$\text{s.t. } (\lambda_1, \dots, \lambda_n, s_1, \dots, s_m, t_1, \dots, t_s) \in S =$$

$$\left\{ (\lambda_1, \dots, \lambda_n) \in R_+^n \left| \begin{array}{l} \sum_{j=1}^n \lambda_j x_{ij} + s_i = x_{i0}, \quad i = 1 \dots m, \\ \sum_{j=1}^n \lambda_j y_{rj} - t_r = y_{r0}, \quad r = 1 \dots s, \\ \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, \quad j = 1 \dots n, \\ s_i \geq 0, \quad i = 1 \dots m, \quad t_r \geq 0, \quad r = 1 \dots s. \end{array} \right. \right\}$$

Now by considering $\theta_i = \frac{x_{i0}-s_i}{x_{i0}}$, $\varphi_r = \frac{y_{r0}+t_r}{y_{r0}}$, and considering 1 for w and w' in objective functions, model 11 is built as follows:

$$\text{Max } E(e_1(u), \dots, e_m(u), e_{m+1}(u), \dots, e_{m+s}(u)) = \left(\frac{-1}{m} \theta_1, \dots, \frac{-1}{m} \theta_m, \frac{1}{s} \varphi_1, \dots, \frac{1}{s} \varphi_s \right) \quad (11)$$

$$\text{s. t. } (\lambda_1, \dots, \lambda_n, \theta_1, \dots, \theta_m, \varphi_1, \dots, \varphi_s) \in S = \left\{ (\lambda_1, \dots, \lambda_n, \theta_1, \dots, \theta_m, \varphi_1, \dots, \varphi_s) \in \mathbb{R}_+^n \left| \begin{array}{l} \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, j = 1 \dots n, \\ \sum_{j=1}^n \lambda_j x_{ij} \leq \theta_i x_{i0}, \theta_i \leq 1, i = 1 \dots m, \\ \sum_{i=1}^n \lambda_i y_{rj} \geq \varphi_r y_{r0}, \varphi_r \geq 1, r = 1 \dots s. \end{array} \right. \right\}$$

It is mentionable that in non-radial model 11, DMU targets are obtained by decreasing inputs ($\theta_i \leq 1, i = 1, \dots, m$) and increasing outputs ($\varphi_r \geq 1, r = 1, \dots, s$), simultaneously. Now, two related interactive stages are described as follows.

3.1.1. The First Interactive Stage

In the first interactive stage, efficient DMUs are determined, and the targets of these DMUs defined. For DMU_o, $o = 1, \dots, n$, variables $x_{i0} = x_{ij}, i = 1, \dots, m, j = 1, \dots, n$, and $y_{r0} = y_{rj}, r = 1, \dots, s, j = 1, \dots, n$ are defined. Non-radial FDH model 12 is then composed for DMU_j, $j = 1, \dots, n$ to distinguish DMUs over efficient frontier as follows.

$$Z_j^o = \text{Max } \sum_{l=1}^{m+s} e_l(u) = \sum_{i=1}^m \frac{-1}{m} \theta_i + \sum_{r=1}^s \frac{1}{s} \varphi_r, \quad \text{s. t.} \quad (12)$$

$$u \in S = \left\{ u \left| \begin{array}{l} \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, j = 1 \dots n, \\ \sum_{j=1}^n \lambda_j x_{ij} \leq \theta_i x_{i0}, 0 \leq \theta_i \leq 1, i = 1 \dots m, \\ \sum_{i=1}^n \lambda_i y_{rj} \geq \varphi_r y_{r0}, \varphi_r \geq 1, r = 1 \dots s. \end{array} \right. \right\}.$$

After the solving of non-radial model 12 for DMU_j, $j = 1, \dots, n$, M (the members of M set are DMUs that lay on the efficient frontier) and M' sets (the process of targets determination of M' set members should be carried out through the second interactive stage of the interactive algorithm) are defined as $M = \{DMU_j | Z_j^o = 0, \forall j\}$ and $M' = \{DMU_j | DMU_j \notin M, \forall j\}$, respectively. It is considerable that DMU_j, $j = 1, \dots, n$ of non-radial model 12 is efficient if $Z_j^o = 0$. The targets of DMUs that are members of M set are then defined as DMU_j Target = $\{DMU_j\}$, $DMU_j \in M$. If $M' = \emptyset$ (in the situations where the process of targets determination for all DMUs has been carried out in the first interactive stage of the interactive algorithm), the interactive algorithm is finished; else (in situation that $M' \neq \emptyset$), the second interactive stage is run.

3.1.2. The Second Interactive Stage

In the second interactive stage, non-efficient DMUs targets are determined by STEM. In this regard, a DMU_j $\in M'$ is considered as DMU_o (a non-proceed DMU_j, $j = 1, \dots, n$) and $x_{i0} = x_{ij}, i = 1, \dots, m, j = 1, \dots, n$, and $y_{r0} = y_{rj}, r = 1, \dots, s, j = 1, \dots, n$, are defined; then the pay-off table (as described in Section 2.2) is constructed by composing and solving the non-radial model 13 for $f_l(u), l = 1, \dots, m + s$ (step 0 of STEM).

$$\text{Max } \{f_1(u) = -x_{10}\theta_1, \dots, f_m(u) = -x_{m0}\theta_m, f_{m+1}(u) = y_{10}\varphi_1, \dots, f_{m+s}(u) = y_{s0}\varphi_s\} \quad \text{s. t.} \quad (13)$$

$$u \in S = \left\{ u \left| \begin{array}{l} \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, j = 1 \dots n, \\ \sum_{j=1}^n \lambda_j x_{ij} \leq \theta_i x_{i0}, 0 \leq \theta_i \leq 1, i = 1 \dots m, \\ \sum_{i=1}^n \lambda_i y_{rj} \geq \varphi_r y_{r0}, \varphi_r \geq 1, r = 1 \dots s. \end{array} \right. \right\}.$$

It is considerable that a DMU in S (in non-radial model 13) is efficient if all $\theta_i^* = 1, i = 1 \dots m$, and all $\varphi_r^* = 1, r = 1 \dots s$. The DM is then asked to suggest a new target for DMU_o. If DM does not suggest a new target for DMU_o, "There is no target for DMU_o in FDH model", then $M' = M' - DMU_o$ (omit DMU_o from M' set), and go to the second interactive stage (to find targets of another non-proceed DMU_j). If DM suggest a new target for DMU_o, extract DM's opinions about the desired value of $f_l, l = 1, \dots, m + s$ from the suggested DMU_o target. This means that the value of $f_l, l = 1, \dots, m + s$ should be as good as the suggested DMU_o target by DM. So the proposed target of DM is considered as DM's opinions for all $m + s$ objective functions ($f_l, l = 1, \dots, m + s$). Attempt are then made to find

the best compromise solution considering DM's opinions using the required iterations of step 1 and 2 of STEM (Section 2.2).

After that, the first iteration of STEM begins (set $p = 1$). In step 1 of STEM (Section 2.2), first $\pi_l, l = 1, \dots, m + s$ is calculated by Equation (4) considering the results of the second interactive stage, then $\beta_l = \frac{\pi_l}{\sum_{d=1}^{m+s} \pi_d}, l = 1, \dots, m + s$ is calculated. If at least one $\beta_l, l = 1, \dots, m + s$ is not calculable, there is no compromise solution (in this situation, obtaining the suggested target of DM for DMU_o is impossible) and the interactive algorithm continues from the start of the second interactive stage (to find another suitable target for DMU_o). In the situation that the value of all $\beta_l, l = 1, \dots, m + s$, are calculable, model 14 is as follows:

$$\begin{aligned} & \text{Min } h \\ & \text{s.t. } h \geq (f_l^* - f_l(u)), \beta_l, l = 1, \dots, m + s, \\ & u \in S^p, h \geq 0. \text{ Where } S^p = \left\{ u \mid \begin{cases} \sum_{j=1}^n \lambda_j = 1, \lambda_j \in \{0,1\}, j = 1 \dots n, \\ \sum_{j=1}^n \lambda_j x_{ij} \leq \theta_i x_{io}, 0 \leq \theta_i \leq 1, i = 1 \dots m, \\ \sum_{j=1}^n \lambda_j y_{rj} \geq \phi_r y_{ro}, \phi_r \geq 1, r = 1 \dots s. \end{cases} \right\} \cup S^{p-1} \end{aligned} \quad (14)$$

It is considerable that S^p denotes the feasible region in the p^{th} iteration. By solving model 14, alternative targets of DMU_o (DMU_o Target = $\{DMU_j | \lambda_j^* = 1, \forall j\}$) are found. The required parameters for Step 2 of STEM (Section 2.2) are then calculated. In this phase, the objective function values of the first step in the p^{th} iteration are considered as $F^p = (f_1(u^p), \dots, f_{m+s}(u^p))$ where u^p denotes the feasible solution in the p^{th} iteration. If all objective function values are satisfied considering the DM's opinion (the situation that the best compromise solution is found), "consider suggested target as the target of DMU_o", $M' = M' - DMU_o$, go to the second interactive stage. If "DM is not satisfied with all objective function values", or "no feasible integer solution is found" (no compromise solution exists in these two situations), algorithm is continued from the start of the second interactive stage. Otherwise $f_l(u^p) = f_l^p$ considering DM's opinions, which means the values of f_l in the p^{th} iteration are satisfied. Consequently, in order to improve the other objective functions, the f_l values are modified by the amounts of Δf_q considering DM's opinions. In this case, DMUs in the feasible region are distinguished in the next iteration.

As mentioned, finding the best compromise solution in the second interactive stage is desired through STEM. If the best compromise solution is obtained using required iterations of step 1 and 2 of STEM, suggested targets are considered as targets of DMU_o. In the situation that no compromise solution is found (in two situations a compromise solution does not exist: 1- at least one $\beta_l, l = 1, \dots, m + s$ is not calculable, 2- there is only one similar unsatisfied objective function in two consecutive steps of STEM), the second interactive stage repeats.

The brief of the interactive algorithm for finding DMU targets is shown in Figure 1. In the interactive stage 1, efficient DMUs and their targets are determined. It is remarkable that by applying the first interactive stage of the interactive algorithm, at least one DMU_j, $j = 1, \dots, n$ is efficient (at least, one member belongs to M set). The number of M' set members are determined by composing and solving of non-radial model 12 for all DMUs. The required number of iterations for the interactive algorithm depends on the number of M' set members. Non-radial model 12 should be constructed and solved n (number of DMUs) times. If the ratio of "the number of M' set members" to "all DMUs" is near to 0, the interactive algorithm needs lesser iterations. If this ratio is near to 1, the interactive algorithm needs more iterations. In the interactive stage 2, non-efficient DMUs targets are determined using STEM.




Figure 1. The interactive algorithm for finding DMU targets in non-radial FDH models.

The required times to construct and find targets of non-radial model 13 (step 0 of STEM) are equal to the number of M' set members. The required times to construct and find targets of non-radial model 14 (steps 1 and 2 of STEM) is dependent on DM's opinions about the suitable target for each considered DMU_o. The required iteration number to apply the second interactive stage is equal to the number of M' set's members, which is defined the first time. As the number of M' set's members is less than n (because at least, one member belongs to M set and M' set is defined as $M' = \{DMU_j | DMU_j \notin M, \forall j\}$ in the first interactive stage), the targets of all DMUs are therefore found in less than n iterations applying the second interactive stage of the interactive algorithm. This means that the interactive algorithm is completed by applying the second interactive stage in less than n iterations.

It is considerable that finding DMU targets of three kinds of non-radial FDH models (models 12–14) is required in the first part of hybrid technique. These targets can be found by solving related non-radial FDH models. RBA is one of the suitable methods to find DMU targets of radial FDH models, without solving of any mathematical models. In the second part of the hybrid technique, extended RBA is proposed to find DMU targets of non-radial FDH models without solving any mathematical models.

3.2. Extended RBA

In this section, extended RBA (the second part of proposed hybrid technique) to find DMU targets in non-radial FDH models of the interactive algorithm is described. As described in Section 3.1, finding DMU targets of three non-radial FDH models of the interactive algorithm (models 12–14) are required. All mentioned models are mixed 0–1 LP and finding DMU targets by solving them is difficult. RBA can find DMU targets of radial FDH models without solving them. Only non-radial model 12 has a lot of similarity to radial FDH models (models 7 and 8). Therefore, finding DMU targets of non-radial model 12 may be possible using RBA with a little modification. However, RBA cannot apply to find DMU targets of non-radial FDH models 13 and 14, because these models have some complicated constraints. In this section, extended RBA is proposed to find DMU targets of non-radial FDH models without solving any mathematical programming models.

Define $x_{io} = x_{ij}$, $i = 1, \dots, m$, and $y_{ro} = y_{rj}$, $r = 1, \dots, s$ for DMU_o , $o = 1, \dots, n$, for $j = 1, \dots, n$. The extended RBA then find DMU targets of non-radial FDH models of the interactive algorithm (models 12–14) through two target finding stages. In the first target finding stage, DMUs of feasible region are found. In this regard, DMUs that have one, two or three conditions are considered as DMUs of feasible region. The number of required conditions for this stage depends on the constraints of the model. The first condition of extended RBA is similar to RBA and can be applied for non-radial FDH models 12–14. The second and third conditions are applied for non-radial model 14. In the second step, the optimum objective function value is calculated and DMU targets are found. Extended RBA are described in the following subsections.

3.2.1. Finding DMU Targets of Non-Radial FDH Models 12 and 13

As the constraints of non-radial FDH models 12 and 13 are exactly the same, finding DMU targets of these models are described simultaneously.

The first target finding stage

First, DMUs of feasible region should be found. Similar to RBA, if the constraints $\frac{x_{ij}}{x_{io}} \leq 1, i = 1, \dots, m$ and $\frac{y_{rj}}{y_{ro}} \geq 1, r = 1, \dots, s$ are satisfied for $DMU_j, j = 1, \dots, n$ (the first condition), $DMU_j, j = 1, \dots, n$ belongs to S . So, the DMUs of feasible region of non-radial FDH models 12 and 13 are determined using Equation (15).

$$S_{\text{step 0 of STEM}} = \{DMU_j | DMU_j \text{ satisfies first condition, } j = 1, \dots, n\}. \quad (15)$$

As mentioned above, the first condition of extended RBA is similar to RBA.

The second target finding stage

The optimum objective function values should be calculated and DMU targets should be found in the second target finding stage. To achieve this, the objective function value of each $DMU_j, j = 1, \dots, n$ in $S_{\text{step 0 of STEM}}$ (in non-radial model 12, $Z_j^0 = \frac{1}{s} \sum_{r=1}^s \frac{y_{rj}}{y_{ro}} - \frac{1}{m} \sum_{i=1}^m \frac{x_{ij}}{x_{io}}$ and in non-radial model 13, $f_1(u) = -x_{10}x_{1j}, \dots, f_m(u) = -x_{m0}x_{mj}, f_{m+1}(u) = y_{1j}y_{10}, \dots, f_{m+s}(u) = y_{sj}y_{s0}$) is calculated. The maximum obtained value for DMUs belonging to $S_{\text{step 0 of STEM}}$, shows optimum objective function value (the optimum objective function value of non-radial FDH models 12 and 13 are Z_j^0 and $f_l(u), l = 1, \dots, m + s$, respectively). Therefore, the optimum objective function values of non-radial FDH models 12 and 13 are calculated by Equations (16) and (17), respectively.

$$Z^{*0} = \max_{DMU_j \in S_{\text{step 0 of STEM}}} \{Z_j^0\}, Z_j^0 = \frac{1}{s} \sum_{r=1}^s \frac{y_{rj}}{y_{ro}} - \frac{1}{m} \sum_{i=1}^m \frac{x_{ij}}{x_{io}} \quad (16)$$

$$f_l^* = \max_{DMU_j \in S_{\text{step 0 of STEM}}} \{f_{lj}\}, f_{lj}(u) = -x_{l0}x_{lj}, l = 1, \dots, m, f_{lj}(u) = y_{(l-m)j}y_{(l-m)0}, l = m + 1, \dots, m + s \quad (17)$$

Then, the DMUs of $S_{\text{step 0 of STEM}}$ which have optimum objective function value show the targets of DMU_0 in each model. Therefore, DMU_0 targets in non-radial FDH models 12 and 13 are determined by Equations (18) and (19), respectively.

$$\begin{aligned} &DMU_0 \text{ Targets in model 12} = \\ &\{DMU_j | DMU_j \in S_{\text{step 0 of STEM}}, j = 1, \dots, n, \text{objective function value} = Z^{*0}\} \end{aligned} \quad (18)$$

$$\begin{aligned} &DMU_0 \text{ Targets in model 13} \\ &= \{DMU_j | DMU_j \in S_{\text{step 0 of STEM}}, j = 1, \dots, n, \text{objective function value} = f_l^*, l = 1, \dots, m + s\} \end{aligned} \quad (19)$$

As described in Section 3.1, both steps 1 and 2 of STEM's model are shown by non-radial model 14. Finding DMU targets of related non-radial FDH models by extended RBA are described in Sections 3.2.2 and 3.2.3.

3.2.2. Finding DMU Targets of Step 1 of STEM's Model

The process of applying extended RBA to find DMU targets of step 1 of STEM's model is described in two target finding stages as follows:

The first target finding stage

In the first target finding stage, finding DMUs of feasible region (in step 1 of STEM model) is desired. In DMUs of feasible region ($DMU_j, j = 1, \dots, n$), two conditions should be satisfied. Firstly, in these DMUs, similar to non-radial FDH models 12 and 13, constraints $\frac{x_{ij}}{x_{io}} \leq 1, i = 1, \dots, m$, and $\frac{y_{rj}}{y_{ro}} \geq 1, r = 1, \dots, s$ should be satisfied for $DMU_j, j = 1, \dots, n$. The method of checking this condition for DMUs of feasible region ($DMU_j, j = 1, \dots, n$) has been described in Section 3.2.1. Secondly, in DMUs of feasible region, the constraint $\max_l \{(f_l^* - f_l(u)) \cdot \beta_l\} \geq 0, l = 1, \dots, m + s$ should be satisfied for $DMU_j, j = 1, \dots, n$. In constraints of the second condition, the values of $f_l^*, l = 1, \dots, m + s$ and $\beta_l, l = 1, \dots, m + s$

are obtained from solving non-radial model 13 and calculated using Equation (4), respectively. For checking the second condition, $(f_i^* + x_{io}x_{ij}) \cdot \beta_i, i = 1, \dots, m$ and $(f_r^* - y_{rj}y_{ro}) \cdot \beta_r, r = 1, \dots, s$ are calculated for $DMU_j, j = 1, \dots, n$ that were compatible in the first condition. Then constraints $\max_i \{(f_i^* - f_i(u)) \cdot \beta_i\} \geq 0, i = 1, \dots, m + s$ are checked for mentioned DMUs. In DMUs belonging to the feasible region of non-radial model 14 (step 1 of STEM), in addition to the first condition, constraints $\max_{i=1, \dots, m, r=1, \dots, s} \{(f_i^* + x_{io}x_{ij}) \cdot \beta_i, (f_r^* - y_{rj}y_{ro}) \cdot \beta_r\} \geq 0$ should be satisfied for $j = 1, \dots, n$ (the second condition), too. In DMUs of feasible region, the first and second conditions should be satisfied in $DMU_j, j = 1, \dots, n$, simultaneously. Therefore, the DMUs of $S_{\text{step 1 of STEM}}^p$ are determined using Equation (20).

$$S_{\text{step 1 of STEM}}^p = \{DMU_j | DMU_j \text{ satisfies first and second conditions, } j = 1, \dots, n\}. \quad (20)$$

The second target finding stage

The optimum objective function values of step 1 of the STEM model should be calculated and DMU targets of the model should be found in the second target finding stage. The optimum objective function value of step 1 of STEM model is calculated by Equation (21).

$$h^* = \min_{DMU_j \in S_{\text{step 1 of STEM}}^p} \left\{ \max_{i=1, \dots, m, r=1, \dots, s} \{(f_i^* + x_{io}x_{ij}) \cdot \beta_i, (f_r^* - y_{rj}y_{ro}) \cdot \beta_r\} \right\} \quad (21)$$

After that, the DMUs of $S_{\text{step 1 of STEM}}^p$ which have optimum objective function value shows the targets of DMU_o in the model. Therefore, the DMU_o targets in step 1 of STEM are determined using Equation (22).

$$\begin{aligned} DMU_o \text{ Targets in step 1 of STEM} = \\ \{DMU_j | DMU_j \in S_{\text{step 1 of STEM}}^p, j = 1, \dots, n, \text{ objective function value} = h^*\}. \end{aligned} \quad (22)$$

3.2.3. Finding DMU Targets of Step 2 of STEM's Model

Finding the optimum solutions of non-radial model 14 (step 2 of STEM), without solving any mathematical models, is described in two target finding stages below.

The first target finding stage

First, DMUs of feasible region in step 2 of the STEM model should be found. In DMUs of feasible region, three conditions should be regarded. The first and second conditions should be tested for $DMU_j, j = 1, \dots, n$ as described in Section 3.2.2, but in the second condition, $\beta_l, l = 1, \dots, m + s$ should be determined again. The third condition relates to $f_q(u) \geq f_q(u^p) - \Delta f_q$ and $f_l(u) \geq f_l(u^p), l = 1, \dots, m + s, l \neq q$ constraints that $f_q(u), \Delta f_q$, and $f_l(u), l = 1, \dots, m + s, l \neq q$ are obtained from suggested target of DM. So in DMUs of feasible region, constraints $f_q(u) \geq -x_{io}x_{ij} - \Delta f_q$, and $f_l(u) \geq -x_{io}x_{ij}, l = 1, \dots, m, l \neq q$ and $f_q(u) \geq y_{lj}y_{lo} - \Delta f_q$, and $f_l(u) \geq y_{lj}y_{lo}, l = m + 1, \dots, m + s, l \neq q$ for $j = 1, \dots, n$ should be satisfied, too. So, the selected DMU_j of $S_{\text{step 2 of STEM}}^p$ set are defined as Equation (23).

$$S_{\text{step 2 of STEM}}^p = \{DMU_j | DMU_j \text{ satisfies first to third conditions, } j = 1, \dots, n\}. \quad (23)$$

The second target finding stage

The optimum objective function value of step 2 of the STEM model should be calculated and DMU targets of mentioned model should be found in the second target finding stage. The optimum objective function value of step 1 of the STEM model is calculated by Equation (24).

$$h^* = \min_{DMU_j \in S_{\text{step 2 of STEM}}^p} \left\{ \max_{i=1, \dots, m, r=1, \dots, s} \{(f_i^* + x_{io}x_{ij}) \cdot \beta_i, (f_r^* - y_{rj}y_{ro}) \cdot \beta_r\} \right\} \quad (24)$$

After that, the DMUs of feasible region which have optimum objective function values show the targets of DMU_o in the model (Equation (25)).

$$DMU_o \text{ Targets in step 2 of STEM} = \{DMU_j | DMU_j \in S_{\text{step 2 of STEM}}^p, j = 1, \dots, n, \text{objective function value} = h^*\}. \quad (25)$$

It is remarkable that DMU_o Target in step 2 of STEM exists, if the best compromise solution is found in step 2 of the STEM model (Section 3.1). The brief of applying extended RBA to find non-radial FDH models targets of the interactive algorithm is shown in Figure 2.




Figure 2. Extended RBA for finding non-radial FDH models targets of the interactive algorithm.

4. Two Real Case Studies

The proposed technique was applied in two real case studies. The description is provided in Sections 4.1 and 4.2.

4.1. The First Case Study: University Departments

The data set of 17 university departments of Islamic Azad University of Mobarakeh Iran (mobarakeh.iau.ir), denoted by $DMU_{01}, \dots, DMU_{17}$, were extracted from reported research [39]. These data were also used by another study [15]. Each DMU_j , $j = 1, \dots, 17$ has two inputs and two outputs.

The names of inputs are “the number of bachelor students” and “the number of (full time and part time) faculty members”. The names of outputs are “the number of graduates” and “the number of research papers”. The data of the first case study are shown in Table 2.

Table 2. Data of the first case study.

DMU _j	I ₁	I ₂	O ₁	O ₂
DMU ₀₁	26	7	12	3
DMU ₀₂	29	6	10	7
DMU ₀₃	40	8	20	6
DMU ₀₄	42	7	12	6
DMU ₀₅	45	9	18	6
DMU ₀₆	92	12	40	2
DMU ₀₇	83	11	58	3
DMU ₀₈	87	14	52	7
DMU ₀₉	149	16	61	4
DMU ₁₀	177	17	54	12
DMU ₁₁	191	19	61	11
DMU ₁₂	185	14	73	4
DMU ₁₃	186	20	85	10
DMU ₁₄	74	12	36	5
DMU ₁₅	164	22	69	8
DMU ₁₆	225	20	80	5
DMU ₁₇	108	10	27	3

As mentioned previously, in DEA models, choosing appropriate inputs, outputs, and DMUs is an important subject. In this case study, the relationship $n \geq 3(m + s)$ existed ($17 \geq 3(2 + 2)$).

4.1.1. Applying the Interactive Algorithm for the First Case Study

To find DMU targets, the interactive algorithm (Figure 1), including two interactive stages, was applied for universities departments.

Applying the first interactive stage

In this stage, efficient DMUs and their targets were determined. For $o = 1, \dots, 17$, parameters $x_{io} = x_{ij}$, $i = 1, 2$, $j = 1, \dots, 17$, and $y_{ro} = y_{rj}$, $r = 1, 2$, $j = 1, \dots, 17$ were defined. Non-radial model 12 was then composed for DMU_j, $j = 1, \dots, 17$. The results of solving of model 12 for DMU_j, $j = \dots, 17$ are shown in Table 3.

Considering Table 3, $M = \{DMU_{01}, DMU_{02}, DMU_{03}, DMU_{04}, DMU_{07}, DMU_{08}, DMU_{09}, DMU_{10}, DMU_{11}, DMU_{12}, DMU_{13}, DMU_{14}, DMU_{15}, DMU_{17}\}$ and $M' = \{DMU_{05}, DMU_{06}, DMU_{16}\}$ were defined. Then DMU_j Target = {DMU_j}, DMU_j ∈ M was also defined. So, obtained results from 17 times (number of DMUs) composing and solving non-radial model 12 (the first interactive stage) showed that 14 DMUs lay on the efficient frontier, and 3 DMUs did not. By applying the second interactive stage, the DMU targets of these three DMUs were distinguished.

Applying the second interactive stage

In the second interactive stage, the targets of DMUs belonging to set M' were found by STEM. First, the pay-off tables for non-proceed DMUs were constructed. As $M' = \{DMU_{05}, DMU_{06}, DMU_{16}\} \neq \emptyset$ (M' has 3 members), the interactive algorithm (Figure 1) should be continued to find targets of M' set considering DMU₀₅, DMU₀₆, and DMU₁₆, one by one. But here, the process of applying the interactive algorithm for DMU₀₅, DMU₀₆ and DMU₁₆ are described simultaneously. So, each of these three DMUs (DMU₀₅, DMU₀₆ and DMU₁₆) was considered as a non-proceed DMU that their targets should be defined.

Table 3. The results of solving model 12 (related to the first interactive stage) in the first case study.

DMU _j	Z_j^0	θ_1^*	θ_2^*	φ_1^*	φ_2^*	DMU _j Target
DMU ₀₁	0	1	1	1	1	DMU ₀₁
DMU ₀₂	0	1	1	1	1	DMU ₀₂
DMU ₀₃	0	1	1	1	1	DMU ₀₃
DMU ₀₄	0	1	1	1	1	DMU ₀₄
DMU ₀₅	0.166	0.89	0.89	1.11	1	DMU ₀₃
DMU ₀₆	0.565	0.9	0.92	1.45	1.5	DMU ₀₇
DMU ₀₇	0	1	1	1	1	DMU ₀₇
DMU ₀₈	0	1	1	1	1	DMU ₀₈
DMU ₀₉	0	1	1	1	1	DMU ₀₉
DMU ₁₀	0	1	1	1	1	DMU ₁₀
DMU ₁₁	0	1	1	1	1	DMU ₁₁
DMU ₁₂	0	1	1	1	1	DMU ₁₂
DMU ₁₃	0	1	1	1	1	DMU ₁₃
DMU ₁₄	0	1	1	1	1	DMU ₁₄
DMU ₁₅	0	1	1	1	1	DMU ₁₅
DMU ₁₆	0.617	0.83	1	1.06	2	DMU ₁₃
DMU ₁₇	0	1	1	1	1	DMU ₁₇

Firstly, $x_{io} = x_{ij}$, $i = 1, 2, j = 1, \dots, 17$, and $y_{ro} = y_{rj}$, $r = 1, 2, j = 1, \dots, 17$ were defined for DMU_o, $o = 5, 6$, and 16 . Then the pay-off tables (Table 4) were constructed by composing and solving of model 13 for $f_l(u)$, $l = 1, \dots, 4$. In model 13, $n = 17$ and $m = s = 2$ are considered for each DMU_o, $o = 5, 6$, and 16 .

It is mentionable that the required times to construct and solve model 13 in the interactive algorithm were equal to 3 (the number of M' set members). In this interactive stage, finding a suitable DMU in feasible region as the target of DMU_o, $o = 5, 6$, and 16 was desired which the value of f_l , $l = 1, \dots, 4$ was as good as the suggested DMU_o target, $o = 5, 6$, and 16 by DM. Supposed that the targets of DMU₀₅, DMU₀₆ and DMU₁₆ according to DM's opinions were DMU₀₃, DMU₀₇ and DMU₁₃, respectively. So, suggested targets of DM were considered as DM's opinions for f_l , $l = 1, \dots, 4$. After that, the first iteration of STEM ($p = 1$) was started. In step 1 of STEM (calculation phase), first π_l , $l = 1, \dots, 4$ was calculated by Equation (4) considering Table 4, then $\beta_l = \frac{\pi_l}{\sum_{d=1}^4 \pi_d}$, $l = 1, \dots, 4$ was calculated as shown in Table 5.

Then model 14 for DMU₀₅, DMU₀₆ and DMU₁₆ (Step 1 of STEM for $p = 1$) were composed. The results of solving these models and DM's opinions about them are shown in Table 6. As seen in the last column of Table 6, the targets of DMU₀₅, DMU₀₆ and DMU₁₆ via solving of model 14 (Step 1 of STEM for $p = 1$) were DMU₀₃, DMU₀₇ and DMU₁₃, respectively.

Table 4. The results of solving model 13 for three DMUs (related to the second interactive stage) in the first case study.

DMU _o	f_l ($l = 1, \dots, 4$)	f^*	f_1	f_2	f_3	f_4	θ_1^*	θ_2^*	φ_1^*	φ_2^*
DMU ₀₅	$f_1 = -45\theta_1$	-40	-40	-8	18	6	0.889	0.889	1.000	1.000
	$f_2 = -9\theta_2$	-8	-40	-8	18	6	0.889	0.889	1.000	1.000
	$f_3 = 18\varphi_1$	20	-40	-8	20	6	0.889	0.889	1.111	1.000
	$f_4 = 6\varphi_2$	6	-40	-8	18	6	0.889	0.889	1.000	1.000
DMU ₀₆	$f_1 = -92\theta_1$	-83	-83	11	40	2	0.902	0.917	1.000	1.000
	$f_2 = -12\theta_2$	-11	-83	11	40	2	0.902	0.917	1.000	1.000
	$f_3 = 40\varphi_1$	58	-83	11	58	2	0.902	0.917	1.450	1.000

	$f_4 = 2\varphi_2$	3	-83	11	40	3	0.902	0.917	1.000	1.500
DMU ₁₆	$f_1 = -225\theta_1$	-186	-186	20	80	5	0.827	1.000	1.000	1.000
	$f_2 = -20\theta_2$	-20	-186	20	80	5	0.827	1.000	1.000	1.000
	$f_3 = 80\varphi_1$	85	-186	20	85	5	0.827	1.000	1.063	1.000
	$f_4 = 5\varphi_2$	10	-186	20	80	10	0.827	1.000	1.000	2.000

Table 5. Calculating parameters for three DMUs to compose model 14 of step 1 of STEM ($p = 1$) (related to the second interactive stage) in the first case study.

DMU _o		l = 1	l = 2	l = 3	l = 4
DMU ₀₅	$f_l^{\min}$	-40	-8.000	18.000	6.000
	f_l^*	-40	-8.000	20.000	6.000
	π_l	0	0.000	0.006	0.000
	β_l	0	0.000	1.000	0.000
DMU ₀₆	$f_l^{\min}$	-83	11.000	40.000	2.000
	f_l^*	-83	-11.000	58.000	3.000
	π_l	0	2.000	0.008	0.167
	β_l	0	0.920	0.004	0.077
DMU ₁₆	$f_l^{\min}$	-186	20.000	80.000	5.000
	f_l^*	-186	-20.000	85.000	10.000
	π_l	0	1.000	0.001	0.100
	β_l	0	0.908	0.001	0.091

As described previously, these targets were acceptable according to DM's opinion, too. As the obtained solutions for DMU₀₅ and DMU₀₆ were satisfied via DM's opinions, the best compromise solutions were obtained for these DMUs.

Then, DMU_o (DMU₀₅ and DMU₀₆) from M' set removed ($M' = \{DMU_{16}\} \neq \emptyset$ (M' has 1 member)). As the value of the fourth target of DMU₁₆ was not satisfied via DM's opinions, the obtained solution was not the best compromise solution. So, composing model 14 related to Step 2 of STEM ($p = 1$) was required for DMU₁₆. The required parameters were calculated as shown in Table 7.

Table 6. The results of solving model 14 (Step 1 of STEM for $p = 1$) and DM's opinions about them for three DMUs (related to the second interactive stage) in the first case study.

DMU _o		Target ₁	Target ₂	Target ₃	Target ₄	Target
DMU ₀₅	Model 14	40	8	20	6	DMU ₀₃
	DM's opinions	40	8	20	6	DMU ₀₃
	Δ_{f_i}	0	0	0	0	-
	Unsatisfied	0	0	0	0	-
DMU ₀₆	Model 14	83	1	58	3	DMU ₀₇
	DM's opinions	83	11	58	3	DMU ₀₇
	Δ_{f_i}	0	-10	0	0	-
	Unsatisfied	0	0	0	0	-
DMU ₁₆	Model 14	86	0	85	0	DMU ₁₃
	DM's opinions	186	20	85	10	DMU ₁₃
	Δ_{f_i}	-100	-20	0	-10	-
	Unsatisfied	0	0	0	1	-

Table 7. Calculated parameters through solving of step 2 of STEM ($p = 1$) models for DMU_{16} (the first case study).

DMU_o	1	i or r	θ_i^* or φ_r^*	f_i	Unsatisfied f_i	Target (DM)	Δ_{f_i}	β_i
DMU_{16}	1	i = 1	0.827	-186	0	186	0.000	0.000
	2	i = 2	1.000	20	0	20	0.000	0.000
	3	r = 1	1.063	85	0	85	0.000	0.000
	4	r = 2	2.000	0	1	10	1.000	0.000

Non-radial model 14 for DMU_{16} was then composed for the first iteration of step 2 of STEM. The results of solving this model showed that there was only one similar unsatisfied objective function (fourth objective function) in two consecutive steps of STEM. Therefore, the target of DMU_{16} could not be considered as DMU_{13} . Therefore, the interactive algorithm was continued by asking DM to suggest a new target for each DMU_o . DM did not suggest new DMUs target for DMU_{16} so there were no targets for DMU_{16} in the FDH model. Then, DMU_o (DMU_{16}) from M' set was removed ($M' = \emptyset$). As there was no non-proceed DMUs and the process of targets determination for all DMUs had been carried out ($M' = \emptyset$), the interactive algorithm was finished. The results of applying the interactive algorithm for all DMUs are shown in Table 8.

As shown in the last column of Table 8, applying the interactive algorithm for university departments are considered as three cases. DMU_{01} , DMU_{02} , DMU_{03} , DMU_{04} , DMU_{07} , DMU_{08} , DMU_{09} , DMU_{10} , DMU_{11} , DMU_{12} , DMU_{13} , DMU_{14} , DMU_{15} , and DMU_{17} were members of M set and the targets of these DMUs were themselves (case 1). As best compromise solutions were found through step 1 of STEM for DMU_{05} , DMU_{06} , these DMUs belonged to case 2. As there was only one similar unsatisfied objective function in two consecutive steps of STEM for DMU_{16} , this DMU belonged to case 3.

Table 8. The brief of applying the interactive algorithm for finding DMU targets (the first case study).

DMU_o	Member of M Set	Obtained Target by Solving of Step 1 of STEM Model	Obtained Target via DM's Opinions	Obtaining DM's Target	Related Interactive Stages of the Interactive Algorithm	Case
DMU_{01}	√	-	-	-	As DMU was a member of M set, target was itself	1
DMU_{02}						
DMU_{03}						
DMU_{04}						
DMU_{07}						
DMU_{08}						
DMU_{09}						
DMU_{10}						
DMU_{11}						
DMU_{12}						
DMU_{05}	×	DMU_{03}	DMU_{03}	√	The best compromise solution via DM's opinions was found in first iteration of step 1 of STEM	2
DMU_{06}		DMU_{07}	DMU_{07}			

DMU ₁₆	×	DMU ₁₃	DMU ₁₃	×	The best compromise solution was not found because there was only one similar unsatisfied objective function in the step 1 and first iteration of step 2 of STEM	3
-------------------	---	-------------------	-------------------	---	--	---

It should be mentioned that in the first case study, 14 DMUs from 17 DMUs (about 82%) lay on the efficient frontier. Moreover, non-radial FDH models 13 and 14 were composed and solved, three and four times through the second interactive stage, respectively. Here, the process of target determination for all DMUs was carried out, and the interactive algorithm finished. In fact, the required repetition times for composing and solving non-radial model 14 depend on DM's opinions about suitable targets for each considered DMU_o.

4.1.2. Applying Extended RBA for the First Case Study

As described in Section 4.1.1, for applying the interactive algorithm for the first case study, solving of non-radial FDH models 12, 13, model 14 of step 1 of STEM ($p = 1$), and model 14 of step 2 of STEM ($p = 1$), 17, 3, 3, and 1 times were needed, respectively. So, solving 24 mixed 0–1 LP models were required to find all DMU targets. The DMU targets of mentioned non-radial FDH models could be found using extended RBA through two target finding stages without solving any mathematical programming models. In this section, applying extended RBA for finding DMU targets of these non-radial FDH models (models 12–14 for DMU₀₅ of university departments) are described. First $x_{i0} = x_{ij}$, $i = 1, 2$, and $y_{r0} = y_{rj}$, $s = 1, 2$ ($j = 1, \dots, 17$) were defined for DMU_o = DMU₀₅ (Figure 2). Target finding of non-radial FDH models 12 and 13 for DMU₀₅ in the first case study

In this regard, DMUs of feasible region should be found in applying the first target finding stage. Firstly, DMUs of feasible region ($S_{\text{step 0 of STEM}}$) should be found using Equation (15) (Figure 2). In DMUs of feasible region, constraints $\frac{x_{ij}}{x_{i0}} \leq 1$, $i = 1, 2$ and $\frac{y_{rj}}{y_{r0}} \geq 1$, $r = 1, 2$ ($j = 1, \dots, 17$) should be satisfied. The results of checking constraints of the first condition are shown in Table 9. As it is observed in the last column of Table 9, only DMU₀₃ and DMU₀₅ satisfied all four constraints of the first condition. So, $S_{\text{step 0 of STEM}} = \{\text{DMU}_{03}, \text{DMU}_{05}\}$.

Table 9. Applying the first target finding stage for DMU₀₅ (the first case study).

DMU _j	I ₁	I ₂	O ₁	O ₂	$\frac{x_{1j}}{x_{10}}$	$\frac{x_{1j}}{x_{10}} \leq 1$	$\frac{x_{2j}}{x_{20}}$	$\frac{x_{2j}}{x_{20}} \leq 1$	$\frac{y_{1j}}{y_{10}}$	$\frac{y_{1j}}{y_{10}} \geq 1$	$\frac{y_{2j}}{y_{20}}$	$\frac{y_{2j}}{y_{20}} \geq 1$	$S_{\text{step 0 of STEM}}$ (Equation (15))
DMU ₀₁	26	7	12	3	0.578	0.578	0.778	0.778	0.667	×	0.500	×	×
DMU ₀₂	29	6	10	7	0.644	0.644	0.667	0.667	0.556	×	1.167	1.167	×
DMU ₀₃	40	8	20	6	0.889	0.889	0.889	0.889	1.111	1.111	1.000	1.000	√
DMU ₀₄	42	7	12	6	0.933	0.933	0.778	0.778	0.667	×	1.000	1.000	×
DMU ₀₅	45	9	18	6	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	√
DMU ₀₆	92	12	40	2	2.044	×	1.333	×	2.222	2.222	0.333	×	×
DMU ₀₇	83	11	58	3	1.844	×	1.222	×	3.222	3.222	0.500	×	×
DMU ₀₈	87	14	52	7	1.933	×	1.556	×	2.889	2.889	1.167	1.167	×
DMU ₀₉	149	16	61	4	3.311	×	1.778	×	3.389	3.389	0.667	×	×
DMU ₁₀	177	17	54	12	3.933	×	1.889	×	3.000	3.000	2.000	2.000	×
DMU ₁₁	191	19	61	11	4.244	×	2.111	×	3.389	3.389	1.833	1.833	×
DMU ₁₂	185	14	73	4	4.111	×	1.556	×	4.056	4.056	0.667	×	×
DMU ₁₃	186	20	85	10	4.133	×	2.222	×	4.722	4.722	1.667	1.667	×

DMU ₁₄	74	12	36	5	1.644	×	1.333	×	2.000	2.000	0.833	×	×
DMU ₁₅	164	22	69	8	3.644	×	2.444	×	3.833	3.833	1.333	1.333	×
DMU ₁₆	225	20	80	5	5.000	×	2.222	×	4.444	4.444	0.833	×	×
DMU ₁₇	108	10	27	3	2.400	×	1.111	×	1.500	1.500	0.500	×	×
DMU ₀ =DMU ₀₅	45	9	18	6	-	-	-	-	-	-	-	-	-

The optimum objective function values and DMU targets should then be found in applying the second target finding stage. At first, the optimum objective function values of models should be calculated by Equations (16) and (17) (Figure 2). The objective function values of DMU₀₃ and DMU₀₅ (in model 12, $Z_j^o = \frac{1}{2} \sum_{r=1}^2 \frac{y_{rj}}{y_{ro}} - \frac{1}{2} \sum_{i=1}^2 \frac{x_{ij}}{x_{io}}$, $j = 1, \dots, 17$, and in model 13, $f_1(u) = -x_{10}x_{1j}$, $f_2(u) = -x_{20}x_{2j}$, $f_3(u) = y_{10}y_{1j}$, $f_4(u) = y_{20}y_{2j}$) are shown in Table 10.

The last row of Table 10 shows the maximum values of Z_j^o , $DMU_j \in S_{\text{step 0 of STEM}}$ and f_l , $l = 1, \dots, 4$, respectively. These values show the optimum objective function value of each model (Z^{*o} , f_l^* , $l = 1, \dots, 4$). The DMU₀ targets were then found by comparing the optimum objective function values of each model with the objective function value of each DMU by Equations (18) and (19) (Figure 2). It is considerable that non-radial model 13 with f_4 objective function had alternative solutions.

Table 10. Applying the second target finding stage for DMU₀₅ (the first case study).

DMU _j	Z_j^o (Equation (16))	Model 12 Targets (Equation (18))	f_1 (Equation (17))	f_1 Targets (Equation (19))	f_2 (Equation (17))	f_2 Targets (Equation (19))	f_3 (Equation (17))	f_3 Targets (Equation (19))	f_4 (Equation (17))	f_4 Targets (Equation (19))
DMU ₀₃	0	DMU ₀₃	-40	DMU ₀₃	-8	DMU ₀₃	20	DMU ₀₃	6	DMU ₀₃
DMU ₀₅	0	×	-45	×	-9	×	18	×	6	DMU ₀₅
Max	0	-	-40	-	-8	-	20	-	6	-

Target finding of non-radial model 14 for DMU₀₅ in the first case study

First, step 1 of STEM ($p = 1$) should be considered. DMUs of feasible region (in step 1 of STEM model) should be found through applying the first target finding stage. First, DMUs belonging to the feasible region should be found. In DMUs of feasible region, two conditions should be satisfied. Firstly, constraints $\frac{x_{ij}}{x_{io}} \leq 1$, $i = 1, 2$, and $\frac{y_{rj}}{y_{ro}} \geq 1$, $r = 1, 2$, ($j = 1, \dots, 17$) should be satisfied (the first condition). As it is seen in Table 9, only DMU₀₃ and DMU₀₅ satisfied these constraints. Moreover, the constraints $\max_l \{(f_l^* - f_l(u)) \cdot \beta_l\} \geq 0$, $l = 1, \dots, 4$ should be satisfied for each DMU of feasible region (the second condition). Checking the second condition for DMU₀₃ and DMU₀₅ are shown in Table 11.

Table 11. Applying target finding stages 1–2 to solve step 1 of STEM ($p = 1$) for DMU₀₅ (the first case study).

DMU _j	$(f_1^* - f_1(u)) \cdot \beta_1$	$(f_2^* - f_2(u)) \cdot \beta_2$	$(f_3^* - f_3(u)) \cdot \beta_3$	$(f_4^* - f_4(u)) \cdot \beta_4$	$S_{\text{step 1 of STEM}}^1$ (Equation (20))	h_j (Equation (21))	Targets (Equation (22))
DMU ₀₃	0	0	0	0	√	0	DMU ₀₃
DMU ₀₅	0	0	2	0	√	2	×
$h^* = \min \max (f_l^* - f_l(u)) \cdot \beta_l$ ($l = 1, \dots, 4$)	-	-	-	-	-	0	-

As seen in Table 11, the constraints of the second condition were also satisfied for DMU₀₃ and DMU₀₅. So, $S_{\text{step 1 of STEM}}^1 = \{\text{DMU}_{03}, \text{DMU}_{05}\}$.

The optimum objective function value of step 1 of STEM model was then calculated, and DMU targets were found through applying the second target finding stage. With respect to Equation (21), the minimum values of the left-hand side of the second condition for DMU₀₃ and DMU₀₅ showed the optimum objective function value of non-radial model 14 (last row of Table 11). As is seen in Equation (22), each DMU with an objective function value equal to the optimum objective function value, showed the targets of DMU₀. So, the target of DMU₀₅ was DMU₀₃, as seen in the last column of Table 11 ($\text{DMU}_{05} \text{ Targets in step 1 of STEM} = \{\text{DMU}_{03}\}$).

It is mentionable that the obtained results from target finding of all non-radial FDH models 12–14 for the first case study for DMU₀₅ using extended RBA are the same as the results obtained from solving the mentioned mathematical models. Here, target finding of three non-radial FDH models of 24 mathematical programming models using extended RBA were described. A similar process should be undertaken to find targets of other 21 non-radial FDH models.

4.2. The Second Case Study: Operations Strategies of Fars Province Pharmaceutical Distributing Companies

By considering expert opinions, 13 active pharmaceutical distributing companies (Fars province of Iran) in 2019 were investigated. These companies were Daroo Pakhsh, Pakhshe Razi, Adora Teb, Mahya Daroo, Alborz, Hejrat, Daroo Gostare Razi, Behestan Pakhsh, Ghasem Iran, Yasin, Ferdos, Elit Daroo, and Soha Helal. The operations strategies of these companies denoted as DMU₀₁, ..., DMU₁₃.

There are five generic performance objectives including cost, delivery speed, quality, dependability, and flexibility. Joining the market requirements through a beneficial method for operations is the purpose of these objectives. These objectives are applicable for different operations. Satisfaction of customers can be obtained through reaching these mentioned objectives [40]. According to expert opinions, these five generic performance objectives were very important in the operation strategies of Fars province pharmaceutical distributing companies. In this regard, cost was defined as the lower cost for producing products and services, and decreasing the cost of distributing medicinal drugs and increasing customers are desirable for pharmaceutical distributing companies. Delivery speed was defined as an elapsed time between the beginning of an operations process and its end, and increasing trucks, vans, and motorcycles numbers could improve delivery speed. Quality was defined as fit-for-purpose, and comparing distributed medicinal drugs with orders and checking the temperature of transportation refrigerators could improve quality. Dependability was defined as keeping delivery promises and distributing medicinal drugs in the promised time and increasing transportation vehicles could improve dependability. Flexibility was defined as treating the operation as a 'black box' and considering the types of flexibility that would contribute to its competitiveness (product or service flexibility, mix flexibility, volume flexibility, and delivery flexibility) and having up to date information about existing medical drugs and having a powerful team to prepare them could improve flexibility.

As mentioned previously, in DEA models, choosing appropriate inputs and outputs is an important subject. As classical DEA models are based on decreasing inputs and increasing outputs, in operations strategies of Fars province pharmaceutical distributing companies, cost and delivery speed were considered as inputs that denoted by I_1 and I_2 . Furthermore, quality, dependability, and flexibility were considered as outputs that were denoted by O_1 , O_2 and O_3 . These inputs and outputs were relevant to the research area and expert(s) confirmed them. All DMUs in the research area were considered as DMUs. So, each DMU_{*j*}, $j = 1, \dots, 13$ had two inputs (I_1 and I_2), and three outputs (O_1 , O_2 and O_3). As mentioned previously, establishing a relationship $n \geq 3(m + s)$ between inputs, out-

puts, and DMU numbers is recommended. Here, all DMUs in the research area were considered and increasing DMUs to reach exact relationship $n \geq 3(m + s)$ was impossible in a practical view (this relationship almost existed ($13 \cong 3(2 + 3)$)). As we will see later, only two DMUs were efficient in the second case study. So, in this case study, decreasing the number of DMUs did not also create difficulty in the theoretical view.

As only experts accessed real data related to inputs and outputs of each DMU, expert judgement was used. With respect to each input and output, the performance of each DMU was scaled by 1 to 9 considering expert opinions (Table 12). The data of inputs and outputs of operations strategies of Fars province pharmaceutical distributing companies (the second case study) are shown in Table 13.

Table 12. Performance scales for inputs and outputs.

Performance Scale	Very Low	Low	Moderate	High	Very High
Outputs	1	3	5	7	9
Inputs	9	7	5	3	1

Table 13. Data of the second case study.

DMU _j	I ₁	I ₂	O ₁	O ₂	O ₃
DMU ₀₁	1	1	8	7	8
DMU ₀₂	1	2	8	6	6
DMU ₀₃	3	4	8	7	4
DMU ₀₄	2	5	5	6	5
DMU ₀₅	1	1	5	5	6
DMU ₀₆	2	3	6	8	7
DMU ₀₇	3	4	4	3	6
DMU ₀₈	3	3	5	5	3
DMU ₀₉	2	2	3	6	4
DMU ₁₀	6	4	7	7	2
DMU ₁₁	5	4	4	7	4
DMU ₁₂	5	5	4	7	6
DMU ₁₃	5	3	6	4	7

As the process of applying the hybrid technique for the second case study had a lot of similarity with the first case study, its main results are described in Sections 4.2.1 and 4.2.2. The details of applying the method for the second case study are shown in Appendix A.

4.2.1. The Main Results of Applying the Interactive Algorithm for the Second Case Study

To find DMU targets for the second case study, the interactive algorithm, including two interactive stages (Figure 1), was applied. In applying the first interactive stage (Appendix A, Appendix A.1), non-radial model 12 was composed and solved 13 times. The results show that only 2 DMUs (about 15%) lay on the efficient frontier, and 11 DMUs did not (Appendix A, Table A1). By applying the second interactive stage (Appendix A, Tables A2–A6), DMU₀₇ target was obtained, but DMU₀₄ and DMU₁₀ target were not obtained. The brief results of applying the interactive algorithm (Figure 1) for all DMUs are shown in Table 14.

As seen in the last column of Table 14, applying the interactive algorithm for the second case study can be categorized into four cases. As shown in Table 14, DMU₀₁ and DMU₀₆ are members of M set and the target of these DMUs are themselves (case 1). The obtained targets by solving step 1 of the STEM model and via DM's opinions of DMU₀₂,

DMU₀₃, DMU₀₅, DMU₀₇, DMU₀₉, and DMU₁₁ are DMU₀₁. By applying step 1 or 2 of STEM, the best compromise solutions for these DMUs are found (DMU₀₁) via DM's opinions (case 2). By solving step 1 of the STEM model, DMU₀₁ is obtained as the target of the rest DMUs. But the target of these DMUs via DM's opinions are DMU₀₆ (case 3 and 4). For DMU₀₄, DMU₀₈, and DMU₁₃, a compromise solution is not found because there is only one similar unsatisfied objective function in step 1 and the first iteration of step 2 of STEM (case 3). The best compromise solution is not obtained for DMU₁₀ because β_l , $l = 1, \dots, 5$ is not calculable for the first iteration of step 2 of STEM (case 4). The required repetition times of composing and solving non-radial model 14, depends on DM's opinions about suitable target for each considered DMU. In Appendix A, finding the target of a DMU in each case are described.

Table 14. The brief of applying the interactive algorithm for finding DMU targets (the second case study).

DMU ₀	Member of M set	Obtained Target by Solving of Step 1 of STEM Model	Obtained Target via DM's Opinions	Obtaining DM's Target	Related Interactive Stages of the Interactive Algorithm	Case
DMU ₀₁	√	-	-	-	As DMU was a member of M set, target was itself	1
DMU ₀₆						
DMU ₀₂						
DMU ₀₃	×	DMU ₀₁	DMU ₀₁	√	The best compromise solution via DM's opinions was found in step 1 or 2 of STEM	2
DMU ₀₅						
DMU ₀₇						
DMU ₀₉						
DMU ₁₁						
DMU ₁₂	×	DMU ₀₁	DMU ₀₆	×	The best compromise solution was not found because there was only one similar unsatisfied objective function in the step 1 and first iteration of step 2 of STEM	3
DMU ₀₄						
DMU ₀₈						
DMU ₁₃	×	DMU ₀₁	DMU ₀₆	×	Compromise solution was not found because calculating value of beta l ($l = 1, \dots, 5$) were not possible for the first cycle of step 2 of STEM	4
DMU ₁₀						

4.2.2. The Main Results of Applying Extended RBA for the Second Case Study

As described in Appendix A.2 (Appendix A), through applying the interactive algorithm for the second case study, non-radial FDH models 12, 13, non-radial FDH model 14 of step 1 of STEM ($p = 1$), and non-radial model 14 of step 2 of STEM ($p = 1$) should be solved 13, 11, 11, and 5 times, respectively. So, finding targets of 40 mixed 0–1 LP models were required. The target of these non-radial FDH models could be obtained using extended RBA through two target finding stages (Figure 2). In Appendix A.2, applying extended RBA for solving three of these non-radial FDH models (models 12–14 for DMU₀₅ of the second case study) are described.

The detail results of applying extended RBA (Figure 2) to solve non-radial FDH models 12–14 for DMU₀₇ of the second case study (related non-radial FDH models in the interactive algorithm) are described in Appendix A.2 (Appendix A). Each target of model was found using two target finding stages for DMU₀₇ of the second case study considering related conditions. $x_{i0} = x_{ij}$, $i = 1, \dots, 3$, and $y_{r0} = y_{rj}$, $s = 1, 2$ ($j = 1, \dots, 13$) were firstly defined for $DMU_0 = DMU_{07}$.

Applying extended RBA for non-radial FDH models 12 and 13 (Appendix A, Appendix A.2), showed that $S_{\text{step 0 of STEM}} = \{\text{DMU}_{01}, \text{DMU}_{02}, \text{DMU}_{05}, \text{DMU}_{06}, \text{DMU}_{07}\}$ and some non-radial FDH models had alternative targets. Applying extended RBA for non-radial FDH model 14 (Appendix A, Appendix A.2), showed that $S_{\text{step 1 of STEM}} = \{\text{DMU}_{01}, \text{DMU}_{02}, \text{DMU}_{05}, \text{DMU}_{06}, \text{DMU}_{07}\}$ and DMU₀₁ was the only target (Table A9).

By solving step 1 of the STEM model (non-radial FDH model 14), DMU₀₁ was obtained as the DMU₀₇ target (Appendix A, Table A10). However, the obtained target via DM's opinions was DMU₀₆. A compromise solution was not found because no feasible integer solution could be found in the first iteration of step 2 of STEM.

It is mentionable that the results obtained from solving of all non-radial FDH models 12–14 for the second case study DMU₀₇ using extended RBA were the same as the results obtained from solving of mentioned models using regular approaches. As described in Appendix A.2 (Appendix A), solving three models of 40 non-radial FDH models were described. A similar process should be undertaken to solve other 37 mathematical models.

As mentioned previously, the suggested hybrid technique contains the interactive algorithm (the first part) and extended RBA (the second part). The interactive algorithm contains two stages. In the interactive stage 1, inputs, outputs, DMUs, and then efficient DMUs and their targets are determined. According to the proposed technique, using real data is preferred rather than gathering them from experts' opinions. In the first case study, inputs, outputs, and DMUs information were real and extracted from a published paper. In the second case study, only experts accessing the real data related to inputs and outputs of each DMU. So, experts' judgements, in situations where experts accessed real data, were used. The number of M' set members are determined by composing and solving non-radial model 12 for all DMUs. The required number of iterations for the interactive algorithm depends on the number of M' set members. Non-radial model 12 should be constructed and solved n (number of DMUs) times. It is considerable that the number of DMUs in the first and second case studies are 17 and 13, respectively. If the ratio of "the number of M' set members" to "all DMUs" is near to 0, the interactive algorithm needs fewer iterations. If this ratio is near to 1, the interactive algorithm needs more iterations. As it is found for applying the first interactive stage in the first case study, if the number of DMUs that lie on the efficient frontier is considerable (14 DMUs from 17 DMUs, about 82%), the interactive requires fewer iterations. However, as it is found for the second case study, if the number of DMUs that lie on the efficient frontier of model 12 is not considerable (2 DMUs from 13 DMUs, about 15%), the interactive algorithm requires more iterations.

In the interactive stage 2, non-efficient DMUs targets are determined using STEM. The required times to construct and find targets of model 13 (step 0 of STEM) are equal to the number of M' set members. The number of M' set members in the first and second case studies were 3 and 11, respectively. The required time for constructing and finding targets of model 14 (steps 1 and 2 of STEM) is dependent on DM's opinions about a suitable target for each considered DMU. These numbers in the first and second case studies were 3 and 11, respectively. Therefore, by applying 3 and 11 times of interactive stage 2, the process of targets determination for non-efficient DMUs of the first and second case studies was carried out, respectively, and the interactive algorithm finished. Moreover, solving 24 and 40 non-radial FDH models were required in the first and second case study, respectively.

5. Conclusions

Finding DMU targets or DMU projections is useful for strategic planning in some organizations. In these organizations, a convexity assumption does not exist, and decisions are made on the basis of pareto solutions. Therefore, these properties should be considered in practical models. Moreover, by eliminating the convexity assumption, FDH models can be applied. As suggesting a technique with mentioned properties has been

neglected in previous research, the main novelty of this paper is in its suggestion of this technique. In this regard, the answers to the two research questions are as follow:

- (a) Is it possible to propose a technique to find all DMU targets in non-radial FDH models based on ASBM using IM? Using IMs such as STEM to find DMU targets in non-radial FDH models on the basis of slack variables can be an important subject and it is applied in the proposed technique.
- (b) Is it possible to find DMU targets of non-radial FDH models in the proposed technique without solving any mathematical models? In this research, in addition to using STEM for finding DMU targets in non-radial FDH models, DMU targets of non-radial FDH models were found by extended RBA without solving any mathematical models.

As has been explained previously, after determining inputs, outputs and DMUs in the first and second case studies, four DMUs from 17 DMUs (76% of DMUs) and two DMUs from 13 DMUs (15% of them) lay on the efficient frontier, respectively. So, in applying the second interactive stage for the first and second case studies, 76% and 20% of DMUs could be selected as targets, respectively. Moreover, the targets of all non-radial FDH models were found using extended RBA without solving any mathematical programming model. In fact, the obtained results can be extended to other cases considering practical and theoretical views.

As mentioned previously, using real data related to inputs, outputs, and DMUs is preferred rather than gathering them from experts' opinions according to the proposed technique. However, if gathering real data was impossible, experts' judgement would be used. In this situation, experts should access real data or have enough information about the problem. The existence of, and access to, such experts can be considered as the first limitation of this study. We assumed that there are no outliers, and this assumption can be considered as the second limitation of this research. Furthermore, all required data existing in deterministic form can be considered as the third limitation of this study.

Detecting outliers and investigating the sensitivity of the modeling approach to outliers in the proposed technique, finding DMU targets using other IMs, using impressive data in supply chain management, and adopting the proposed technique in situations where data are fuzzy can be considered as future studies.

Author Contributions: Conceptualization, M.A. and M.R.M.; methodology, M.A. and M.R.M.; software, M.A. and M.R.M.; validation, M.A. and M.R.M.; formal analysis, M.A. and M.R.M.; investigation, M.A., M.R.M. and P.F.W.; resources, M.A. and M.R.M.; data curation, M.A.K. and M.A.; writing—original draft preparation, M.A., M.R.M. and P.F.W.; writing—review and editing, M.A. and M.R.M.; visualization, M.A., M.R.M. and P.F.W.; supervision, M.A. and M.R.M.; project administration, M.A.; funding acquisition, P.F.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Fundação Carlos Chagas Filho de Amparo à Pesquisa (FAPERJ).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: This project (Determining Appropriate Operations Strategies of Organizations and improving their Operations Performance using Data Envelopment Analysis method: A Case Study in Fars Province Pharmaceutical Distributing Companies) was financially supported by Research Council of Shiraz Branch, Islamic Azad University, Iran.

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A

The application of the main results of the interactive algorithm for the second case study are described in Section 4.2. The detailed results of the application of the interactive algorithm (Figure 1) for the second case study are described here.

Appendix A.1. The Detailed Results of Applying the Interactive Algorithm for the Second Case Study

For determination of DMU targets, the interactive algorithm, including two interactive stages (Figure 1), was applied for the second case study. For the determination of efficient DMUs and their targets, the interactive stage was applied. For $o = 1, \dots, 13$, parameters $x_{io} = x_{ij}$, $i = 1, 2, j = 1, \dots, 13$, and $y_{ro} = y_{rj}$, $r = 1, 2, 3, j = 1, \dots, 13$ were defined. Then non-radial FDH model 12 was made for DMU_j , $j = 1, \dots, 13$. These non-radial FDH models were solved, and the obtained results shown in Table A1.

Table A1. The results of solving model 12 (related to the first interactive stage) in the second case study.

DMU _j	Z_j^o	θ_1^*	θ_2^*	φ_1^*	φ_2^*	φ_3^*	DMU _j Target (Model 12)
DMU ₀₁	0.000	1.000	1.000	1.000	1.000	1.000	DMU ₀₁
DMU ₀₂	0.416	1.000	0.500	1.000	1.170	1.330	DMU ₀₁
DMU ₀₃	1.041	0.330	0.250	1.000	1.000	2.000	DMU ₀₁
DMU ₀₄	1.105	0.500	0.200	1.600	1.170	1.600	DMU ₀₁
DMU ₀₅	0.444	1.000	1.000	1.600	1.400	1.330	DMU ₀₁
DMU ₀₆	0.000	1.000	1.000	1.000	1.000	1.000	DMU ₀₆
DMU ₀₇	1.597	0.330	0.250	2.000	2.330	1.330	DMU ₀₁
DMU ₀₈	1.555	0.330	0.330	1.600	1.400	2.670	DMU ₀₁
DMU ₀₉	1.444	0.500	0.500	2.670	1.170	2.000	DMU ₀₁
DMU ₁₀	1.839	0.170	0.250	1.140	1.000	4.000	DMU ₀₁
DMU ₁₁	1.441	0.200	0.250	2.000	1.000	2.000	DMU ₀₁
DMU ₁₂	1.244	0.200	0.200	2.000	1.000	1.330	DMU ₀₁
DMU ₁₃	1.142	0.200	0.330	1.330	1.750	1.140	DMU ₀₁

$M = \{ DMU_{01}, DMU_{06} \}$ and $M' = \{ DMU_{02}, DMU_{03}, DMU_{04}, DMU_{05}, DMU_{07}, DMU_{08}, DMU_{09}, DMU_{10}, DMU_{11}, DMU_{12}, DMU_{13} \}$ are defined with consideration to Table A1. Then DMU_j Target = $\{ DMU_j \}$, $DMU_j \in M$ was also defined. To apply the second interactive stage, the pay-off table for non-proceed DMUs was constructed. As $M' \neq \emptyset$ (M' has 11 members), the algorithm should be continued (to find targets of M' set). Targets of all members of M' should be determined individually. Suppose that only DMU_{04} , DMU_{07} , and DMU_{10} were non-proceed DMUs that defined their targets as desired. Therefore, suppose $M' = \{ DMU_{04}, DMU_{07}, DMU_{10} \}$ (a sample DMU in cases 2–4 in Table 14). Also suppose that the finding process of DMUs targets were carried out simultaneously ($DMU_o = DMU_{04}, DMU_{07}$, and DMU_{10}).

At first, $x_{io} = x_{ij}$, $i = 1, 2, j = 1, \dots, 13$, and $y_{ro} = y_{rj}$, $r = 1, 2, 3, j = 1, \dots, 13$ were defined for DMU_o , $o = 4, 7$, and 10 . Then the pay-off tables (Table A2) were created by composing and solving non-radial model 13 for $f_l(u)$, $l = 1, \dots, 5$. $N = 13$, $m = 2$ and $s = 3$ were considered for each DMU_o in model 13. The calculation and decision phases of STEM were then applied. Finding the suitable DMUs in feasible region were suitable targets of DMU_o in which the value of f_l , $l = 1, \dots, 5$ were as good as the suggested DMU_o target by DM. Suppose that DM suggests DMU_{06} as the target of DMU_{04}, DMU_{07} , and DMU_{10} . So, suggested targets of DM were considered as DM's opinions for f_l , $l = 1, \dots, 5$ for each DMU_o . After that, the first iteration of STEM ($p = 1$) began. With respect to step 1 of STEM, $\pi_l, l =$

1, ..., 5 was calculated by Equation (3), and then $\beta_l = \frac{\pi_l}{\sum_{d=1}^5 \pi_d}$, $l = 1, \dots, 5$ was obtained as seen in Table A3. Then model 14 (Step 1 of STEM for $p = 1$) was composed for DMU₀₄, DMU₀₇, and DMU₁₀. The results of this model solving and DM's opinions about them are shown in Table A4. As seen in the last column of Table A4, the obtained target through solving of non-radial model 14 (step 1 of STEM ($p = 1$)) for all DMU₀₄, DMU₀₇, and DMU₁₀ was DMU₀₁. As described previously, DM suggested DMU₀₆ as the target of DMU₀₄, DMU₀₇, and DMU₁₀. Therefore, the interactive algorithm should be continued for these DMUs one by one. Here, these three DMUs were considered as DMU₀, simultaneously.

Table A2. The results of solving model 13 for three DMUs (related to the second interactive stage) in the second case study.

DMU ₀	f_l ($l = 1, \dots, 5$)	f^*	f_1	f_2	f_3	f_4	f_5	θ_1^*	θ_2^*	φ_1^*	φ_2^*	φ_3^*	Target
DMU ₀₄	$f_1 = -1\theta_1$	-1	-1	-1	5	6	5	0.500	0.200	1.000	1.000	1.000	DMU ₀₁
	$f_2 = -2\theta_2$	-1	-1	-1	5	6	5	0.500	0.200	1.000	1.000	1.000	DMU ₀₁
	$f_3 = 8\varphi_1$	8	-1	-1	8	6	5	0.500	0.200	1.600	1.000	1.000	DMU ₀₁
	$f_4 = 6\varphi_2$	8	-2	-3	5	8	5	1.000	0.600	1.000	1.333	1.000	DMU ₀₆
	$f_5 = 6\varphi_3$	8	-1	-1	5	6	8	0.500	0.200	1.000	1.000	1.600	DMU ₀₁
DMU ₀₇	$f_1 = -3\theta_1$	-1	-1	-1	4	3	6	0.333	0.250	1.000	1.000	1.000	DMU ₀₁
	$f_2 = -4\theta_2$	-1	-1	-1	4	3	6	0.333	0.250	1.000	1.000	1.000	DMU ₀₁
	$f_3 = 4\varphi_1$	8	-1	-1	8	3	6	0.333	0.250	2.000	1.000	1.000	DMU ₀₁
	$f_4 = 3\varphi_2$	8	-2	-3	4	8	6	0.667	0.750	1.000	2.667	1.000	DMU ₀₆
	$f_5 = 6\varphi_3$	8	-1	-1	4	3	8	0.333	0.250	1.000	1.000	1.333	DMU ₀₁
DMU ₁₀	$f_1 = -6\theta_1$	-1	-1	-1	7	7	2	0.167	0.250	1.000	1.000	1.000	DMU ₀₁
	$f_2 = -4\theta_2$	-1	-1	-1	7	7	2	0.167	0.250	1.000	1.000	1.000	DMU ₀₁
	$f_3 = 7\varphi_1$	8	-1	-1	8	7	2	0.167	0.250	1.143	1.000	1.000	DMU ₀₁
	$f_4 = 7\varphi_2$	7	-1	-1	7	7	2	0.167	0.250	1.000	1.000	1.000	DMU ₀₁
	$f_5 = 2\varphi_3$	8	-1	-1	7	7	8	0.167	0.250	1.000	1.000	4.000	DMU ₀₁

As shown in Table A4, the value of f_4 for DMU₀₄, DMU₀₇ and DMU₁₀ was not satisfied via DM's opinions. Therefore, the obtained solutions for considered DMUs were not the best compromise solutions and composing model 14 (the first iteration of step 2 of STEM) for these three DMUs were required.

The required parameters for step 2 of STEM ($p = 1$) considering DMU₀₇ and DMU₁₀ were calculated as shown in Table A5.

Table A3. Calculating parameters for three DMUs to compose model 14 of step 1 of STEM ($p = 1$) in the second case study.

DMU ₀		$l = 1$	$l = 2$	$l = 3$	$l = 4$	$l = 5$
DMU ₀₄	$f_l^{\min}$	-2.000	-3.000	5.000	6.000	5.000
	f_l^*	-1.000	-1.000	8.000	8.000	8.000
	π_l	0.250	0.133	0.075	0.042	0.075
	β_l	0.435	0.232	0.130	0.072	0.130
DMU ₀₇	$f_l^{\min}$	-2.000	-3.000	4.000	3.000	6.000
	f_l^*	-1.000	-1.000	8.000	8.000	8.000
	π_l	0.167	0.167	0.125	0.208	0.042
	β_l	0.235	0.235	0.176	0.294	0.059
DMU ₁₀	$f_l^{\min}$	-1.000	-1.000	7.000	7.000	2.000
	f_l^*	-1.000	-1.000	8.000	7.000	8.000
	π_l	0.000	0.000	0.018	0.000	0.375
	β_l	0.000	0.000	0.045	0.000	0.955

Table A4. The results of solving model 14 (Step 1 of STEM for $p = 1$) and DM's opinions about them for three DMUs in the second case study.

DMU ₀		Target ₁	Target ₂	Target ₃	Target ₄	Target ₅	Target
DMU ₀₄	Model 14	1.000	1.000	8.000	7.000	8.000	DMU ₀₁
	DM's opinions	2.000	3.000	6.000	8.000	7.000	DMU ₀₆
	Δ_{f_i}	-1.000	-2.000	2.000	-1.000	1.000	-
	Unsatisfied	0	0	0	1	0	-
DMU ₀₇	Model 14	1.000	1.000	8.000	7.000	8.000	DMU ₀₁
	DM's opinions	2.000	3.000	6.000	8.000	7.000	DMU ₀₆
	Δ_{f_i}	-1.000	-2.000	2.000	-1.000	1.000	-
	Unsatisfied	0	0	0	1	0	-
DMU ₁₀	Model 14	1.000	1.000	8.000	7.000	8.000	DMU ₀₁
	DM's opinions	2.000	3.000	6.000	8.000	7.000	DMU ₀₆
	Δ_{f_i}	-1.000	-2.000	2.000	-1.000	1.000	-
	Unsatisfied	0	0	0	1	0	-

As can be seen in the last column of Table A5, all β_l , $l = 1, \dots, 5$ were not calculable (n.c.) for DMU₁₀ (because $\frac{0}{0}$ occurs in calculating β_l , $l = 1, \dots, 5$). So, composing non-radial FDH model 14 for the second iteration of step 2 of STEM for DMU₁₀ was not possible. This means that the best compromise solution for DMU₁₀ could not be obtained. As DM suggested no new target for DMU₁₀ in restarting the second interactive stage, no target existed in FDH model for DMU₁₀ according to DM's opinion, $M' = M' - \text{DMU}_{10}$ ($M' = \{\text{DMU}_{04}, \text{DMU}_{07}\}$).

Table A5. Calculated parameters through solving step 2 of the STEM model ($p = 1$) considering two DMUs (the second case study).

DMU ₀	1	i or r	θ_i^* or φ_r^*	f_l	Unsatisfied f_l	Target (DM)	Δ_{f_i}	β_l
DMU ₀₄	1	i = 1	0.500	-1.000	0	2.000	-1.000	0.000
	2	i = 2	0.200	-1.000	0	3.000	-2.000	0.000
	3	r = 1	1.600	8.000	0	6.000	2.000	0.000
	4	r = 2	1.167	7.000	<u>1</u>	8.000	-1.000	1.000
	5	r = 3	1.600	8.000	0	7.000	1.000	0.000
DMU ₀₇	1	i = 1	0.333	-1.000	0	2.000	-1.000	0.000
	2	i = 2	0.250	-1.000	0	3.000	-2.000	0.000
	3	r = 1	2.000	8.000	0	6.000	2.000	0.000
	4	r = 2	2.333	7.000	1	8.000	-1.000	1.000
	5	r = 3	1.333	8.000	0	7.000	1.000	0.000
DMU ₁₀	1	i = 1	0.167	-1.000	0	2.000	-1.000	<u>n.c.</u>
	2	i = 2	0.250	-1.000	0	3.000	-2.000	<u>n.c.</u>
	3	r = 1	1.143	8.000	0	6.000	2.000	<u>n.c.</u>
	4	r = 2	1.000	7.000	1	8.000	-1.000	<u>n.c.</u>
	5	r = 3	4.000	8.000	0	7.000	1.000	<u>n.c.</u>

Then, non-radial FDH model 14 for DMU₀₄ and DMU₀₇ was composed for the first iteration of step 2 of STEM. The results of solving model 14 (Step 2 of STEM for $p = 2$) and DM's opinions about them for DMU₀₄ and DMU₀₇ are shown in Table A6. As all objective functions for DMU₀₇ were satisfied via DM's opinions (the value of all targets of unsatisfied row for DMU₀₇, that are underlined, were zero), the best compromise solution was obtained, $M' = M' - \text{DMU}_{07}$ ($M' = \{\text{DMU}_{10}\}$). As there was only one similar unsatisfied

objective function (fourth objective function) in two consecutive steps of STEM for DMU₀₄, the target of this DMU could not be considered as DMU₀₆. So, the interactive algorithm was continued by asking DM to suggest a new target for each DMU. Suppose that DM did not suggest new DMUs target for DMU₀₄. So, there was no target for DMU₀₄ in the non-radial FDH model. This DMU was then removed from M' set ($M' = \emptyset$). As there was no non-proceed DMUs and the process of targets determination for all DMUs had been carried out ($M' = \emptyset$), the interactive algorithm was finished.

As shown in the last row of Table A7, the proposed target of DM for DMU₀₇ was obtained (DMU₀₆), but the proposed targets of DM for DMU₀₄ and DMU₁₀ were not obtained.

Appendix A.2. The Detail Results of Applying Extended RBA for the Second Case Study

Applying extended RBA (Section 3.2) is described to find targets of the non-radial FDH model 12–14 for DMU₀₇ of the second case study (related non-radial FDH models in the interactive algorithm). Each target of the non-radial FDH model is found using two target finding stages. A similar process should be carried out to find the target of the mentioned non-radial FDH models for other DMUs. First $x_{i0} = x_{ij}$, $i = 1, 2$, and $y_{r0} = y_{rj}$, $r = 1, 2, 3$ ($j = 1, \dots, 13$) for DMU₀ = DMU₀₇ should be defined.

Table A6. The results of solving model 14 (Step 2 of STEM for $p = 2$) and DM's opinions about them for two DMUs in the second case study.

DMU ₀		Target ₁	Target ₂	Target ₃	Target ₄	Target ₅	Target
DMU ₀₄	Model 14	1	1	8	7	8	DMU ₀₁
	DM's opinions	2	3	6	8	7	DMU ₀₆
	Δ_{fj}	-1	-2	2	-1	1	-
	Unsatisfied	0	0	0	1	0	-
DMU ₀₇	Model 14	1	1	8	7	8	DMU ₀₆
	DM's opinions	2	3	6	8	7	DMU ₀₆
	Δ_{fj}	-1	-2	2	-1	1	-
	Unsatisfied	0	0	0	0	0	-

Table A7. Applying the second interactive stage for three considered DMUs (the second case study).

STEM Phases	DMU ₀₄	DMU ₀₇	DMU ₁₀	Table
Construct pay-off table	√	√	√	A.2
Calculating β_l ($l = 1, \dots, 5$) for proposing step 1 of STEM ($p = 1$) model	√	√	√	A.3
Solving of step 1 of STEM ($p = 1$) model	√	√	√	A.4
Calculating β_l ($l = 1, \dots, 5$) for proposing step 2 of STEM ($p = 1$) model	√	√	n.a.	A.5
Solving of step 2 of STEM ($p = 1$) model	√	√	×	A.6
Obtaining target regards DM's opinions	×	√	×	A.6

Target finding of non-radial FDH models 12 and 13 for DMU₀₇ in the second case study

By applying the first target finding stage, DMUs of feasible region were found. As described in Section 3.2.1, firstly, DMUs of feasible region ($S_{\text{step 0 of STEM}}$) should be found using Equation (15) (Table A8).

As is observed in the last column of Table A8, only five DMUs (DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇) satisfied all five constraints. So, these DMUs belonged to the feasible region ($S_{\text{step 0 of STEM}} = \{\text{DMU}_{01}, \text{DMU}_{02}, \text{DMU}_{05}, \text{DMU}_{06}, \text{DMU}_{07}\}$).

By applying the second target finding stage, the optimum objective function values were then calculated, and DMU targets were found. In this regard, firstly, the optimum

objective function values of models should be calculated by Equations (16) and (17). The objective function value of DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆, and DMU₀₇ (in non-radial FDH model 12, $Z_j^o = \frac{1}{3} \sum_{r=1}^3 \frac{y_{rj}}{y_{ro}} - \frac{1}{2} \sum_{i=1}^2 \frac{x_{ij}}{x_{io}}$, $j = 1, \dots, 13$, and in non-radial FHD model 13, $f_1(u) = -x_{10}x_{1j}$, $f_2(u) = -x_{20}x_{2j}$, $f_3(u) = y_{10}y_{1j}$, $f_4(u) = y_{20}y_{2j}$, $f_5(u) = y_{30}y_{3j}$) are shown in Table A9.

The last row of Table A9 shows the maximum value of Z_j^o and f_l , $l = 1 \dots 5$, respectively. These values show the optimum objective function value (optimum value of Z_j^o and f_l , $l = 1, \dots, 5$) of each non-radial FDH model.

DMU₀ targets are found by comparing the optimum objective function value (optimum value of Z^{*o} and f_l^* , $l = 1, \dots, 5$) with the objective function value of each DMU (Z_j^o and f_l , $l = 1, \dots, 5$) by Equations (15) and (16) (Table A9). It is considerable that the objective functions f_1 , f_2 , and f_3 of model 14 had alternative optimum solutions.

Table A8. Applying the first target finding stage for DMU₀₇ (the second case study).

DMU _j	$\frac{x_{1j}}{x_{10}}$	$\frac{x_{1j}}{x_{10}} \leq 1$	$\frac{x_{2j}}{x_{20}}$	$\frac{x_{2j}}{x_{20}} \leq 1$	$\frac{y_{1j}}{y_{10}}$	$\frac{y_{1j}}{y_{10}} \geq 1$	$\frac{y_{2j}}{y_{20}}$	$\frac{y_{2j}}{y_{20}} \geq 1$	$\frac{y_{3j}}{y_{30}}$	$\frac{y_{3j}}{y_{30}} \geq 1$	S _{step 0 of STEM} (Equation (15))
DMU ₀₁	0.33	0.33	0.25	0.25	2.00	2.00	2.33	2.33	1.33	1.33	√
DMU ₀₂	0.33	0.33	0.50	0.50	2.00	2.00	2.00	2.00	1.00	1.00	√
DMU ₀₃	1.00	1.00	1.00	1.00	2.00	2.00	2.33	2.33	0.67	×	×
DMU ₀₄	0.67	0.67	1.25	×	1.25	1.25	2.00	2.00	0.83	×	×
DMU ₀₅	0.33	0.33	0.25	0.25	1.25	1.25	1.67	1.67	1.00	1.00	√
DMU ₀₆	0.67	0.67	0.75	0.75	1.50	1.50	2.67	2.67	1.17	1.17	√
DMU ₀₇	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	√
DMU ₀₈	1.00	1.00	0.75	0.75	1.25	1.25	1.67	1.67	0.50	×	×
DMU ₀₉	0.67	0.67	0.50	0.50	0.75	×	2.00	2.00	0.67	×	×
DMU ₁₀	2.00	×	1.00	1.00	1.75	1.75	2.33	2.33	0.33	×	×
DMU ₁₁	1.67	×	1.00	1.00	1.00	1.00	2.33	2.33	0.67	×	×
DMU ₁₂	1.67	×	1.25		1.00	1.00	2.33	2.33	1.00	1.00	×
DMU ₁₃	1.67	×	0.75	0.75	1.50	1.50	1.33	1.33	1.17	1.17	×

Table A9. Applying the second target finding stage for DMU₀₇ (the second case study).

DMU _j	Z_j^o (Equation (16))	Model 12 Targets (Equation (18))	f_1 (Equation (17))	f_1 Targets (Equation (19))	f_2 (Equation (17))	f_2 Targets (Equation (19))	f_3 (Equation (17))	f_3 Targets (Equation (19))	f_4 (Equation (17))	f_4 Targets (Equation (19))	f_5 (Equation (17))	f_5 Targets (Equation (19))
DMU ₀₁	2	DMU ₀₁	−1	DMU ₀₁	−1	DMU ₀₁	8	DMU ₀₁	7	×	8	DMU ₀₁
DMU ₀₂	1	×	−1	DMU ₀₂	−2	×	8	DMU ₀₂	6	×	6	×
DMU ₀₅	1	×	−1	DMU ₀₅	−1	DMU ₀₅	5	×	5	×	6	×
DMU ₀₆	1	×	−2	×	−3	×	6	×	8	DMU ₀₆	7	×
DMU ₀₇	0	×	−3	×	−4	×	4	×	3	×	6	×
Max	2	−	−1	−	−1	−	8	−	8	−	8	−

Target finding of non-radial FDH model 14 for DMU₀₇ in the second case study

First, step 1 of STEM ($p = 1$) should be considered. By applying the first target finding stage, DMUs of feasible region (in step 1 of STEM model) were found. As described in Section 3.2.2, DMUs belonging to the feasible region should be found. In DMUs of feasible regions, two conditions should be satisfied. At first, constraints $\frac{x_{ij}}{x_{io}} \leq 1$, $i = 1, 2$, and

$\frac{y_{rj}}{y_{ro}} \geq 1, r = 1, 2, 3 (j = 1, \dots, 13)$ should be satisfied (the first condition). As is observable in Table A8, only DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇ satisfied these constraints.

Moreover, the constraints $\max_l \{(f_l^* - f_l(u)) \cdot \beta_l\} \geq 0, l = 1, \dots, 5$ should be satisfied for each DMU of feasible region (the second condition). Checking the second condition for DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇ are shown in Table A10. As shown in Table A10, the constraints of the second condition were also satisfied for DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇. So, $S_{\text{step 1 of STEM}}^1 = \{\text{DMU}_{01}, \text{DMU}_{02}, \text{DMU}_{05}, \text{DMU}_{06}, \text{DMU}_{07}\}$.

Table A10. Applying the first and second target finding stages to find targets of step 1 of STEM ($p = 1$) for DMU₀₇ (the second case study).

DMU _j	$(f_1^* - f_1(u)) \cdot \beta_1$	$(f_2^* - f_2(u)) \cdot \beta_2$	$(f_3^* - f_3(u)) \cdot \beta_3$	$(f_4^* - f_4(u)) \cdot \beta_4$	$(f_5^* - f_5(u)) \cdot \beta_5$	$S_{\text{step 1 of STEM}}^1$ (Equation (20))	h_j (Equation (21))	Targets (Equation (22))
DMU ₀₁	0.000	0.000	0.000	0.294	0.000	√	0.294	DMU ₀₁
DMU ₀₂	0.000	0.235	0.000	0.588	0.118	√	0.588	×
DMU ₀₅	0.000	0.000	0.529	0.882	0.118	√	0.882	×
DMU ₀₆	0.235	0.471	0.353	0.000	0.059	√	0.471	×
DMU ₀₇	0.471	0.706	0.706	1.471	0.118	√	1.471	×
$h^* = \min \max$ $(f_l^* - f_l(u)) \cdot \beta_l$ ($l = 1, \dots, 5$)	-	-	-	-	-	-	0.294	-

By applying the second target finding stage, the optimum objective function value (in step 1 of STEM model) was calculated and DMU targets were found. By considering Equation (21), the minimum value of the left-hand side of the second condition for DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇ showed the optimum objective function value of non-radial model 14 (last row of Table A10). By considering Equation (22), each DMU that the objective function value was equal to optimum objective function value, showed the targets of DMU₀. So, the target of DMU₀₇ was DMU₀₁ (the last column of Table A10).

For target finding in step 2 of the STEM model ($p = 1$), first, DMUs belonging to feasible region should be found. By considering Equation (22), DMUs of $S_{\text{step 2 of STEM}}^1$ should satisfy three conditions. In this regard, DMU₀₁, DMU₀₂, DMU₀₅, DMU₀₆ and DMU₀₇ satisfied the first and second conditions. Moreover, the constraints $\max_l \{(f_l^* - f_l(u)) \cdot \beta_l\} \geq 0, l = 1, \dots, 5$ (the third condition) should be checked for DMUs of feasible regions. As DMU₀₁ and DMU₀₆ satisfied the third condition, these DMUs belong to $S_{\text{step 2 of STEM}}^1$ and two optimum benchmarks existed.

References

- Farrell, M.J. The measurement of productive efficiency. *J. R. Stat. Soc. Ser. A (General)* **1957**, *120*, 253–281. <https://doi.org/10.2307/2343100>.
- Charnes, A.; Cooper, W.W.; Rhodes, E. Measuring the efficiency of decision making units. *Eur. J. Oper. Res.* **1978**, *2*, 429–444. <https://doi.org/10.1016/0377-221790138-8>.
- Banker, R.D.; Charnes, A.; Cooper, W.W. Some models for estimating technical and scale inefficiencies in data envelopment analysis. *Manag. Sci.* **1984**, *30*, 1078–1092. <https://doi.org/10.1287/mnsc.30.9.1078>.
- Charnes, A.; Cooper, W.W.; Golany, B.; Seiford, L.; Stutz, J. Foundations of data envelopment analysis for Pareto-Koopmans efficient empiric functions. *J. Econom.* **1985**, *30*, 91–107. <https://doi.org/10.1016/0304-407690133-2>.
- Tone, K. A slack-based measure of efficiency in data envelopment analysis. *Eur. J. Oper. Res.* **2001**, *130*, 498–509. <https://doi.org/10.1016/S0377-221700407-5>.
- Green, R.H.; Cook, W.D.; Doyle, J. A note on the additive data envelopment analysis model. *J. Oper. Res. Soc.* **1997**, *48*, 446–448. <https://doi.org/10.1057/palgrave.jors.2600386>.
- Emrouznejad, A.; Parker, B.R.; Tavares, G. Evaluation of research in efficiency and productivity: A survey and analysis of the first 30 years of scholarly literature in DEA. *Socio-Econ. Plan. Sci.* **2008**, *42*, 151–157. <https://doi.org/10.1016/j.seps.2007.07.002>.
- Cook, W.D.; Seiford, L.M. Data envelopment analysis (DEA)—Thirty years on. *Eur. J. Oper. Res.* **2009**, *192*, 1–17. <https://doi.org/10.1016/j.ejor.2008.01.032>.

9. Ramón, N.; Ruiz, J.L.; Sirvent, I. Two-step benchmarking: Setting more realistically achievable targets in DEA. *Expert Syst. Appl.* **2018**, *92*, 124–131. <https://doi.org/10.1016/j.eswa.2017.09.044>.
10. Tsolas, I.E.; Charles, V.; Gherman, T. Supporting Better Practice Benchmarking: A DEA-ANN Approach to Bank Branch Performance Assessment. *Expert Syst. Appl.* **2020**, *160*, 113599. <https://doi.org/10.1016/j.eswa.2020.113599>.
11. Chen, K.; Zhu, J. Additive slacks-based measure: Computational strategy and extension to network DEA. *Omega* **2020**, *91*, 102022. <https://doi.org/10.1016/j.omega.2018.12.011>.
12. Deprins, D.; Simar, L.; Tulkens, H. Measuring labor efficiency in post offices. In *The Performance of Public Enterprises: Concepts and Measurement*; Center for Operations Research and Econometrics (CORE), Amsterdam, North Holland, The Netherlands, 1984.
13. Soleimani-Damaneh, M.; Jahanshahloo, G.R.; Reshadi, M. On the estimation of returns-to-scale in FDH models. *Eur. J. Oper. Res.* **2006**, *174*, 1055–1059. <https://doi.org/10.1016/j.ejor.2005.01.051>.
14. Soleimani-Damaneh, M.; Reshadi, M. A polynomial-time algorithm to estimate returns to scale in FDH models. *Comput. Oper. Res.* **2007**, *34*, 2168–2176. <https://doi.org/10.1016/j.cor.2005.08.004>.
15. Mostafaei, A.; Soleimani-damaneh, M. Global sub-increasing and global sub-decreasing returns to scale in free disposal hull technologies: Definition, characterization and calculation. *Eur. J. Oper. Res.* **2020**, *280*, 230–241. <https://doi.org/10.1016/j.ejor.2019.07.005>.
16. Fukuyama, H.; Leth Hougaard, J.; Sekitani, K.; Shi, J. Efficiency measurement with a non-convex free disposal hull technology. *J. Oper. Res. Soc.* **2016**, *67*, 9–19. <https://doi.org/10.1057/jors.2015.4>.
17. Manzari Tavakoli, I.; Mostafaei, A. Free disposal hull efficiency scores of units with network structures. *Eur. J. Oper. Res.* **2019**, *277*, 1027–1036. <https://doi.org/10.1016/j.ejor.2019.03.023>.
18. Arfa, C.; Leleu, H.; Romdhane, H.B.; van Mosseveld, C.; Sadraoui, T. Measuring the Efficiency of Hospital's Cardiology Wards Using the Free Disposal Hull Approach. *Int. J.* **2020**, *8*, 37–45. <https://doi.org/10.12691/ijefm-8-2-1>.
19. Kerstens, K.; Van De Woestyne, I. Solution methods for nonconvex free disposal hull models: A review and some critical comments. *Asia-Pac. J. Oper. Res.* **2014**, *31*, 1450010. <https://doi.org/10.1142/S0217595914500109>.
20. Soleimani-Damaneh, M.; Mostafaei, A. Identification of the anchor points in FDH models. *Eur. J. Oper. Res.* **2015**, *246*, 936–943. <https://doi.org/10.1016/j.ejor.2015.05.051>.
21. Mirmozaffari, M.; Azeem, G.; Boskabadi, A.; Aranizadeh, A.; Vaishnav, A.; John, J. A Novel Improved Data Envelopment Analysis Model Based on SBM and FDH Models. *Eur. J. Electr. Eng. Comput. Sci.* **2020**, *4*, 1–7. <https://doi.org/10.24018/ejece.2020.4.3.214>.
22. Joro, T.; Korhonen, P. *Extension of Data Envelopment Analysis with Preference Information*; Springer: Berlin/Heidelberg, Germany, 2015.
23. Hwang, C.L.; Masud AS, M. *Multiple Objective Decision Making—Methods and Applications: A State-of-the-Art Survey*; Springer Science & Business Media: Berlin, Germany, 1979; Volume 164.
24. Henriques, C.; Luque, M.; Marcenaro-Gutierrez, O. Coupling distinct MOLP interactive approaches with a novel DEA hybrid model. *Int. Trans. Oper. Res.* **2020**, 1–22. <https://doi.org/10.1111/itor.12879>.
25. Hakanen, J.; Miettinen, K.; Matković, K. Task-based visual analytics for interactive multiobjective optimization. *J. Oper. Res. Soc.* **2020**, *72*, 2073–2090. <https://doi.org/10.1080/01605682.2020.1768809>.
26. Golany, B. An interactive MOLP procedure for the extension of DEA to effectiveness analysis. *J. Oper. Res. Soc.* **1988**, *39*, 725–734. <https://doi.org/10.1057/jors.1988.127>.
27. Benayoun, R.; De Montgolfier, J.; Tergny, J.; Laritchev, O. Linear programming with multiple objective functions: Step method (STEM). *Math. Program.* **1971**, *1*, 366–375. <https://doi.org/10.1007/BF01584098>.
28. Izadikhah, M.; Alikhani, S. An improvement on STEM method in multi-criteria analysis. *J. Math. Ext.* **2012**, *6*, 21–39.
29. Perić, T.; Babić, Z.; Matejaš, J. Comparative analysis of application efficiency of two iterative multi objective linear programming methods (MP method and STEM method). *Cent. Eur. J. Oper. Res.* **2018**, *26*, 565–583. <https://doi.org/10.1007/s10100-018-0543-x>.
30. Gerami, J. An interactive procedure to improve estimate of value efficiency in DEA. *Expert Syst. Appl.* **2019**, *137*, 29–45. <https://doi.org/10.1016/j.eswa.2019.06.061>.
31. Borovička, A. Algorithmic Improvements of the KSU-STEM Method Verified on a Fund Portfolio Selection. *Information* **2020**, *11*, 262. <https://doi.org/10.3390/info11050262>.
32. Wang, C.N.; Dang, T.T. Location Optimization of Wind Plants Using DEA and Fuzzy Multi-Criteria Decision Making: A Case Study in Vietnam. *IEEE Access* **2021**, *9*, 116265–116285. <https://doi.org/10.1109/ACCESS.2021.3106281>.
33. Wang, C.N.; Dang, T.T.; Tibo, H.; Duong, D.H. Assessing renewable energy production capabilities using DEA window and fuzzy TOPSIS model. *Symmetry* **2021**, *13*, 334. <https://doi.org/10.3390/sym13020334>.
34. Wang, C.N.; Dang, T.T.; Bayer, J. A Two-Stage Multiple Criteria Decision Making for Site Selection of Solar Photovoltaic (PV) Power Plant: A Case Study in Taiwan. *IEEE Access* **2021**, *9*, 75509–75525. <https://doi.org/10.1109/ACCESS.2021.3081995>.
35. Rousseeuw, P.J.; Leroy, A.M. *Robust Regression and Outlier Detection*; Wiley: New York, NY, USA, 1987.
36. Han, J.; Kamber, M.; Pei, J. *Data Mining: Concepts and Techniques*; Morgan Kaufmann: Waltham, MA, USA, 2012.
37. Yang, M.; Wan, G.; Zheng, E. A predictive DEA model for outlier detection. *J. Manag. Anal.* **2014**, *1*, 20–41. <https://doi.org/10.1080/23270012.2014.889911>.
38. Cooper, W.W.; Seiford, L.M.; Tone, K. *Data Envelopment Analysis: A Comprehensive Text with Models, Applications, References and DEA Solver Software*; Springer: Berlin/Heidelberg, Germany, 2007.

-
39. Hadi-Vencheh, A.; Foroughi, A.A.; Soleimani-damaneh, M. A DEA model for resource allocation. *Econ. Model.* **2008**, *25*, 983–993. <https://doi.org/10.1016/j.econmod.2008.01.003>.
 40. Slack, N.; Lewis, M. *Operations Strategy*; Pearson Education: London, UK, 2017.