

Editorial

Special Issue on Advances in Marine Biotechnology: Exploitation of Halophyte Plants

Catarina Guerreiro Pereira 

Centre of Marine Sciences—CCMAR, Faculty of Sciences and Technology, Ed. 7, Campus of Gambelas, University of Algarve, 8005-139 Faro, Portugal; cagpereira@ualg.pt

Halophyte (salt tolerant) plants encompass roughly 1% of the world's plant species that can thrive in a multitude of saline biotopes, where glycophytes (non-salt tolerant) cannot [1,2]. They are highly resistant to the abiotic constraints characteristic of saline ecosystems, such as salinity, drought, extreme temperatures, and UV radiation, having evolved specialized morphological and physiological adaptations in response to the challenges of living in such harsh conditions [1,3]. The response mechanisms to handle abiotic stress include the synthesis and accumulation of bioactive metabolites, like phenolic compounds, terpenoids, vitamins, etc. These molecules are usually powerful antioxidants but can display other biological activities such as enzyme inhibition, antimicrobial, anti-inflammatory, and antitumoral activities, among others, thus granting halophytes with potential biotechnological applications for the food, pharmaceutical and cosmetic industries [4–17].

Halophyte plants have long been used for food, forage and ethno-medicinal/veterinary purposes and are nowadays raising scientific and commercial interest particularly in the agri-food sector due to their nutritional, functional and organoleptic properties [2,4,6,18–20]. Being naturally adapted to salinity, halophytes appear pivotal within the context of soil and water salinization and freshwater scarcity for agriculture. They are promising sustainable alternatives to the traditional glycophytic crops, since they can be cultivated under saline conditions using brackish and salt water for irrigation [1,15,19,21–23]. In fact, several halophyte species are emerging as candidates for alternative crops with applications in various commercial segments, from human and/or animal nutrition to pharmaceutical and/or cosmetic industries, being an approach to the sustained management of saline areas, revegetation, soil-remediation, or even ecosystem-engineering purposes [1–6,18,21,24–27]. In the scientific literature, halophytes are increasingly proven to be an important source of relevant compounds, ingredients, and/or raw material with multiple potential biotechnological applications [7–17,19,20,22,23,28–36]. Moreover, several halophyte species are still traditionally used in many parts of the world as food and folk remedies [4], which stands to prove their great potential.

Altogether, halophyte plants represent an outstanding (and underexplored) reservoir of natural biologically active compounds and a promising source for a new generation of sustainable bioproducts (such as foods, drugs and/or health-promoting products) awaiting imaginative exploration. Hence, this Special Issue “Advances in Marine Biotechnology: Exploitation of Halophyte Plants” aimed at providing a platform for current studies on the exploitation of halophytes in view of their vast potential biotechnological applications in all fields of science.

A total of five articles, three research papers and two review papers, are presented in this Special Issue. The three research articles concerned halophyte cultivation, but two of them focused on halophyte culture coupled with Integrated Multi-Trophic Aquaculture (IMTA) systems. Custódio et al. [37] assessed the conditions that benefit the performance of the sea purslane *Halimione portulacoides* in extracting dissolved inorganic nutrients from saline aquaculture effluents under IMTA systems by investigating the influence of lighting



Citation: Pereira, C.G. Special Issue on Advances in Marine Biotechnology: Exploitation of Halophyte Plants. *Appl. Sci.* **2022**, *12*, 10573. <https://doi.org/10.3390/app122010573>

Received: 14 October 2022

Accepted: 18 October 2022

Published: 19 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

conditions and planting density on its vegetative development and nutrient (nitrogen and phosphorous) extraction capacity. The authors discovered that plant growth was unaffected by artificial lighting type, although LED systems proved more energy-efficient, and that high-density planting produced more biomass per unit of area and extracted more dissolved inorganic nitrogen and phosphorus. They state that *H. portulacoides* can be hydroponically cultivated using nutrient-rich saline effluents, employing LEDs as an alternative to fluorescent lighting, and using high-density planting to promote higher yields and extraction efficiencies. Marques et al. [38] investigated the fatty acids (FA) profiles of three halophyte species (*H. portulacoides*, *Salicornia ramosissima* and *Sarcocornia perennans*) whose nutritional properties for human consumption have been documented, comparing wild specimens retrieved from donor sites with conspecifics cultivated in aquaponics coupled with an IMTA system. The authors found that the FA profiles of halophytes from aquaponics differed significantly from those of the wild specimens, displaying higher content of omega-3 FAs in edible parts, and mostly in the form of α -linolenic acid. Their results suggest that halophytes were not nutrient-limited under aquaponic conditions and they highlight the nutritional value of these species, reinforcing the potential of aquaponics coupled with IMTA to produce high-quality halophytes for human consumption. The third research article considered the role of different salt types on specific salinity stress conditions for plants [39]. Nissim et al. [39] compared the effects of seawater at moderate and high concentrations and analogous NaCl solutions on salt-tolerant (*Tetragonia tetragonoides*) and salt-sensitive (*Lactuca sativa*) crops grown in hydroponics, theorising that, due to its mineral composition, diluted seawater would result in a less stressful growing medium than NaCl. The authors confirmed this hypothesis and found significant differences between the halophyte and the glycophyte species: the growth rates in *T. tetragonoides* did not vary between the two types of stress, but in *L. sativa* the diluted seawater led to higher biomass yield in both fresh weigh and dry weight. Their results in terms of physiological parameters and concentration of minerals, phenolics, and proline also showed that seawater is a less detrimental growing medium than a similar NaCl solution, potentially due to different mineral composition.

From the two review articles, Hulkko et al. [40] addressed the contents of various types of compounds in halophyte biomass, while Karkanis et al. [41] focused on *Crithmum maritimum* (sea fennel) as an alternative crop for saline agriculture. Hulkko et al. [40] encompassed proanthocyanidins (condensed tannins), total phenolic compounds, photosynthetic pigments (chlorophyll and carotenoids), and vitamins in various halophytes, also considering extraction and analytical methods, and showed that several species have potential for nutrient-rich food production in salt-affected areas as well as to produce nutraceuticals and ingredients for cosmetics and pharmaceutical industries. They also postulate that waste repurposing could help identify new bio-derived sources for valuable compounds, while improving the sustainability and economic feasibility of existing production systems. Karkanis et al. [41] compiles recent literature on farming practices that may allow the establishment of sea fennel cultivation protocols for farmers, while also presenting the main constraints that hinder its further exploitation. Additionally, the authors gathered up-to-date information on the chemical composition (e.g., volatile and polyphenolic compounds, fatty acids, amino acids, minerals) and health-related properties (e.g., antimicrobial, insecticidal, anticholinesterase activities) of different plant organs to help uncover potential alternative uses that can increase the sea fennel's added value, contributing to its commercial exploration. The authors conclude that cultivation of halophytes as cash crops allows the use of salinized soils and/or salinized water in coastal areas to produce high-added-value products, and that sea fennel possesses features that fit this context.

Overall, the published studies provided advances on the biotechnological exploitation of halophytes, but more research in the field will continue to address the potential applications of these salt-tolerant plant species to uncover all that they have to offer.

Funding: This research was funded by Portuguese national funds from FCT—Foundation for Science and Technology through projects UIDB/04326/2020, UIDP/04326/2020, LA/P/0101/2020, and PTDC/BAAAGR/1391/2020 (GreenValue).

Acknowledgments: The author is grateful for the opportunity to join *Applied Sciences* as a Guest Editor of this Special Issue, with a special thanks to *Applied Sciences* editorial team. The author expresses her gratitude to the authors and peer reviewers for their valuable contributions.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Hasanuzzaman, M.; Nahar, K.; Alam, M.M.; Bhowmik, P.C.; Hossain, M.A.; Rahman, M.M.; Prasad, M.N.V.; Ozturk, M.; Fujita, M. Potential use of halophytes to remediate saline soils. *Biomed. Res. Int.* **2014**, *2014*, 589341. [[CrossRef](#)] [[PubMed](#)]
2. Oliveira, M.; Hoste, H.; Custódio, L. A systematic review on the ethnoveterinary uses of mediterranean salt-tolerant plants: Exploring its potential use as fodder, nutraceuticals or phytotherapeutics in ruminant production. *J. Ethnopharmacol.* **2021**, *267*, 113464. [[CrossRef](#)] [[PubMed](#)]
3. Grigore, M.-N.; Toma, C. *Anatomical Adaptations of Halophytes. A Review of Classic Literature and Recent Findings*; Springer International Publishing: Cham, Switzerland, 2017.
4. Ksouri, R.; Ksouri, W.M.; Jallali, I.; Debez, A.; Magné, C.; Hiroko, I.; Abdelly, C. Medicinal halophytes: Potent source of health promoting biomolecules with medical, nutraceutical and food applications. *Crit. Rev. Biotechnol.* **2012**, *32*, 289–326. [[CrossRef](#)] [[PubMed](#)]
5. Petropoulos, S.A.; Karkanis, A.; Martins, N.; Ferreira, I.C.F.R. Halophytic herbs of the Mediterranean basin: An alternative approach to health. *Food Chem. Toxicol.* **2018**, *114*, 155–169. [[CrossRef](#)]
6. Petropoulos, S.A.; Karkanis, A.; Martins, N.; Ferreira, I.C.F.R. Edible halophytes of the Mediterranean basin: Potential candidates for novel food products. *Trends Food Sci. Technol.* **2018**, *74*, 69–84. [[CrossRef](#)]
7. Lopes, A.; Rodrigues, M.J.; Pereira, C.; Oliveira, M.; Barreira, L.; Varela, J.; Trampetti, F.; Custódio, L. Natural products from extreme marine environments: Searching for potential industrial uses within extremophile plants. *Ind. Crops Prod.* **2016**, *94*, 299–307. [[CrossRef](#)]
8. Oliveira, M.; Rodrigues, M.J.; Neng, N.R.; Nogueira, J.M.F.; Bessa, R.J.B.; Custódio, L. Seasonal variations of the nutritive value and phytotherapeutic potential of *Cladum mariscus* L. (Pohl.) targeting ruminant's production. *Plants* **2021**, *10*, 556. [[CrossRef](#)]
9. Rodrigues, M.J.; Soszynski, A.; Martins, A.; Rauter, A.P.; Neng, N.R.; Nogueira, J.M.F.; Varela, J.; Barreira, L.; Custódio, L. Unravelling the antioxidant potential and the phenolic composition of different anatomical organs of the marine halophyte *Limonium algarvense*. *Ind. Crops Prod.* **2015**, *77*, 315–322. [[CrossRef](#)]
10. Castañeda-Loaiza, V.; Placines, C.; Rodrigues, M.J.; Pereira, C.; Zengin, G.; Uysal, A.; Jekö, J.; Cziáky, Z.; Reis, C.P.; Gaspar, M.M.; et al. If you cannot beat them, join them: Exploring the fruits of the invasive species *Carpobrotus edulis* (L.) N.E. Br as a source of bioactive products. *Ind. Crops Prod.* **2020**, *144*, 112005. [[CrossRef](#)]
11. Pereira, C.G.; Locatelli, M.; Innosa, D.; Cacciagrano, F.; Polesná, L.; Santos, T.F.; Rodrigues, M.J.; Custódio, L. Unravelling the potential of the medicinal halophyte *Eryngium maritimum* L.: In vitro inhibition of diabetes-related enzymes, antioxidant potential, polyphenolic profile and mineral composition. *S. Afr. J. Bot.* **2019**, *120*, 204–212. [[CrossRef](#)]
12. Rodrigues, M.J.; Jekö, J.; Cziáky, Z.; Pereira, C.G.; Custódio, L. The medicinal halophyte *Frankenia laevis* L. (sea heath) has in vitro antioxidant activity, α -glucosidase inhibition, and cytotoxicity towards hepatocarcinoma cells. *Plants* **2022**, *11*, 1353. [[CrossRef](#)] [[PubMed](#)]
13. Placines, C.; Castañeda-Loaiza, V.; Rodrigues, M.J.; Pereira, C.G.; Stefanucci, A.; Mollica, A.; Zengin, G.; Llorent-Martínez, E.J.; Castilho, P.C.; Custódio, L. Phenolic profile, toxicity, enzyme inhibition, in silico studies, and antioxidant properties of *Cakile maritima* Scop. (Brassicaceae) from southern Portugal. *Plants* **2020**, *9*, 142. [[CrossRef](#)] [[PubMed](#)]
14. Rodrigues, M.J.; Gangadhar, K.N.; Vizzeto-Duarte, C.; Wubshet, S.G.; Nyberg, N.T.; Barreira, L.; Varela, J.; Custódio, L. Maritime halophyte species from Southern Portugal as sources of bioactive molecules. *Mar. Drugs* **2014**, *12*, 2228–2244. [[CrossRef](#)] [[PubMed](#)]
15. Rodrigues, M.J.; Castañeda-Loaiza, V.; Monteiro, I.; Pinela, J.; Barros, L.; Abreu, R.M.V.; Oliveira, M.C.; Reis, C.; Soares, F.; Pousão-Ferreira, P.; et al. Metabolomic profile and biological properties of sea lavender (*Limonium algarvense* Erben) plants cultivated with aquaculture wastewaters: Implications for its use in herbal formulations and food additives. *Foods* **2021**, *10*, 3104. [[CrossRef](#)] [[PubMed](#)]
16. Pereira, C.G.; Barreira, L.; Bijttebier, S.; Pieters, L.; Marques, C.; Santos, T.F.; Rodrigues, M.J.; Varela, J.; Custódio, L. Health promoting potential of herbal teas and tinctures from *Artemisia campestris* subsp. *maritima*: From traditional remedies to prospective products. *Sci. Rep.* **2018**, *8*, 4689. [[CrossRef](#)]
17. Pereira, C.G.; Barreira, L.; Neng, N.R.; Nogueira, J.M.F.; Marques, C.; Santos, T.F.; Varela, J.; Custódio, L. Searching for new sources of innovative products for the food industry within halophyte aromatic plants: In vitro antioxidant activity and phenolic and mineral contents of infusions and decoctions of *Crithmum maritimum* L. *Food Chem. Toxicol.* **2017**, *107*, 581–589. [[CrossRef](#)]
18. Custódio, L.; Rodrigues, M.J.; Pereira, C.G.; Castañeda-Loaiza, V.; Fernandes, E.; Standing, D.; Neori, A.; Shpigel, M.; Sagi, M. A review on *Sarcococca* species: Ethnopharmacology, nutritional properties, phytochemistry, biological activities and propagation. *Foods* **2021**, *10*, 2778. [[CrossRef](#)]

19. Castañeda-Loaiza, V.; Oliveira, M.; Santos, T.; Schüler, L.; Lima, A.R.; Gama, F.; Salazar, M.; Neng, N.R.; Nogueira, J.M.F.; Varela, J.; et al. Wild vs cultivated halophytes: Nutritional and functional differences. *Food Chem.* **2020**, *333*, 127536. [[CrossRef](#)]
20. Barreira, L.; Resek, E.; Rodrigues, M.J.; Rocha, M.I.; Pereira, H.; Bandarra, N.; Silva, M.M.; Varela, J.; Custódio, L. Halophytes: Gourmet food with nutritional health benefits? *J. Food Compos. Anal.* **2017**, *59*, 35–42. [[CrossRef](#)]
21. Peña, R.J.H.; Hernández, D.M.; Ghasemi, M.; Puente, E.O.R. Salt tolerant plants as a valuable resource for sustainable food production in arid and saline coastal zones. *Acta Biol. Colomb.* **2021**, *26*, 116–126. [[CrossRef](#)]
22. Rodrigues, M.J.; Monteiro, I.; Placines, C.; Castañeda-Loaiza, V.; Ślusarczyk, S.; Matkowski, A.; Pereira, C.; Pousão, P.; Custódio, L. The irrigation salinity and harvesting affect the growth, chemical profile and biological activities of *Polygonum maritimum* L. *Ind. Crops Prod.* **2019**, *139*, 111510. [[CrossRef](#)]
23. Rodrigues, M.J.; Monteiro, I.; Castañeda-Loaiza, V.; Placines, C.; Oliveira, M.C.; Reis, C.; Caperta, A.D.; Soares, F.; Pousão-Ferreira, P.; Pereira, C.; et al. Growth performance, in vitro antioxidant properties and chemical composition of the halophyte *Limonium algarvense* Erben are strongly influenced by the irrigation salinity. *Ind. Crops Prod.* **2020**, *143*, 111930. [[CrossRef](#)]
24. Buhmann, A.; Papenbrock, J. An economic point of view of secondary compounds in halophytes. *Funct. Plant Biol.* **2013**, *40*, 952–967. [[CrossRef](#)] [[PubMed](#)]
25. Flowers, T.J.; Galal, H.K.; Bromham, L. Evolution of halophytes: Multiple origins of salt tolerance in land plants. *Funct. Plant Biol.* **2010**, *37*, 604–612. [[CrossRef](#)]
26. Santos, J.; Al-Azzawi, M.; Aronson, J.; Flowers, T.J. EHALOPH a database of salt-tolerant plants: Helping put halophytes to work. *Plant Cell Physiol.* **2015**, *57*, e10. [[CrossRef](#)]
27. Ventura, Y.; Sagi, M. Halophyte crop cultivation: The case for *Salicornia* and *Sarcocornia*. *Environ. Exp. Bot.* **2013**, *92*, 144–153. [[CrossRef](#)]
28. Custódio, L.; Ferreira, A.C.; Pereira, H.; Silvestre, L.; Vizotto-Duarte, C.; Barreira, L.; Rauter, A.P. The marine halophytes *Carpobrotus edulis* L. and *Arthrocnemum macrostachyum* L. are potential sources of nutritionally important PUFAs and metabolites with antioxidant, metal chelating and anticholinesterase inhibitory activities. *Bot. Mar.* **2012**, *55*, 281–288. [[CrossRef](#)]
29. Oliveira, M.; Junior, P.A.S.; Rodrigues, M.J.; DellaGreca, M.; Barreira, L.; Murta, S.M.F.; Romanha, A.J.; Custódio, L. Unlocking the in vitro anti-*Trypanosoma cruzi* activity of halophyte plants from the southern Portugal. *Asian Pac. J. Trop. Med.* **2016**, *9*, 735–741. [[CrossRef](#)]
30. Oliveira, M.; Rodrigues, M.J.; Pereira, C.; Neto, R.L.M.; Junior, P.A.S.; Neng, N.R.; Nogueira, J.M.F.; Varela, J.; Barreira, L.; Custódio, L. First report of the in vitro antileishmanial properties of extremophile plants from the Algarve Coast. *Nat. Prod. Res.* **2018**, *32*, 600–604. [[CrossRef](#)]
31. Pereira, C.G.; Custódio, L.; Rodrigues, M.J.; Neng, N.R.; Nogueira, J.M.F.; Carlier, J.; Costa, M.C.; Varela, J.; Barreira, L. Profiling of antioxidant potential and phytoconstituents of *Plantago coronopus*. *Braz. J. Biol.* **2017**, *77*, 632–641. [[CrossRef](#)]
32. Pereira, C.G.; Barreira, L.; Blijtebier, S.; Pieters, L.; Neves, V.; Rodrigues, M.J.; Rivas, R.; Varela, J.; Custódio, L. Chemical profiling of infusions and decoctions of *Helichrysum italicum* subsp. *picardii* by UHPLC-PDA-MS and in vitro biological activities comparatively with green tea (*Camellia sinensis*) and rooibos tisane (*Aspalathus linearis*). *J. Pharm. Biomed. Anal.* **2017**, *145*, 593–603. [[CrossRef](#)] [[PubMed](#)]
33. Rodrigues, M.J.; Gangadhar, K.N.; Zengin, G.; Mollica, A.; Varela, J.; Barreira, L.; Custódio, L. *Juncaceae* species as sources of innovative bioactive compounds for the food industry: In vitro antioxidant activity, neuroprotective properties and in silico studies. *Food Chem. Toxicol.* **2017**, *107B*, 590–596. [[CrossRef](#)] [[PubMed](#)]
34. Rodrigues, M.J.; Custódio, L.; Lopes, A.; Oliveira, M.; Neng, N.R.; Nogueira, J.M.F.; Martins, A.; Rauter, A.P.; Varela, J.; Barreira, L. Unlocking the in vitro anti-inflammatory and antidiabetic potential of *Polygonum maritimum*. *Pharm. Biol.* **2017**, *55*, 1348–1357. [[CrossRef](#)] [[PubMed](#)]
35. Rodrigues, M.J.; Ślusarczyk, S.; Pecio, Ł.; Matkowski, A.; Salmas, R.E.; Durgadi, S.; Pereira, C.; Varela, J.; Barreira, L.; Custódio, L. In vitro and in silico approaches to appraise *Polygonum maritimum* L. as a source of innovative products with anti-ageing potential. *Ind. Crops Prod.* **2018**, *111*, 391–399. [[CrossRef](#)]
36. Rodrigues, M.J.; Pereira, C.A.; Oliveira, M.; Neng, N.R.; Nogueira, J.M.F.; Zengin, G.; Mahomoodally, M.F.; Custódio, L. Sea rose (*Armeria pungens* (Link) Hoffmanns. & Link) as a potential source of innovative industrial products for anti-ageing applications. *Ind. Crops Prod.* **2018**, *121*, 250–257. [[CrossRef](#)]
37. Custódio, M.; Cartaxana, P.; Villasante, S.; Calado, R.; Lillebø, A.I. LED Lighting and high-density planting enhance the cost-efficiency of *Halimione portulacoides* extraction units for integrated aquaculture. *Appl. Sci.* **2021**, *11*, 4995. [[CrossRef](#)]
38. Marques, B.; Maciel, E.; Domingues, M.R.; Calado, R.; Lillebø, A.I. Halophyte plants cultured in aquaponics hold the same potential for valorization as wild conspecifics from donor sites. *Appl. Sci.* **2021**, *11*, 11586. [[CrossRef](#)]
39. Nissim, W.N.; Masi, E.; Pandolfi, C.; Mancuso, S.; Atzori, G. The response of halophyte (*Tetragonia tetragonoides* (Pallas) Kuntz.) and glycophyte (*Lactuca sativa* L.) crops to diluted seawater and NaCl solutions: A comparison between two salinity stress types. *Appl. Sci.* **2021**, *11*, 6336. [[CrossRef](#)]
40. Hulkko, L.S.S.; Chaturvedi, T.; Thomsen, M.H. Extraction and quantification of chlorophylls, carotenoids, phenolic compounds, and vitamins from halophyte biomass. *Appl. Sci.* **2022**, *12*, 840. [[CrossRef](#)]
41. Karkanis, A.; Polyzos, N.; Kompocholi, M.; Petropoulos, S.A. Rock samphire, a candidate crop for saline agriculture: Cropping practices, chemical composition and health effects. *Appl. Sci.* **2022**, *12*, 737. [[CrossRef](#)]