applied sciences

MDPI

Article

Do You Train Like You Compete? A Comparison of Training Tasks and Competition in Elite Basketball Based on Biomechanical and External Physiological Load

Carlos Sosa Marín [1,2], Enrique Alonso-Pérez-Chao [1,3], Xavier Schelling [4] and Alberto Lorenzo [2,*]

[1] Department of Sports Sciences, Faculty of Medicine, Health and Sports, Universidad Europea de Madrid, 28001 Villaviciosa de Odón, Spain
[2] Facultad de Ciencias de la Actividad Física y del Deporte, Universidad Politécnica de Madrid, 28040 Madrid, Spain
[3] Facultad de Ciencias Biomédicas y de la Salud, Universidad Alfonso X el Sabio (UAX), 28691 Villanueva de la Cañada, Spain
[4] Institute for Health and Sport (iHeS), Victoria University, Melbourne, VIC 8001, Australia
* Correspondence: alberto.lorenzo@upm.es

Abstract

Basketball is an intermittent sport with high neuromuscular and metabolic demands. To optimize specificity, training tasks should replicate competitive loads, but little is known about how drills compare to official matches. This study compared the physiological and biomechanical load of training tasks with official competition in elite U18 basketball players. Twelve male players (16.9 ± 0.8 years) were monitored across two seasons (179 training sessions, 21 matches). A total of 3136 individual records were collected using Catapult Vector S7 LPS units. Training drills were classified by specificity (0–5). Physiological (distance and intensity zones) and biomechanical variables (accelerations, decelerations, jumps, explosive efforts, PlayerLoad™) were analyzed using cluster analysis and linear mixed models. Competition imposed the highest physiological and biomechanical loads. Non-opposition drills (1v0–5v0) showed limited transfer, though 1v0–2v0 accumulated higher jump density. Among opposition formats, 3v3 full-court was the best at replicating match demands. Continuous opposition tasks (3v3v3, 4v4v4, 5v5v5) elicited lower physiological but comparable biomechanical load. Small-sided formats, particularly 3v3 and 4v4, are the most effective training tools for reproducing competition demands, while non-opposition drills are better suited for technical or rehabilitation purposes.

Keywords: team sports; ball drills; small-sided games; scrimmage games; load monitoring; drill-design

Check for updates

Academic Editor: Argyris Toubekis

Received: 12 November 2025
Revised: 12 January 2026
Accepted: 15 January 2026
Published: 19 January 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Basketball is an open skills sport in which players are required to react and make decisions in an unpredictable and changing external rhythm environment [1–4]; intermittent between offensive and defensive actions [5–7], with frequent changes in movement types and intensity [8–10], alternating periods of high intentionality with periods of medium and low intensity [7,11,12]. From a physical-demand standpoint, it could be said that, in competition, there is an alternation of aerobic and anaerobic demands consecutively [13], which implies high neuromuscular and metabolic demands [14].

The principle of training specificity states that the greatest performance gains of training are obtained when the training stimulus is matched to the physiological and motor

requirements of the sport during competition [15]. As Dawson et al. [16] state with reference to a well-known aphorism in the training world, 'you should train as you compete'. For this reason, training strategies that replicate the performance demands of competition should be employed throughout the season [17–19]. Attention needs to be paid to task design, as if the training does not match the load endured and does not meet the specific demands of competition, this will result in lower competitive performance [20]. Knowledge and understanding of the movement patterns and activities performed in competitions by players is essential to plan relevant and effective training [16].

The findings from research examining competition demands in basketball are highly variable and inconsistent, as they are influenced by multiple factors that determine both external and internal load. In summary, and in relation to competition and peak demands observed in competition, the following table provides the most relevant data in this regard (Table 1).

Table 1. Absolute average values, average values relative to time and values relative to time at peak demands for 1 min windows.

Variable	Absolute Average	Normalized Average (per Minute)	Peak Demand Relative to Time (per Minute) in a 1 min Window
PL	240–423 ua	6–10 ua/min	18.45–19.92 ua/min
Distance	3209 m–3325 m (LPS) 4404 m–7558 m (TMA)	70–95 m/min 93.1–133.1 m/min	120–144 m/min X
Distance < 7 Km/h	1259 m–1455 m (LPS) 404 m–1838 m (TMA)	X	X
Distance 7–14 Km/h	1122.72 m–1199.01 m (LPS) 1424 m–2208 m (TMA)	X	X
Distance 14–18 Km/h	483.89 m–451.78 m (LPS) 928 m–2845 m (TMA)	X	X
Distance > 18 Km/h	139.83 m–195.29 m (LPS) 70 m–1096 m (TMA)	2–4 m/min X	20–30 m/min X
Accelerations	100–140 n	2.5–4.2 n/min	6.4–9.2 n/min
Decelerations	84–124 n	2.1–3.1 n/min	6.3–8.4 n/min
Jumps	32–52 n	1.1–1.6 n/min	X
Activity change (frequency)	1–3 sg	X	X
Lactate concentration	2.7–6.8 mmol/L^{-1}	X	X
HR Max	170–198 lpm	X	X
HR Avg	136–158 lpm	X	X
% time < 85% HR Max	19.6–35%	X	X
% time > 85% HR Max	80.4–65%	X	X

Notes: Sosa et al. (2021) [21]; Stojanović et al. (2018) [7]; Torres-Ronda et al. (2016) [22].

In basketball, there is research that has compared training with official competition [20,23,24], with friendly competition [22,25,26] or with simulated competition contexts [27]. In addition, there are studies that, although they have monitored training and official competition, have not compared the demands, but have quantified the weekly percentage change in external load and related it to internal load [28,29] or have attempted to relate training load to injury risk [30–33]. Given the time and resources devoted to planning training tasks and programs with the aim of simulating the demands of competition, this information enables the design of training protocols that are appropriately tailored to both individual and team needs related to competition [34]. However, as the same authors point out, most research to date has examined the load experienced by players using the traditional distinction between external and internal load (e.g., [35,36]). However, it is necessary to go beyond

this classification, distinguishing more precisely between physiological and biomechanical loading [14,37].

Therefore, the aim of the present study is to compare training drills with competition in elite basketball players from the perspective of physiological and biomechanical load experienced.

2. Materials and Methods

2.1. Sample

Eighteen elite male basketball players [38] from the same team, competing in the highest regional division of the U18 Spanish basketball league, were included in this investigation (n = 18; mean ± SD: age 16.9 ± 0.8 years, height 196.6 ± 9.4 cm, body mass 91.7 ± 8.2 kg). Data collection was conducted across 179 training sessions and 21 official matches, all of which followed standard FIBA regulations. This guaranteed uniformity in match duration, court size, player numbers and overall game structure throughout the entire sample. Data collection took place at the same training facility over two consecutive seasons (2018–2019 and 2019–2020). To qualify for inclusion, athletes were required to participate in at least 50% of training sessions (≥90 of 179) and a minimum of 35% of official games (≥8 of 21) across both seasons. Players failing to meet these minimum thresholds were not considered in the final analyses.

At the session level, any training entry in which an athlete completed less than 80% of the planned duration was discarded; nonetheless, these players remained part of the study sample provided they satisfied the general inclusion criteria. In competition, game-level data were omitted when players accumulated fewer than 15 min of effective playing time, as recorded by the tracking devices. These 15 min reflected the actual time spent on the court, including periods of live play and stoppages (e.g., fouls, free throws, out-of-bounds actions), but not time spent on the bench or during official breaks such as timeouts or quarter intervals.

After applying all exclusion criteria, six players who initially met the recruitment criteria were removed from the final dataset because they did not fulfill one or more of the predefined inclusion criteria. The resulting sample comprised 3136 valid observations, 141 from official games and 2995 from training sessions, collected from a total of 12 players. All procedures were carried out following the guidelines of the Declaration of Helsinki [39].

2.2. Procedures

This observational study spanned two complete competitive seasons (2019–2020 and 2020–2021). During every monitored activity, both training sessions and official games, athletes wore Vector S7™ devices (Catapult Sports, Melbourne, Australia) secured within a custom-fitted vest featuring a pocket positioned over the upper thoracic spine, between the scapulae. Each unit integrated a tri-axial accelerometer (±16 g, 100 Hz), tri-axial magnetometer (±4900 µT, 100 Hz), tri-axial gyroscope (up to 2000 deg/s, 100 Hz) and a local positioning system (LPS). Positional data were collected through the ClearSky system (ClearSky S7, Catapult Sports, Melbourne, Australia, firmware 5.6) operating at 10 Hz and based on ultra-wideband technology (4 GHz). To guarantee optimal signal reception and accurate spatial reconstruction, 24 antennas were strategically installed around the court's perimeter. Previous research has verified the validity of this technology for accurately quantifying distance-related variables [40–43], such as speed, accelerations, decelerations [40,41] and Player LoadTM [44], and has demonstrated high reliability (coefficient of variation < 5%) for distance and speed measurements [43]. All athletes were familiar with this monitoring equipment, as the same tracking units had been routinely used throughout both training sessions and competition during the previous season. Before each official match, the devices

were powered on approximately 20–40 min before warm-up initiation. To reduce potential variability between units in the quantification of external load, each player was assigned a specific device and used the same unit consistently throughout the entire study [45].

Activity editing was performed both in real time and after each session to maintain consistency in the dataset. To limit inter-observer variation, all editing procedures were conducted by the same researcher. For training sessions, active duration was defined as the time during which a player was directly participating in the task, excluding transitions between drills, water breaks, or any period during which the player was present but not actively engaged. A player was considered inactive when positioned off the court without taking part in the exercise (for example, during a 5v5 drill while waiting to substitute a teammate). Once data collection was finalized, player-specific information for each training session, organized by drill type, was exported using Catapult Sports' OpenField cloud software (version 1.22.0). After applying predefined exclusion criteria, processed data were transferred to a Microsoft Excel spreadsheet (version 16.0, Microsoft Corporation, Redmond, WA, USA) for subsequent analysis. Training drills were then categorized based on level of specificity, using a 0-to-5 classification based on the framework proposed by Schelling and Torres [46]. Activities classified as levels 0–1 involved non-basketball work performed outside the court (e.g., cycling), whereas level 5 represented formal competitive play (Table 2).

Table 2. Drill classification based on their specificity from 5 to 0 [46].





4	5v5 full-court	A 5v5 drill involves 10 players on the court simultaneously. The number of continuous actions and the work-to-rest ratio depend on the coach's instructions and the pauses introduced during the drill.	
4	5v5v5	In this 5v5v5 format, the team coming off a defensive action transitioned directly into an offensive phase, progressing to the basket located on the opposite side.	
4	4v4 full-court	The 4v4 drill involves eight players with continuous transitions: the defending team converts to offense and attacks the same opponent at the opposite basket. The number of consecutive actions (typically 3–6) and work-to-rest ratios vary according to coaching instructions and integrated rest intervals.	
4	4v4v4	In this 4v5v5 format, the team coming off a defensive action transitioned directly into an offensive phase, progressing to the basket located on the opposite side.	

Table 2. *Cont.*












3	3v3 full-court	The 3v3 drill involves six players with continuous transitions: the defending team converts to offense and attacks the same opponent at the opposite basket. The number of consecutive actions (typically 3–6) and work-to-rest ratios vary according to coaching instructions and integrated rest intervals.	
3	3v3v3	In the 3v3v3 drill, the defending team transitions to offense and attacks the opposite basket against the third team awaiting in defensive position.	
3	Eleven Player Break	A continuous 3v2 drill is performed. Among the five players involved, the player who secures possession at the end of the action, whether through a made basket, rebound, or turnover, initiates the attack on the opposite side of the court, supported by two teammates positioned in the corners and facing two defenders already waiting on that end.	
3	2v2 full-court	In the 2v2 drill, teams transition immediately from offense to defense at the opposite basket, then initiate a counterattack by passing to either of two teammates positioned on the far side of the court.	
2	1v1 in longitudinal half-court (28 × 7.5 m)	The drill begins with the attacker seeking to penetrate and score in a 1v1 situation. When the action ends, the attacker rotates into the defensive position.	
2	5v0 full-court	A 5v0 drill is executed starting from midcourt, followed by another 5v0 sequence at the opposite end of the court. Once both drills are completed, a new group of five players rotates onto the court.	
2	4v0 full-court	A 4v0 drill is executed from midcourt and then repeated at the opposite end. Once both sequences are completed, a new group of four players rotates onto the court.	

Table 2. *Cont.*

2	4v0 full-court	A 4v0 drill is executed from midcourt and then repeated at the opposite end. Once both sequences are completed, a new group of four players rotates onto the court.	
2	3v0 full-court	A 3v0 drill is carried out from midcourt and then repeated at the opposite end. After completing both sequences, a new group of three players enters the court.	
2	2v0 (individual technical-tactical) half-court	Several individual technical-tactical tasks were undertaken in a controlled, no-opposition two-player format (2v0)	
2	1v0 (individual technical-tactical) half-court	Several individual technical-tactical tasks were undertaken in a controlled, no-opposition one-player format (1v0)	

2.3. Physical Variables

The selected physical indicators were divided into physiological and biomechanical categories [3,37]. The variables were processed and reported as normalized per-minute accumulation metrics.

2.3.1. Physiological Variables

Five physiological variables were analyzed: total distance covered per minute (TD, $m \cdot min^{-1}$) and distance per minute across four intensity zones: standing–walking (S-W, <7 $km \cdot h^{-1}$), jogging (JOG, 7–14 $km \cdot h^{-1}$), running (RUN, 14.01–18 $km \cdot h^{-1}$), and high-speed running (HSR, >18 $km \cdot h^{-1}$), consistent with established basketball research protocols [47].

To characterize the external physiological demands of each drill, a two-step cluster analysis (average silhouette = 0.5) was applied using distance-based metrics, including total distance per minute and distance per minute within each speed zone (Table 3). This procedure yielded four discrete clusters representing ascending levels of external physiological load: low (assigned value 1), medium (2), high (3) and very high (4).

Table 3. Clusters of training drills based on external physiological load profiles.

Variables	External Physiological Load			
	Low	Medium	High	Very High
Distance per minute (m)	18.56	62.09	75.28	80.50
Standing–walking (<7 km·h^{-1})	15.53	31.75	65.40	34.35
Jogging (7–14 km·h^{-1})	3.13	21.36	51.81	28.05
Running (14.01–18 km·h^{-1})	0.85	6.37	17.32	12.29
High-speed running (>18 km·h^{-1})	0.22	2.42	3.77	6.02
Sample size (N)	141	1831	122	1042
Sample proportion (%)	4.5%	58.4%	3.9%	32.2%
Bayesian information criterion (BIC)	9214.44			
Average silhouette	0.5			

Note: Data for each external physiological load variable are presented as mean ± standard deviation, with sample sizes representing the number of tasks classified within each category.

To determine the external physiological load associated with each drill type, an average load value was calculated based on the assigned cluster scores (1–4). For example, if 100 recordings from official games were classified evenly between cluster 4 and cluster 3 (50 each), the resulting average would be 3.5, indicating an external physiological load value of 3.5 for that drill category.

2.3.2. Biomechanical Variables

Five biomechanical metrics were analyzed: jumps per minute (JUMPS) exceeding 20 cm; the number of accelerations per minute (ACC) greater than 2 m·s^{-2} (dwell time: 0.3 s); the number of decelerations per minute (DEC) below −2 m·s^{-2} (dwell time: 0.3 s); PlayerLoad™ per minute (PL, arbitrary units); and explosive efforts per minute (EE). The selected dwell time was set at 0.3 s, as durations between 0.3 and 0.4 s are commonly applied in basketball monitoring contexts [47–49].

PL was calculated as the square root of the sum of the instantaneous rate of change in acceleration in the three movement planes (x-, y- and z-axes) using the following formula [47,50]: $PlayerLoad^{TM} = \left[\sqrt{(a_{y1} - a_{y-1})^2} + \sqrt{(a_{x1} - a_{x-1})^2} + \sqrt{(a_{z1} - a_{z-1})^2}\right]/100$.

In this equation, fwd denotes movement along the anterior–posterior axis, side corresponds to medial–lateral displacement and up reflects vertical motion; t indicates time. EE were quantified as the number of inertial actions performed per minute (n·min^{-1}), based on the detection of medium- and high-intensity inertial events such as accelerations, decelerations and changes in direction.

A two-step cluster analysis (average silhouette = 0.5) was applied to categorize drills according to biomechanical load demands, incorporating five input variables: jumps (JUMPS), accelerations (ACC), decelerations (DEC), player load (PL) and eccentric efforts (EE) (Table 4). Two biomechanical load clusters emerged—low (1) and high (2)—with drill-specific load profiles determined by averaging the cluster assignments across repetitions.

2.4. Statistical Analysis

To analyze differences in external load orientation (physiological or biomechanical), a Linear Mixed Model (LMM) was applied. The comparative framework positioned training drills against competitive matches rather than examining inter-drill differences. Training exposure included 14 practice configurations: individual technical-tactical work on half-court (1v0, 2v0), constrained-space games (1v1 in longitudinal half-court, 28 × 7.5 m),

balanced small-sided games across various formats (2v2, 3v3, 4v4, 5v5 on full-court), numerical advantage situations (3v0, 4v0, 5v0 full-court), continuous wave exercises (3v3v3, 4v4v4, 5v5v5) and eleven-player transition drills. Each task was treated as a categorical predictor in the model, while "player" was included as a random effect to account for repeated measures across individuals. Each of the 14 training situations was included as a categorical fixed effect, and 'player' was entered as a random effect to account for repeated measures across individuals. This structure allowed the model to control for within-player variability while estimating condition-related differences. Cohen's effect sizes and 95% confidence intervals were subsequently derived from the estimated marginal means.

Table 4. Clusters of training drills based on biomechanical load profiles.

Variables	Biomechanical Load	
	Low	**High**
Accelerations per minute	1.35	2.71
Decelerations per minute	1.20	3.38
Explosive efforts per minute	1.56	3.26
PlayerLoad per minute	5.91	8.42
Jumps per minute	0.65	0.73
Sample size (N)	128	2124
Sample proportion (%)	47.4%	67.7%
Bayesian Information Criterion (BIC)	10,677.49	
Average silhouette	0.5	

Note: Data for each external physiological load variable are presented as mean ± standard deviation, with sample sizes representing the number of tasks classified within each category.

Cohen's effect size (ES) and the mean differences with their corresponding 95% confidence intervals (CI) were calculated for each pairwise comparison. Effect sizes were interpreted according to the following qualitative thresholds: <0.20 trivial; 0.20–0.59 small; 0.60–1.19 moderate; 1.20–1.99 large; and ≥2.00 very large [51]. Statistical analyses were performed using IBM SPSS Statistics for Windows (version 23; IBM Corporation, Armonk, NY, USA), whereas ES values were derived using a customized Microsoft Excel spreadsheet (version 16.0; Microsoft Corporation, Redmond, WA, USA).

3. Results

Figure 1 graphically shows the classification of the different exercises analyzed according to their physical orientation and specificity. Regarding the comparison between training tasks for **external** physiological load (Figure 2), official games showed significantly higher values than 5v5v5 (ES: 2.07), 4v4v4 (ES: 2.04), 1v1 in longitudinal half-court 28 × 7.5 m-(ES: 1.62), 2v0 individual technical-tactical half-court (ES: 1.38) and 1v0 individual technical-tactical half-court (ES: 1.43); all of which represent large effect sizes (Cohen's d > 0.8). Regarding the comparison between training tasks for biomechanical load (Figure 3), official competition showed significantly higher values ($p < 0.05$) than 5v0 full-court (ES: 1.74), 4v0 full-court (ES: 2.07), 3v0 full-court (ES: 2.51), 2v0 individual technical-tactical half-court (ES: 2.47) and 1v0 individual technical-tactical half-court (ES: 2.09), all of which represent large effect sizes (Cohen's d > 0.8).




Figure 1. Classification of Drills According to Physical Orientation and Specificity. Note: The color scale indicates specificity as low (blue), medium (yellow) and high (red).




Figure 2. External Physiological load. Effect Size Estimates (Cohen's d) with 95% CI for Training–Match Differences in Load Orientation. Notes: The dotted line represents the effect size threshold from large to very large.




Figure 3. Biomechanical load. Effect size estimates (Cohen's d) with 95% CI for training–match differences in load orientation. Notes: The dotted line represents the effect size threshold from large to very large.

4. Discussion

The aim of the present study has been to compare training tasks with competition in elite basketball players from the perspective of physiological and biomechanical load experienced.

Previously, several studies have compared training and competition demands in the same group of players. The training analysis has been compared with official competition [23,24,52–55], with friendly competition [22,25,26] or with simulated competition contexts [27]. Moreover, some of the studies that have compared training with competition have not specified the type of tasks used in the practices, just the training load [20,21,25,27,54]. Only in the studies by Svilar et al. [26], in which 5v5 (regular-stops and no-stops) was analyzed, and by Torres-Ronda et al. [22], in which BDs (2v0, 3v0, 4v0 and 5v0), SSGs (1v1, 2v2, 3v3, 4v4) and SGs (5v5) were analyzed, the type of tasks has been specified, although in both studies the tasks were compared with friendly competition. To date, to our knowledge, there are no studies that have compared training tasks with official competition over several seasons in the same group of players.

4.1. Considerably Inferior to Competitive Benchmarks Across Biomechanical and Physiological Dimensions

Regarding non-opposition drills, it has been observed that official competition obtains higher values in biomechanical variables such as PL/min and explosive efforts/min than all non-opposition activities (1v0, 2v0, 3v0, 4v0 and 5v0). However, 1v0 and 2v0 tasks obtain more jumps/min than the competition and the rest of the tasks, data that coincides with the study of Torres-Ronda et al. [22], which suggests that when the work–rest ratio is

considered, non-opposition tasks have the highest density of actions. The fact that in 1v0 and 2v0 tasks there is a high density of jumps/min and low running values in all zones compared to competition, could be justified on the basis of the following: (a) in these tasks it is not necessary to perform certain actions such as dribbling or change of direction to progress, which limits the ability to accumulate other explosive actions; (b) these tasks are usually set at half-court, so there is no possibility to reach or stay long in high speed bands and accumulate distance and PL given the limited space available; and (c) they are usually tasks with a high technical orientation where a high number of repetitions are performed with little displacement, such as shooting drills. Although these drills allow for a high accumulation of jump-related load, caution is advised for players with patellar tendinopathy, ankle issues, or low tolerance to high jump volumes, as excessive jump density may be counterproductive.

However, when the competition is compared with BDs with a higher tactical orientation (3v0, 4v0 and 5v0), and that could be organized as full-court, no differences were observed in terms of total distance/min or in high intensity running zones (Runing zone 3 and 4), only significantly higher values were obtained for competition in zone 1 and 2 compared to 3v0, 4v0 and 5v0. These data are compatible with those obtained by Sosa et al. [21], in which it was observed that, during competition, 80.09% of the distance covered and 93.65% of the time is spent in zone 1 and 2. Therefore, 3v0, 4v0 and 5v0 accumulate distances in zone 3 and 4 similar to those of competition, but do not accumulate high biomechanical load. These tasks could serve as a valuable tool for improving physiological parameters with reduced risk of joint stress and injury due to the absence of external opposition and lower biomechanical loading requirements. Concurrently, these tasks represent ideal activities for implementation during the initial phase of sports injury rehabilitation, particularly for articular injuries, due to their low biomechanical requirements.

4.2. Progressing Toward Competitive Benchmarks Across Biomechanical and Physiological Dimensions

At specific times during the season, it is desirable that training mimics competition demands in terms of physical demands and technical-tactical specificity to the competition [21]. For this reason, the SSGs and SGs seem to be a tool that prepares players better than the BDs. While non-opposition tasks could generate a high density of actions, tasks with opposition have proven to be the most intense. The 1v1 half-court longitudinal 1v1 has shown a high biomechanical load, being the task that accumulates more explosive efforts/min, accelerations/min and decelerations/min, above any task and official competition. Our data coincide with another study [22] in which it was observed that 1v1 full-court represented the highest intensity ball exercise. However, these authors found that the next task in intensity was 2v2 half-court, whereas in the present study, this task is far from the biomechanical load competition values.

As with the non-opposition tasks, if we compare official competition with the set of SSGs, we observe that in total distance/min there are no significant differences with any task, but there are differences in the different zones, especially in zones 1 and 2, which are where more time is spent in competition [21].

To date, there are no studies that have compared SSGs with continuous opposition (3v3v3, 4v4v4 and 5v5v5) or Eleven Player Break with competition, so the data of the present investigation could not be compared with other studies. It can be observed that the tasks 4v4v4 and 5v5v5 accumulate significantly less external physiological load than official competition, and 3v3v3 accumulate significantly less distance than competition in zones 1, 2 and 3. In addition, there is a higher PL/min in competition compared to 4v4v4 and 5v5v5. These differences are aligned with the idea that with a lower m^2/player ratio, less external physiological load is accumulated [14], and given that PL has a high

correlation with distance traveled [56,57], it seems logical to think that in situations with similar opposition but with greater space, a greater load will be accumulated. For this reason, continuous opposition tasks could be an interesting tool to work on tactical aspects without accumulating a large external physiological load. On the other hand, the Eleven Player Break task accumulates less biomechanical load than competition but accumulates a similar external physiological load, so that it could be used with the objective of improving the athlete's conditioning, including opposition, and with a lower biomechanical load than competition.

The data obtained in this study are consistent with other studies [20,22,58], which reflect that competition is more intense than the rest SSGs and SGs; however, there are differences. Feu et al. [59] found that the external workload variables, such as distance covered, high-intensity displacements, accelerations and decelerations, tend to be lower in 5v5 in training sessions compared to competitive matches. Additionally, internal workload measures, such as heart rate, were higher during competition, reflecting greater physiological stress. Torres-Ronda [22] observed that the intensity, measured by mean HR and Peak HR reached, is higher in competition, 158 bpm-198 ppm, respectively, compared to the rest of SSGs (5v5, 4v4, 3v3 and 2v2). These authors found that 5v5 (152 ppm and 182 ppm, respectively) was more intense than 3v3 (145 ppm and 179 ppm, respectively) and 4v4 (143 ppm and 174 ppm, respectively), contrasting with the present study, in which it has been observed that official competition is the most intense activity, followed by 3v3, 4v4 and 5v5, respectively. As reflected by other authors [60], it has been found that 3v3 and 4v4 tasks do not differ significantly. Although the 3v3, 4v4 and 5v5 tasks have very similar biomechanical load values, the cumulative external physiological load is higher in 3v3, resembling that of official competition. The results of Montgomery et al. [58], in which they compared SSGs and SGs with friendly competition (Peak FC reached, % Maximal FC and VO_2Max), contrast with the present study because, although they found that friendly competition was more intense than any drill, the most intense drill was 5v5, as its demands most resembled those of competition. In fact, in their study, they reflected that the peak HR reached was similar between 5v5 tasks and friendly competition, although there was a greater time in % HR max in favor of competition. It seems logical to think, that since 5v5 tasks are the most similar to the game, they are the ones that generate load values closer to competition [14,26,27], but observing the results obtained in the present study, it seems that 3v3 and 4v4, when played in full-court, are the best option to achieve a high work intensity similar to that of official competition, especially biomechanical load, given the high frequency of movements. In this aspect, McCormick et al. [61] observed that the number of high-intensity actions in 3v3 is higher than in friendly competition (33 movements per minute), both in half-court (46 movements per minute) and full-court (53 movements per minute) situations. Thus, based on the data obtained in the present study, the 3v3 full-court task seems to be the best solution, both at the biomechanical and physiological levels. Although no tasks of 3v2, 4v3 or 5v4 have been analyzed in this study, it seems that these tasks may show higher demands and a greater similarity with friendly competition [22].

4.2.1. Contradictory Findings: When Training Exceeds Competition Intensity

In contrast to the data obtained in this study, there are authors who have observed that basketball training demands can exceed those of friendly competition. In their study, Svilar et al. [26] found that 5v5 no-stops elicited higher values of PL/min, decelerations/min, jumps/min, high intensity jumps/min and COD/min than friendly competition. However, when 5v5 was performed with the regular-stop game, it did not outperform friendly competition in any variable except accelerations/min. These results may be justi-

fied by the intentional intensification of the 5v5 when rest time during the task is eliminated. Similarly, Fox, Stanton, and Scanlan [25] found that external demands, both absolute and relative, were greater during pre-season training tasks compared to friendly competition. It seems logical that pre-season training load would be more demanding for the purpose of generating physiological adaptations than during the in-season [62,63]. The authors Fox, Stanton, and Scanlan [25] posit that it is possible that in-season training loads may not present higher workloads compared to competition, as the goal of training would reflect fitness maintenance rather than development, which could justify the results of their study that indicated that training was more intense than competition. These findings contrast with data from other authors [22], in which friendly matches showed a higher cardiovascular load than training. In the case of the study by Brandao et al. [27], in which no significant differences were observed between training and competition, four training sessions were monitored, and the competitive reference was a match simulation between components of the same team. These factors could cause the competition to drop in intensity compared with official competition. Observing these data, it seems that the moment of the season (pre-season/in-season) and schedule could modulate the training and competition loads [20,25].

4.2.2. Factors Influencing Load Variability Between Training and Competition

In this sense, there are several factors that can influence a match in relation to the increase in maximal HR values, such as duration, physical fitness of the athletes, motivation or the tactical approach of the teams [64]. Some authors [65] have also suggested that the exclusion of rest periods from the analyses may overestimate the external load in competition.

The available literature is unclear as to the variations that may exist during basketball training sessions. The inconsistency in results when comparing training with competition may derive from different factors that have to do with the intentional modulation of training tasks with the aim of increasing intensity and eliciting a greater number of movements, which increases cumulative Training Load [25]. This can be achieved through task design: court size, space restriction, number of participants, coach involvement or duration [34]; or through task orientation: tactically oriented tasks involve more interruptions for coach instructions and require lower intensity periods to learn systems and plays [23,66]. However, if the task objective has a greater physical, rather than tactical, developmental focus, higher relative intensity values appear to be obtained [25]. So, coaches should keep in mind that the excessive use of tactical tasks during the training session could compromise the physical fitness of the players, unless they are designed with a specific conditioning objective. In addition, it should be assessed that, if this type of exercise is characterized by a high tactical content, it could increase the cognitive load [22,46].

Other factors, not depending on the task design, that could modify the load are the monitoring methodology: different methods of filtering and data processing can modify the results [65,66]; or the time of the season: preseason training is typically characterized by higher workloads than those observed during the in-season period, as this phase is designed to elicit substantial physiological adaptations [67]. In contrast, in-season training generally does not exceed the demands imposed by competition, given that the primary objective during this period is the maintenance, rather than further development, of physiological capacities [25]. Differences could also be attributed to the fact that in competition there are prolonged and frequent stoppages, and the available playing time is restricted to only five players simultaneously [25].

4.3. Limitations of the Study

The limitations encountered in completing this study should be considered when interpreting our results. First, the sample size we employed was small, given that it was indicative of a single elite, junior, male basketball team. In turn, further research is needed to identify if our findings hold consistent in other teams across wider samples of basketball players encompassing male and female play. Undoubtedly, the analysis of training and competition is complex, and the values of a chosen variable may be different in one context or another, which could cause the results to vary between studies [21]. Secondly, before analyzing, we should reflect on how no single variable is determinant or capable of describing the complexity or demand of an activity; rather, it is the way in which the different variables interact that determines the nature or orientation of the task. As posited by Weaving et al. [68], one variable cannot show the significant differences shown by a set of variables. For this reason, it seems that a data collection adjusted to the needs and capabilities of the team context and sufficiently broad so as to allow a global view of the load received by the athlete should be performed. Therefore, it seems advisable to assess a set of variables systematically to increase the sensitivity of the changes. Future studies may integrate additional wearable sensors, such as leg-mounted inertial units or combined motion-analysis systems, to directly examine lower-limb biomechanics during cutting, landing and deceleration actions. This approach could provide valuable insights into joint-level mechanical demands and their potential relationship with common basketball injuries.

5. Conclusions

The results of this study provide practical insights into how different training tasks replicate the physical demands of competition. The following conclusions can help coaches better tailor task selection based on desired intensity and workload:

- Official competition consistently represents the most demanding context in both biomechanical and external physiological load, exceeding all training tasks analyzed.
- Non-opposition drills (e.g., 1v0 to 5v0) show limited transferability to the physical demands of competition, suggesting they may be insufficient to replicate match intensity.
- Among the training formats, the 3v3 task most closely mirrors the physiological and biomechanical demands of official competition.
- The 3v3 and 4v4 full-court tasks appear to be highly effective for eliciting elevated intensities, particularly in biomechanical load due to the high frequency of movements. These formats may be especially useful during training phases aimed at accumulating a high external workload.
- In small-sided games with constrained space and continuous opposition (e.g., 3v3v3, 4v4v4, 5v5v5), external physiological load is lower than in official matches, although biomechanical load remains comparable. This suggests these tasks may be well-suited for emphasizing technical-tactical development while minimizing excessive external load.
- Despite the greater overall load values observed in competition, a substantial portion of match time consists of low- or very low-intensity actions. This implies that decisive moments in competition are significantly more intense than what is typically elicited during training.
- The findings highlight the critical importance of players' abilities to accelerate, decelerate, change direction and jump, as these actions underpin performance in basketball's high-intensity, intermittent context.

Author Contributions: Conceptualization, C.S.M. and E.A.-P.-C.; methodology, C.S.M., E.A.-P.-C. and A.L.; formal analysis, C.S.M. and E.A.-P.-C.; investigation, C.S.M. and E.A.-P.-C.; data curation, C.S.M. and E.A.-P.-C.; writing—original draft preparation, C.S.M., E.A.-P.-C. and A.L.; writing—review and editing, A.L. and X.S.; supervision, A.L. and X.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The datasets presented in this article are not readily available because are the property of the basketball team in which they were recorded and are not publicly available. The authors do not have permission to share the data publicly. However, should anyone be interested in learning more, please contact the authors directly. Requests to access the datasets should be directed to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Baker, J.; Côté, J.; Abernethy, B. Learning from the experts: Practice activities of expert decision makers in sport. *Res. Q. Exerc. Sport* **2003**, *74*, 342–347. [CrossRef]
2. Bishop, D.C.; Wright, C.A. Time-motion analysis of professional basketball to determine the relationship between three activity profiles: High, medium and low intensity and the length of the time spent on court. *Int. J. Perform. Anal. Sport* **2006**, *6*, 130–139. [CrossRef]
3. Coyne, J.O.C.; Gregory Haff, G.; Coutts, A.J.; Newton, R.U.; Nimphius, S. The current state of subjective training load monitoring—A practical perspective and call to action. *Sports Med. Open* **2018**, *4*, 58. [CrossRef] [PubMed]
4. Scanlan, A.T.; Tucker, P.S.; Dascombe, B.J.; Berkelmans, D.M.; Hiskens, M.I.; Dalbo, V.J. Fluctuations in activity demands across game quarters in professional and semiprofessional male basketball. *J. Strength Cond. Res.* **2015**, *29*, 3006–3015. [CrossRef]
5. Aschendorf, P.F.; Zinner, C.; Delextrat, A.; Engelmeyer, E.; Mester, J. Effects of basketball-specific high-intensity interval training on aerobic performance and physical capacities in youth female basketball players. *Physician Sportsmed.* **2019**, *47*, 65–70. [CrossRef]
6. Maggioni, M.A.; Bonato, M.; Stahn, A.; Torre, A.L.; Agnello, L.; Vernillo, G.; Castagna, C.; Merati, G. Effects of ball drills and repeated-sprint-ability training in basketball players. *Int. J. Sports Physiol. Perform.* **2019**, *14*, 757–764. [CrossRef]
7. Stojanović, E.; Stojiljković, N.; Scanlan, A.T.; Dalbo, V.J.; Berkelmans, D.M.; Milanović, Z. The activity demands and physiological responses encountered during basketball match-play: A systematic review. *Sports Med.* **2018**, *48*, 111–135. [CrossRef]
8. Matthew, D.; Delextrat, A. Heart rate, blood lactate concentration, and time-motion analysis of female basketball players during competition. *J. Sports Sci.* **2009**, *27*, 813–821. [CrossRef]
9. McInnes, S.E.; Carlson, J.S.; Jones, C.J.; McKenna, M.J. The physiological load imposed on basketball players during competition. *J. Sports Sci.* **1995**, *13*, 387–397. [CrossRef] [PubMed]
10. Taylor, J.B.; Wright, A.A.; Dischiavi, S.L.; Townsend, M.A.; Marmon, A.R. Activity demands during multi-directional team sports: A systematic review. *Sports Med.* **2017**, *47*, 2533–2551. [CrossRef]
11. Doma, K.; Leicht, A.; Sinclair, W.; Schumann, M.; Damas, F.; Burt, D.; Woods, C. Impact of exercise-induced muscle damage on performance test outcomes in elite female basketball players. *J. Strength Cond. Res.* **2018**, *32*, 1731–1738. [CrossRef]
12. Edwards, T.; Spiteri, T.; Piggott, B.; Bonhotal, J.; Haff, G.G.; Joyce, C. Monitoring and managing fatigue in basketball. *Sports* **2018**, *6*, 19. [CrossRef]
13. Scanlan, A.; Dascombe, B.; Reaburn, P. A Comparison of the activity demands of elite and sub-elite australian men's basketball competition. *J. Sports Sci.* **2011**, *29*, 1153–1160. [CrossRef] [PubMed]
14. Sosa, C.; Alonso-Pérez-Chao, E.; Ribas, C.; Schelling, X.; Lorenzo, A. Description and classification of training drills, based on biomechanical and physiological load, in elite basketball. *Sensors* **2025**, *25*, 262. [CrossRef] [PubMed]
15. Robertson, S.; Joyce, D. Evaluating strategic periodisation in team sport. *J. Sports Sci.* **2018**, *36*, 279–285. [CrossRef]
16. Dawson, B.; Hopkinson, R.; Appleby, B.; Stewart, G.; Roberts, C. Comparison of training activities and game demands in the australian football league. *J. Sci. Med. Sport* **2004**, *7*, 292–301. [CrossRef]
17. Roberts, S.P.; Trewartha, G.; Higgitt, R.J.; El-Abd, J.; Stokes, K.A. The physical demands of elite English rugby union. *J. Sports Sci.* **2008**, *26*, 825–833. [CrossRef]
18. Smith, D.J. A framework for understanding the training process leading to elite performance. *Sports Med.* **2003**, *33*, 1103–1126. [CrossRef]

19. Champion, L.; Middleton, K.; MacMahon, C. Many pieces to the puzzle: A new holistic workload approach to designing practice in sports. *Sports Med. Open* **2023**, *9*, 38. [CrossRef]
20. Román, M.R.; García-Rubio, J.; Feu, S.; Ibáñez, S.J. Training and competition load monitoring and analysis of women's amateur basketball by playing position: Approach study. *Front. Psychol.* **2019**, *9*, 2689. [CrossRef] [PubMed]
21. Sosa, C.; Lorenzo, A.; Trapero, J.; Ribas, C.; Alonso, E.; Jimenez, S.L. Specific absolute velocity thresholds during male basketball games using local positional system; differences between age categories. *Appl. Sci.* **2021**, *11*, 4390. [CrossRef]
22. Torres-Ronda, L.; Ric, A.; Llabres-Torres, I.; de las Heras, B.; Schelling i del Alcazar, X. Position-dependent cardiovascular response and time-motion analysis during training drills and friendly matches in elite male basketball players. *J. Strength Cond. Res.* **2016**, *30*, 60–70. [CrossRef]
23. Fox, J.L.; O'Grady, C.J.; Scanlan, A.T. Game schedule congestion affects weekly workloads but not individual game demands in semi-professional basketball. *Biol. Sport* **2020**, *37*, 59–67. [CrossRef] [PubMed]
24. Fox, J.L.; Scanlan, A.T.; Stanton, R.; Sargent, C. The effect of game-related contextual factors on sleep in basketball players. *J. Strength Cond. Res.* **2022**, *36*, 2942–2945. [CrossRef] [PubMed]
25. Fox, J.L.; Stanton, R.; Scanlan, A.T. A comparison of training and competition demands in semiprofessional male basketball players. *Res. Q. Exerc. Sport* **2018**, *89*, 103–111. [CrossRef]
26. Svilar, L.; Castellano, J.; Jukic, I. Comparison of 5vs5 training games and match-play using microsensor technology in elite basketball. *J. Strength Cond. Res.* **2019**, *33*, 1897–1903. [CrossRef] [PubMed]
27. Brandão, F.M.; Junior, D.B.R.; da Cunha, V.F.; Meireles, G.B.; Filho, M.G.B. Differences between training and game loads in young basketball players. *Rev. Bras. Cineantropometria E Desempenho Hum.* **2019**, *21*, e59840. [CrossRef]
28. Conte, D.; Kolb, N.; Scanlan, A.T.; Santolamazza, F. Monitoring training load and well-being during the in-season phase in national collegiate athletic association division I men's basketball. *Int. J. Sports Physiol. Perform.* **2018**, *13*, 1067–1074. [CrossRef]
29. Peterson, K.D.; Quiggle, G.T. Tensiomyographical responses to accelerometer loads in female collegiate basketball players. *J. Sports Sci.* **2017**, *35*, 2334–2341. [CrossRef]
30. Weiss, K.J.; Allen, S.V.; McGuigan, M.R.; Whatman, C.S. The relationship between training load and injury in men's professional basketball. *Int. J. Sports Physiol. Perform.* **2017**, *12*, 1238–1242. [CrossRef]
31. Piedra, A.; Peña, J.; Ciavattini, V.; Caparrós, T. Relationship between injury risk, workload, and rate of perceived exertion in professional women's basketball. *Apunt. Sports Med.* **2020**, *55*, 71–79. [CrossRef]
32. Caparrós, T.; Casals, M.; Solana, Á.; Peña, J. External workloads are related to higher injury risk in professional male basketball games. *J. Sports Sci. Med.* **2018**, *17*, 289–297.
33. Ferioli, D.; La Torre, A.; Tibiletti, E.; Dotto, A.; Rampinini, E. Determining the relationship between load markers and non-contact injuries during the competitive season among professional and semi-professional basketball players. *Res. Sports Med.* **2021**, *29*, 265–276. [CrossRef]
34. Sosa Marín, C.; Alonso-Pérez-Chao, E.; Schelling, X.; Lorenzo, A. How to optimize training design? A narrative review of load modulators in basketball drills. *Appl. Sci.* **2025**, *15*, 3816. [CrossRef]
35. Sansone, P.; Gasperi, L.; Makivic, B.; Gomez-Ruano, M.; Tessitore, A.; Conte, D. An ecological investigation of average and peak external load intensities of basketball skills and game-based training drills. *Biol. Sport* **2023**, *40*, 649–656. [CrossRef] [PubMed]
36. Russell, J.L.; McLean, B.D.; Impellizzeri, F.M.; Strack, D.S.; Coutts, A.J. Measuring physical demands in basketball: An explorative systematic review of practices. *Sports Med.* **2021**, *51*, 81–112. [CrossRef]
37. Vanrenterghem, J.; Nedergaard, N.J.; Robinson, M.A.; Drust, B. Training load monitoring in team sports: A novel framework separating physiological and biomechanical load-adaptation pathways. *Sports Med.* **2017**, *47*, 2135–2142. [CrossRef]
38. McKay, A.K.A.; Stellingwerff, T.; Smith, E.S.; Martin, D.T.; Mujika, I.; Goosey-Tolfrey, V.L.; Sheppard, J.; Burke, L.M. Defining training and performance caliber: A participant classification framework. *Int. J. Sports Physiol. Perform.* **2022**, *17*, 317–331. [CrossRef] [PubMed]
39. Harriss, D.J.; Atkinson, G. Ethical standards in sport and exercise science research: 2014 Update. *Int. J. Sports Med.* **2014**, *34*, 1025–1029. [CrossRef] [PubMed]
40. Hodder, R.W.; Ball, K.A.; Serpiello, F.R. Criterion validity of catapult clearsky t6 local positioning system for measuring inter-unit distance. *Sensors* **2020**, *20*, 3693. [CrossRef]
41. Luteberget, L.; Spencer, M.; Gilgien, M. Validity of the catapult clearsky t6 local positioning system for team sports specific drills, in indoor conditions. *Front. Physiol.* **2018**, *9*, 115. [CrossRef]
42. Serpiello, F.R.; Hopkins, W.G.; Barnes, S.; Tavrou, J.; Duthie, G.M.; Aughey, R.J.; Ball, K. Validity of an ultra-wideband local positioning system to measure locomotion in indoor sports. *J. Sports Sci.* **2018**, *36*, 1727–1733. [CrossRef] [PubMed]
43. Hoppe, M.W.; Baumgart, C.; Polglaze, T.; Freiwald, J. Validity and reliability of gps and lps for measuring distances covered and sprint mechanical properties in team sports. *PLoS ONE* **2018**, *13*, e0192708. [CrossRef] [PubMed]
44. Luteberget, L.; Holme, B.; Spencer, M. Reliability of wearable inertial measurement units to measure physical activity in team handball. *Int. J. Sports Physiol. Perform.* **2017**, *13*, 467–473. [CrossRef]

45. Castellano, J.; Casamichana, D.; Calleja-González, J.; Román, J.S.; Ostojic, S. Reliability and accuracy of 10 hz gps devices for short-distance exercise. *J. Sports Sci. Med.* **2011**, *10*, 233–234.
46. Schelling, X.; Torres, L. Accelerometer load profiles for basketball-specific drills in elite players. *J. Sports Sci. Med.* **2016**, *15*, 585–591.
47. Alonso Pérez-Chao, E.; Gómez, M.Á.; Lisboa, P.; Trapero, J.; Jiménez, S.L.; Lorenzo, A. Fluctuations in external peak demands across quarters during basketball games. *Front. Psychol.* **2022**, *13*, 868009. [CrossRef] [PubMed]
48. Alonso Perez-Chao, E.; Lorenzo, A.; Scanlan, A.; Lisboa, P.; Sosa, C.; Gómez, M.A. Higher playing times accumulated across entire games and prior to intense passages reduce the peak demands reached by elite, junior, male basketball players. *Am. J. Mens. Health* **2021**, *15*, 15579883211054353. [CrossRef]
49. Svilar, L.; Castellano, J.; Jukic, I. Load monitoring system in top-level basketball team: Relationship between external and internal training load. *Kinesiology* **2018**, *50*, 25–33. [CrossRef]
50. Alonso, E.; Miranda, N.; Zhang, S.; Sosa, C.; Trapero, J.; Lorenzo, J.; Lorenzo, A. Peak match demands in young basketball players: Approach and applications. *Int. J. Environ. Res. Public Health* **2020**, *17*, 2256. [CrossRef]
51. Hopkins, W.; Marshall, S.; Batterham, A.; Hanin, J. Progressive statistics for studies in sports medicine and exercise science. *Med. Sci. Sports Exerc.* **2009**, *41*, 3–12. [CrossRef] [PubMed]
52. Reina, M.; García-Rubio, J.; Pino-Ortega, J.; Ibáñez, S.J. The acceleration and deceleration profiles of U-18 women's basketball players during competitive matches. *Sports* **2019**, *7*, 165. [CrossRef] [PubMed]
53. Williams, M.N.C.; Dalbo, V.J.; Fox, J.L.; O'Grady, C.J.; Scanlan, A.T. Comparing weekly training and game demands according to playing position in a semiprofessional basketball team. *Int. J. Sports Physiol. Perform.* **2021**, *16*, 772–778. [CrossRef] [PubMed]
54. Fox, J.L.; Stanton, R.; O'Grady, C.J.; Teramoto, M.; Sargent, C.; Scanlan, A.T. Are acute player workloads associated with in-game performance in basketball? *Bio. Sport* **2022**, *39*, 95–100. [CrossRef]
55. Williams, M.N.C.; Fox, J.L.; O'Grady, C.J.; Gardner, S.; Dalbo, V.J.; Scanlan, A.T. Weekly training demands increase, but game demands remain consistent across early, middle, and late phases of the regular season in semiprofessional basketball players. *Int. J. Sports Physiol. Perform.* **2022**, *17*, 350–357. [CrossRef]
56. Bredt, S.D.G.T.; Chagas, M.H.; Peixoto, G.H.; Menzel, H.J.; Andrade, A.G.P. Understanding player load: Meanings and limitations. *J. Hum. Kinet.* **2020**, *71*, 5–9. [CrossRef]
57. Heishman, A.D.; Daub, B.D.; Miller, R.M.; Freitas, E.D.S.; Bemben, M.G. Monitoring External Training Loads and Neuromuscular Performance for Division I Basketball Players over the Preseason. *J. Sports Sci. Med.* **2020**, *19*, 204–212.
58. Montgomery, P.G.; Pyne, D.B.; Minahan, C.L. The physical and physiological demands of basketball training and competition. *Int. J. Sports Physiol. Perform.* **2010**, *5*, 75–86. [CrossRef]
59. Feu, S.; García-Ceberino, J.M.; López-Sierra, P.; Ibáñez, S.J. Training to compete: Are basketball training loads similar to competition achieved? *Appl. Sci.* **2023**, *13*, 12512. [CrossRef]
60. Sampaio, J.; Abrantes, C.; Leite, N. Power, heart rate and perceived exertion responses to 3 × 3 and 4 × 4 basketball small-sided games. *Rev. De Psicol. Del Deporte* **2009**, *18*, 463–467.
61. Mccormick, B.T.; Hannon, J.C.; Newton, M.; Shultz, B.; Miller, N.; Young, W.; Arias, J.; Jenkins, D. Comparison of physical activity in small-sided basketball games versus full-sided games. *Int. J. Sports Sci. Coach.* **2012**, *7*, 689–697. [CrossRef]
62. Dupont, G.; Akakpo, K.; Berthoin, S. The effect of in-season, high-intensity interval training in soccer players. *J. Strength Cond. Res.* **2004**, *18*, 584–589. [PubMed]
63. Clark, N.A.; Edwards, A.M.; Morton, R.H.; Butterly, R.J. Season-to-season variations of physiological fitness within a squad of professional male soccer players. *J. Sports Sci. Med.* **2008**, *7*, 157–165.
64. Ben Abdelkrim, N.; Castagna, C.; Jabri, I.; Battikh, T.; El Fazaa, S.; Ati, J. El Activity profile and physiological requirements of junior elite basketball players in relation to aerobic-anaerobic fitness. *J. Strength Cond. Res.* **2010**, *24*, 2330–2342. [CrossRef]
65. O'grady, C.J.; Dalbo, V.J.; Teramoto, M.; Fox, J.L.; Scanlan, A.T. External Workload Can be anticipated during 5 vs. 5 games-based drills in basketball players: An exploratory study. *Int. J. Environ. Res. Public Health* **2020**, *17*, 2103. [CrossRef]
66. Malone, S.; Owen, A.; Newton, M.; Mendes, B.; Collins, K.D.; Gabbett, T.J. The acute:chonic workload ratio in relation to injury risk in professional soccer. *J. Sci. Med. Sport* **2017**, *20*, 561–565. [CrossRef] [PubMed]
67. Narazaki, K.; Berg, K.; Stergiou, N.; Chen, B. Physiological demands of competitive basketball. *Scand. J. Med. Sci. Sports* **2009**, *19*, 425–432. [CrossRef]
68. Weaving, D.; Jones, B.; Till, K.; Abt, G.; Beggs, C. The case for adopting a multivariate approach to optimize training load quantification in team sports. *Front. Physiol.* **2017**, *8*, 1024. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.