

Emotional Components of Pain Perception in Borderline Personality Disorder and Major Depression – a repetitive peripheral magnetic stimulation (rPMS) study

¹Kathrin Malejko, ²André Huss, ¹Carlos Schönfeldt-Lecuona, ¹Maren Braun, ¹Heiko Graf

¹Department of Psychiatry and Psychotherapy III, University of Ulm, Germany

²Department of Neurology, University of Ulm, Germany

Table S1: Characteristics of patients with Borderline personality disorder (BPD) and comorbid major depression, patients with major depressive disorder (MD) without BPD and healthy controls (HC).

		BPD (n=10)	MD (n=12)	HC (n=12)
Age of onset NSSI:				
	< 20 years	7	0	0
	> 20 years	3	0	0
Years of NSSI:				
	< 5 years	3	0	0
	> 5 years	7	0	0
Frequency of NSSI:				
	Daily	2	0	0
	Weekly	8	0	0
Substance abuse history:				
	Alcohol	1	1	0
	Psychotropic drugs	1	0	0
Medication:				
	Antidepressants	8	12	0
	Mood stabilizers	5	0	0
	Antiepileptic drugs	5	0	0
	Antipsychotic drugs	5	4	0
	Antihypertensive drugs	3	0	0
	Benzodiazepines	0	4	0
Dissociation Tension scale (mean±SD)	Before rPMS	2.80±2.07	1.88±1.59	n.a.
	After rPMS	2.35±2.23	1.80±1.86	n.a.
BDI (mean±SD)		39.56±12.20	29.67±11.23	n.a.

NSSI=nonsuicidal self-injury; SD=standard deviation; n.a.=not assessed