



Article

Influence of the Ripening Stage and Extraction Conditions on the Phenolic Fingerprint of 'Corbella' Extra-Virgin Olive Oil

Anallely López-Yerena ^{1,†}, Antonia Ninot ^{2,†}, Núria Jiménez-Ruiz ¹, Julián Lozano-Castellón ^{1,3},
Maria Pérez ^{1,4}, Elvira Escribano-Ferrer ^{3,5}, Agustí Romero-Aroca ², Rosa M. Lamuela-Raventós ^{1,3}
and Anna Vallverdú-Queralt ^{1,3,*}

¹ Department of Nutrition, Food Science and Gastronomy, XIA, Faculty of Pharmacy and Food Sciences, Institute of Nutrition and Food Safety (INSA-UB), University of Barcelona, 08028 Barcelona, Spain; naye.yerena@gmail.com (A.L.-Y.); nuriajimenru@gmail.com (N.J.-R.); julian.lozano@ub.edu (J.L.-C.); mariaperez@ub.edu (M.P.); lamuela@ub.edu (R.M.L.-R.)

² Institute of Agrifood Research and Technology (IRTA), Fruit Science Program, Olive Growing and Oil Technology Research Team, 43120 Constantí, Spain; Antonia.Ninot@irta.cat (A.N.); agusti.romero@irta.cat (A.R.-A.)

³ CIBER Physiopathology of Obesity and Nutrition (CIBEROBN), Institute of Health Carlos III, 28029 Madrid, Spain; escribano@ub.edu

⁴ Laboratory of Organic Chemistry, Faculty of Pharmacy and Food Sciences, University of Barcelona, 08028 Barcelona, Spain

⁵ Pharmaceutical Nanotechnology Group I+D+I Associated Unit to CSIC, Biopharmaceutics and Pharmacokinetics Unit, Department of Pharmacy and Pharmaceutical Technology and Physical Chemistry, Institute of Nanoscience and Nanotechnology (IN2UB), Faculty of Pharmacy and Food Sciences, University of Barcelona, 08028 Barcelona, Spain

* Correspondence: avallverdu@ub.edu; Tel.: +34-934-024-508

† These authors equally contributed to the work.



Citation: López-Yerena, A.; Ninot, A.; Jiménez-Ruiz, N.; Lozano-Castellón, J.; Pérez, M.; Escribano-Ferrer, E.; Romero-Aroca, A.; Lamuela-Raventós, R.M.; Vallverdú-Queralt, A. Influence of the Ripening Stage and Extraction Conditions on the Phenolic Fingerprint of 'Corbella' Extra-Virgin Olive Oil. *Antioxidants* **2021**, *10*, 877. <https://doi.org/10.3390/antiox10060877>

Academic Editors: Monica Rosa Loizzo and Rosa Tundis

Received: 6 May 2021

Accepted: 26 May 2021

Published: 30 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The ancient 'Corbella' olive variety from the center-north of Catalonia is being recovered to obtain quality extra-virgin olive oil (EVOO) with unique organoleptic properties. The aim of this work was to determine the effect of agronomic and technical factors on the phenolic fingerprint of EVOO and to establish the optimum harvesting time and crushing and malaxation conditions for 'Corbella' olives. Therefore, three different ripening indices (0.3, 1.2, and 3.2) and three crushing temperatures (10, 18, and 25 °C) were studied. Additionally, a factorial design to optimize the phenolic concentration of the EVOO was developed, applying a range of sieve diameters (4 and 6 mm), and malaxation time (30 and 60 min) and temperature (27, 32, and 37 °C). The phenolic profile was analyzed by ultra-high performance liquid chromatography coupled to mass spectrometry in a tandem detector. The level of secoiridoids, the major phenolic compounds in the oil, was higher when using olives harvested earlier. Oleuropein aglycone and ligstroside aglycone were degraded during crushing at high temperatures, resulting in the formation of oleacein and oleocanthal. The best processing conditions in terms of total phenolic content were found to be 30 min of malaxation at 37 °C, the crushing size not having any affect.

Keywords: polyphenols; malaxation; crushing size; oleocanthal; oleacein

1. Introduction

The demand for extra virgin olive oil (EVOO) is increasing due to growing consumer awareness of the health benefits of the Mediterranean diet, in which it is the main source of dietary fat [1]. The nutritional and health-promoting properties of EVOO [2,3] are mainly correlated with its highly bioactive components, such as monounsaturated fatty acids (55–83%), unsaponifiable compounds (1–2%), and soluble or hydrophilic compounds, including α -tocopherol, phenolic compounds, and other compounds with antioxidant properties [4]. In 2012, the European Food Safety Authority (EFSA) recognized that a daily

consumption of 5 mg hydroxytyrosol and/or its derivatives (e.g., the oleuropein complex and tyrosol) per 20 g of olive oil afforded protection of LDL particles from oxidative damage and maintenance of a normal blood HDL-cholesterol concentration [5].

Since this recognition, the scientific community and olive oil industry have focused considerable efforts on optimizing the EVOO extraction process, searching for a balance between yield and quality [6]. To date, many studies have shown that the phenolic composition of EVOO depends on a highly complex multivariate interaction between agronomic, biochemical, and technological factors [7–10]. Firstly, the composition of the olives depends on their genetic origin, the geographical area of cultivation and its climate, and agronomic practices such as fertilization and water availability [11,12]. Then, the parameters associated with EVOO quality can be controlled by regulating the main variables of mechanical extraction (temperature, oxygen, enzymatic activities, and the disruption level of olive cell tissues) [13,14]. In this context, crushing is a determinant factor in the release and activation of endogenous enzymes (β -glycosidases and oxidoreductases), whereas malaxation is crucial for the modulation of the enzymatic activity [14–17] and production of high quality EVOO. During malaxation, the chemical composition of the oil undergoes considerable changes due to partition phenomena between oil and water, as well as reactions catalyzed by the fruit enzymes released during crushing [13].

Given the above, the main novelty of this work lies in the study of both agronomic and technological variables to obtain EVOO with a high content of phenolic compounds. For this purpose, the objectives of this study were (i) to determine the effect of three different ripening indices (RI, 0.3, 1.2, and 3.2) on the phenolic profile of EVOO, (ii) to evaluate changes in polyphenol content of EVOO obtained at three crushing temperatures (10, 18, and 25 °C), and (iii) to optimize the malaxation parameters (time and temperature) and crushing size to obtain an EVOO with higher phenolic compounds content. Regarding this last objective, a factorial design was developed to optimize the time (30 and 60 min) and temperature (27, 32, and 37 °C) of malaxation, applying two sieve diameters (4 and 6 mm). The study was carried out with olives from the ‘Corbella’ cultivar, native to Bages in the center-north of Catalonia, an ancient variety that was abandoned and at risk of extinction. EVOO produced exclusively with ‘Corbella’ olives has a powerful, fruity, and balanced taste. In recent years, centenary ‘Corbella’ olive trees have been recovered to promote the cultivation of this autochthonous variety and offer the market ecological products with added value.

2. Materials and Methods

2.1. Chemicals and Reagents

Regarding the secoiridoids, oleocanthal ($\geq 95\%$ purity) was purchased from Merck (Darmstadt, Germany), while oleacein and oleuropein aglycone (≥ 90 and 95% purity, respectively) from Toronto Research Chemical Inc. (ON, Canada). Luteolin was acquired from Sigma-Aldrich (Madrid, Spain), ferulic acid and apigenin from Fluka (Buchs, Switzerland), and hydroxytyrosol from Extrasynthese (Genay, France). Hexane, methanol, and acetonitrile were purchased from Sigma-Aldrich. Ultrapure water was obtained using a Milli-Q purification system (Millipore, Bedford, MA, USA).

2.2. Plant Material

Olive fruits were hand collected from mature olive trees of the ‘Corbella’ cultivar in the Oil Migjorn farm in Bages, Barcelona Province, Spain. The olives used to study the crushing temperature, crushing size, and malaxation were cultivated under controlled deficit irrigation conditions, whereas those used for the ripening study were obtained from non-irrigated trees. On the day of harvesting, the olives were sent to the IRTA laboratories in Constantí, Tarragona, where, after a health assessment, they were treated as described below.

2.3. Oil Extraction

All the fruit samples were processed using an Abencor analyser (Abengoa S.A., Seville, Spain), which reproduces the industrial process on a laboratory scale. The fruits were milled using a stainless-steel rotatory hammer mill operating at 3000 rpm and equipped with different sieves, depending on the experiment. The resulting olive paste was immediately passed into a malaxation unit with blades spinning at 50 rpm. The time and temperature of this step were set depending on each experiment. A basket centrifuge (3000 rpm for 3 min) was used to centrifuge the kneaded olive paste without addition of hot water or talcum. After centrifugation, the oil was decanted and stored in amber glass bottles at $-20\text{ }^{\circ}\text{C}$ in darkness and without headspace until analysis.

2.3.1. Ripening Study

The fruits were collected at 3 different stages of maturity and, following the method described by Uceda and Frías [18], the 3 RI were calculated to be 0–1, 1–2, and 2–3. For each RI, the olives were crushed in a 5 mm diameter sieve, and the olive paste was placed in a $28\text{ }^{\circ}\text{C}$ malaxator for 30 min without the addition of talcum powder or hot water. The centrifugation and storage conditions were as described above (Section 2.2).

2.3.2. Crushing Temperature Study

Healthy olives were stored for two hours in a fridge ($7\text{ }^{\circ}\text{C}$) and then washed using clean water heated to three different temperatures: (10, 18, and $25\text{ }^{\circ}\text{C}$) in a water bath. Once the desired temperature was reached, fruits were crushed in a 5 mm diameter sieve, and the resulting paste was placed in a malaxator at $18\text{--}19\text{ }^{\circ}\text{C}$ for 30 min without adding either talcum or water. It was then centrifuged at 3000 rpm for 3 min without adding water. The EVOO obtained was separated and stored under the conditions described above (Section 2.2).

2.3.3. Crushing Size and Malaxation Study

The olive oil was extracted following a factorial model with three factors: crushing size (4 and 6 mm), malaxation temperature (27, 32, and $37\text{ }^{\circ}\text{C}$), and malaxation time (30 and 60 min). The trials were performed using a 3.3 kg batch of drupes. Three representative olive samples, each weighing a minimum of one kilogram, were processed, and the corresponding EVOOs were obtained using an Abencor analyser (Abengoa S.A., Seville, Spain). The olive fruits were milled using a stainless-steel hammer mill equipped with a 4- and 6-mm sieve that was operated at 3000 rpm. The resulting olive paste was immediately kneaded in a mixer at 50 rpm for 30 or 60 min at 27, 32, or $37\text{ }^{\circ}\text{C}$. The centrifugation and storage conditions were as described above (Section 2.2).

2.4. Phenolic Compound Composition

After a liquid–liquid extraction, the identification and quantification of phenolic compounds were carried out by liquid chromatography coupled to mass spectrometry in tandem mode (LC-MS/MS), as described in our previous studies [11,19]. Samples were analyzed using an AcquityTM UPLC (Waters; Milford, MA, EUA) system with an API 3000 triple-quadrupole mass spectrometer (PE Sciex, Concord, Ontario, Canada). The system was equipped with a turbo ion spray source. Separation of phenolic compounds was conducted using an Acquity UPLC[®] BEH C₁₈ column (2.1 × 50 mm, i.d., 1.7 μm particle size) and Acquity UPLC[®] BEH C₁₈ Pre-Column (2.1 × 5 mm, i.d., 1.7 μm particle size) (Waters Corporation[®], Wexford, Ireland).

The liquid–liquid extraction was carried out following procedures described previously [19]. For this purpose, 0.5 g of EVOO was dissolved in hexane (oil/hexane 1:2, *w/v*) in a 10 mL centrifuge tube and shaken for 1 min. For the phenolic compound extraction, 2 mL of MeOH: H₂O (4:1 *v/v*) was added and the samples were homogenized for 1 min. The two phases were separated by centrifugation (3000 rpm, $4\text{ }^{\circ}\text{C}$, 4 min). The methanolic fraction was separated and subjected to a second cleaning using hexane, while the remain-

ing phenolic compounds in the hexane fraction were extracted by a further addition of MeOH: H₂O (4:1 *v/v*). The extracts were shaken and centrifuged under the same conditions as previously described. The two methanolic fractions were pooled and reduced to dryness with N₂, redissolved in 800 µL of ACN, and stored at −80 °C prior to analysis.

The separation of oleocanthal, oleacein, ligstroside, and oleuropein aglycone was performed following the procedure proposed by Lozano–Castellon [19]. MeOH (A) and H₂O (B) were used as mobile phases, both with 0.1% of formic acid. An increasing linear gradient (*v/v*) of B was used (t (min), %B), as follows: (0, 100); (2, 100); (4.75, 46.4); (4.9, 100); (5.9, 0); (6, 100); (6.5, 100), at 0.6 mL/min, 5 µL, and 50 °C of constant flow rate, injection volume, and column temperature, respectively. The separation of other polyphenols was achieved following procedures described previously [11] using H₂O with 0.2% acetic acid (A) and ACN (B) as mobile phases. An increasing linear gradient (*v/v*) of B was used (t (min), %B), as follows: (0, 5); (2.5, 5); (12.5, 40); (12.6, 100); (13.5, 100); (13.6, 5); (15, 5), at 0.4 mL/min, 5 µL, and 40 °C of constant flow rate, injection volume, and column temperature, respectively.

The ionization was achieved using an electrospray (ESI) interface operating in negative ionization mode (M[−]H), and the compounds were monitored in the multiple monitoring mode (MRM). The system was controlled by Analyst version 1.4.2 software supplied by Applied Biosystems. The calibration curves were prepared in refined olive oil and were linear over the concentration range 0–20 mg/mL using hydroxytyrosol, apigenin, luteolin, oleocanthal, oleacein, and oleuropein aglycone as standards.

2.5. Statistical Analyses

The normality of the data was tested by the Q–Q plot diagnostic graphic. After viewing the results, all subsequent tests were non-parametric. In the ripening and crushing study, to assess differences between groups, a Kruskal–Wallis test was performed, and the post-hoc test was the pairwise Mann–Whitney U-test. To investigate the main effects of the investigated factors, as well as to determine possible interactions between them, phenolic concentration analysis data were subjected to three-way analysis of variance (ANOVA), with 2 crushing size, malaxation temperature, and malaxation duration as factors (3 malaxation temperatures × 2 malaxation durations × 2 crushing size × 3 replicates). Means were compared by least significant difference (LSD) test, at the level of $p < 0.05$. All the statistical analyses were performed with R Project for Statistical Computing 3.6.0.

3. Results and Discussion

3.1. Ripening Study

The effects of ripening on the phenolic concentrations are shown in Figure 1. The total phenolic content was affected negatively by ripening stage, decreasing up to 49% at 3.2 of RI (from 566.74 ± 51.45 to 289.48 ± 17.43 mg·kg^{−1}) ($p \leq 0.05$). The same behaviour was observed for the secoiridoids' and phenolic alcohols; concentration. In contrast, the maximum concentration of flavones and lignans was reached at 3.2 of RI, whereas the highest phenolic acid content was obtained with olives from the second harvest with an RI of 1.2 (2.16 ± 0.06 mg·kg^{−1}).

The RI of the olive fruits affected the total amount of phenolic compounds in the EVOO, which was 43.64% lower ($p \leq 0.05$) when produced from the third harvest. This could be caused by endogenous enzymes in the olives, such as esterases, polyphenol oxidases, and β-glucosidases, which degrade phenols during the maturation process [13,14,16]. These results are in agreement with the literature [16,20–22], including a study carried out with 'Frantoio' and 'Manzanilla' EVOO, in which a gradual decrease in phenolic compounds was observed from the first to the fifth harvest [20]. In another study, however, this tendency did not begin below an RI of 2.5–3 [16]. It has also been suggested that the phenolic content is more influenced by the cultivar than the degree of fruit maturity [23]. It should be mentioned that the fruit of some olive cultivars ripens at an uneven rate, and therefore the

same harvest will contain olives of variable ripeness [24], rendering the RI a poor descriptor for analysis. The ‘Corbella’ cultivar exhibits a very uniform ripening pattern [25].




Figure 1. Effect of the ripening index on the total phenolic content and the phenolic composition of EVOO. TP: total phenols; SEC: secoiridoids. * indicates significant differences between phenolic concentration ($p < 0.05$).

The RI effect on the concentrations of individual phenolic compounds is shown in Table 1. The concentration of secoiridoids, the prevalent phenolic group found in EVOO and associated with many of its benefits [26], was greatly affected by the harvest time, as in other studies [13,27,28]. All the secoiridoids decreased as the RI increased, except for hydroxyoleuropein aglycone. The explanation may be that the enzymatic degradation of oleuropein and ligstroside results in a higher concentration of the aglycone.

Table 1. Phenolic compound contents ($\text{mg} \cdot \text{kg}^{-1}$) of EVOO produced from ‘Corbella’ olives with different crushing temperatures.

| Group/Compound | Ripening Index | | |
|----------------------------|-----------------------------|-----------------------------|-----------------------------|
| | 0.3 ± 0.07 | 1.2 ± 0.17 | 3.2 ± 0.27 |
| Total Polyphenols | 566.74 ± 51.45 ^a | 483.72 ± 26.65 ^b | 289.48 ± 17.43 ^c |
| Secoiridoids | 556.81 ± 51.45 ^a | 473.07 ± 26.17 ^b | 275.72 ± 17.43 ^c |
| Elenolic acid | 272.12 ± 9.71 ^a | 228.82 ± 16.30 ^b | 125.52 ± 9.89 ^c |
| Hydroxyelenolic acid | 8.68 ± 0.38 ^a | 8.79 ± 0.29 ^a | 4.32 ± 0.21 ^b |
| Hydroxyoleuropein aglycone | 1.11 ± 0.09 ^b | 1.16 ± 0.07 ^b | 2.03 ± 0.07 ^a |
| Ligstroside aglycone | 15.07 ± 0.84 ^a | 11.92 ± 0.47 ^b | 9.00 ± 0.51 ^c |
| Methyloleuropein aglycone | 0.92 ± 0.04 ^a | 0.88 ± 0.02 ^a | 0.87 ± 0.02 ^a |
| Oleacein | 94.25 ± 6.29 ^a | 78.72 ± 4.48 ^b | 34.06 ± 2.06 ^c |
| Oleocanthal | 9.14 ± 1.20 ^a | 6.84 ± 0.31 ^b | 2.36 ± 0.21 ^c |
| Oleuropein aglycone | 155.51 ± 5.71 ^a | 135.95 ± 2.98 ^a | 97.56 ± 3.75 ^b |
| Phenolic alcohols | 1.34 ± 0.21 ^a | 0.78 ± 0.11 ^b | 0.66 ± 0.04 ^c |
| Hydroxytyrosol | 1.27 ± 0.08 ^a | 0.68 ± 0.06 ^b | 0.46 ± 0.05 ^c |
| Hydroxytyrosol acetate | 0.08 ± 0.02 ^b | 0.11 ± 0.02 ^b | 0.21 ± 0.02 ^a |
| Flavones | 0.90 ± 0.09 ^c | 1.15 ± 0.08 ^b | 2.92 ± 0.41 ^a |
| Apigenin | 0.49 ± 0.01 ^c | 0.82 ± 0.06 ^b | 2.23 ± 0.2 ^a |
| Luteolin | 0.41 ± 0.02 ^b | 0.32 ± 0.04 ^b | 0.69 ± 0.04 ^a |
| Phenolic acids | 2.12 ± 0.32 ^{ab} | 2.16 ± 0.06 ^a | 1.93 ± 0.15 ^b |
| <i>m</i> -coumaric acid | 1.53 ± 0.04 ^a | 1.47 ± 0.03 ^a | 1.42 ± 0.03 ^b |
| <i>p</i> -coumaric acid | 0.59 ± 0.05 ^{ab} | 0.69 ± 0.04 ^a | 0.50 ± 0.04 ^b |
| Lignans | 5.57 ± 0.76 ^b | 6.56 ± 0.75 ^b | 8.24 ± 0.85 ^a |
| Pinosresinol | 5.57 ± 0.76 ^b | 6.56 ± 0.75 ^b | 8.24 ± 0.85 ^a |

Different letters within the same row mean significant differences (p -value < 0.05) according to the Kruskal–Wallis test.

Among the phenolic alcohols, hydroxytyrosol decreased with maturation. In contrast, hydroxytyrosol acetate increased, probably because it can form part of more complex compounds such as oleacein, oleuropein, and verbascoside, and is released when they are

broken down [29]. Due to its powerful radical scavenging activity, hydroxytyrosol acetate plays an important role in the anti-inflammatory effect of EVOO and is able to downregulate cyclooxygenase-2 expression [30]. In the case of hydroxytyrosol, which is also a degradation product, its reduced concentration could be caused by its higher lability. This effect is of interest, as according to EFSA, the consumption of at least 5 mg of hydroxytyrosol and its derivatives is required to protect LDL particles from oxidative damage [5]. Therefore, to preserve the health benefits of EVOO, a higher RI is not recommended, as the resulting reduction of hydroxytyrosol far exceeds the increase in the acetate derivative.

In the case of flavones, an upward trend was observed, as in other studies [31,32]. Bengana et al., 2013 [33] reported an increase in the flavonoid concentration in EVOO produced from olives in the first period of maturation, although levels began to decline when the harvest was in December. This tendency was not observed in our work, as all the olives were collected before the end of October. The glycosidic forms of flavones found in the drupelet become more soluble in oil during ripening, when the bond with the sugar is broken down by β -glucosidase. Thus, with flavones in oil being retained to a higher degree than in water during the extraction, oils from late season olives have a higher flavone content. However, other studies report that while luteolin increased, apigenin levels remained unchanged in 'Hojiblanca', 'Picual', and 'Picudo' oil varieties [34] or even decreased in the 'Arbequina' variety [35]. Such variable maturation behaviour can be attributed to differences in composition and endogenous enzymes among varieties [23,36]. Additionally, accurate comparisons of results from different studies are not always possible, as some refer only to the harvest date but not the RI [23].

The level of phenolic acids showed a slight decrease from the first to the second RI, as reported elsewhere [32]. In contrast, other studies have found an increase, associated with the activity of hydrolytic enzymes in the drupelet [28,37].

Among the lignans, pinoresinol increased by approximately 40.75% over the three ripening stages, possibly because of the degradation of the coexisting acetoxypinoresinol [38], whose levels were below the detection limit. Pinoresinol is a very stable compound and is preserved during ripening. Lignans are reported to have an inhibitory activity against cancer cell growth and antiestrogenic effects [39,40].

3.2. Crushing Temperature Study

The crushing of the olive fruit is the first operation in the extraction of olive oil. The disruption of fruit tissues releases the oil and triggers the enzymatic reactions that shape the profile and concentration of phenolic and volatile compounds [41]. Table 2 presents the effect of the three different crushing temperatures on the phenolic profile of EVOO. As shown in Figure 2, the total phenols decreased with higher temperatures, with a significant difference observed between 10 and 18 °C (360.70 ± 15.00 to 337.55 ± 17.00 mg·kg⁻¹) ($p \leq 0.05$), but not between 18 and 25 °C. All the phenolic groups (secoiridoids, phenolic alcohols, flavones, phenolic acids, and lignans) followed the same trend ($p \leq 0.05$). Similarly, a study on olive oil of the 'Cima di Bitonto' cultivar, applying three crushing temperatures (12, 16, and 20 °C), reported a gradual decrease in the phenolic content [42]. The overall quality of EVOO, attributed to its content of minor compounds, can therefore be modulated by using the appropriate crushing conditions [43]. A study carried out by Caponio and Catalano [44] showed that higher temperatures in the crusher during olive processing led to a shorter preservation of the oils, mainly attributed to a higher degree of auto-oxidation.

The secoiridoids oleuropein aglycone and ligstroside aglycone underwent enzymatic degradation at higher crushing temperatures, resulting in the formation of oleacein and oleocanthal. During crushing and malaxation, β -glucosidases and esterases are released from the olive flesh and pit and can readily interact with phenols, the reaction kinetics increasing with the temperature [45]. A high concentration of oleocanthal and oleacein in EVOO might be desirable due to their health properties [8,26].

Table 2. Phenolic compound contents ($\text{mg}\cdot\text{kg}^{-1}$) of EVOO produced from ‘Corbella’ olives using different crushing temperatures.

| Group/Compound | Crushing Temperature ($^{\circ}\text{C}$) | | |
|----------------------------|---|----------------------------------|---------------------------------|
| | 10 | 18 | 25 |
| Total Polyphenols | 360.70 \pm 15.00 ^a | 337.55 \pm 17.00 ^b | 331.23 \pm 20.00 ^b |
| Secoiridoids | 346.03 \pm 14.00 ^a | 325.30 \pm 17.00 ^b | 321.28 \pm 19.00 ^b |
| Elenolic acid | 191.00 \pm 8.00 ^a | 175.00 \pm 10.00 ^{ab} | 171.00 \pm 4.00 ^b |
| Hydroxyelenolic acid | 7.44 \pm 0.50 ^a | 6.06 \pm 0.50 ^b | 6.13 \pm 0.30 ^b |
| Hydroxyoleuropein aglycone | 2.14 \pm 0.08 ^a | 1.72 \pm 0.10 ^b | 1.74 \pm 0.08 ^b |
| Ligstroside aglycone | 10.00 \pm 0.20 ^a | 8.96 \pm 0.50 ^b | 8.47 \pm 0.10 ^c |
| Methyloleuropein aglycone | 0.87 \pm 0.01 ^a | 0.87 \pm 0.01 ^a | 0.87 \pm 0.00 ^a |
| Oleacein | 16.50 \pm 0.60 ^c | 29.70 \pm 2.00 ^b | 35.20 \pm 1.00 ^a |
| Oleocanthal | 0.95 \pm 0.09 ^c | 2.05 \pm 0.1 ^b | 2.42 \pm 0.20 ^a |
| Oleuropein aglycone | 117.10 \pm 2.96 ^a | 97.20 \pm 5.00 ^b | 95.10 \pm 2.00 ^b |
| Phenolic alcohols | 0.79 \pm 0.10 ^a | 0.45 \pm 0.05 ^b | 0.39 \pm 0.04 ^b |
| Hydroxytyrosol | 0.59 \pm 0.04 ^a | 0.28 \pm 0.03 ^b | 0.29 \pm 0.04 ^b |
| Hydroxytyrosol acetate | 0.20 \pm 0.02 ^a | 0.16 \pm 0.01 ^b | 0.10 \pm 0.01 ^c |
| Flavones | 3.74 \pm 0.20 ^a | 3.04 \pm 0.20 ^b | 1.91 \pm 0.09 ^c |
| Apigenin | 2.85 \pm 0.20 ^a | 2.28 \pm 0.10 ^b | 1.43 \pm 0.07 ^c |
| Luteolin | 0.89 \pm 0.09 ^a | 0.67 \pm 0.04 ^b | 0.49 \pm 0.05 ^c |
| Phenolic acids | 2.08 \pm 0.09 ^a | 1.79 \pm 0.20 ^b | 1.65 \pm 0.01 ^b |
| <i>m</i> -coumaric acid | 1.49 \pm 0.04 ^a | 1.43 \pm 0.04 ^{ab} | 1.38 \pm 0.03 ^b |
| <i>p</i> -coumaric acid | 0.60 \pm 0.06 ^a | 0.33 \pm 0.02 ^b | 0.27 \pm 0.04 ^b |
| Lignans | 8.06 \pm 1.00 ^a | 6.97 \pm 0.80 ^{ab} | 5.99 \pm 0.80 ^b |
| Pinoresinol | 8.06 \pm 1.00 ^a | 6.97 \pm 0.80 ^{ab} | 5.99 \pm 0.80 ^b |

Different letters within the same row mean significant differences (p -value $<$ 0.05) according to the Kruskal–Wallis test.




Figure 2. Effect of crushing temperature on the total phenolic content and the phenolic composition of EVOO. TP: total phenols; Sec: secoiridoids. * indicates significant differences between phenolic concentration ($p <$ 0.05).

The levels of the phenolic alcohols hydroxytyrosol and hydroxytyrosol acetate decreased by 50.77 and 49.02% at 18 and 25 $^{\circ}\text{C}$, respectively. Hydroxytyrosol underwent a great change between 10 and 18 $^{\circ}\text{C}$, whereas the acetate derivative declined progressively. The EVOO concentration of flavones at 25 $^{\circ}\text{C}$ was 48.93% lower than at 10 $^{\circ}\text{C}$. Similarly, in a study of ‘Ayvalik’ oils, malaxation at 27 $^{\circ}\text{C}$ produced oils with a higher concentration of luteolin than at 37 and 47 $^{\circ}\text{C}$ [46]. Accordingly, higher processing temperatures reduce the content of flavones in olive oil.

The effect on the phenolic acids was comparable to that observed in total phenols, with the most substantial degradation recorded at 18 $^{\circ}\text{C}$, and only a slight further reduction at 25 $^{\circ}\text{C}$. These results reflect that at lower temperatures phenolic acids are stable or degrade very little. Finally, although less drastically, lignans also declined with higher temperatures, in contrast with another study that found their concentration was unaltered by changes in temperature during malaxation [47].

3.3. Crushing Size and Malaxation Study

Considerable research has been carried out to clarify which factors most affect the concentration of phenolic compounds in EVOO, with the aim of maximizing its health properties. In this context, many studies have investigated the impact of malaxation temperature and time, together with either the time of harvest [48], extraction techniques [49], or the degree of ripeness [27,50], on the concentration of total and individual phenols. Crushing size has been evaluated to a lesser extent, often together with the hammer mill rotor speed [41,51]. Therefore, by analyzing the influence of the temperature and time of malaxation, as well as the crushing size, our study opens a new three-factor approach to optimizing the phenol content of EVOO.

The effect of malaxation temperature (27, 32, and 37 °C), malaxation time (30 and 60 min) and crushing size (4 and 6 mm) on the total and individual phenol content is presented in Table 3. In general, the malaxation temperature affected the content of total and individual phenols, except for liguistroside aglycone and luteolin ($p > 0.05$), whereas the duration was the only factor that modified the concentration of all the compounds analyzed and consequently the total polyphenol concentration ($p < 0.05$). Crushing size did not affect the concentration of liguistroside and oleuropein aglycone, luteolin, and pinoreosinol ($p > 0.05$).

Concerning the factorial interaction effects, temperature*time modified the concentration of hydroxyelenolic acid, oleocanthal, hydroxytyrosol, and apigenin, but to a low extent ($p < 0.05$). Mixing temperature * crushing size affected the final concentration of up to 70% of phenolic compounds and the total polyphenols. Finally, malaxation time* crushing size significantly affected the concentration of total polyphenols, hydroxyelenolic acid, oleocanthal, oleacein, oleuropein aglycone, and hydroxytyrosol ($p < 0.05$).

The temperature*time*size interaction affected the content of total phenols and oleocanthal, oleacein, oleuropein aglycone, hydroxytyrosol, and apigenin. The factorial design showed that the EVOOs with the highest phenolic contents were obtained after 30 min of malaxation at 37 °C, the amount being $513.43 \pm 70.89 \text{ mg}\cdot\text{kg}^{-1}$ for a crushing size of 4 mm and $503.21 \pm 66.72 \text{ mg}\cdot\text{kg}^{-1}$ for 6 mm; conversely, the lowest content was obtained at 32 °C (Figure 3). The final concentration of phenolic compounds in EVOO probably depends on the equilibrium between their transfer to the oil phase and oxidative degradation [52]. A similar positive influence of a higher malaxation temperature has been reported [17,49,52], although some studies found that temperatures beyond 30 °C resulted in a lower phenolic concentration due to an enhanced activity of endogenous peroxidase (POD) and polyphenol oxidase (PPO) enzymes [17,47,53–56]. In addition, a short malaxation time increased the total phenol content (Figure 3), as in other studies [49,56,57], whereas a long malaxation time is rarely reported to have a positive effect [58]. The benefits of a brief malaxation can be attributed to a reduced activity of the hydrolytic and oxidative enzymes (POD and PPO) and less diffusion of the phenols to the water phase [56,57]. In our study, it was also demonstrated that the crushing size (4 and 6 mm) barely affected the total phenol content when the malaxation conditions were 37 °C for 30 min, in accordance with results in the literature [41,43,50].

Table 3. Effect of malaxation temperature, malaxation time, crushing size, and their interaction on phenolic compound contents ($\text{mg}\cdot\text{kg}^{-1}$) in EVOO produced from ‘Corbella’ olives.

| Compound | MTE | | MT | | CS | | MTE \times MT | | MTE \times CS | | MT \times CS | | MTE \times MT \times CS | |
|----------------------|----------------|---------|----------------|---------|----------------|---------|-----------------|---------|-----------------|---------|----------------|---------|-----------------------------|---------|
| | <i>p</i> Value | F Value | <i>p</i> Value | F Value | <i>p</i> Value | F Value | <i>p</i> Value | F Value | <i>p</i> Value | F Value | <i>p</i> Value | F Value | <i>p</i> Value | F Value |
| Total Polyphenols | <0.001 | 218.2 | <0.001 | 96.7 | 0.0013 | 11.39 | 0.1351 | 2.29 | <0.001 | 21.93 | <0.001 | 32.83 | <0.001 | 32.67 |
| Elenolic acid | <0.001 | 34.4 | <0.001 | 76.14 | 0.0179 | 5.92 | 0.1762 | 1.87 | <0.001 | 37.95 | 0.2157 | 1.57 | 0.1909 | 1.75 |
| Hydroxyelenolic acid | <0.001 | 103.58 | <0.001 | 89.65 | 0.0019 | 10.59 | 0.0005 | 13.46 | 0.0445 | 4.21 | 0.0013 | 11.36 | 0.3988 | 0.72 |
| Ligstroside aglycone | 0.1559 | 2.07 | <0.001 | 57.75 | 0.2206 | 1.53 | 0.4396 | 0.61 | 0.6632 | 0.19 | 0.3395 | 0.93 | 0.113 | 2.59 |
| Oleacein | <0.001 | 1271.08 | <0.001 | 147.99 | <0.001 | 35.66 | 0.4081 | 0.69 | 0.0001 | 18.05 | <0.001 | 66.25 | <0.001 | 60.79 |
| Oleocanthal | <0.001 | 432.54 | <0.001 | 45.1 | <0.001 | 30.78 | 0.0008 | 12.38 | 0.0001 | 18.95 | <0.001 | 64.67 | <0.001 | 71.49 |
| Oleuropein aglycone | 0.0002 | 15.88 | <0.001 | 50.93 | 0.6893 | 0.16 | 0.6681 | 0.19 | 0.8331 | 0.04 | 0.0075 | 7.65 | 0.0053 | 8.39 |
| Hydroxytyrosol | <0.001 | 46.77 | <0.001 | 21.08 | 0.0032 | 9.44 | 0.023 | 5.44 | <0.001 | 266.07 | 0.0036 | 9.16 | 0.0002 | 15.34 |
| Apigenin | 0.0007 | 12.92 | 0.0001 | 18.33 | 0.0191 | 5.8 | 0.0036 | 9.16 | 0.3479 | 0.89 | 0.2888 | 1.15 | 0.0103 | 7.02 |
| Luteolin | 0.1896 | 1.76 | 0.0171 | 6.02 | 0.3164 | 1.02 | 0.6117 | 0.26 | <0.001 | 25.01 | 0.1389 | 4.69 | 0.1310 | 2.34 |
| Pinoresinol | 0.0012 | 11.49 | 0.0223 | 5.5 | 0.1407 | 2.23 | 0.9473 | <0.001 | <0.001 | 32.07 | 0.3674 | 0.82 | 0.323 | 0.99 |

MTE: malaxation temperature; MT: malaxation time and CS: crushing size.




Figure 3. Effect of malaxation temperature (27, 32, and 37 °C), malaxation time (30 and 60 min), and crushing size (4 and 6 mm) on the total phenol content and the phenolic profile of EVOO. * indicates significant differences between phenolic concentration ($p < 0.05$).

Regarding the secoiridoids, the highest concentration of oleuropein aglycone was obtained after 30 min of malaxation at 37 °C applying a crushing size of 4 mm ($183.27 \pm 18.99 \text{ mg} \cdot \text{kg}^{-1}$). On the other hand, three treatments provided the best conditions to obtain an EVOO rich in ligstroside aglycone and oleacein. In the case of ligstroside aglycone, two treatments with the highest temperature (37 °C, 30 min, and 4 or 6 mm) and one with the lowest (27 °C, 30 min, and 6 mm) gave the best results. These findings are in line with the analysis of the factorial and interaction effects, where only the malaxation time significantly modified the concentration of this aglycone ($p < 0.05$). In the case of oleacein, its EVOO concentration was favored by the highest temperature (37 °C, 30 or 60 min, and 4 or 6 mm) (Figure 3). The optimal conditions for oleocanthal were 60 min at 37 °C with a crushing size of 6 mm. Finally, the highest concentration of elenolic acid ($135.52 \pm 11.33 \text{ mg} \cdot \text{kg}^{-1}$) was obtained using conditions of 27 °C, 30 min, and 4 mm and hydroxyelenolic acid with 37 °C, 30 min, and 4 or 6 mm.

Other studies have reported that higher malaxation temperatures may (i) increase the partition coefficient between oil and water phases in olive paste [59], (ii) reduce the activity of PPO enzymes [17], (iii) boost the release of phenols from the cell wall of polysaccharides

and other olive tissues [15,17], and (iv) increase enzymatic activity of β -glucosidases and esterases. β -glucosidases catalyze the hydrolysis of oleuropein and ligstroside, leading to their aglycone forms, and also contribute to the formation of oleacein and oleocanthal in combination with esterases [60]. The observed increase in concentration of most of the secoiridoids can be attributed to the reasons mentioned above.

Regarding duration, a longer malaxation noticeably reduced the content of secoiridoids, except for oleocanthal. An excessive malaxation time can result in undesired reactions catalyzed by PPO and phenoloxidasases [17,57,61] by increasing olive paste exposure to air. Moreover, prolonged contact between the paste and water favors the diffusion of phenols into the aqueous phase [57,61]. Previous studies have reported a similar increase in oleocanthal, with other secoiridoids showing either a weak dependence on the malaxation duration [27,50] or a decreasing trend over time [48,57]. In summary, the oils with the highest secoiridoid content, except for elenolic acid, were produced by a malaxation temperature of 37 °C. These results are in agreement with Boselli et al. (2009), but again they disagree with the conclusions of other studies that describe a negative impact of higher temperatures on secoiridoids, probably due to an enhanced rate of degradation [52].

The hydroxytyrosol content was also affected by extraction conditions (Figure 3), the highest concentration ($10.45 \pm 0.98 \text{ mg}\cdot\text{kg}^{-1}$) being obtained by malaxation at 37 °C for 60 min and a crushing size of 6 mm. Contradictory results have been reported in the literature, in that a longer olive paste malaxation reduced the hydroxytyrosol concentration [57], as did a higher malaxation temperature [27].

Flavones (apigenin and luteolin) and lignans (pinoresinol) followed a similar pattern, concentrations being lowest at 32 °C and highest at 27 °C, in both cases with a malaxation time of 30 min and crushing size of 4 mm. Similar results have been recently reported for flavones [31,52,62], whereas other authors describe that the duration of malaxation has little effect on flavone and lignan content [57]. Other studies have found opposite trends, reporting that the pinoresinol content increased with temperature and malaxation time [31].

4. Conclusions

The total phenol content in the EVOO extracted from 'Corbella' olives decreased with fruit maturity, but flavones, hydroxyoleuropein aglycone, hydroxytyrosol acetate, and lignans increased. A negative correlation was found between the total phenols and higher crushing temperatures, although individual phenolic compounds such as oleocanthal and oleacein increased. In summary, to obtain a 'Corbella' EVOO rich in phenolic compounds, an early harvest and low crushing temperature should be employed. On the contrary, if a higher concentration of specific phenols such as oleocanthal or oleacein is desired, due to their health properties, a moderate crushing temperature should be used.

Regarding malaxation conditions, the results suggest that a higher temperature (37 °C) increased the total phenol content and the concentration of the most relevant phenols. A short malaxation time (30 min) enhanced the total phenols and oleuropein and ligstroside aglycone, whereas a longer process (60 min) increased oleocanthal concentration. Finally, a clear trend was not observed with respect to crushing size.

Based on these results, future research could be focused on studying the correlation between the enzymatic activity and ripening index or how the malaxation temperature affects olives collected at different times.

Author Contributions: Conceptualization, A.N., A.R.-A., and A.V.-Q.; methodology, A.N. and A.V.-Q.; software, J.L.-C. and M.P.; validation, E.E.-F., A.R.-A., and R.M.L.-R.; formal analysis, A.L.-Y., N.J.-R., J.L.-C., and M.P.; investigation, A.L.-Y., A.N., N.J.-R., J.L.-C., and A.V.-Q.; resources, E.E.-F., A.R.-A., and R.M.L.-R.; data curation, A.L.-Y., N.J.-R., J.L.-C., and M.P.; writing—original draft preparation, A.L.-Y., A.N., and N.J.-R.; writing—review and editing, J.L.-C., M.P., E.E.-F., A.R.-A., R.M.L.-R., and A.V.-Q.; visualization, A.L.-Y., A.N., and N.J.-R.; supervision, A.V.-Q.; project administration, E.E.-F., A.R.-A., and R.M.L.-R.; funding acquisition, E.E.-F., A.R.-A., and R.M.L.-R. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by CICYT [AGL2016-75329-R], CIBEROBN from the Instituto de Salud Carlos III, ISCIII from the Ministerio de Ciencia, Innovación y Universidades, (AEI/FEDER, UE) and Generalitat de Catalunya (GC) [2017SGR 196]. A.L-Y wishes to thank the Consejo Nacional de Ciencia y Tecnología (CONACYT) of Mexico for the doctoral scholarship. J. L-C thanks the Ministry of Science Innovation and Universities for the FPI contract (BES-2017-080017). A. V-Q thanks the Ministry of Science Innovation and Universities for the Ramon y Cajal contract (RYC-2016-19355). A. R-A and A.N. acknowledge financial support from the CERCA Program of the Generalitat of Catalonia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article.

Acknowledgments: The authors wish to thank the CCiT-UB for use of the mass spectrometry equipment.

Conflicts of Interest: R.M.L.-R. reports receiving lecture fees from Cerveceros de España and receiving lecture fees and travel support from Adventia. The other authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Salas-Salvadó, J.; Becerra-Tomás, N.; García-Gavilán, J.F.; Bulló, M.; Barrubés, L. Mediterranean Diet and Cardiovascular Disease Prevention: What Do We Know? *Prog. Cardiovasc. Dis.* **2018**, *61*, 62–67. [[CrossRef](#)] [[PubMed](#)]
2. García-Gavilán, J.F.; Bulló, M.; Canudas, S.; Martínez-González, M.A.; Estruch, R.; Giardina, S.; Fitó, M.; Corella, D.; Ros, E.; Salas-Salvadó, J. Extra virgin olive oil consumption reduces the risk of osteoporotic fractures in the PREDIMED trial. *Clin. Nutr.* **2018**, *37*, 329–335. [[CrossRef](#)]
3. Estruch, R.; Ros, E.; Salas-Salvadó, J.; Covas, M.-I.; Corella, D.; Arós, F.; Gómez-Gracia, E.; Ruiz-Gutiérrez, V.; Fiol, M.; Lapetra, J.; et al. Primary Prevention of Cardiovascular Disease with a Mediterranean Diet. *N. Engl. J. Med.* **2013**, *368*, 1279–1290. [[CrossRef](#)] [[PubMed](#)]
4. Yubero-Serrano, E.M.; Lopez-Moreno, J.; Gomez-Delgado, F.; Lopez-Miranda, J. Extra virgin olive oil: More than a healthy fat. *Eur. J. Clin. Nutr.* **2019**, 8–17. [[CrossRef](#)] [[PubMed](#)]
5. The European Commission. *Commission Regulation (EU) No 1018/2013 of 23 October 2013 Amending Regulation (EU) No 432/2012 Establishing a List of Permitted Health Claims Made on Foods Other Than Those Referring to the Reduction of Disease Risk and to Children's Development and Health*; 2013; Vol. L 113, pp. 6–12. Available online: http://eur-lex.europa.eu/pri/en/oj/dat/2003/L_285/L_28520031101en00330037.pdf (accessed on 20 September 2020).
6. Pérez, M.; López-yerena, A.; Lozano-castellón, J.; Olmo-cunillera, A.; Lamuela-raventós, R.M.; Martin-belloso, O.; Vallverdú-queralt, A. Impact of novel technologies on virgin olive oil processing, consumer acceptance, and the valorization of olive mill wastes. *Antioxidants* **2021**, *10*, 417. [[CrossRef](#)] [[PubMed](#)]
7. López-Yerena, A.; Ninot, A.; Lozano-Castellón, J.; Escribano-Ferrer, E.; Romero-Aroca, A.J.; Belaj, A.; Vallverdú-Queralt, A.; Lamuela-Raventós, R.M. Conservation of native wild ivory-white olives from the Medes islands natural reserve to maintain virgin olive oil diversity. *Antioxidants* **2020**, *9*, 1009. [[CrossRef](#)] [[PubMed](#)]
8. Lozano-Castellón, J.; López-Yerena, A.; Rinaldi de Alvarenga, J.F.; Romero del Castillo-Alba, J.; Vallverdú-Queralt, A.; Escribano-Ferrer, E.; Lamuela-Raventós, R.M. Health-promoting properties of oleocanthal and oleacein: Two secoiridoids from extra-virgin olive oil. *Crit. Rev. Food Sci. Nutr.* **2019**, *60*, 2532–2548. [[CrossRef](#)] [[PubMed](#)]
9. Vallverdú-Queralt, A.; Regueiro, J.; Rinaldi de Alvarenga, J.F.; Torrado, X.; Lamuela-Raventós, R.M. Home Cooking and Phenolics: Effect of Thermal Treatment and Addition of Extra Virgin Olive Oil on the Phenolic Profile of Tomato Sauces. *J. Agric. Food Chem.* **2014**. [[CrossRef](#)]
10. Borges, T.H.; Serna, A.; López, L.C.; Lara, L.; Nieto, R.; Seiquer, I. Composition and antioxidant properties of Spanish extra virgin olive oil regarding cultivar, harvest year and crop stage. *Antioxidants* **2019**, *8*, 217. [[CrossRef](#)] [[PubMed](#)]
11. López-Yerena, A.; Lozano-Castellón, J.; Olmo-Cunillera, A.; Tresserra-Rimbau, A.; Quifer-Rada, P.; Jiménez, B.; Pérez, M.; Vallverdú-Queralt, A. Effects of organic and conventional growing systems on the phenolic profile of extra-virgin olive oil. *Molecules* **2019**, *24*, 1986. [[CrossRef](#)]
12. Taticchi, A.; Selvaggini, R.; Esposito, S.; Sordini, B.; Veneziani, G.; Servili, M. Physicochemical characterization of virgin olive oil obtained using an ultrasound-assisted extraction at an industrial scale: Influence of olive maturity index and malaxation time. *Food Chem.* **2019**, *289*, 7–15. [[CrossRef](#)] [[PubMed](#)]
13. Ben Brahim, S.; Kelebek, H.; Ammar, S.; Abichou, M.; Bouaziz, M. LC-MS phenolic profiling combined with multivariate analysis as an approach for the characterization of extra virgin olive oils of four rare Tunisian cultivars during ripening. *Food Chem.* **2017**, *229*, 9–19. [[CrossRef](#)]

14. Clodoveo, M.L.; Hbaieb, R.H.; Kotti, F.; Mugnozza, G.S.; Gargouri, M. Mechanical strategies to increase nutritional and sensory quality of virgin olive oil by modulating the endogenous enzyme activities. *Compr. Rev. Food Sci. Food Saf.* **2014**, *13*, 135–154. [[CrossRef](#)] [[PubMed](#)]
15. Clodoveo, M.L. Malaxation: Influence on virgin olive oil quality. Past, present and future—An overview. *Trends Food Sci. Technol.* **2012**, *25*, 13–23. [[CrossRef](#)]
16. Peres, F.; Martins, L.L.; Ferreira-Dias, S. Influence of enzymes and technology on virgin olive oil composition. *Crit. Rev. Food Sci. Nutr.* **2017**, *57*, 3104–3126. [[CrossRef](#)] [[PubMed](#)]
17. Taticchi, A.; Esposito, S.; Veneziani, G.; Urbani, S.; Selvaggini, R.; Servili, M. The influence of the malaxation temperature on the activity of polyphenoloxidase and peroxidase and on the phenolic composition of virgin olive oil. *Food Chem.* **2013**, *136*, 975–983. [[CrossRef](#)] [[PubMed](#)]
18. Uceda, M.; Frías, L. Seasons of harvest. Changes on fruit oil content, oil composition and oil quality. In *Proceedings of the II Seminario Oleicola International*; IOOC: Cordoba, Spain, 1975; pp. 25–46.
19. Lozano-Castellón, J.; López-Yerena, A.; Olmo-Cunillera, A.; Jáuregui, O.; Pérez, M.; Lamuela-Raventós, R.M.; Vallverdú-Queralt, A. Total analysis of the major secoiridoids in extra virgin olive oil: Validation of an uhplc-esi-ms/ms method. *Antioxidants* **2021**, *10*, 540. [[CrossRef](#)]
20. Alowairesh, B.; Singh, Z.; Fang, Z.; Kailis, S.G. Harvest time impacts the fatty acid compositions, phenolic compounds and sensory attributes of Frantoio and Manzanilla olive oil. *Sci. Hort.* **2018**, *234*, 74–80. [[CrossRef](#)]
21. Matos, L.C.; Pereira, J.A.; Andrade, P.B.; Seabra, R.M.; Oliveira, M.B.P.P. Evaluation of a numerical method to predict the polyphenols content in monovarietal olive oils. *Food Chem.* **2007**, *102*, 976–983. [[CrossRef](#)]
22. Gómez-Rico, A.; Salvador, M.D.; La Greca, M.; Fregapane, G. Phenolic and volatile compounds of extra virgin olive oil (*Olea europaea* L. Cv. Cornicabra) with regard to fruit ripening and irrigation management. *J. Agric. Food Chem.* **2006**, *54*, 7130–7136. [[CrossRef](#)]
23. Bilušić, T.; Žanetić, M.; Ljubenković, I.; Generalić Mekinić, I.; Štambuk, S.; Bojović, V.; Soldo, B.; Magiatis, P. Molecular characterization of Dalmatian cultivars and the influence of the olive fruit harvest period on chemical profile, sensory characteristics and oil oxidative stability. *Eur. Food Res. Technol.* **2018**, *244*, 281–289. [[CrossRef](#)]
24. Romero, A.; Díaz, I.; Tous, J. Optimal harvesting period for “arbequina” olive cultivar in catalonia (spain). *Acta Hort.* **2002**, *586*, 393–396. [[CrossRef](#)]
25. Tous, J.; Romero, A. Variedades del olivo. In *Fundación La Caixa*; AEDOS: Barcelona, Spain, 1993.
26. Castejón, M.L.; Montoya, T.; Alarcón, C.; Lastra, D. Potential Protective Role Exerted by Secoiridoids from *Olea europaea* L. in Cancer, Cardiovascular, Neurodegenerative, Aging—Related, and Immunoinflammatory Diseases. *Antioxidants* **2020**, *9*, 149. [[CrossRef](#)]
27. Lukić, I.; Krapac, M.; Horvat, I.; Godena, S.; Kosić, U.; Brkić Bubola, K. Three-factor approach for balancing the concentrations of phenols and volatiles in virgin olive oil from a late-ripening olive cultivar. *LWT Food Sci. Technol.* **2018**, *87*, 194–202. [[CrossRef](#)]
28. Amanpour, A.; Kelebek, H.; Selli, S. LC-DAD-ESI-MS/MS-based phenolic profiling and antioxidant activity in Turkish cv. Nizip Yaglik olive oils from different maturity olives. *J. Mass Spectrom.* **2019**, *54*, 227–238. [[CrossRef](#)] [[PubMed](#)]
29. Vilaplana-Pérez, C.; Auñón, D.; García-Flores, L.A.; Gil-Izquierdo, A. Hydroxytyrosol and Potential Uses in Cardiovascular Diseases, Cancer, and AIDS. *Front. Nutr.* **2014**, *1*, 1–11. [[CrossRef](#)]
30. Aparicio-Soto, M.; Sánchez-Fidalgo, S.; González-Benjumea, A.; Maya, I.; Fernández-Bolaños, J.G.; Alarcón-de-la-Lastra, C. Naturally occurring hydroxytyrosol derivatives: Hydroxytyrosyl acetate and 3,4-dihydroxyphenylglycol modulate inflammatory response in murine peritoneal macrophages. potential utility as new dietary supplements. *J. Agric. Food Chem.* **2015**, *63*, 836–846. [[CrossRef](#)]
31. De Torres, A.; Espinola, F.; Moya, M.; Alcalá, S.; Vidal, A.M.; Castro, E. Assessment of phenolic compounds in virgin olive oil by response surface methodology with particular focus on flavonoids and lignans. *LWT Food Sci. Technol.* **2018**, *90*, 22–30. [[CrossRef](#)]
32. Jiménez, B.; Sánchez-Ortiz, A.; Lorenzo, M.L.; Rivas, A. Influence of fruit ripening on agronomic parameters, quality indices, sensory attributes and phenolic compounds of Picudo olive oils. *Food Res. Int.* **2013**, *54*, 1860–1867. [[CrossRef](#)]
33. Bengana, M.; Bakhouch, A.; Lozano-Sánchez, J.; Amir, Y.; Youyou, A.; Segura-Carretero, A.; Fernández-Gutiérrez, A. Influence of olive ripeness on chemical properties and phenolic composition of Chemlal extra-virgin olive oil. *Food Res. Int.* **2013**, *54*, 1868–1875. [[CrossRef](#)]
34. Brenes, M.; García, A.; García, P.; Rios, J.J.; Garrido, A. Phenolic compounds in Spanish olive oils. *J. Agric. Food Chem.* **1999**, *47*, 3535–3540. [[CrossRef](#)]
35. Artajo, L.S.; Romero, M.P.; Motilva, M.J. Transfer of phenolic compounds during olive oil extraction in relation to ripening stage of the fruit. *J. Sci. Food Agric.* **2006**, *86*, 518–527. [[CrossRef](#)]
36. Ye, J.H.; Wijesundera, C.; Shi, M. Effects of Agronomic and Oil Processing Conditions on Natural Antioxidative Phenolics in Olive (*Olea europaea* L.). *Austin J. Nutr. Food Sci.* **2014**, *2*.
37. Baccouri, O.; Guerfel, M.; Baccouri, B.; Cerretani, L.; Bendini, A.; Lercker, G.; Zarrouk, M.; Daoud Ben Miled, D. Chemical composition and oxidative stability of Tunisian monovarietal virgin olive oils with regard to fruit ripening. *Food Chem.* **2008**, *109*, 743–754. [[CrossRef](#)] [[PubMed](#)]

38. Olmo-Cunillera, A.; López-Yerena, A.; Lozano-Castellón, J.; Tresserra-Rimbau, A.; Vallverdú-Queralt, A.; Pérez, M. NMR spectroscopy: A powerful tool for the analysis of polyphenols in extra virgin olive oil. *J. Sci. Food Agric.* **2020**, *100*, 1842–1851. [[CrossRef](#)] [[PubMed](#)]
39. Fini, L.; Hotchkiss, E.; Fogliano, V.; Graziani, G.; Romano, M.; De Vol, E.B.; Qin, H.; Selgrad, M.; Boland, C.R.; Ricciardiello, L. Chemopreventive properties of pinoresinol-rich olive oil involve a selective activation of the ATM-p53 cascade in colon cancer cell lines. *Carcinogenesis* **2008**, *29*, 139–146. [[CrossRef](#)]
40. Owen, R.W.; Mier, W.; Giacosa, A.; Hull, W.E.; Spiegelhalder, B.; Bartsch, H. Identification of lignans as major components in the phenolic fraction of olive oil. *Clin. Chem.* **2000**, *46*, 976–988. [[CrossRef](#)] [[PubMed](#)]
41. Polari, J.J.; Wang, S.C. Hammer Mill Sieve Design Impacts Olive Oil Minor Component Composition. *Eur. J. Lipid Sci. Technol.* **2019**, *121*, 1900168. [[CrossRef](#)]
42. Caponio, F.; Pasqualone, A.; Gomes, T.; Catalano, P. Use of HPSEC analysis of polar compounds to assess the influence of crushing temperature on virgin olive oil's quality. *Eur. Food Res. Technol.* **2002**, *215*, 534–537. [[CrossRef](#)]
43. Inarejos-García, A.M.; Fregapane, G.; Salvador, M.D. Effect of crushing on olive paste and virgin olive oil minor components. *Eur. Food Res. Technol.* **2011**, *232*, 441–451. [[CrossRef](#)]
44. Caponio, F.; Catalano, P. Hammer crushers vs disk crushers: The influence of working temperature on the quality and preservation of virgin olive oil. *Eur. Food Res. Technol.* **2001**, *213*, 219–224. [[CrossRef](#)]
45. Abbattista, R.; Losito, I.; De Ceglie, C.; Basile, G.; Calvano, C.D.; Palmisano, F.; Cataldi, T.R.I. Structural characterization of the ligstroside aglycone isoforms in virgin olive oils by liquid chromatography–high-resolution Fourier-transform mass spectrometry and H/Dexchange. *J. Mass Spectrom.* **2019**, *54*, 843–855. [[CrossRef](#)] [[PubMed](#)]
46. Jolayemi, O.S.; Tokatli, F.; Ozen, B. Effects of malaxation temperature and harvest time on the chemical characteristics of olive oils. *Food Chem.* **2016**, *211*, 776–783. [[CrossRef](#)] [[PubMed](#)]
47. Parenti, A.; Spugnoli, P.; Masella, P.; Calamai, L. The effect of malaxation temperature on the virgin olive oil phenolic profile under laboratory-scale conditions. *Eur. J. Lipid Sci. Technol.* **2008**, *110*, 735–741. [[CrossRef](#)]
48. Diamantakos, P.; Giannara, T.; Skarkou, M.; Melliou, E.; Magiatis, P. Influence of harvest time and malaxation conditions on the concentration of individual phenols in extra virgin olive oil related to its healthy properties. *Molecules* **2020**, *25*, 2449. [[CrossRef](#)] [[PubMed](#)]
49. Stefanoudaki, E.; Koutsaftakis, A.; Harwood, J.L. Influence of malaxation conditions on characteristic qualities of olive oil. *Food Chem.* **2011**, *127*, 1481–1486. [[CrossRef](#)]
50. Lukić, I.; Žanetić, M.; Jukić Špika, M.; Lukić, M.; Koprivnjak, O.; Brkić Bubola, K. Complex interactive effects of ripening degree, malaxation duration and temperature on Oblica cv. virgin olive oil phenols, volatiles and sensory quality. *Food Chem.* **2017**, *232*, 610–620. [[CrossRef](#)]
51. Polari, J.J.; Wang, S.C. Comparative Effect of Hammer Mill Screen Size and Cell Wall-Degrading Enzymes during Olive Oil Extraction. *ACS Omega* **2020**, *5*, 6074–6081. [[CrossRef](#)] [[PubMed](#)]
52. Boselli, E.; Di Lecce, G.; Strabbioli, R.; Pieralisi, G.; Frega, N.G. Are virgin olive oils obtained below 27 °C better than those produced at higher temperatures? *LWT Food Sci. Technol.* **2009**, *42*, 748–757. [[CrossRef](#)]
53. Servili, M.; Selvaggini, R.; Esposto, S.; Taticchi, A.; Montedoro, G.F.; Morozzi, G. Health and sensory properties of virgin olive oil hydrophilic phenols: Agronomic and technological aspects of production that affect their occurrence in the oil. *J. Chromatogr. A* **2004**, *1054*, 113–127. [[CrossRef](#)]
54. Ryan, D.; Antolovich, M.; Prenzler, P.; Robards, K.; Lavee, S. Biotransformations of phenolic compounds in *Olea europaea* L. *Sci. Hortic.* **2002**, *92*, 147–176. [[CrossRef](#)]
55. Ranalli, A.; Contento, S.; Schiavone, C.; Simone, N. Malaxing temperature affects volatile and phenol composition as well as other analytical features of virgin olive oil. *Eur. J. Lipid Sci. Technol.* **2001**, *103*, 228–238. [[CrossRef](#)]
56. Angerosa, F.; Mostallino, R.; Basti, C.; Vito, R. Influence of malaxation temperature and time on the quality of virgin olive oils. *Food Chem.* **2001**, *72*, 19–28. [[CrossRef](#)]
57. Jiménez, B.; Sánchez-Ortiz, A.; Rivas, A. Influence of the malaxation time and olive ripening stage on oil quality and phenolic compounds of virgin olive oils. *Int. J. Food Sci. Technol.* **2014**, *49*, 2521–2527. [[CrossRef](#)]
58. Polari, J.J.; Garcí-Aguirre, D.; Olmo-García, L.; Carrasco-Pancorbo, A.; Wang, S.C. Interactions between hammer mill crushing variables and malaxation time during continuous olive oil extraction. *Eur. J. Lipid Sci. Technol.* **2018**, *120*, 1–19. [[CrossRef](#)]
59. Gomez-Rico, A.; Inarejos-Garcia, A.M.; Salvador, M.D.; Fregapane, G. Effect of malaxation conditions on phenol and volatile profiles in olive paste and the corresponding virgin olive oils (*Olea europaea* L. Cv. Cornicabra). *J. Agric. Food Chem.* **2009**, *57*, 3587–3595. [[CrossRef](#)] [[PubMed](#)]
60. Miho, H.; Moral, J.; López-González, M.A.; Díez, C.M.; Priego-Capote, F. The phenolic profile of virgin olive oil is influenced by malaxation conditions and determines the oxidative stability. *Food Chem.* **2020**, *314*, 126183. [[CrossRef](#)] [[PubMed](#)]
61. Ranalli, A.; Pollastri, L.; Contento, S.; Iannucci, E.; Lucera, L. Effect of olive paste kneading process time on the overall quality of virgin olive oil. *Eur. J. Lipid Sci. Technol.* **2003**, *105*, 57–67. [[CrossRef](#)]
62. Marx, Í.M.G.; Casal, S.; Rodrigues, N.; Pinho, T.; Veloso, A.C.A.; Pereira, J.A.; Peres, A.M. Impact of the malaxation temperature on the phenolic profile of cv. Cobrançosa olive oils and assessment of the related health claim. *Food Chem.* **2021**, *337*, 127726. [[CrossRef](#)]