

Profiling dopamine-induced oxidized proteoforms of β -synuclein by top-down mass spectrometry

Arianna Luise¹, Elena De Cecco², Erika Ponzini³, Martina Sollazzo², PierLuigi Mauri⁴, Frank Sobott⁵, Giuseppe Legname², Rita Grandori¹, and Carlo Santambrogio^{1*}

¹ Department of Biotechnology and Biosciences, University of Milano-Bicocca, Piazza della Scienza 2, 20126 Milan, Italy; a.luise1@campus.unimib.it (A.L.), rita.grandori@unimib.it (R.G.), carlo.santambrogio@unimib.it (C.S.)

² Department of Neuroscience, Scuola Internazionale Superiore di Studi Avanzati (SISSA) and ELETTRA-Sincrotrone Trieste S.C.p.A, 34136 Trieste, Italy; edececco@sissa.it (E.D.C.), legname@sissa.it (G.L.)

³ Department of Materials Science, University of Milano-Bicocca, Via Roberto Cozzi 55, I-20125 Milan, Italy; erika.ponzini@unimib.it (E.P.)

⁴ Institute of Biomedical Technologies, National Research Council of Italy, Segrate, 20090 Milan, Italy; pierluigi.mauri@itb.cnr.it (P.M.)

⁵ Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds LS2 9JT, United Kingdom; F.Sobott@leeds.ac.uk (F.S.)

* Correspondence: carlo.santambrogio@unimib.it; Tel.: +39 02 6448 3395

SUPPLEMENTARY MATERIALS




Figure S1. Dependence of OE values on charge state in MS spectra. Relative abundance of OE values vs charge state in MS spectra under denaturing conditions, after 24 h incubation of BS in the presence of DA.