

Supplementary material

Table S1. Gene primers used in the PCR assay

Gene( NCBI No.)	Primer direction	Primer (5' → 3')
Srebf1 (78968)	forward	CATGGATTGCACATTGAAGACA
	reverse	CAGAGAACAGAAGAGAACG
Cyp7a1 (13122)	forward	ACAGCTAAGGAGGACTTCACTCT
	reverse	TTCATCAAGGTACCGGTCGTATT
Ppara (19013)	forward	CTCCACCTGCAGAGCAACCA
	reverse	CGTCAGACTCGGTCTTCTTGAT
Cyp51 (13121)	forward	GAGAGAACGTTGCCTATGTGCC
	reverse	TGTAACGGATTACTGGGTTTCT
Scd1 (20249)	forward	AGGCTACATTGAGGTCTGGT
	reverse	AGGAAGGAGTGAGACTGAGC
Fasn (14104)	forward	GCAAAGTCCTGTCCAGGTA
	reverse	TGCTTGGCTTGGTAGCCG
Slc2a4 (20528)	forward	TTCTATTGCCGTCCCTCTGCTT
	reverse	TCATTCTCATCTGGCCCTAAAGTAT
Adipoq (11450)	forward	GGAGAGAACGGAGAGAAAGGAG
	reverse	GAACGCTGAGCGATACACATAA
36b4 (11837)	forward	TCTAGGACCCGAGAACGACCTC
	reverse	GTTGTCAAACACACTGCTGGAT




Figure S1. Concentrations of plasma inflammatory cytokines, n = 8/group. Top edge of the box, 75th percentile; bottom edge, 25th percentile; horizontal bar within box, median; top horizontal bar outside box, maximum concentration; bottom horizontal bar outside box, minimum concentration. Boxes with different letters were significantly different ( $p < 0.05$ ).