

Supplementary Material

Identification of a novel inhibitor of TfR1 from designed and synthesized Muriceidine A derivatives

Yu Wu ^{1,⊥}, Zongchen Ma ^{1,⊥}, Xiaoyuan Mai ¹, Xiaoling Liu¹, Pinglin Li ¹, Xin Qi¹,
Guoqiang Li ^{1,2,*} and Jing Li ^{1,2,*}

¹ Key Laboratory of Marine Drugs, Chinese Ministry of Education, School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, P. R. China

² Laboratory for Marine Drugs and Bioproducts, Open Studio for Druggability Research of Marine Natural Products, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, P. R. China.

***Corresponding Author**

E-mail address: lijing_ouc@ouc.edu.cn (J. L.) and liguoqiang@ouc.edu.cn (G. L.)

[⊥]These authors contributed equally to this work

^1H -NMR spectra and ^{13}C -NMR spectra of compounds **Muriceidine A. 12a-12d** and **16**




Figure S1: The ^1H -NMR spectra of **Muriceidine A**




Figure S2: The ^1H -NMR spectra of **12a**




Figure S3: The ^{13}C -NMR spectra of **12a**




Figure S4: The ^1H -NMR spectra of **12b**




Figure S5: The ^{13}C -NMR spectra of **12b**




Figure S6: The ^1H -NMR spectra of **12c**




Figure S7: The ^{13}C -NMR spectra of **12c**




Figure S8: The ^1H -NMR spectra of **12d**




Figure S9: The ^{13}C -NMR spectra of **12d**




Figure S10: The ^1H -NMR spectra of **16**




Figure S11: The ^{13}C -NMR spectra of **16**