

## Article

# Sinapic acid Inhibits Group IIA Secretory Phospholipase A<sub>2</sub> and Its Inflammatory Response in Mice

Aladahalli S. Giresha <sup>1,2</sup>, Deepadarshan Urs <sup>1</sup>, Sophiya Pundalik <sup>1</sup>, Rajkumar S. Meti <sup>1</sup>, Siddanakoppalu N. Pramod <sup>3</sup>, Ballenahalli H. Supreetha <sup>1</sup>, Madhusudana Somegowda <sup>4</sup>, Kattapura K. Dharmappa <sup>1,\*</sup>, Ahmed M. El-Shehawi <sup>5</sup>, Sarah Albogami <sup>5</sup>, Mona M. Elseehy <sup>6</sup>, Abdullah Alaklabi <sup>7</sup>, Hosam O. Elansary <sup>8,\*</sup>, Alanoud Omur A. Mehder <sup>9</sup> and Eman A. Mahmoud <sup>10</sup>

- <sup>1</sup> Inflammation Research Laboratory, Department of Studies and Research in Biochemistry, Mangalore University, Jnana Kaveri Post Graduate Campus, Chikka Aluvara, Kodagu 571232, India; girishas25@mangaloreuniversity.ac.in (A.S.G.); deepadarshanurs@mangaloreuniversity.ac.in (D.U.); p.sophiya89@mangaloreuniversity.ac.in (S.P.); rsmeti@gmail.com (R.S.M.); suprithabh5985@gmail.com (B.H.S.)
  - <sup>2</sup> Department of Biochemistry, School of Science, Jain (Deemed-to-Be University), JC Road, Bangalore 560027, India
  - <sup>3</sup> Department of Studies in Food Technology, Davangere University, Shivagangothri, Davangere 577007, India; pramodsn@davangereuniversity.ac.in
  - <sup>4</sup> Department of Plant Biochemistry, University of Agriculture and Horticulture Science, Shivamogga 577204, India; ysmadhu84@gmail.com
  - <sup>5</sup> Department of Biotechnology, College of Science, Taif University, Taif 21944, Saudi Arabia; a.elshehawi@tu.edu.sa (A.M.E.-S.); dr.sarah@tu.edu.sa (S.A.)
  - <sup>6</sup> Department of Genetics, Faculty of Agriculture, University of Alexandria, Alexandria 21545, Egypt; monaahmedma@yahoo.com
  - <sup>7</sup> Department of Biology, Faculty of Science, University of Bisha, Bisha 61922, Saudi Arabia; alaklabia@gmail.com
  - <sup>8</sup> Plant Production Department, College of Food & Agriculture Sciences, King Saud University, Riyadh 11451, Saudi Arabia
  - <sup>9</sup> Family Education Department, Education College, Umm AL-Qura University, Mecca 24382, Saudi Arabia; aomehder@uqu.edu.sa
  - <sup>10</sup> Department of Food Industries, Faculty of Agriculture, Damietta University, Damietta 34511, Egypt; emanmail2005@yahoo.com
- \* Correspondence: dharmappa@mangaloreuniversity.ac.in (K.K.D.); helansary@ksu.edu.sa (H.O.E.)



**Citation:** Giresha, A.S.; Urs, D.; Pundalik, S.; Meti, R.S.; Pramod, S.N.; Supreetha, B.H.; Somegowda, M.; Dharmappa, K.K.; El-Shehawi, A.M.; Albogami, S.; et al. Sinapic acid Inhibits Group IIA Secretory Phospholipase A<sub>2</sub> and Its Inflammatory Response in Mice. *Antioxidants* **2022**, *11*, 1251. <https://doi.org/10.3390/antiox11071251>

Academic Editor: Erick Leite Bastos

Received: 12 May 2022

Accepted: 22 June 2022

Published: 25 June 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


**Abstract:** Human Group IIA secreted phospholipase A<sub>2</sub> (sPLA<sub>2</sub>-IIA) enzyme plays a crucial role in several chronic inflammatory diseases such as asthma, atherosclerosis, gout, bronchitis, etc. Several studies showed that the antioxidants exert an anti-inflammatory function by inhibiting the sPLA<sub>2</sub>-IIA enzyme. Hence, the present study evaluated an antioxidant molecule, sinapic acid, for sPLA<sub>2</sub>-IIA inhibition as an anti-inflammatory function. Initially, the antioxidant efficacy of sinapic acid was evaluated, and it showed greater antioxidant potency. Further, sinapic acid inhibited 94.4 ± 4.83% of sPLA<sub>2</sub>-IIA activity with an IC<sub>50</sub> value of 4.16 ± 0.13 μM. The mode of sPLA<sub>2</sub>-IIA inhibition was examined by increasing the substrate concentration from 30 to 120 nM and the calcium concentration from 2.5 to 15 mM, which did not change the level of inhibition. Further, sinapic acid altered the intrinsic fluorescence and distorted the far UltraViolet Circular Dichroism (UV-CD) spectra of the sPLA<sub>2</sub>-IIA, indicating the direct enzyme-inhibitor interaction. Sinapic acid reduced the sPLA<sub>2</sub>-IIA mediated hemolytic activity from 94 ± 2.19% to 12.35 ± 2.57% and mouse paw edema from 171.75 ± 2.2% to 114.8 ± 1.98%, demonstrating the anti-inflammatory efficiency of sinapic acid by in situ and in vivo methods, respectively. Finally, sinapic acid reduced the hemorrhagic effect of *Vipera russelli* venom hemorrhagic complex-I (VR-HC-I) as an anti-hemorrhagic function. Thus, the above experimental results revealed the sinapic acid potency to be an antioxidant, anti-inflammatory and anti-hemorrhagic molecule, and therefore, it appears to be a promising therapeutic agent.

**Keywords:** secreted phospholipase A<sub>2</sub>; chronic inflammatory diseases; proinflammatory mediators; anti-inflammatory drugs; sinapic acid; hemorrhagic activity

### 1. Introduction

Inflammation is a defensive process and a necessary prerequisite to healing the tissue injury that occurs due to physical, chemical, or biological agents. However, if the inflammation remains beyond its defensive role, it leads to serious consequences such as systemic shock, circulatory collapse, and local tissue injury [1]. Studies have shown that secreted phospholipase A<sub>2</sub> group IIA (sPLA<sub>2</sub>-IIA) enzymes play a significant role in oxidative stress [2] and inflammatory diseases [3,4]. In healthy people, the concentration of sPLA<sub>2</sub>-IIA is minimal (3 ng/mL) but increases significantly (250–500 ng/mL) during infection and injuries [5]. The sPLA<sub>2</sub>-IIA concentration has been elevated in most inflammatory fluids of patients with rheumatoid arthritis [6], asthma [7], atherosclerosis [8], and acute respiratory distress syndrome [9], as well as a biomarker for cardiovascular complications [10,11], sepsis [12] and transplant rejection [13].

The sPLA<sub>2</sub>-IIA enzyme catalyzes membrane phospholipid into arachidonic acid and lysophosphatidic acid. Arachidonic acid is converted into inflammatory mediators such as thromboxane, leukotriene, prostaglandins, and prostacyclins. Lysophosphatidic acid is catalyzed to a platelet activating factor (PAF) that further intensifies the inflammatory condition (Figure 1). Furthermore, the arachidonic acid pathway produces loads of reactive oxygen species (ROS), which contribute to the defensive function by destroying inflowing pathogens [14,15]. However, the persistence of ROS after the defensive role causes deleterious complications [16]. Furthermore, they play an important role in several inflammatory diseases such as ARDS, COPD, chronic bronchitis, asthma [17], rheumatoid arthritis [18], and Alzheimer’s disease [19].



**Figure 1.** Schematic representation of sPLA<sub>2</sub>-IIA mediated inflammatory pathways, ROS production and check points of anti-inflammatory molecule/drugs.

The arachidonic acid pathway-mediated ROS production modulates the cPLA<sub>2</sub> and iPLA<sub>2</sub> functions that enhance the production of arachidonic acid and free radicals [20]. Interestingly, the ROS increases the sPLA<sub>2</sub>-IIA activity and lipid peroxidation that modulate the downstream reactions, which further increase the proinflammatory mediators. Therefore, a single bioactive molecule with both sPLA<sub>2</sub>-IIA inhibitory and antioxidant activities may become a more effective anti-inflammatory agent. Till today, Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) are widely used to control chronic inflammatory disorders [21,22]. NSAIDs limit the COX-1/2 enzymes but have no effect on the generation of leukotrienes and PAF [23]; they continue to cause inflammation (Figure 1). Furthermore, the prolonged use of NSAIDs leads to several complications such as hepatotoxicity, renal injury, hypertension, cardiovascular risks, and gastrointestinal toxicity [24–27]. The specific sPLA<sub>2</sub>-IIA inhibitors such as varespladib (LY315920) and varespladib-methyl (LY333013) were examined in clinical trials, where they were used to treat patients with cardiovascular complications [28,29], but they failed to demonstrate the therapeutic effects. Drugs such as LY315920NA, ginkgetin and petrosaspongiolide M were not successful even though they limit the sPLA<sub>2</sub>-IIA activity at nanomolar concentrations. The unsuccessfulness of these sPLA<sub>2</sub>-IIA inhibitors may be due to the problem associated with formulation or their cytotoxic nature [30,31]. As a result, there is an urgent need for safe and effective sPLA<sub>2</sub>-IIA inhibitors from natural resources with minimal or no adverse effects [32].

Antioxidants such as flavonoids, phenols, and retinoids scavenge ROS and prevent lipid peroxidation, and they further limit the sPLA<sub>2</sub>-IIA-mediated arachidonic acid cascade [14]. In our initial study on pharmaceutically important bioactive molecules, sinapic acid, an antioxidant found in dietary sources [33], has been shown to interfere with the pathways connected to inflammation. Sinapic acid plays a protective role against oxidative stress disorders, as shown in [34], and another study has shown the anti-inflammatory effect by down regulating the synthesis of iNOS and COX-2 in murine macrophage cell lines [35]. Sinapic acid is also documented for its anti-inflammatory effects by inhibiting IL-1 $\beta$  [36], NF- $\kappa$ B [35], reducing the risk of inflammatory colitis in mice by suppressing malondialdehyde, TNF- $\alpha$  and myeloperoxidase expression [37], and reducing carrageenan-induced edema [35]. Therefore, we hypothesized our research to evaluate the potency of sinapic acid for neutralizing the PLA<sub>2</sub>-IIA enzyme and its inflammation responses.

## 2. Materials and Methods

### 2.1. Materials

Sinapic acid, gallic acid (GA), thiobarbituric acid (TBA), Sephadex (G25, G50, and G75), CM-Sephadex-25, 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH), Ultima Gold Scintillation Cocktail, and dimethyl sulfoxide (DMSO) are products of Sigma-Aldrich, St. Louis, MO, USA. <sup>14</sup>C-oleic acid was procured from Perkin Elmer Life Sciences Inc. in Boston, MA, USA. The venom of *Viper russelli* was purchased from Irula Cooperative Society Ltd., Tamil Nadu, India. All reagents and chemicals used in the investigation were of superior quality.

### 2.2. Animals

Swiss albino mice (weighing around 20–25 g, males) were procured from the University Animal House Facility (AHF), Mangalore University, Mangalore, India. Animals were maintained and handled according to the guidelines of the Indian National Regulations for Animal Research. In the present study, we conducted the experiments according to the guidelines of Mangalore University's Institutional Animal Ethical Committee (No: MU/AZ/504(a)/IAEC/2015-2016).

### 2.3. Human Biological Fluid

Institutional Human Ethical Committee (IHEC), Mangalore University, Mangalore, India, permitted the usage of human blood samples (IHEC-No. MU/IHEC/2018/7). The blood samples were collected from volunteers after obtaining the consent letter.

#### 2.4. Purification of sPLA<sub>2</sub>-IIA

The sPLA<sub>2</sub>-IIA was purified from *Viper russelli* venom as per the protocol of Kasturi and Gowda [38]. The purity of sPLA<sub>2</sub>-IIA was tested by sodium dodecyl-sulfate polyacrylamide gel electrophoresis [39]. The sPLA<sub>2</sub>-IIA of *Vipera russelli* venom was generally used to study the mode of action of human inflammatory sPLA<sub>2</sub>-IIA because of the simple purification procedure, availability, close structural similarities, and catalytic action compared to human sPLA<sub>2</sub> IIA [40].

The human and snake venom sPLA<sub>2</sub> enzymes share similar functional and biological properties such as edema, pain, muscle injury and leukocyte influx [41]. It was also reported that the binding pattern of a known inhibitor with human and venom phospholipase A<sub>2</sub> was very similar [42]. Hence, it is suggested the use of snake venom PLA<sub>2</sub> as a tool for investigating a new pharmacological inhibitor of human sPLA<sub>2</sub>-IIA [42].

#### 2.5. Molecular Docking

The structures of phospholipase A<sub>2</sub> (PLA<sub>2</sub>) were downloaded from the Protein Data Bank (PDB ID: 1POE and 3H1X). Structures of sinapic acid and genistein were drawn and analyzed with ChemDraw Ultra 12.0. The three-dimensional coordination was derived through the PRODRG online server [43]. The potential active pockets for PLA<sub>2</sub> protein were determined and identified from both of CASTp server and reference [44]. During the process, intermediary steps, such as grid box creation, energy optimization, protein and ligand preparations, were established through the Graphical User Interface program of AutoDock Tools (ADT). AutoDock prepared the data and saved the prepared file in the required PDBQT format. Using the available information about chosen protein and ligand, AutoDock Vina was used for the docking process along with grid box characteristics in the configuration file. AutoDock Vina employs an iterated local search global optimizer to process the submitted data [45]. During the docking procedure, the option was selected to consider both the ligand and protein as rigid. Following the completion of the scheduled docking runs, the variable conformations of the protein with ligands were obtained as binding methods with their respective binding affinity. The stable confirmation mode with the optimum best interaction was selected and was the one that represented the lowest binding affinity; the same was picked and aligned alongside the receptor structure for further investigation [46].

#### 2.6. Estimation of Antioxidant Activity

The antioxidant activity of sinapic acid was estimated by DPPH radical scavenging activity, as described by Blois [47], anti-lipid peroxidation activity, as described by Gutteridge [48], and reducing power activity, as described by Oyaizu [49]. The antioxidant activity was expressed as percent radical scavenging activity.

#### 2.7. Secreted Phospholipase A<sub>2</sub> Assay (sPLA<sub>2</sub>-IIA)

Autoclaved *E. coli* cells labeled with <sup>14</sup>C-oleic acid were used as a substrate for estimation of sPLA<sub>2</sub>-IIA activity [50,51]. Briefly, the reaction mixture (350 µL) consists of 3.18 × 10<sup>9</sup> autoclaved *E. coli* cells, calcium (5 mM), Tris-HCl buffer (100 mM), enzyme, and water. The 30 µL *E-coli* substrate was added and incubated at 37 °C for 60 min. In total, 2N HCl (100 µL) and 100 µL fatty acid-free BSA (10%) were added, vortexed and centrifuged at 20,000 × g for 5 min. A total of 140 µL supernatant containing <sup>14</sup>C-oleic acid was collected with caution and was added to a scintillation cocktail, and the <sup>14</sup>C radioactivity was measured.

#### 2.8. Inhibition of sPLA<sub>2</sub>-IIA Activity

Sinapic acid was dissolved in a small amount of DMSO and made up to the appropriate concentration with Tris-HCl buffer. sPLA<sub>2</sub>-IIA inhibition was performed by taking different concentrations of sinapic acid. Genistein was used as the standard molecule as it was a proven sPLA<sub>2</sub>-IIA inhibitor and an anti-inflammatory molecule [52]. The maxi-

mum concentration of DMSO used in the experiment was 0.022%. The Graph pad Prism version 5.0 (GraphPad Software, San Diego, CA, USA) software was employed to calculate the IC<sub>50</sub> value.

#### 2.9. The Effect of Concentrations of Substrate and Calcium on sPLA<sub>2</sub>-IIA Inhibition

The assay was carried out with and without IC<sub>50</sub> concentration of sinapic acid as described above. The effect of substrate concentration on sPLA<sub>2</sub>-IIA inhibition was studied by increasing its concentration from 30 to 120 nmoles. The effect of calcium concentrations on sPLA<sub>2</sub>-IIA inhibition was examined by increasing its concentration from 2.5 to 15 mM.

#### 2.10. Intrinsic Fluorescence Study

The fluorescence intensity of the sPLA<sub>2</sub>-IIA enzyme was measured with and without sinapic acid using the Horiba Jobin Yvon Fluorolog-3 spectrofluorometer. The standard reaction mixture (2.0 mL) in a 1 cm path length cuvette consists of sPLA<sub>2</sub>-IIA (20 µg/mL) and sinapic acid concentrations ranging from 0.02 to 0.10 µM. The spectra of the fluorescence were measured between the wavelength of 300 and 380 nm. The tryptophan standard was employed to correct the measurements empirically [53].

#### 2.11. The Far UV-Circular Dichroism Study

The UV-CD spectra of sPLA<sub>2</sub>-IIA (30 µg/mL) were recorded with/without sinapic acid in a reaction mixture using a Jasco J-810 spectropolarimeter. The quartz cuvette was used to record the spectra of sPLA<sub>2</sub>-IIA between 200 and 240 nm at room temperature. The bandwidth was 1 nm, and the response time was set for 2 s. Ten scans in total were carried out to obtain the final spectrum. The spectrum of the blank solution containing the standard reaction mixture was subtracted to correct the protein spectra. The secondary structure of sPLA<sub>2</sub>-IIA was calculated using the K2D3 software (<http://cbdm-01.zdv.uni-mainz.de/~andrade/k2d3/>).

#### 2.12. Study of Reversibility of sPLA<sub>2</sub>-IIA Inhibition

The sPLA<sub>2</sub>-IIA with IC<sub>50</sub> concentration of sinapic acid in 350 µL standard reaction mixture was preincubated and then subjected to dialysis (MW cut off of bag is 3000–6000) for twenty-four hours by changing two buffers. The sPLA<sub>2</sub>-IIA activity was determined before and following the dialysis procedure.

#### 2.13. Neutralization of Indirect Haemolytic Activity

The experiment was conducted as per the method of Boman and Kaletta [54]. The human RBC (1 mL) and egg yolk (1 mL) in 8 mL of PBS were mixed fresh as a substrate for indirect hemolytic activity. The inhibitor (sinapic acid) was preincubated with sPLA<sub>2</sub>-IIA (30 µg) at 37 °C for 30 min, and 1 mL of a substrate was added and allowed for the reaction for 45 min at 37 °C. In total, 9 mL of ice-cold PBS was added to halt the reaction. The suspension was vortexed and centrifuged for 20 min at 1500 × g. The hemolytic activity in terms of released hemoglobin was measured at 530 nm. The sPLA<sub>2</sub>-IIA enzyme without sinapic acid was the positive control.

#### 2.14. Neutralization of Edema Inducing Activity of sPLA<sub>2</sub>-IIA

The assay was performed as per the method of Yamakawa et al. [55], slightly modified by Vishwanath et al. [56]. The sPLA<sub>2</sub>-IIA (5 µg) with different concentrations of sinapic acid, making up a total of 20 µL, was injected into the plantar surface of the right hind footpad of mice (weighing 20–25 g). The saline was injected to the respective left hind limb as negative controls. The animals were euthanized after 45 min by administering anesthesia (30 mg/kg of pentobarbitone i.p.), and hind limbs were amputated at the ankle

joint and were weighed individually. The edema percentage was calculated using the following formula

$$\text{Edema ratio} = \frac{\text{Weight of the edematous leg}}{\text{Weight of normal leg (saline injected)}} \times 100$$

### 2.15. Hemorrhagic Activity of sPLA<sub>2</sub>-IIA

The hemorrhagic activity of sPLA<sub>2</sub>-IIA was estimated as described by Kondo H and Venkatesh M [57,58]. Briefly, 10 µg of the hemorrhagic complex containing 5:2 ratio of sPLA<sub>2</sub>-IIA enzyme and nonenzymatic peptide (Vipera neurotoxin-II, VNTx-II) was injected subcutaneously (s.c.). The mice were euthanized after three hours, the skin was removed, and hemorrhagic spots were measured on the dorsal surface. Saline alone was injected as the control. The preincubated hemorrhagic complex was injected with different concentrations of sinapic acid for the inhibition studies.

### 2.16. Statistical Analysis

The test results were given as the mean standard deviation of three determinations. Graph Pad Prism Version 5.0 was used to calculate IC<sub>50</sub> values. Percent inhibition was calculated from the difference between the control receiving vehicle and the inhibitor-treated animals.

## 3. Results

### 3.1. Molecular Docking


The molecular docking study was carried out to analyze the enzyme–inhibitor interaction. The sinapic acid interacted with the human sPLA<sub>2</sub>-IIA (1POE) enzyme, and it showed the binding energy of −7.6 (E-value). The sinapic acid interacted with the active site conserved amino acid Asp48 through hydrogen bonding and hydrophobic interaction with Cys124, Val45, Cys49, Thr121, Pro122, Lys52, Gly32, and Gly31 (Figure 2B and Table 1). Similarly, the binding energy of standard genistein was −7.2, which interacted with active site Asp48 and Lys52 through hydrogen bonding and showed hydrophobic interaction with Val45, Cys49, Thr121, and Cys124 residues (Figure 2A and Table 1).

**Table 1.** The binding energy, H-bonds and hydrophobic interactions of genistein and sinapic acid with human sPLA<sub>2</sub>-IIA (PDB ID: 1POE).

No.	Ligands	Enzyme Name	Binding Energy (kcal/Mol)	No. of H Bonds	RMSD	Inhibition Constant (Ki) µM	Amino Acids Involved in Interactions
1	Genistein	1POE (Human PLA <sub>2</sub> )	−7.2	02	80.96	1.95	Val45, Asp48, Cys49, Lys52, Thr121, Cys124
2	Sinapic acid		−7.6	01	83.44	2.71	Gly31, Gly32, Asp48, Cys49, Lys52, Thr121, Pro122, Cys124,

### 3.2. Estimation of Antioxidant Activities

The antioxidant potency of sinapic acid was examined by the DPPH method, reducing power assay and anti-lipid peroxidation activity. The sinapic acid scavenged the DPPH radicals to 89% ± 2.13, whereas the ascorbic acid (standard) showed 96% ± 1.78. The reducing power of sinapic acid was 63.5% ± 2.05, compared to quercetin (standard), which showed 66.5% ± 1.82. The anti-lipid peroxidation was 80.6% ± 1.07, whereas α-lipoic acid (standard) exhibited 89% ± 1.69 (Table 2).



**Figure 2.** Docked images of sPLA<sub>2</sub>-IIA (1POE) with genistein (standard) and sinapic acid: 2D Ligplot analysis and the crystal structure of sPLA<sub>2</sub>-IIA (1POE) with standard genistein (A) and sinapic acid (B). Hydrogen bonds are presented by the green dashed line, and the unit of distance was set as Å.

**Table 2.** The antioxidant activity of sinapic acid.


Antioxidant Assays	Sinapic Acid (25 µM)	Ascorbic Acid (25 µM)	Quercetin (25 µM)	α-Lipoic Acid (25 µM)
DPPH free radical	89% ± 2.13	96% ± 1.78	NT	NT
Reducing power	63.5% ± 2.05	NT	66.5% ± 1.82	NT
Anti-lipid peroxidation	80.6% ± 1.07	NT	NT	89% ± 1.69

NT—not tested.

The antioxidant activity of sinapic acid was evaluated by DPPH radical scavenging, reducing power assay and anti-lipid peroxidation.

### 3.3. Inhibition of sPLA<sub>2</sub>-IIA

Further, the sinapic acid was employed to inhibit the inflammatory sPLA<sub>2</sub>-IIA enzyme. It potentially inhibited the sPLA<sub>2</sub>-IIA enzyme to the extent of 94.4% ± 4.83 at 16 μM concentration with an F-statistic value of 0.0031 and *p*-value of 0.9969 (Figure 3). The IC<sub>50</sub> value of sinapic acid was calculated by the software Graphpad Prism 5.0, and it was shown to be 4.16 ± 0.13 μM, whereas the standard genistein was 11.75 μM (historical IC<sub>50</sub> value) (Table 3) [52].



**Figure 3.** Inhibition of sPLA<sub>2</sub>-IIA enzyme by sinapic acid: sPLA<sub>2</sub>-IIA enzyme was incubated with sinapic acid (2–16 μM) at 37 °C for 60 min. Enzyme activity was measured by measuring the radiation of <sup>14</sup>C by liquid scintillation spectrometer. Enzyme inhibition was noted as a percentage of control. The data represents mean ± SD (n = 3).


**Table 3.** IC<sub>50</sub> value of sinapic acid for sPLA<sub>2</sub>-IIA enzyme.

Enzyme	Specific Activity* (nmol/mg/min at 37 °C)	IC <sub>50</sub> (μM) #		Type of Inhibition
		Sinapic Acid	Genistein	
sPLA <sub>2</sub> -IIA	160.0	4.16 ± 0.13	11.75 [58]	Both are competitive inhibitors bound to the active site of the sPLA <sub>2</sub> -IIA.


\* The specific activity is defined as nmoles of fatty acid released/mg of protein/min at 37 °C. # IC<sub>50</sub> value is the concentration that inhibits 50% sPLA<sub>2</sub>-IIA enzyme activity.

### 3.4. Effect of Calcium and Substrate Concentration on sPLA<sub>2</sub> IIA Inhibition

The sPLA<sub>2</sub>-IIA activity was measured with and without sinapic acid (IC<sub>50</sub> concentration) by increasing the calcium concentration from 2.5 to 15 mM, the activity of the enzyme was increased linearly and maintained the constant inhibition of 49.34% ± 1.35 over all the ranges of the calcium concentrations (Figure 4). Furthermore, sPLA<sub>2</sub>-IIA activity was measured with and without IC<sub>50</sub> concentration of sinapic acid by increasing substrate concentration from 30 to 120 nmoles; the enzyme activity increased linearly and maintained the constant inhibition of 48.43% ± 1.76 over all the ranges of substrate concentrations (Figure 5).




**Figure 4.** Effect of calcium concentration on sPLA2-IIA inhibition: The sPLA2 IIA activity was measured with calcium concentration ranges from 2.5 to 15 mM with (■) and without (□) IC<sub>50</sub> concentration of sinapic acid. The sPLA<sub>2</sub>-IIA inhibition is shown in the inlet. The data are expressed as mean ± standard deviation (n = 3).



**Figure 5.** Effect of substrate concentration on sPLA2-IIA inhibition: The sPLA2 IIA activity was measured with the substrate concentrations ranging from 30 to 120 μL with (■) and without (□) IC<sub>50</sub> concentration of sinapic acid. The sPLA<sub>2</sub>-IIA inhibition is shown in the inlet. The data are expressed as mean ± standard deviation (n = 3).

The inhibition constant ( $K_i$ ) was determined by fitting the data to the competitive inhibition model in GraphPad Prism 5.0 via nonlinear regression analysis of competitive enzyme kinetics [59] (Figure 6). The  $K_i$  of sinapic acid for sPLA<sub>2</sub>-IIA inhibition was found to be  $2.711 \pm 1.19$ . The  $K_i$  and  $IC_{50}$  values are often used to compare the relative potency of inhibitors. Smaller  $K_i$  values denote tight binding, and if the  $K_i$  value is less than the  $IC_{50}$  value, it indicates competitive inhibition [60].



**Figure 6.** Inhibition constant ( $K_i$ ) of sinapic acid for sPLA<sub>2</sub>-IIA. The primary data fit the competitive inhibition model, and  $K_i$  was determined by increasing the concentrations of sinapic acid by using Graph pad prism 5.0. Values plotted mean  $\pm$  standard deviation ( $n = 3$ ).

### 3.5. Intrinsic Fluorescence Study

The altered intrinsic fluorescence spectrum of the enzyme indicates the structural changes due to interaction with the inhibitor. Sinapic acid altered the relative intrinsic fluorescence of the sPLA<sub>2</sub>-IIA enzyme in accordance with the inhibitor concentration (0.02 to 0.1  $\mu$ M). The maximum fluorescence intensity of sPLA<sub>2</sub>-IIA was noted at 338 nm and shifted to the higher wavelength of 344 nm in the presence of sinapic acid at 0.1  $\mu$ M concentration (Figure 7I,II).


### 3.6. Circular Dichroism (CD) Study

The change in the secondary structure of the enzyme implies the direct interaction with the inhibitor. The CD spectrum of sPLA<sub>2</sub>-IIA with and without  $IC_{50}$  concentration of sinapic acid was recorded, which exhibited two major negative bands at 210 and 222 nm (red line). In the presence of sinapic acid ( $IC_{50}$  concentration), the negative bands were significantly reduced and abruptly shifted to longer wavelengths of 220 and 224 nm, respectively (Figure 8). The sPLA<sub>2</sub>-IIA spectra were corrected by subtracting spectra of the blank solution containing 100 mM Tris-HCl buffer (pH 7.4) and 5 mM calcium. The K2D3 software was used to determine the secondary structure of the sPLA<sub>2</sub>-IIA enzyme (Table 4).


**Table 4.** Secondary structure of sPLA<sub>2</sub>-IIA enzyme upon addition of sinapic acid.

* Secondary Structure of sPLA <sub>2</sub> -IIA	sPLA <sub>2</sub> -IIA Alone	sPLA <sub>2</sub> -IIA + Sinapic Acid ( $IC_{50}$ )
$\alpha$ -helix	43.66%	21.47%
$\beta$ -strand	10.62%	31.65%
Random coil	45.72%	46.88%

\* Secondary structure of sPLA<sub>2</sub> IIA was calculated with K2D3.



**Figure 7.** Intrinsic fluorescence spectra of sPLA<sub>2</sub>-IIA enzyme: (I) Fluorescence spectra of sPLA<sub>2</sub>-IIA enzyme (20 µg/mL) alone (a), with sinapic acid concentrations of 0.02 µM (b), 0.04 µM (c), 0.06 µM (d), 0.08 µM (e) and 0.1 µM (f). (II) The graph showed maximum fluorescence intensity of sPLA<sub>2</sub>-IIA enzyme of each concentration of sinapic acid.




**Figure 8.** Circular Dichroism spectra of sPLA<sub>2</sub>-IIA with and without sinapic acid: The far-UV CD spectra of sPLA<sub>2</sub>-IIA alone (red line) and with sinapic acid (IC<sub>50</sub> concentration) (blue line) were recorded between 200 and 240 nm using a Jasco J-810 spectropolarimeter.

### 3.7. Determination of Binding Characteristics

The reversibility of sPLA<sub>2</sub>-IIA inhibition was studied by subjecting the preincubated reaction mixture to dialysis. The sPLA<sub>2</sub>-IIA activity was measured before and after the dialysis. The percentage of sPLA<sub>2</sub>-IIA inhibition before and after the dialysis was found to be  $50.2\% \pm 2.3$  and  $47.8\% \pm 1.55$ , respectively.

### 3.8. Neutralization of Indirect Haemolytic Activity

Sinapic acid was subjected to neutralizing indirect hemolytic activity of the sPLA<sub>2</sub>-IIA enzyme. Sinapic acid reduced the indirect hemolytic activity of sPLA<sub>2</sub>-IIA in a concentration-dependent manner. The sPLA<sub>2</sub>-IIA (30 µg) alone causes erythrocyte lysis to  $94\% \pm 2.19$ , which was reduced to  $12.35\% \pm 2.57$  by sinapic acid at a concentration of 16 µM (Figure 9). Distilled water served as a positive control (100% lysis).




**Figure 9.** Neutralization of sPLA<sub>2</sub>-IIA induced indirect hemolytic activity: The reaction was initiated by adding 1 mL of substrate to preincubated sPLA<sub>2</sub>-IIA with indicated concentration sinapic acid and incubated at 37 °C for 30 min. The released hemoglobin was measured by reading the optical density at 540 nm. Data represent the mean standard deviation (n = 3).

### 3.9. Neutralization of sPLA<sub>2</sub> IIA Induced Mouse Paw Edema


The different doses of sinapic acid (3–18 µM) were preincubated with sPLA<sub>2</sub>-IIA and injected into the right hind paw of mice, and saline injected to the left hind paw served as the control. Sinapic acid reduced the edema from  $171.75\% \pm 2.2$  to  $114.8\% \pm 1.98$  at 18 µM concentration, and the reduced percentage of sPLA<sub>2</sub>-IIA-induced edema was  $79.12\% \pm 1.52$  (Figure 10).

### 3.10. Neutralization of Haemorrhagic Activity

This study reveals the synergistic effect of sPLA<sub>2</sub>-IIA and nonenzymatic peptides. The *Vipera russellii* sPLA<sub>2</sub>-IIA and *Vipera russellii* neurotoxic nonenzymatic peptide (VNTx-II) in the 5:2 molar ratio is called *V. russelli* Hemorrhagic Complex-I (VR-HC-I) [58] and was administered to mice intradermally. VR-HC-I induced the hemorrhage at the injection site (Figure 11c). Neither sPLA<sub>2</sub>-IIA nor VNTx-II independently showed the hemorrhagic effect (Figure 11a,b respectively). The mice were injected with preincubated VR-HC-I with sinapic acid (5, 10 and 15 µM), which reduced the hemorrhagic potential (Figure 11d–f respectively). Sinapic acid significantly neutralized the hemorrhagic activity at 15 µM concentration.



**Figure 10.** Neutralization of sPLA<sub>2</sub>-IIA induced edema: The preincubated PLA<sub>2</sub>-IIA enzyme (5 µg) with the indicated concentrations of sinapic acid injected to the right hind footpad of mice (20 µL). The mice were euthanized after 45 min and legs were amputated at the ankle, and the edema ratio was calculated. The data are shown in mean ± standard deviation (n = 3).



**Figure 11.** Reducing VR-HC-I mediated hemorrhagic activity by sinapic acid: Mice injected with 10 µg of sPLA<sub>2</sub>-IIA (a), 10 µg of VNTx-II alone (b), 10 µg of VR-HC-I (c). The injection of preincubated 10 µg of VR-HC-I with sinapic acid of 5 µM (d), 10 µM (e), 15 µM (f). The mice were sacrificed after 3 h, and hemorrhagic spots were recorded using the graph sheet.

#### 4. Discussion

Sinapic acid is rich in fruits such as orange, mango, avocado, strawberries, and raspberries [61–63], vegetables such as garlic, onions, cabbage [64,65], and legumes such as horse grams [66]. Among them, avocado, garlic, and horse gram are well documented for their anti-inflammatory activity [67,68]. The sinapic acid reported no cytotoxic effect on V79 cells [69], and there was no effect on lactate dehydrogenase activity and serum creatine kinase in broilers, suggesting that there are no effects on the brain, liver, kidneys, and cardiac muscle [70]. Thus, sinapic acid from the food items was demonstrated as a non-toxic and therapeutically important molecule.

The *in silico* molecular docking study is important at the early stage of drug discovery because it provides basic knowledge of binding energy, pattern and binding affinity. The docking results of sinapic acid with sPLA<sub>2</sub>-IIA (1POE) exhibited greater binding energy (E value −7.6), which was slightly higher than the binding energy of standard genistein (E value −7.2). Most of the sPLA<sub>2</sub>-IIA inhibitors interfere with the catalytic site by binding to His 47/Asp 48 and decreases the catalytic activity by weakening Ca<sup>2+</sup> interaction [71,72]. The sinapic acid interacted with the active site amino acid Asp48 through hydrogen bonding and showed hydrophobic interaction with few amino acids. Therefore, sinapic acid was assumed to be a potent inhibitor of sPLA<sub>2</sub>-IIA enzyme (Table 1).

The reactive oxygen species (ROS) and their role in human disease has become an important aspect of disease management. The sPLA<sub>2</sub>-IIA-mediated ROS generation through activation of NADPH oxidases [73] is an important pathway as it is known to be involved in the activation of cPLA<sub>2</sub> and ERK1/2 [74], leading to the release of arachidonic acid. Furthermore, the hydroxyl radicals that formed during inflammation attack membrane glycerophospholipids and initiate lipid peroxidation [73]. Hence, sinapic acid was evaluated for its antioxidant efficacy. Sinapic acid effectively scavenged the DPPH radical, reduced the ferric ions, and demonstrated its ability to protect lipid peroxidation. Thus, it was concluded that sinapic acid, if developed as an anti-inflammatory drug, limits free radicals and their intermediates released during inflammatory pathologies.

The results illustrated that the sinapic acid exhibited greater binding energy (docking study) and better antioxidant potency and, hence, was further examined for its anti-inflammatory activity. Sinapic acid potentially inhibited the sPLA<sub>2</sub>-IIA enzyme in a concentration-dependent manner and with a comparatively low IC<sub>50</sub> value. The genistein was taken as a standard molecule in this study as it is a well-known anti-inflammatory and antioxidant molecule [75–77]. The kinetics study of sPLA<sub>2</sub>-IIA inhibition was performed using Graph pad prism software 5.0, which suggested that the sinapic acid is a competitive inhibitor of sPLA<sub>2</sub>-IIA.

Many sPLA<sub>2</sub>-IIA inhibitors limit the enzyme activity either by chelating calcium (metal ion) or binding to substrates. Inhibitors such as lipocortin I and II bind sPLA<sub>2</sub>-IIA enzymes non-specifically and affect the quality of the lipid interface [78]. Therefore, we examined the effect of substrate and calcium concentrations on sPLA<sub>2</sub>-IIA inhibition. The findings exhibit that the inhibition of sPLA<sub>2</sub>-IIA by sinapic acid was not dependent on calcium or substrate concentrations.

Many sPLA<sub>2</sub>-IIA inhibitors were reported to alter fluorescence spectra [79]. The structural change in the enzyme upon interaction with the inhibitor alters the intrinsic fluorescence. The aromatic amino acids of protein such as tryptophan, tyrosine, and phenylalanine are responsible for intrinsic fluorescence. The quantum yield, intensity, and wavelength of maximum fluorescence emission depend upon the microenvironment of these aromatic amino acids. The sinapic acid shifts the maximum fluorescence spectrum of sPLA<sub>2</sub>-IIA towards the shorter wavelength and increases the fluorescence intensity as the polarity of the solvent surrounding the aromatic amino acids decreases [80,81]. Sinapic acid alone does not show any fluorescence, indicating that sinapic acid interacts directly with the sPLA<sub>2</sub>-IIA enzyme.

To substantiate fluorimetry results, a circular dichroism (CD) study was carried out. The studies showed that significant changes occur in the secondary structure of sPLA<sub>2</sub>-IIA

upon inhibitors binding [82]. In the present study, the sinapic acid interaction with the sPLA<sub>2</sub>-IIA enzyme caused significant changes in the secondary structure (Figure 8). Hence, it is indisputably concluded that sinapic acid inhibits sPLA<sub>2</sub>-IIA by irreversibly binding to the active site.

The reversibility of sPLA<sub>2</sub>-IIA inhibition was examined by measuring the percentage of inhibition before and after the dialysis of the reaction mixture. The inhibition percentages before and after the dialysis were almost the same. Hence, it again is implicit that sinapic acid irreversibly binds to the sPLA<sub>2</sub>-IIA enzyme.

The indirect hemolytic assay is an indirect way of estimating sPLA<sub>2</sub>-IIA activity using egg yolk phospholipid and washed erythrocytes as substrates [83]. Sinapic acid efficiently neutralized the sPLA<sub>2</sub>-IIA-mediated hemolysis in a dose-dependent way. Thus, sinapic acid neutralizes sPLA<sub>2</sub>-IIA enzyme activity irrespective of the nature of the substrate because the sinapic acid binds to the enzyme irreversibly.

It has been observed that the in vitro experiments show positive results but fail to show efficiency in the in vivo studies. This could be due to the heterogeneity of the environment in the in vivo models. Animal experiments are important for researchers as they provide peer knowledge of pharmacodynamics and pharmacokinetics in the early stages of drug discovery [84]. Therefore, the effectiveness of the sinapic acid in neutralizing the sPLA<sub>2</sub>-IIA-induced inflammatory response in the Swiss albino mice was evaluated. Sinapic acid reduced inflammatory edema to a greater extent. Thus, sinapic acid demonstrated the in vivo efficacy by neutralizing the sPLA<sub>2</sub>-IIA mediated inflammatory response.

In the living system, protein–protein interactions lead to pharmacological damage due to the synergistic effect [85]. For example, the interaction of human sPLA<sub>2</sub>-IIA and vimentin (an intracellular protein) further exacerbates inflammatory pathologies. Interestingly, the addition of LY311727 (sPLA<sub>2</sub>-IIA inhibitor) causes substantial structural displacement in the amino terminus of the sPLA<sub>2</sub>-IIA enzyme, and that is sufficient to minimize its interaction with the vimentin. The interaction between sPLA<sub>2</sub>-IIA and nonenzymatic peptides is synergistic in snake bites and leads to increased hemorrhage [58]. In the present study, sinapic acid significantly reduced the synergistic hemorrhagic effect of *V. russelli* Haemorrhagic Complex-I (sPLA<sub>2</sub>-IIA and *V. russelli* neurotoxic nonenzymatic peptide) (Figure 10).

## 5. Conclusions

Activated sPLA<sub>2</sub>-IIA generates proinflammatory lipid mediators and oxygen-free radicals that intensify the status of oxidative stress disorders and chronic inflammatory diseases. The present study evaluated the sinapic acid from a dietary source for both antioxidant potency and sPLA<sub>2</sub>-IIA inhibition as an anti-inflammatory function, and the result showed that sinapic acid exhibit both the potencies to a better extent. Further, sPLA<sub>2</sub>-IIA inhibition was not dependent on either calcium or substrate concentration. The altered fluorescence intensity and shifted negative bands of the circular dichroism spectrum suggest the direct interaction of the sinapic acid with the active site of the sPLA<sub>2</sub>-IIA enzyme. Furthermore, sinapic acid neutralized sPLA<sub>2</sub>-IIA induced erythrolysis, mouse paw edema and the hemorrhagic effect. As a result, sinapic acid is a potential therapeutic candidate for both inflammatory diseases and snakebite envenomation. However, more clinical studies are needed to claim sinapic acid as an anti-inflammatory drug.

**Author Contributions:** Conceptualization, A.S.G., B.H.S. and K.K.D.; methodology, A.S.G., D.U. and S.P.; software, S.N.P.; result and validation, A.S.G. and K.K.D.; formal analysis, R.S.M.; writing—original draft preparation, A.S.G.; review and editing, K.K.D. and H.O.E.; visualization, M.S.; supervision, K.K.D. and H.O.E.; project administration, A.M.E.-S., S.A. and M.M.E.; funding acquisition, A.A., A.O.A.M. and E.A.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** Taif University Researchers Supporting Project number (TURSP-2020/75).

**Institutional Review Board Statement:** The studies on edema and hemorrhage using Swiss albino mice were conducted according to the guidelines of the Declaration of Indian National Regulations for Animal Research, India, and approved by the Institutional Ethical Committee of Mangalore University, India. (No: MU/AZ/504(a)/IAEC/2015–2016).

**Informed Consent Statement:** All tests were performed according to AVMA's guidelines 55. Human blood samples were collected from healthy volunteers after receiving ethical approval and informed consent letters (IHEC-No. MU/IHEC/2018/7).

**Data Availability Statement:** Data is contained within the article.

**Acknowledgments:** The current work was funded by Taif University Researchers Supporting Project number (TURSP-2020/75), Taif University, Taif, Saudi Arabia. The authors are owed the Indian Council for Medical Research (ICMR), New Delhi, for financial support through Sanctions a Senior Research Fellowship. We also thank Mangalore University for providing laboratory facilities.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Chen, L.; Deng, H.; Cui, H.; Fang, J.; Zuo, Z.; Deng, J.; Li, Y.; Wang, X.; Zhao, L. Inflammation-associated diseases in organs. *Oncotarget* **2018**, *23*, 7204. [[CrossRef](#)]
- Rathee, P.; Chaudhary, H.; Rathee, S.; Rathee, D.; Kumar, V.; Kohli, K. Mechanism of action of flavonoids as anti-inflammatory agents: A review. *Inflamm. Allergy-Drug Targets* **2009**, *8*, 229–235. [[CrossRef](#)]
- Tietge, U.J.; Pratico, D.; Ding, T.; Funk, C.D.; Hildebrand, R.B.; Van Berkel, T.; Van Eck, M. Macrophage-specific expression of group IIA sPLA2 results in accelerated atherogenesis by increasing oxidative stress. *J. Lipid Res.* **2005**, *46*, 1604–1614. [[CrossRef](#)]
- Bidgood, M.J.; Jamal, O.S.; Cunningham, A.M.; Brooks, P.M.; Scott, K.F. Type IIA secreted phospholipase A2 up-regulates cyclooxygenase-2 and amplifies cytokine-mediated prostaglandin production in human rheumatoid synoviocytes. *J. Immunol.* **2000**, *165*, 2790–2797. [[CrossRef](#)]
- Van Hensbergen, V.P.; Wu, Y.; van Sorge, N.M.; Touqui, L. Type IIA secreted phospholipase A2 in host defense against bacterial infections. *Trends Immunol.* **2020**, *41*, 313–326. [[CrossRef](#)]
- Stefanski, E.; Pruzanski, W.; Sternby, B.; Vadas, P. Purification of a soluble phospholipase A2 from synovial fluid in rheumatoid arthritis. *J. Biochem.* **1986**, *100*, 1297–1303. [[CrossRef](#)]
- Bowton, D.L.; Seeds, M.C.; Fasano, M.B.; Goldsmith, B.; Bass, D.A. Phospholipase A2 and arachidonate increase in bronchoalveolar lavage fluid after inhaled antigen challenge in asthmatics. *Am. J. Respir. Crit. Care Med.* **1997**, *155*, 421–425. [[CrossRef](#)]
- Menschikowski, M.; Hagelgans, A.; Siegert, G. Secreted phospholipase A2 of group IIA: Is it an offensive or a defensive player during atherosclerosis and other inflammatory diseases? *Prostaglandins Other Lipid Mediat.* **2006**, *79*, 1–33. [[CrossRef](#)]
- Styles, L.A.; Schalkwijk, C.G.; Aarsman, A.J.; Vichinsky, E.P.; Lubin, B.H.; Kuypers, F.A. Phospholipase A2 levels in acute chest syndrome of sickle cell disease. *Blood* **1996**, *87*, 2573–2578. [[CrossRef](#)]
- Mallat, Z.; Lambeau, G.; Tedgui, A. Lipoprotein-associated and secreted phospholipases A2 in cardiovascular disease: Roles as biological effectors and biomarkers. *Circulation* **2010**, *122*, 2183–2200. [[CrossRef](#)]
- Tietge, J.F. Extracellular phospholipases: Role in inflammation and atherosclerotic cardiovascular. *Atheroscler. Risks Mech. Ther.* **2015**, *10*, 279.
- Tan, T.L.; Goh, Y.Y. The role of group IIA secretory phospholipase A2 (sPLA2-IIA) as a biomarker for the diagnosis of sepsis and bacterial infection in adults—A systematic review. *PLoS ONE* **2017**, *12*, e0180554. [[CrossRef](#)] [[PubMed](#)]
- Annema, W.; de Boer, J.F.; Dijkers, A.; Dimova, L.G.; van der Giet, M.; Bakker, S.J.; Tietge, U.J. Group IIA Secretory Phospholipase A2 Predicts Graft Failure and Mortality in Renal Transplant Recipients by Mediating Decreased Kidney Function. *J. Clin. Med.* **2020**, *9*, 1282. [[CrossRef](#)]
- Nanda, B.L.; Nataraju, A.; Rajesh, R.; Rangappa, K.S.; Shekar, M.A.; Vishwanath, B.S. PLA2 mediated arachidonate free radicals: PLA2 inhibition and neutralization of free radicals by anti-oxidants—a new role as anti-inflammatory molecule. *Curr. Top. Med. Chem.* **2007**, *7*, 765–777. [[CrossRef](#)]
- Nethery, D.; Stofan, D.; Callahan, L.; DiMarco, A.; Supinski, G. Formation of reactive oxygen species by the contracting diaphragm is PLA(2) dependent. *J. Appl. Physiol.* **1999**, *87*, 792–800. [[CrossRef](#)]
- Odinga, E.S.; Waigi, M.G.; Gudda, F.O.; Wang, J.; Yang, B.; Hu, X.; Li, S.; Gao, Y. Occurrence, formation, environmental fate and risks of environmentally persistent free radicals in biochars. *Environ. Int.* **2020**, *134*, 105172. [[CrossRef](#)] [[PubMed](#)]
- Zinellu, A.; Fois, A.G.; Sotgia, S.; Zinellu, E.; Bifulco, F.; Pintus, G.; Mangoni, A.A.; Carru, C.; Pirina, P. Plasma protein thiols: An early marker of oxidative stress in asthma and chronic obstructive pulmonary disease. *Eur. J. Clin. Investig.* **2016**, *46*, 181–188. [[CrossRef](#)] [[PubMed](#)]
- Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.T.; Mazur, M.; Telser, J. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* **2007**, *39*, 44–84. [[CrossRef](#)]
- Bhatt, S.; Puli, L.; Patil, C.R. Role of reactive oxygen species in the progression of Alzheimer's disease. *Drug Discov. Today* **2021**, *26*, 794–803. [[CrossRef](#)]

20. Zhu, D.; Lai, Y.; Shelat, P.B.; Hu, C.; Sun, G.Y.; Lee, J.C. Phospholipases A2 mediate amyloid-beta peptide-induced mitochondrial dysfunction. *J. Neurosci.* **2006**, *26*, 11111–11119. [[CrossRef](#)]
21. Göttsche, P.C. Non-steroidal anti-inflammatory drugs. *BMJ* **2000**, *320*, 1058–1061. [[CrossRef](#)] [[PubMed](#)]
22. Warner, D.C.; Schnepf, G.; Barrett, M.S.; Dian, D.; Swigonski, N.L. Prevalence, attitudes, and behaviours related to the use of nonsteroidal anti-inflammatory drugs (NSAIDs) in student athletes. *J. Adolesc. Health* **2002**, *30*, 150–153. [[CrossRef](#)]
23. Burnett, B.P.; Levy, R.M. 5-Lipoxygenase metabolic contributions to NSAID-induced organ toxicity. *Adv. Ther.* **2012**, *29*, 79–98. [[CrossRef](#)] [[PubMed](#)]
24. Ghosh, R.; Alajbegovic, A.; Gomes, A.V. NSAIDs and Cardiovascular Diseases: Role of Reactive Oxygen Species. *Oxid. Med. Cell. Longev.* **2015**, *2015*, 536962. [[CrossRef](#)]
25. Fitzgerald, G.A. Coxibs and cardiovascular disease. *N. Engl. J. Med.* **2004**, *351*, 1709–1711. [[CrossRef](#)]
26. Vane, J.R.; Botting, R.M. Anti-inflammatory drugs and their mechanism of action. *Inflamm. Res.* **1998**, *47*, 78–87. [[CrossRef](#)]
27. Bavry, A.A.; Khaliq, A.; Gong, Y.; Handberg, E.M.; Cooper-DeHoff, R.M.; Pepine, C.J. Harmful effects of NSAIDs among patients with hypertension and coronary artery disease. *Am. J. Med.* **2011**, *124*, 614–620. [[CrossRef](#)]
28. Nicholls, S.J.; Cavender, M.A.; Kastelein, J.J.; Schwartz, G.; Waters, D.D.; Rosenson, R.S.; Bash, D.; Hislop, C. Inhibition of secreted phospholipase A2 in patients with acute coronary syndromes: Rationale and design of the vascular inflammation suppression to treat acute coronary syndrome for 16 weeks (VISTA-16) trial. *Cardiovasc. Drugs Ther.* **2012**, *26*, 71–75. [[CrossRef](#)]
29. Nicholls, S.J.; Kastelein, J.J.; Schwartz, G.G.; Bash, D.; Rosenson, R.S.; Cavender, M.A.; Brennan, D.M.; Koenig, W.; Jukema, J.W.; Nambi, V.; et al. Varespladib and cardiovascular events in patients with an acute coronary syndrome: The VISTA-16 randomized clinical trial. *JAMA* **2014**, *311*, 252–262. [[CrossRef](#)]
30. Zeiher, B.G.; Steingrub, J.; Laterre, P.F.; Dmitrienko, A.; Fukiishi, Y.; Abraham, E.; EZZI Study Group. LY315920NA/S-5920, a selective inhibitor of group IIA secretory phospholipase A2, fails to improve clinical outcome for patients with severe sepsis. *Crit. Care Med.* **2005**, *33*, 1741–1748. [[CrossRef](#)]
31. Garcia-Pastor, P.; Randazzo, A.; Gomez-Paloma, L.; Alcaraz, M.J.; Paya, M. Effects of petrosaspongolide M, a novel phospholipase A2 inhibitor, on acute and chronic inflammation. *J. Pharmacol. Exp. Ther.* **1999**, *289*, 166–172. [[PubMed](#)]
32. Reid, R.C. Inhibitors of secreted phospholipase A2 group IIA. *Curr. Med. Chem.* **2005**, *12*, 3011–3026. [[CrossRef](#)] [[PubMed](#)]
33. Gaspar, A.; Martins, M.; Silva, P.; Garrido, E.M.; Garrido, J.; Firuzi, O.; Miri, R.; Saso, L.; Borges, F. Dietary phenolic acids and derivatives. Evaluation of the antioxidant activity of sinapic acid and its alkyl esters. *J. Agric. Food Chem.* **2010**, *58*, 11273–11280. [[CrossRef](#)]
34. Chen, C. Sinapic Acid and Its Derivatives as Medicine in Oxidative Stress-Induced Diseases and Aging. *J. Oxid. Med. Cell. Longev.* **2016**, *2016*, 3571614. [[CrossRef](#)] [[PubMed](#)]
35. Yun, K.J.; Koh, D.J.; Kim, S.H.; Park, S.J.; Ryu, J.H.; Kim, D.G.; Lee, J.Y.; Lee, K.T. Anti-inflammatory effects of sinapic acid through the suppression of inducible nitric oxide synthase, cyclooxygenase-2, and proinflammatory cytokines expressions via nuclear factor- $\kappa$ B inactivation. *J. Agric. Food Chem.* **2008**, *56*, 10265–10272. [[CrossRef](#)]
36. Huang, X.; Pan, Q.; Mao, Z.; Zhang, R.; Ma, X.; Xi, Y.; You, H. Sinapic acid inhibits the IL-1 $\beta$ -induced inflammation via MAPK downregulation in rat chondrocytes. *Inflammation* **2018**, *41*, 562–568. [[CrossRef](#)]
37. Lee, J.Y. Anti-inflammatory effects of sinapic acid on 2,4,6-trinitrobenzenesulfonic acid-induced colitis in mice. *Arch. Pharm. Res.* **2018**, *41*, 243–250. [[CrossRef](#)]
38. Kasturi, S.; Gowda, T.V. Purification and characterization of a major phospholipase A2 from Russell's viper (*Viperarusselli*) venom. *Toxicon* **1989**, *27*, 229–237. [[CrossRef](#)]
39. Laemmli, U.K. SDS-page Laemmli method. *Nature* **1970**, *227*, 680–685. [[CrossRef](#)]
40. Davidson, F.F.; Dennis, E.A. Evolutionary relationships and implications for the regulation of phospholipase A2 from snake venom to human secreted forms. *J. Mol. Evol.* **1990**, *31*, 228–238. [[CrossRef](#)]
41. Teixeira, C.D.; Landucci, E.C.; Antunes, E.; Chacur, M.; Cury, Y. Inflammatory effects of snake venom myotoxic phospholipases A2. *Toxicon* **2003**, *42*, 947–962. [[CrossRef](#)] [[PubMed](#)]
42. Sales, T.A.; Marcussi, S.; Da Cunha, E.F.; Kuca, K.; Ramalho, T.C. Can inhibitors of snake venom phospholipases A2 lead to new insights into anti-inflammatory therapy in humans? A theoretical study. *Toxins* **2017**, *9*, 341. [[CrossRef](#)] [[PubMed](#)]
43. Schüttelkopf, W.; van Aalten, D.M.F. PRODRG: A tool for high-throughput crystallography of protein-ligand complexes. *Acta Crystallogr. D Biol. Crystallogr.* **2004**, *60*, 1355–1363. [[CrossRef](#)] [[PubMed](#)]
44. Tian, W.; Chen, C.; Lei, X.; Zhao, J.; Liang, J. CASTp 3.0: Computed atlas of surface topography of proteins. *Nucleic Acids Res.* **2018**, *46*, W363–W367. [[CrossRef](#)]
45. Morris, G.M.; Huey, R.; Lindstrom, W.; Sanner, M.F.; Belew, R.K.; Goodsell, D.S.; Olson, A.J. Autodock4 and AutoDockTools4: Automated docking with selective receptor flexibility. *J. Comput. Chem.* **2009**, *30*, 2785–2791. [[CrossRef](#)]
46. Jinadatta, P.; Rajshekarappa, S.; Rao, K.S.R.; Subbaiah, S.G.P.; Shastri, S. In silico, in vitro: Antioxidant and antihepatotoxic activity of gnetol from *Gnetum ula* Brongn. *Bioimpacts* **2019**, *9*, 239–249. [[CrossRef](#)]
47. Blois, M.S. Antioxidant determinations by the use of a stable free radical. *Nature* **1958**, *181*, 1199–1200. [[CrossRef](#)]
48. Gutteridge, J.M.; Halliwell, B. The measurement and mechanism of lipid peroxidation in biological systems. *Trends Biochem. Sci.* **1990**, *15*, 129–135. [[CrossRef](#)]
49. Oyaizu, M. Studies on products of browning reaction. Antioxidative activities of products of browning reaction prepared from glucosamine. *Jpn. J. Nutr. Diet.* **1986**, *44*, 307–315. [[CrossRef](#)]

50. Patriarca, P.; Beckerdite, S.; Pettis, P.; Elsbach, P. Phospholipid metabolism by phagocytic cells: VII. The degradation and utilization of phospholipids of various microbial species by rabbit granulocytes. *Biochim. Biophys. Acta BBA Lipids Lipid Metab.* **1972**, *280*, 45–56. [[CrossRef](#)]
51. Vishwanath, B.S.; Frey, F.J.; Bradbury, M.J.; Dallman, M.F.; Frey, B.M. Glucocorticoid deficiency increases phospholipase A2 activity in rats. *J. Clin. Investig.* **1993**, *92*, 1974–1980. [[CrossRef](#)] [[PubMed](#)]
52. Dharmappa, K.K.; Mohamed, R.; Shivaprasad, H.V.; Vishwanath, B.S. Genistein, a potent inhibitor of secreted phospholipase A2: A new insight in down regulation of inflammation. *Inflammopharmacology* **2010**, *18*, 25–31. [[CrossRef](#)] [[PubMed](#)]
53. Prigent-Dachary, J.; Boffa, M.C.; Boisseau, M.R.; Dufourcq, J. Snake venom phospholipases A2. A fluorescence study of their binding to phospholipid vesicles correlation with their anticoagulant activities. *J. Biol. Chem.* **1980**, *255*, 7734–7739. [[CrossRef](#)]
54. Boman, H.G.; Kaletta, U. Chromatography of rattlesnake venom: A separation of three phosphodiesterases. *Biochim. Biophys. Acta* **1957**, *24*, 619–631. [[CrossRef](#)]
55. Yamakawa, M.; Nozaki, M.; Hokama, Z. Fractionation of sakishima habu (*Trimeresurus elegans*) venom and lethal, hemorrhagic and edema-forming activities of the fractions. *Anim. Plant Microb. Toxins* **1976**, *1*, 97–109.
56. Vishwanath, B.S.; Gowda, T.V. Interaction of aristolochic acid with Viperarusselli phospholipase A2: Its effect on enzymatic and pathological activities. *Toxicon* **1987**, *25*, 929–937. [[CrossRef](#)]
57. Kondo, H.; Kondo, S.; Ikezawa, H.; Murata, R.; Ohsaka, A. Studies on the quantitative method for determination of hemorrhagic activity of Habu snake venom. *Jpn. J. Med. Sci. Biol.* **1960**, *13*, 43–51. [[CrossRef](#)]
58. Venkatesh, M.; Gowda, V. Synergistically acting PLA2: Peptide hemorrhagic complex from Daboia russelii venom. *Toxicon* **2013**, *73*, 111–120. [[CrossRef](#)]
59. Figueiredo, A.C.; de Sanctis, D.; Pereira, P.J.B. The tick-derived anticoagulant madanin is processed by thrombin and factor Xa. *PLoS ONE* **2013**, *8*, e71866. [[CrossRef](#)]
60. Burlingham, B.T.; Widlanski, T.S. An intuitive look at the relationship of Ki and IC50: A more general use for the Dixon plot. *J. Chem. Educ.* **2003**, *80*, 214. [[CrossRef](#)]
61. Nićiforović, N.; Abramović, H. Sinapic acid and its derivatives: Natural sources and bioactivity. *Compr. Rev. Food Sci. Food Saf.* **2014**, *13*, 34–51. [[CrossRef](#)] [[PubMed](#)]
62. Wang, Y.C.; Chuang, Y.C.; Ku, Y.H. Quantitation of bioactive compounds in citrus fruits cultivated in Taiwan. *Food Chem.* **2007**, *102*, 1163–1171. [[CrossRef](#)]
63. Russell, W.R.; Labat, A.; Scobbie, L.; Duncan, G.J.; Duthie, G.G. Phenolic acid content of fruits commonly consumed and locally produced in Scotland. *Food Chem.* **2009**, *115*, 100–104. [[CrossRef](#)]
64. Gorinstein, S.; Leontowicz, H.; Leontowicz, M.; Namiesnik, J.; Najman, K.; Drzewiecki, J.; Cvikrová, M.; Martinová, O.; Katrich, E.; Trakhtenberg, S. Comparison of the main bioactive compounds and antioxidant activities in garlic and white and red onions after treatment protocols. *J. Agric. Food Chem.* **2008**, *56*, 4418–4426. [[CrossRef](#)]
65. Ferreres, F.; Sousa, C.; Vrchovska, V.; Valentão, P.; Pereira, J.A.; Seabra, R.M.; Andrade, P.B. Chemical composition and antioxidant activity of tronchuda cabbage internal leaves. *Eur. Food Res. Technol.* **2006**, *222*, 88–98. [[CrossRef](#)]
66. Kawsar, S.M.; Huq, E.; Nahar, N.; Ozeki, Y. Identification and quantification of phenolic acids in *Macrotyloma uniflorum* by reversed phase-HPLC. *Am. J. Plant Physiol.* **2010**, *5*, 204–211. [[CrossRef](#)]
67. Dabas, D.; Ziegler, G.R.; Lambert, J.D. Anti-Inflammatory Properties of a Colored Avocado Seed Extract. *Adv. Food Technol. Nutr. Sci. Open J.* **2019**, *5*, 8–12. [[CrossRef](#)]
68. Giresha, A.S.; Narayanappa, M.; Joshi, V.; Vishwanath, B.S.; Dharmappa, K.K. Human secreted phospholipase A2 (spla2) inhibition by aqueous extract of *Macrotyloma uniflorum* (seed) as anti-inflammatory activity. *Int. J. Pharm. Pharm. Sci.* **2015**, *7*, 217–222.
69. Hameed, H.; Aydin, S.; Başaran, A.A.; Başaran, N. Assessment of cytotoxic properties of sinapic acid in vitro. *Turk. J. Pharm. Sci.* **2016**, *13*, 225–232. [[CrossRef](#)]
70. Qiao, H.Y.; Dahiya, J.P.; Classen, H.L. Nutritional and physiological effects of dietary sinapic acid (4-hydroxy-3, 5-dimethoxy-cinnamic acid) in broiler chickens and its metabolism in the digestive tract. *Poult. Sci.* **2008**, *87*, 719–726. [[CrossRef](#)]
71. Ntie-Kang, F.; Lifongo, L.L.; Mbah, J.; Owono, L.C.O.; Megnassan, E.; Mbaze, L.M.; Judson, P.N.; Sippl, W.; Efange, S.M.N. In silico drug metabolism and pharmacokinetic profiles of natural products from medicinal plants in the Congo basin. *Silico Pharmacol.* **2013**, *1*, 12. [[CrossRef](#)]
72. Kim, R.R.; Chen, Z.; Mann, T.; Bastard, K.F.; Scott, K.; Church, W.B. Structural and functional aspects of targeting the secreted human group IIA phospholipase A2. *Molecules* **2020**, *25*, 4459. [[CrossRef](#)] [[PubMed](#)]
73. Adibhatla, R.M.; Hatcher, J.F. Phospholipase A2, reactive oxygen species, and lipid peroxidation in CNS pathologies. *BMB Rep.* **2008**, *41*, 560. [[CrossRef](#)] [[PubMed](#)]
74. Shelat, P.B.; Chalimoniuk, M.; Wang, J.H.; Strosznajder, J.B.; Lee, J.C.; Sun, A.Y.; Simonyi, A.; Sun, G.Y. Amyloid  $\beta$  peptide and NMDA induce ROS from NADPH oxidase and AA release from cytosolic phospholipase A2 in cortical neurons. *J. Neurochem.* **2008**, *106*, 45–55. [[CrossRef](#)] [[PubMed](#)]
75. Zhao, J.H.; Arao, Y.; Sun, S.J.; Kikuchi, A.; Kayama, F. Oral administration of soy-derived genistin suppresses lipopolysaccharide-induced acute liver inflammation but does not induce thymic atrophy in the rat. *Life Sci.* **2006**, *78*, 812–819. [[CrossRef](#)] [[PubMed](#)]
76. Park, Y.J.; Ko, J.W.; Jeon, S.; Kwon, Y.H. Protective effect of genistein against neuronal degeneration in ApoE<sup>-/-</sup> mice fed a high-fat diet. *Nutrients* **2016**, *8*, 692. [[CrossRef](#)]

77. Rumman, M.; Pandey, S.; Singh, B.; Gupta, M.; Ubaid, S.; Mahdi, A.A. Genistein Prevents Hypoxia-Induced Cognitive Dysfunctions by Ameliorating Oxidative Stress and Inflammation in the Hippocampus. *Neurotox. Res.* **2021**, *39*, 1123–1133. [[CrossRef](#)]
78. Bastian, B.C.; Sellert, C.; Seekamp, A.; Römisch, J.; Pâques, E.P.; Bröcker, E.B. Inhibition of human skin phospholipase A2 by “lipocortins” is an indirect effect of substrate/lipocortin interaction. *J. Investig. Dermatol.* **1993**, *101*, 359–363. [[CrossRef](#)]
79. Nemeč, K.N.; Pande, A.H.; Qin, S.; Urbauer, R.J.B.; Tan, S.; Moe, D.; Tatulian, S.A. Structural and functional effects of tryptophans inserted into the membrane-binding and substrate-binding sites of human group IIA phospholipase A2. *Biochemistry* **2006**, *45*, 12448–12460. [[CrossRef](#)]
80. Matveeva, E.G.; Morisseau, C.; Goodrow, M.H.; Mullin, C.; Hammock, B.D. Tryptophan fluorescence quenching by enzyme inhibitors as a tool for enzyme active site structure investigation: Epoxide hydrolase. *Curr. Pharm. Biotechnol.* **2009**, *10*, 589–599. [[CrossRef](#)]
81. Joshi, B.N.; Sainani, M.N.; Bastawade, K.B.; Deshpande, V.V.; Gupta, V.S.; Ranjekar, P.K. Pearl millet cysteine protease inhibitor: Evidence for the presence of two distinct sites responsible for anti-fungal and anti-feedentactivities. *Eur. J. Biochem.* **1999**, *265*, 556–563. [[CrossRef](#)] [[PubMed](#)]
82. Kelly, S.M.; Jess, T.J.; Price, N.C. How to study proteins by circular dichroism. *Biochim. Biophys. Acta BBA Proteins Proteom.* **2005**, *1751*, 119–139. [[CrossRef](#)] [[PubMed](#)]
83. Yousefpour, A.; Amjad Iranagh, S.; Nademi, Y.; Modarress, H. Molecular dynamics simulation of nonsteroidal antiinflammatory drugs, naproxen and relafen, in a lipid bilayer membrane. *Int. J. Quantum Chem.* **2013**, *113*, 1919–1930. [[CrossRef](#)]
84. Badyal, D.K.; Desai, C. Animal use in pharmacology education and research: The changing scenario. *Indian J. Pharmacol.* **2014**, *46*, 257–265. [[CrossRef](#)] [[PubMed](#)]
85. Hakes, L.; Pinney, J.W.; Robertson, D.L.; Lovell, S.C. Protein-protein interaction networks and biology—What’s the connection? *Nat. Biotechnol.* **2008**, *26*, 69–72. [[CrossRef](#)]